Title Dementia Risk and Dynamic Response to Exercise: Methodology for an Acute Exercise

Clinical Trial

Running Head Acute exercise protocol

Word Count 3976

Authors Dreu White¹, Casey S John², Ashley Kucera², Bryce Truver², Rebecca J Lepping^{2,3}, Phil Lee^{3,4}, Laura Martin^{3,5}, Sandra A Billinger^{1,2}, Jeffrey M Burns², Jill K Morris² and Eric D Vidoni²

Author affiliations ¹ Department of Physical Therapy and Rehabilitation Science, University of Kansas Medical Center, Kansas City, KS, USA

² University of Kansas Alzheimer's Disease Center, Fairway, KS, USA

³ Hoglund Biomedical Imaging Center, University of Kansas Medical Center, Kansas City, KS, USA

⁴ Department of Radiology, University of Kansas Medical Center, Kansas City, KS, USA

⁵ Department of Population Health, University of Kansas Medical Center, Kansas City, KS, USA

Corresponding author Eric D. Vidoni, University of Kansas Alzheimer's Disease Center, 4350

Shawnee Mission Parkway, Fairway, KS 60205, MS6002

Email: evidoni@kumc.edu

Trial registration ClinicalTrials.gov NCT04009629

Abstract

Background: Exercise likely has numerous, meaningful benefits for brain and cognition.

However, those benefits and their causes remain imprecisely defined, especially in the context of cognitive disorders associated with aging, such as Alzheimer's disease (AD). If the brain does benefit from exercise it does so primarily through exposure to brief, "acute" exposures to

exercise over a lifetime. Methods: The Dementia Risk and Dynamic Response to Exercise

(DYNAMIC) clinical trial seeks to characterize the acute exercise response in cerebral

perfusion, and circulating neurotrophic factors in older adults with and without the apolipoprotein e4 genotype (*APOE4*), the strongest genetic predictor of sporadic, late onset AD. DYNAMIC will

enroll 60 older adults into a single moderate intensity bout of exercise intervention. We will

measure pre- and post-exercise cerebral blood flow using arterial spin labeling, and

neurotrophic factors. We expect that *APOE4* carriers will have poor CBF regulation, i.e. slower

return to baseline perfusion after exercise, and will demonstrate blunted neurotrophic response

to exercise, with concentrations of neurotrophic factors positively correlating with CBF

regulation. If exercise-induced changes in perfusion and circulating factors can be detected,

DYNAMIC will contribute to our understanding of exercise-induced brain change and potential

biomarker outcomes of exercise interventions. Results: Preliminary proof-of-concept findings on

7 older adults and 9 younger adults. We have found that this experimental method can capture

CBF and neurotrophic response over a time course, and best practices following exercise.

Conclusions: This methodology will provide important insight into acute exercise response and

potential directions for clinical trial outcomes. ClinicalTrials.gov NCT04009629

Keywords

arterial spin labeling, aerobic exercise, cerebrovascular reserve, acute

Abbreviations

AD - Alzheimer's disease, *APOE4* - apolipoprotein e4, *CBF* - cerebral blood flow, *TCD* - transcranial Doppler, *BDNF* - Brain Derived Neurotrophic Factor, *VEGF* - Vascular Endothelia Growth Factor, *IGF1* - Insulin-like Growth Factor 1, *AUC* - area under the curve, *KU ADC* - University of Kansas Alzheimer's Disease Center, *MRI* - magnetic resonance imaging, *TGSE* - turbo gradient spin echo, *pCASL* - pseudo-continuous arterial spin labeling, *MPRAGE* - magnetization prepared rapid gradient echo, *CTCAE* - Common Terminology Criteria for Adverse Events, *DSMC* - Data and Safety Monitoring Committee, *PET* - positron emission tomography

Background

The number of Americans 65 years and older will double in size over the next 40 years.(1) Aging is often associated with increased cognitive decline.(2) Alzheimer's disease (AD) is of particular concern to the health care system, with an expected two-fold increase in prevalence over the next 30 years, and high direct and indirect costs of care.(3, 4) We must find effective interventions to reduce the burden of cognitive decline and AD on our society.

There is evidence that risk of age-related cognitive decline, including mild cognitive impairment and dementia, can be reduced by health behavior interventions such as exercise.(5-12) Although the literature is not conclusive,(13-16) there is a growing consensus that common healthy behaviors, and especially exercise, support brain health and cognitive function.(17, 18) A number of potential mechanisms may link exercise with brain health. Increased brain volume,(19) regional neurogenesis,(20) circulating neurotrophic factors,(21) and cerebrovascular reserve (i.e. capacity for response to a stimulus challenge) (22) all have been implicated as mediators of exercise benefits for the brain. If the brain does benefit from exercise it does so primarily through brief, "acute" exposures to exercise over a lifetime.(23)

Due to the considerable benefits of aerobic exercise on cardiovascular function, there is interest in precisely defining cerebrovascular adaptations to aerobic exercise and how those adaptations may support cognitive function.(24, 25) We know that sedentary older adults demonstrate decreasing cerebral blood flow (CBF) over time.(26-29) Older athletes who take time off of training quickly experience reductions in CBF,(30) whereas habitual exercise appears to increase CBF.(26, 28, 31) Heart disease and stroke patients improve CBF while completing their cardiac rehab program.(32, 33) Yet, evidence from prior aerobic exercise intervention trials with otherwise healthy older adults have been mixed, some reporting increased CBF,(9) and others reporting no difference.(34)

In general, these studies have used passive "resting" conditions when measuring CBF and blood biomarkers, rather than employing tasks or experimental challenge (e.g. exercise,

task-based fMRI, neuropsychological test) meant to approximate the ecological stressors of daily life. Our own work and others' demonstrates the importance of measuring the response to challenge, especially to exercise.(22, 35-42) For example, using transcranial Doppler (TCD), we recently described the dynamic change in a proxy measure of CBF with onset of aerobic exercise in young and older individuals.(35) Our work demonstrates that older adults have a noticeable blunting of CBF during a dynamic condition like exercise that was less appreciably different from younger adults during a static, resting, measure.

There is evidence that the Apolipoprotein epsilon4 (*APOE4*) single nucleotide polymorphism, the strongest genetic risk factor for late-onset, sporadic AD,(43) leads to modified neurovascular coupling, a leaky blood–brain barrier, angiopathy, and disrupted nutrient transport.(44) *APOE* appears key to maintaining cerebrovascular integrity independent of β -amyloid deposition.(45) *APOE4* carriers also demonstrate altered (46-48) and generally lower resting CBF especially in regions associated with AD-related change.(29, 49-53) Because exercise has such a strong and reliable benefit for the vascular system *APOE4* carriers, who are at greater vascular risk than non-carriers,(12) may preferentially benefit.(54-56)

There are well-documented differences in cerebral blood flow (CBF) and tissue oxygenation based on age or *APOE* genotype,(29, 48, 49, 57) with deleterious consequences for cognition.(29, 58) The exercise stimulus may counter this through cerebral oxygenation and stimulation of neurotrophics, among other mechanisms. Cerebrovascular endothelial cells respond to the shear force stress on the vessel walls by releasing Brain Derived Neurotrophic Factor (BDNF).(59, 60) Peripheral Vascular Endothelia Growth Factor (VEGF) appears to be essential for running-induced neurogenesis and benefits acute exercise performance and brain blood flow in mice.(61, 62) Measurable increases in VEGF are seen after acute exercise.(38, 39) But to date, the literature has not connected CBF, neurotrophins, and *APOE* genotype in exercising humans, despite convergent data pointing to their intimate involvement in cognitive

decline. Our project will extend the prior work by directly assessing the relationship of these factors in response to an acute exercise challenge.

We set out to quantify the CBF response to exercise which has the potential to be a valuable measure of cerebrovascular health.(22) This manuscript details the clinical trial methodology and proof-of-concept preliminary data of the Dementia Risk and Dynamic Response to Exercise study (DYNAMIC - NCT04009629). The scientific premise underlying this project is that CBF and blood-based biomarkers such as VEGF, BDNF, and Insulin-like Growth Factor 1 (IGF1) are interrelated mechanisms driving chronic aerobic exercise effects on brain health and cognition.(9, 20, 63) Our single visit clinical trial seeks to characterize the relationship of *APOE4* carrier status with CBF and blood-based biomarkers of brain health. We capture dynamic fluctuations in CBF and blood-based biomarkers in a time-sensitive manner before and after an acute bout of moderate intensity aerobic exercise. Our working hypothesis is that individuals at genetic risk for AD have poor CBF regulation and altered neurotrophic response and that this methodological approach will inform future clinical trial biomarker protocols in the future.

A secondary goal of this manuscript is to detail the protocol design and adaptations of a single visit experimental study as a clinical trial. Changes in 2018 to the Federal Policy for the Protection of Human Subjects ('Common Rule') has expanded the definition of a clinical trial to include any investigation in which human subjects are prospectively assigned to an intervention to evaluate the effect on a biomedical or behavioral outcome.(64) As a result, most experimental exercise manipulations are now classified as clinical trials even if not traditionally considered an intervention, resulting in increased scrutiny and more strict standards for protocol and reporting. This manuscript also serves to detail clinical trial adaptations for single visit or short experimental studies that have previously fallen outside the aegis of clinical intervention regulation.

Methods

Summary of design

DYNAMIC is a single site, non-randomized, prospectively enrolling trial testing *APOE4*-related response differences to a single, 15-minute bout of moderate intensity aerobic exercise. The study plans to enroll 60 older adults (≥ 65 years), approximately balanced for E4 carriage, and up to 20 younger individuals to serve as normative cohort. We hypothesize that *APOE4* carriers will have poor CBF regulation, i.e. slower return to baseline perfusion (reduced area under the curve) after exercise, and will demonstrate blunted neurotrophic response to exercise, with concentrations of neurotrophic factors positively correlating with CBF regulation. We will also explore the relationship of the CBF and neurotrophic responses to cognitive performance.

Outcomes

As a registered clinical trial, we have identified a single primary outcome and several secondary outcomes of interest. Our primary outcome of interest is global CBF area under the curve (AUC) of the cumulative cerebral blood flow before and after our exercise intervention. Secondary outcomes are change in IGF1, change in VEGF, and change in BDNF from baseline to post intervention. Our exploratory cognitive measures and associated cognitive domains are episodic memory (NIH Toolbox Picture Sequence Memory Test 8+), processing speed (Pattern Comparison Test 7+), and attention and executive function (Flanker Inhibitory Control and Attention Test 12+).

Recruitment and Eligibility

We significantly reduce participant burden by heavily leveraging the infrastructure of the University of Kansas Alzheimer's Disease Center (KU ADC), an NIH-designated Alzheimer's disease research center. The KU ADC follows a Clinical Cohort of 400 individuals with annual cognitive evaluations and prior genetic testing.⁽⁶⁵⁾ We recruit younger participants from the

local area via social networks and fliers. All individuals provide written, institutionally approved written informed consent according to the Declaration of Helsinki guidance either on the day of the visit or in advance through electronic consenting.

Participants have no changes in memory or thinking, or diagnosis of cognitive impairment. Additional inclusion criteria are: 1) Age 18-85 (inclusive); 2) English speaking; 3) corrected hearing or vision; 4) willingness to have genotyping performed if necessary. Exclusion criteria are: 1) health care provider recommended activity restrictions; 2) prior diagnosis of clinically significant cognitive decline judged on Clinical Dementia Rating(66) or Quick Dementia Rating Scale(67) and AD8(68) equivalent of non-impaired, or similar clinical determination in the prior 6 months; 3) anti-coagulant use; 4) high cardiovascular risk without physician clearance for exercise(69); 5) exercise-limiting pain, musculoskeletal, or metabolic condition; 6) MRI contraindications; 7) clinically significant psychiatric illness or other neurological disorders that have the potential to impair cognition (e.g., Parkinson's disease, stroke defined as a clinical episode with neuroimaging evidence in an appropriate area to explain the symptoms); 8) myocardial infarction or symptomatic coronary artery disease in the prior 2 years.

Secondary enrollment considerations are sex and E4 carriage. Within the older adult Clinical Cohort 95% already have *APOE* genotype determined from prior study activities and have consented to share that information with collaborating studies such as DYNAMIC. Genotype is not disclosed to the participant. Study staff with direct participant contact remain blinded to genotype. Because *APOE4* does not have equal penetrance in the Clinical Cohort, the KU ADC continually monitors enrollment rates for DYNAMIC based on sex and E4 carriage and an unblinded study team member not involved in recruitment, consent, or study visit execution reviews and provides enriched contact lists to blinded staff to support balanced participation. Participants who have not previously had their E4 characterized consent to have genotyping performed for the purpose of the study. We exclude individuals with an E2 allele. We make efforts to preferentially match *APOE4* genotype groups based on sex.

Procedures

Participants attend a single study visit. Because our focus is on assessment of dynamic time- and intervention-related changes in our outcomes, precise study timing and short transitions between study activities is critical. Our procedures have been planned to minimize waiting and transition time between study events.

Participants first change into provided magnetic resonance imaging (MRI) compatible clothes (scrubs) and remove all MRI-incompatible dental appliances, jewelry, etc. The exercise bicycle ergometer (Corival, Lode B.V., www.lode.nl) is adjusted so that the knee achieves near but not complete extension. Participants practice pedaling between 60-70 rpm until they report feeling comfortable with the movement and equipment.

Participants then complete approximately 20 minutes of NIH Toolbox-based neuropsychological testing on an iPad (Apple Inc.) in the following order, Pattern Comparison, Picture Sequence Memory, and Flanker Inhibitory Control and Attention tests. Two seated blood pressure and pulse readings and taken one minute apart are averaged as a baseline measure of vitals (Welch Allyn ProBP3400).

Next, participants are escorted to the adjacent MRI suite. They are fitted with a continuous blood pressure monitoring cuff (Caretaker 4, Caretaker Medical N.A. caretakermedical.net) on the finger which is calibrated to the baseline vitals. The MRI technologist fits ear plugs and headphones on the participant, lays them on the MRI table positioning the cuffed finger on the abdomen and begins scanning. Rapid transition from preparation to scanning is emphasized.

Imaging data are collected with a 3 Tesla whole-body scanner (Siemens Skyra, Erlangen, Germany) fitted with a 20-channel head and neck receiver coil. The MRI session is split into two parts: pre-exercise, and post-exercise. Each portion of the session begins with automated scout image acquisition and shimming procedures to optimize field homogeneity.

The pre-exercise portion consists of two 3D turbo gradient spin echo (TGSE) pseudo-continuous arterial spin labeling (pCASL) sequences,(70-73) yielding 9 minutes and 56 seconds of pre-exercise pCASL data. All pCASL sequences are collected with the same acquisition parameters, designed for full brain coverage, and optimized for cerebral blood flow (CBF) measurements in older adults (repetition time/echo time (TR/TE) = 4300/22.42 ms, flip angle = 120 deg, field of view (FOV) = 300 x 300 mm², matrix = 96 x 96 voxels, voxel in-plane resolution = 3.125 x 3.125 mm², slice thickness = 2.5 mm, 48 axial slices, 20 volumes, acquisition time = 5:48). Positioning of the pCASL sequences is prescribed using the automated scout image, aligning the top of the acquisition box to the top of the brain and the angle of acquisition to the base of the corpus callosum as landmarks. The two pre-exercise pCASL sequences are followed by a T₁-weighted, 3D magnetization prepared rapid gradient echo (MPRAGE) structural scan(74) (TR/TE = 2300/2.95 ms, inversion time (TI) = 900 ms, flip angle = 9 deg, FOV = 253 x 270 mm², matrix = 240 x 256 voxels, voxel in-plane resolution = 1.05 x 1.05 mm², slice thickness = 1.2 mm, 176 sagittal slices, in-plane acceleration factor = 2, acquisition time = 5:09), used for alignment and normalization of the pCASL data.

Participants return to the testing room. An optical heart rate sensor (OH1 Polar Electro, Inc. polar.com) is secured to the forearm with self-adhering wrap. Blood pressure is taken again. Then, a flexible intravenous catheter is placed and 10mL of blood is collected in vacutainer tubes containing EDTA as an anti-coagulant. If the genotype is not available, an additional 3mL of blood is collected in a single vacutainer tube containing acid citrate dextrose and stored for future genotyping.

Participants then remount the cycle ergometer and begin a 5-minute warm up. During the initial 5 minutes, study staff gradually increase resistance with a goal of achieving the target heart rate of 45-55% of HR reserve in minutes 4 and 5. Participants begin pedaling at 60-70 rpm and a resistance based on an age-dependent decision algorithm (Figure 1). The 15-minute aerobic exercise bout begins immediately following the warm-up. Study staff check heart rate

every 1-minute and adjust cycle resistance to maintain the heart rate in the target zone. After 15 minutes of exercise resistance is reduced to 10W and participants pedal at a self-selected cadence for 3 minutes to cool down and drink 100mL of water to reduce potential perspiration-related changes in blood volume. Post-exercise vitals are taken immediately at the beginning of the cool down period. An additional 10mL of blood is collected during cool down. The heart rate monitor is removed, and the participant is quickly escorted back to the MRI room.

Once back in the MRI room, the same preparatory procedures for MRI are repeated and 4 consecutive pCASL sequences are acquired, yielding 23 minutes and 12 seconds of post-exercise pCASL data. Finally, the participant is escorted back to the testing room where vitals and 10mL of blood are taken one more time, and neuropsychological testing is repeated. Participants are compensated \$100 upon completion of the visit.

Blood Processing and Assessment

We optimized sample collection and processing procedures for accurate measurement of plasma neurotrophins in 5 samples independent of this study. As demonstrated in our proof-of-concept results, when platelets remain in a blood sample, a freeze thaw can greatly increase the concentration of such factors and may not accurately reflect levels that were circulating in the plasma at the time of acute exercise. Consequently, our protocol emphasizes immediate processing. Our optimized protocol is as follows: plasma is generated immediately upon collection by centrifugation by processing at 1500 RCF (g) (2800 RPM) at 4°C for ten minutes. Platelet-rich plasma is then centrifuged in four, 1.5mL aliquots at 1700 x g (4500 RPM) (4C) for 15 minutes. The resulting platelet-poor plasma is separated from the pellet and snap frozen in liquid nitrogen until stored at -80 C at the end of the visit.

Imaging Analysis

Planned pCASL data analyses include using the USC Laboratory of Functional MRI Technology CBF Preprocess and Quantify packages,(75, 76) and Statistical Parametric Mapping CAT12 package.(77) We motion correct labeled and control pCASL images separately, calculate the CBF timeseries via simple subtraction, and coregister with the T₁-weighted anatomical image. Mean CBF in the entire gray matter mask is extracted from each label-control subtraction image in the timeseries in units of mL*100g tissue⁻¹*min⁻¹. Imaging analysis may change as the field improves methods.

Cognitive Test Assessment

Cognitive assessments scores are calculated automatically by the NIH Toolbox software (<https://www.healthmeasures.net/>). We plan to use change in T-Score for each domain which is age, education, gender, and ethno-racial identification corrected and provides a score based on a normative mean of 50 with a standard deviation of 10.

Data Collection

All data are collected and organized in a custom designed REDCap(78) database. Project access is role based. APOE4 genotypes are kept in a separate database and the linking list is kept by a designated, unblinded investigator.

Sample Size

To our knowledge, there are currently no peer-reviewed reports of genotype-based CBF differences in response to acute exercise. However, perfusion measures in genetic risk for AD (APOE4) have been performed previously and can form the basis of a reasonable power analysis. Two prior cross-sectional estimates of the relationship of perfusion and E4 carriage

have delivered similar effect sizes ($d=1.0$). (48, 49) Given this effect size, we expect to be able to discern differences based on *APOE4* in a sample of 60 older adults.

Safety

Adverse Events. Adverse events are defined as any untoward medical occurrence in study participants, which does not necessarily have a causal relationship with the study treatment. The seriousness of the adverse event is determined using the National Cancer Institute Common Terminology Criteria for Adverse Events (CTCAE) v3.0, and need for hospital admission. Adverse events are assessed only at the visit, but the consent form has contact information should the participant need to contact the study team regarding delayed development of an AE. Serious Adverse Events are reported per institutional and NIH requirements.

Monitoring. Safety of the study is monitored in an ongoing manner by a chartered Data and Safety Monitoring Committee (DSMC) and Independent Safety Officer according to a Safety Plan. The Independent Safety Officer advises the NIH and the Principal Investigator regarding participant safety, participant risks and benefits, scientific integrity and ethical conduct of a study. The DSMC provides additional support and guidance for the Principal Investigator.

Response to SARS-CoV-2

Data collection began prior to the SARS-CoV-2 novel coronavirus pandemic, was paused between March 11 and June 1, 2020, and resumed with additional safety procedures. The blood processing centrifuge and staff member were moved to a separate, nearby testing room to allow for physical distancing. All participants are screened via telephone a maximum of 72 hours before the day of visit, and are screened again (including temperature) outside the imaging facility on the day of the visit. All staff wear level 1 surgical masks, gloves, and face

shields. Participants are provided a surgical mask to be worn at all times except during exercise and in the MRI. The staff member in charge of exercise and neuropsychological testing minimizes time within 6 feet of the participant and MRI tech staff. Electronic, advance consenting has been implemented following the coronavirus pandemic to reduce the amount of close contact time between staff and participant.

Proof-of-Concept Results

To date, we have enrolled 16 participants into the study. Demographics are provided in Table 1. To demonstrate proof-of-concept, we provide preliminary analyses and comparisons of individuals above and below 65 years of age. E4 carriage has not been unmasked to date and is not included in this report. There were more women in the older adult group. Figure 2 depicts the study flow and average time for each study event. Figure 3 demonstrates our ability to capture dynamic blood flow changes post-exercise. The area under the CBF curve (Figure 3A) is greater in the younger individuals. Additionally, even when we normalize CBF to the average of the second pre-exercise pCASL sequence (a stabilized resting CBF), the area under the curve of younger adults remains greater. Notably, these dynamic changes do not appear to be heavily driven by blood pressure which is plotted underneath the CBF curves.

We also provide evidence for our decision to process blood collection immediately. In plasma processed with one centrifugation, mean plasma levels of BDNF were $17,506 \pm 4031$ pg/mL. Duplicate samples that underwent an immediate second centrifugation to remove the platelet pellet, followed by an immediate snap-freeze, were measured as having mean levels to 29 ± 10.6 pg/mL. A delay of 15 minutes prior to each centrifugation step, which may allow increased time for platelet release when activated by the shear stress of centrifugation, increased levels to 245 ± 108 pg/mL.

Adverse Events

To date there has been one adverse event, nausea, upon IV placement. Symptoms resolved with a light snack and rest.

Discussion

We have designed and implemented a single visit clinical trial to test the effect of *APOE4* carriage on brain blood flow response to an acute, 15-minute, moderate intensity exercise challenge. We have also refined optimal blood collection and processing procedures to characterize circulating, bio-available neurotrophic responses to exercise. We expect that our strict and time sensitive protocol will allow us to identify CBF and blood-based neurotrophic responses that are obscured during typical resting conditions. We will also explore whether CBF or neurotrophic responses are related to performance changes on neuropsychological tests. To be clear, we do not expect measurable vascularization, neurogenesis, or other benefits immediately following exercise. Nor do we suggest that any neurotrophic increases are causal of *ad hoc* cognitive change or CBF response. Rather, we seek to index the transient changes and relationships that are hypothesized to mediate these benefits with chronic exposure to exercise.

Our proof-of-concept results suggest that we can capture a dynamic CBF recovery response following exercise that can differentiate groups. We found that older adults have lower overall blood flow and slower return to resting blood flow as measured by area under the CBF curve. Our blood processing results provide a clear case for immediate post-processing to identify the bio-available circulating neurotrophic factors following exercise. Optimization of pre- and post- processing techniques revealed higher BDNF levels in plasma collected without removal of platelets compared to plasma where platelets had been removed. In addition, sitting time after the centrifugation step to remove platelets, which may result in platelet activation, also affected neurotrophic factor levels.

Despite the many strengths and carefully constructed protocol, there are notable limitations. First, our pre-post exercise measures are proxies of CBF during exercise as we are not imaging during exercise. Some protocols for MRI during exercise are beginning to emerge, though concerns about motion artifact remain. CBF can also be measured using contrast-enhanced MRI, TCD, or positron emission tomography (PET). TCD has the advantages of temporal resolution and ease of use during exercise, but can only index blood velocity. Xenon PET imaging can produce whole brain CBF but requires a radioactive isotope. pCASL provides both the whole brain spatial resolution, and potentially improved temporal resolution compared to PET, using only magnetically labeled arterial blood water, an endogenous tracer that is highly reproducible.(70) Additionally, optimal imaging parameters to capture CBF response to acute exercise using have been investigated previously.(79) Pseudo-continuous ASL sequences have improved signal-to-noise ratio while maintaining high labeling efficiency.(80) Common to all ASL sequences are pairs of images with and without labeling, that allow for CBF quantification. Typically, the subtraction of these pairs is averaged over a sequence to calculate CBF. However, the pairs viewed as a timeseries, may also capture information about transient CBF changes, for example following exercise, as we have done here. We believe this timecourse will provide important additional information about dynamic responsiveness, or cerebrovascular reserve, of the system.

A second limitation is our focus on three neurotrophic factors and the *APOE4* genotype. Recent evidence suggests that there are myriad important biochemical changes in response to exercise.(81) Similarly, thousands of genes are responsible for exercise response.(82) We have chosen to focus on those that are most consistently linked to brain health and Alzheimer's disease in the literature. While important for our own research line, it leaves the potential for uninvestigated responses that may be important.

Despite these limitations the DYNAMIC study offers a blueprint for a unique and innovative methodology to capture acute exercise response in individuals at risk for Alzheimer's disease.

Aerobic exercise is among the most important and cost effective tools available for chronic disease management.(83) Our work along with others' strongly suggests that moderate-intensity aerobic exercise benefits brain health and cognition.(10, 11, 84-87) However, the field continue to struggle with adequate methods for capturing mechanistic drivers of exercise benefits on the brain. New protocols such as DYNAMIC are necessary to move forward our mechanistic explorations of exercise effects on brain health and cognition.

Declarations

Ethics approval and consent to participate All participants have provided written informed consent approved by the University of Kansas Institutional Review Board, consistent with the Declaration of Helsinki.

Consent for publication Not applicable.

Availability of data and materials Data may be obtained by contacting the corresponding author. Data will not be released prior to the closing of the trial.

Competing interests No authors have a perceived or real conflict of interest.

Funding Research reported in this publication was supported by the National Institute on Aging of the National Institutes of Health under award number R21AG061548 and P30 AG035982, an equipment grant S10 RR29577, and the Leo and Anne Albert Charitable Trust. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the Leo and Anne Albert Trust, and they had no involvement in the acquisition or dissemination of the results.

Authors' contributions EDV, JKM, PL, SAB, JMB, LM and RJL, contributed to the conception of the work. All authors contributed to the design of the work. DW, CSJ, AK, BT, RJL, EDV, JKM contributed to the acquisition or analysis of the work. CSJ, RJL, SAB, JMB, JKM, EDV contributed to the interpretation of data. All authors have contributed to the drafting and revision of the work. All authors have approved submission and agreed to be personally accountable for the work

References

1. Colby SL, Ortman JM. Projections for the Size and Composition of the U. S. Population: 2014 to 2060. In: US Census Bureau, editor. Washington, DC2014.

2. National Institute on Aging. Growing Older in America: the Healthy and Retirement Study. Bethesda, MD: National Institutes of Health; 2007.
3. Jutkowitz E, Kane RL, Gaugler JE, MacLehose RF, Dowd B, Kuntz KM. Societal and Family Lifetime Cost of Dementia: Implications for Policy. J Am Geriatr Soc. 2017;65(10):2169-75.
4. Alzheimer's A. 2016 Alzheimer's disease facts and figures. Alzheimer Dementia. 2016;12(4):459-509.
5. Yaffe K, Barnes D, Nevitt M, Lui LY, Covinsky K. A prospective study of physical activity and cognitive decline in elderly women: women who walk. Arch Intern Med. 2001;161(14):1703-8.
6. Weuve J, Kang JH, Manson JE, Breteler MM, Ware JH, Grodstein F. Physical activity, including walking, and cognitive function in older women. JAMA. 2004;292(12):1454-61.
7. Nagamatsu LS, Chan A, Davis JC, Beattie BL, Graf P, Voss MW, et al. Physical activity improves verbal and spatial memory in older adults with probable mild cognitive impairment: a 6-month randomized controlled trial. J Aging Res. 2013;2013:861893.
8. Ngandu T, Lehtisalo J, Solomon A, Levalahti E, Ahtiluoto S, Antikainen R, et al. A 2 year multidomain intervention of diet, exercise, cognitive training, and vascular risk monitoring versus control to prevent cognitive decline in at-risk elderly people (FINGER): a randomised controlled trial. Lancet. 2015;385(9984):2255-63.
9. Maass A, Duzel S, Goerke M, Becke A, Sobieray U, Neumann K, et al. Vascular hippocampal plasticity after aerobic exercise in older adults. Molecular psychiatry. 2015;20(5):585-93.
10. Erickson KI, Voss MW, Prakash RS, Basak C, Szabo A, Chaddock L, et al. Exercise training increases size of hippocampus and improves memory. Proc Natl Acad Sci U S A. 2011;108(7):3017-22.

11. Vidoni ED, Johnson DK, Morris JK, Van Sciver A, Greer CS, Billinger SA, et al. Dose-Response of Aerobic Exercise on Cognition: A Community-Based, Pilot Randomized Controlled Trial. *PLoS One*. 2015;10(7):e0131647.
12. Rosenberg A, Ngandu T, Rusanen M, Antikainen R, Backman L, Havulinna S, et al. Multidomain lifestyle intervention benefits a large elderly population at risk for cognitive decline and dementia regardless of baseline characteristics: The FINGER trial. *Alzheimer Dementia*. 2017.
13. Sink KM, Espeland MA, Castro CM, Church T, Cohen R, Dodson JA, et al. Effect of a 24-Month Physical Activity Intervention vs Health Education on Cognitive Outcomes in Sedentary Older Adults: The LIFE Randomized Trial. *JAMA*. 2015;314(8):781-90.
14. Brasure M, Desai P, Davila H, Nelson VA, Calvert C, Jutkowitz E, et al. Physical Activity Interventions in Preventing Cognitive Decline and Alzheimer-Type Dementia: A Systematic Review. *Ann Intern Med*. 2018;168(1):30-8.
15. Barreto PS, Demougeot L, Vellas B, Rolland Y. Exercise training for preventing dementia, mild cognitive impairment, and clinically meaningful cognitive decline: a systematic review and meta-analysis. *J Gerontol A Biol Sci Med Sci*. 2017.
16. Daviglus ML, Bell CC, Berrettini W, Bowen PE, Connolly ES, Jr., Cox NJ, et al. NIH state-of-the-science conference statement: Preventing Alzheimer's disease and cognitive decline. *NIH Consens State Sci Statements*. 2010;27(4):1-30.
17. Global Council on Brain Health. The Brain-Body Connection: GCBH Recommendations on Physical Activity and Brain Health. 2016.
18. Erickson KI, Hillman C, Stillman CM, Ballard RM, Bloodgood B, Conroy DE, et al. Physical Activity, Cognition, and Brain Outcomes: A Review of the 2018 Physical Activity Guidelines. *Med Sci Sports Exerc*. 2019;51(6):1242-51.

19. Erickson KI, Raji CA, Lopez OL, Becker JT, Rosano C, Newman AB, et al. Physical activity predicts gray matter volume in late adulthood: The Cardiovascular Health Study. *Neurology*. 2010;75(16):1415-22.
20. Pereira AC, Huddleston DE, Brickman AM, Sosunov AA, Hen R, McKhann GM, et al. An in vivo correlate of exercise-induced neurogenesis in the adult dentate gyrus. *Proc Natl Acad Sci U S A*. 2007;104(13):5638-43.
21. Erickson KI, Miller DL, Roecklein KA. The aging hippocampus: interactions between exercise, depression, and BDNF. *Neuroscientist*. 2012;18(1):82-97.
22. Davenport MH, Hogan DB, Eskes GA, Longman RS, Poulin MJ. Cerebrovascular reserve: the link between fitness and cognitive function? *Exerc Sport Sci Rev*. 2012;40(3):153-8.
23. Buchman AS, Boyle PA, Yu L, Shah RC, Wilson RS, Bennett DA. Total daily physical activity and the risk of AD and cognitive decline in older adults. *Neurology*. 2012;78(17):1323-9.
24. Tyndall AV, Argourd L, Sajobi TT, Davenport MH, Forbes SC, Gill SJ, et al. Cardiometabolic risk factors predict cerebrovascular health in older adults: results from the Brain in Motion study. *Physiol Rep*. 2016;4(8).
25. Leeuwis AE, Hooghiemstra AM, Amier R, Ferro DA, Franken L, Nijveldt R, et al. Design of the ExCersion-VCI study: The effect of aerobic exercise on cerebral perfusion in patients with vascular cognitive impairment. *Alzheimer dementia*. 2017;3(2):157-65.
26. Rogers RL, Meyer JS, Mortel KF. After reaching retirement age physical activity sustains cerebral perfusion and cognition. *J Am Geriatr Soc* 1990;38(2):123-8.
27. Bailey DM, Marley CJ, Brugniaux JV, Hodson D, New KJ, Ogoh S, et al. Elevated aerobic fitness sustained throughout the adult lifespan is associated with improved cerebral hemodynamics. *Stroke*. 2013;44(11):3235-8.
28. Thomas BP, Yezhuvath US, Tseng BY, Liu P, Levine BD, Zhang R, et al. Life-long aerobic exercise preserved baseline cerebral blood flow but reduced vascular reactivity to CO₂. *Journal of magnetic resonance imaging : JMRI*. 2013;38(5):1177-83.

29. Kim SM, Kim MJ, Rhee HY, Ryu CW, Kim EJ, Petersen ET, et al. Regional cerebral perfusion in patients with Alzheimer's disease and mild cognitive impairment: effect of APOE epsilon4 allele. *Neuroradiology*. 2013;55(1):25-34.
30. Alfini AJ, Weiss LR, Leitner BP, Smith TJ, Hagberg JM, Smith JC. Hippocampal and Cerebral Blood Flow after Exercise Cessation in Master Athletes. *Frontiers Aging Neurosci*. 2016;8:184.
31. Xu X, Jerskey BA, Cote DM, Walsh EG, Hassenstab JJ, Ladino ME, et al. Cerebrovascular perfusion among older adults is moderated by strength training and gender. *Neurosci Lett*. 2014;560:26-30.
32. Anazodo UC, Shoemaker JK, Suskin N, Ssali T, Wang DJ, St Lawrence KS. Impaired Cerebrovascular Function in Coronary Artery Disease Patients and Recovery Following Cardiac Rehabilitation. *Frontiers Aging Neurosci* e. 2015;7:224.
33. Robertson AD, Crane DE, Rajab AS, Swardfager W, Marzolini S, Shirzadi Z, et al. Exercise intensity modulates the change in cerebral blood flow following aerobic exercise in chronic stroke. *Exp Brain Res*. 2015;233(8):2467-75.
34. Flodin P, Jonasson LS, Riklund K, Nyberg L, Boraxbekk CJ. Does Aerobic Exercise Influence Intrinsic Brain Activity? An Aerobic Exercise Intervention among Healthy Old Adults. *Frontiers Aging Neurosci*. 2017;9:267.
35. Billinger SA, Craig JC, Kwapiszeski SJ, Sisante JV, Vidoni ED, Maletsky R, et al. Dynamics of middle cerebral artery blood flow velocity during moderate-intensity exercise. *J Appl Physiol* (1985). 2017;122(5):1125-33.
36. Ogoh S, Ainslie PN. Cerebral blood flow during exercise: mechanisms of regulation. *J Appl Physiol* (1985). 2009;107(5):1370-80.
37. Jorgensen LG, Perko G, Secher NH. Regional cerebral artery mean flow velocity and blood flow during dynamic exercise in humans. *J Appl Physiol* (1985). 1992;73(5):1825-30.

38. Asano M, Kaneoka K, Nomura T, Asano K, Sone H, Tsurumaru K, et al. Increase in serum vascular endothelial growth factor levels during altitude training. *Acta Physiol Scand*. 1998;162(4):455-9.
39. Schobersberger W, Hobisch-Hagen P, Fries D, Wiedermann F, Rieder-Scharinger J, Villiger B, et al. Increase in immune activation, vascular endothelial growth factor and erythropoietin after an ultramarathon run at moderate altitude. *Immunobiology*. 2000;201(5):611-20.
40. Zoladz JA, Pilc A, Majerczak J, Grandys M, Zapart-Bukowska J, Duda K. Endurance training increases plasma brain-derived neurotrophic factor concentration in young healthy men. *J Physiol Pharmacol*. 2008;59 Suppl 7:119-32.
41. Rasmussen P, Brassard P, Adser H, Pedersen MV, Leick L, Hart E, et al. Evidence for a release of brain-derived neurotrophic factor from the brain during exercise. *Exp Physiol*. 2009;94(10):1062-9.
42. Seifert T, Brassard P, Wissenberg M, Rasmussen P, Nordby P, Stallknecht B, et al. Endurance training enhances BDNF release from the human brain. *Am J Physiol*. 2010;298(2):R372-7.
43. Bertram L, McQueen MB, Mullin K, Blacker D, Tanzi RE. Systematic meta-analyses of Alzheimer disease genetic association studies: the AlzGene database. *Nature Genetics*. 2007;39(1):17-23.
44. Tai LM, Thomas R, Marottoli FM, Koster KP, Kanekiyo T, Morris AW, et al. The role of APOE in cerebrovascular dysfunction. *Acta Neuropathol*. 2016;131(5):709-23.
45. Bell RD, Winkler EA, Singh I, Sagare AP, Deane R, Wu Z, et al. Apolipoprotein E controls cerebrovascular integrity via cyclophilin A. *Nature*. 2012;485(7399):512-6.
46. Scarmeas N, Habeck CG, Stern Y, Anderson KE. APOE genotype and cerebral blood flow in healthy young individuals. *JAMA*. 2003;290(12):1581-2.

47. Fleisher AS, Podraza KM, Bangen KJ, Taylor C, Sherzai A, Sidhar K, et al. Cerebral perfusion and oxygenation differences in Alzheimer's disease risk. *Neurobiol Aging*. 2009;30(11):1737-48.
48. Zlatar ZZ, Bischoff-Grethe A, Hays CC, Liu TT, Meloy MJ, Rissman RA, et al. Higher Brain Perfusion May Not Support Memory Functions in Cognitively Normal Carriers of the ApoE epsilon4 Allele Compared to Non-Carriers. *Frontiers Aging Neurosci*. 2016;8:151.
49. Thambisetty M, Beason-Held L, An Y, Kraut MA, Resnick SM. APOE epsilon4 genotype and longitudinal changes in cerebral blood flow in normal aging. *Arch Neurology*. 2010;67(1):93-8.
50. Farrer LA, Cupples LA, Haines JL, Hyman B, Kukull WA, Mayeux R, et al. Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease. A meta-analysis. APOE and Alzheimer Disease Meta Analysis Consortium. *JAMA*. 1997;278(16):1349-56.
51. Corder E, Saunders A, Strittmatter W, Schmechel DE, Gaskell PC, Small GW, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science*. 1993;261:921-3.
52. Myers RH, Schaefer EJ, Wilson PW, D'Agostino R, Ordovas JM, Espino A, et al. Apolipoprotein E epsilon4 association with dementia in a population-based study: The Framingham study. *Neurology*. 1996;46(3):673-7.
53. Seshadri S, Fitzpatrick AL, Ikram MA, DeStefano AL, Gudnason V, Boada M, et al. Genome-wide analysis of genetic loci associated with Alzheimer disease. *JAMA*. 2010;303(18):1832-40.
54. Pizzie R, Hindman H, Roe CM, Head D, Grant E, Morris JC, et al. Physical activity and cognitive trajectories in cognitively normal adults: the adult children study. *Alzheimer Disease Assoc Disord*. 2014;28(1):50-7.

55. Head D, Bugg JM, Goate AM, Fagan AM, Mintun MA, Benzinger T, et al. Exercise Engagement as a Moderator of the Effects of APOE Genotype on Amyloid Deposition. Arch Neurology. 2012;69(5):636-43.
56. Schultz SA, Boots EA, Darst BF, Zetterberg H, Blennow K, Edwards DF, et al. Cardiorespiratory fitness alters the influence of a polygenic risk score on biomarkers of AD. Neurology. 2017;88(17):1650-8.
57. Viticchi G, Falsetti L, Vernieri F, Altamura C, Altavilla R, Luzzi S, et al. Apolipoprotein E genotype and cerebrovascular alterations can influence conversion to dementia in patients with mild cognitive impairment. J Alzhiemer Dis. 2014;41(2):401-10.
58. Tarumi T, Dunskey DI, Khan MA, Liu J, Hill C, Armstrong K, et al. Dynamic cerebral autoregulation and tissue oxygenation in amnesic mild cognitive impairment. J Alzhiemer Dis. 2014;41(3):765-78.
59. Prigent-Tessier A, Quirie A, Maguin-Gate K, Szostak J, Mossiat C, Nappey M, et al. Physical training and hypertension have opposite effects on endothelial brain-derived neurotrophic factor expression. Cardiovascular Res. 2013;100(3):374-82.
60. Banoujaafar H, Van Hoecke J, Mossiat CM, Marie C. Brain BDNF levels elevation induced by physical training is reduced after unilateral common carotid artery occlusion in rats. J Cerebral Blood Flow Metabol. 2014;34(10):1681-7.
61. Fabel K, Fabel K, Tam B, Kaufer D, Baiker A, Simmons N, et al. VEGF is necessary for exercise-induced adult hippocampal neurogenesis. Eur J Neurosci. 2003;18(10):2803-12.
62. Rich B, Scadeng M, Yamaguchi M, Wagner PD, Breen EC. Skeletal myofiber vascular endothelial growth factor is required for the exercise training-induced increase in dentate gyrus neuronal precursor cells. J Physiol. 2017;595(17):5931-43.
63. Borror A. Brain-derived neurotrophic factor mediates cognitive improvements following acute exercise. Med Hypotheses. 2017;106:1-5.
64. Subjects PoH. 46 CFR. 2018.

65. Graves RS, Mahnken JD, Swerdlow RH, Burns JM, Price C, Amstein B, et al. Open-source, Rapid Reporting of Dementia Evaluations. *J Registry Manag.* 2015;42(3):111-4.
66. Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology.* 1993;43(11):2412b-4.
67. Galvin JE. The Quick Dementia Rating System (Qdrs): A Rapid Dementia Staging Tool. *Alzheimers Dement (Amst).* 2015;1(2):249-59.
68. Galvin JE, Roe CM, Powlishta KK, Coats MA, Muich SJ, Grant E, et al. The AD8: a brief informant interview to detect dementia. *Neurology.* 2005;65(4):559-64.
69. ACSM ACoSM. ACSM's guidelines for exercise testing and prescription. 8th ed. Thompson WR, editor. Philadelphia: Lippincott Williams & Wilkins; 2010.
70. Kilroy E, Apostolova L, Liu C, Yan L, Ringman J, Wang DJ. Reliability of 2D and 3D Pseudo-Continuous Arterial Spin Labeling Perfusion MRI in Elderly Populations—Comparison with 15O-water PET. *JMRI.* 2014;39(4):931.
71. Wang Y, Moeller S, Li X, Vu AT, Krasileva K, Ugurbil K, et al. Simultaneous multi-slice Turbo-FLASH imaging with CAIPIRINHA for whole brain distortion-free pseudo-continuous arterial spin labeling at 3 and 7 T. *Neuroimage.* 2015;113:279-88.
72. Wang DJ, Alger JR, Qiao JX, Gunther M, Pope WB, Saver JL, et al. Multi-delay multi-parametric arterial spin-labeled perfusion MRI in acute ischemic stroke - Comparison with dynamic susceptibility contrast enhanced perfusion imaging. *Neuroimage Clin.* 2013;3:1-7.
73. Yan L, Wang S, Zhuo Y, Wolf RL, Stiefel MF, An J, et al. Unenhanced dynamic MR angiography: high spatial and temporal resolution by using true FISP-based spin tagging with alternating radiofrequency. *Radiology.* 2010;256(1):270-9.
74. Shah M, Ross JS, Tkach J, Modic MT. 3D MPRAGE evaluation of the internal auditory canals. *J Comput Assist Tomogr.* 1993;17(3):442-5.
75. Jann K. LOFT_CBFpreprocess.m.
76. Jann K, Smith R, Piedra ER, Xiao X, Wang DJ. LOFT_CBFquantify.m. 3.0 ed2017.

77. Dahnke R, Yotter RA, Gaser C. Cortical thickness and central surface estimation. *Neuroimage*. 2013;65:336-48.
78. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)--a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform*. 2009;42(2):377-81.
79. Smith JC, Paulson ES, Cook DB, Verber MD, Tian Q. Detecting changes in human cerebral blood flow after acute exercise using arterial spin labeling: implications for fMRI. *J Neurosci Method*. 2010;191(2):258-62.
80. Wu WC, Fernandez-Seara M, Detre JA, Wehrli FW, Wang J. A theoretical and experimental investigation of the tagging efficiency of pseudocontinuous arterial spin labeling. *Magn Reson Med*. 2007;58(5):1020-7.
81. Contrepolis K, Wu S, Moneghetti KJ, Hornburg D, Ahadi S, Tsai MS, et al. Molecular Choreography of Acute Exercise. *Cell*. 2020;181(5):1112-30 e16.
82. Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, et al. Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA*. 1995;273(5):402-7.
83. Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, King AC, et al. Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Med Sci Sports Exerc*. 2007;39(8):1435-45.
84. Defina LF, Willis BL, Radford NB, Gao A, Leonard D, Haskell WL, et al. The association between midlife cardiorespiratory fitness levels and later-life dementia: a cohort study. *Ann Intern Med*. 2013;158(3):162-8.
85. Voss MW, Prakash RS, Erickson KI, Basak C, Chaddock L, Kim JS, et al. Plasticity of brain networks in a randomized intervention trial of exercise training in older adults. *Frontiers Aging Neurosci*. 2010;2.

86. Muller J, Chan K, Myers JN. Association Between Exercise Capacity and Late Onset of Dementia, Alzheimer Disease, and Cognitive Impairment. Mayo Clin Proc. 2017;92(2):211-7.
87. Billinger SA, Vidoni ED, Morris JK, Thyfault JP, Burns JM. Exercise Test Performance Reveals Support of the Cardiorespiratory Fitness Hypothesis. J Aging Phys Act. 2016:1-17.

Age $\leq$ 60

No

Yes

Begin at 40W
2 minutes at steps
from above

Begin at 60W
2 minutes at steps
from above

After 3 warm-up minutes have elapsed check HR

No

In Target
HR Zone?

Yes

Increase or decrease
resistance in ~5W
steps. Check HR
again in 15 seconds

Maintain Watts. Check
HR in again in 15
seconds

No

In Target
HR Zone?

Yes

Increase or decrease
resistance in ~10W
steps. Check again in
15 seconds

Maintain Watts.
Check HR again in
15 seconds

Continue to adjust resistance based on HR
for 2 minutes

No

In Target
HR Zone?

Yes

Increase or
decrease
resistance in ~5W
steps.
Check HR again in
1 minute

Maintain Watts.
Check HR again in 1
minute

No

In Target
HR Zone?

Yes

Increase or decrease
resistance in ~5W
steps.
Check again in 1
minute

Maintain Watts.
Check HR again in 1
minute

Exercise for 15 minutes.

Table 2. Demographics and Preliminary Results		
	Younger (n=9)	Older (n=7)
Age, yrs	25.9 [23-29]	76.6 [71-81]
Female, %	5 (56%)	6 (86%)
Resting Heart Rate, bpm	81.4 [68-95]	62 [54-71]
Resting MAP, mmHg	99.6 [83.8-128.0]	113 [90.2-142.0]
Mean Workload	82.3W [49.3-114]	29.7W [0-49.3]
Peak Workload	89.3W [60-120]	31.4 [0-50]
Pre-exercise MAP AUC	1480 [1476-1485]	1673 [1443-1913]
Pre-exercise MAP AUC	3397 [3192-3602]	3856 [3514-4207]
Pre-exercise CBF AUC	1571 [920-2241]	1339 [1043-1591]
Post-exercise CBF AUC	2973 [1893-4027]	2413 [2035-2897]
<p>Values mean [range] unless otherwise noted. bpm= beats per minute;</p> <p>CBF=cerebral blood flow; MAP = mean arterial pressure; AUC=area under curve.</p>		