Title: Pathogenic *SPTBN1* variants cause a novel autosomal dominant neurodevelopmental syndrome

Authors: Margot A. Cousin^{1,2*}, Keith A. Breau³, Blake A. Creighton³, Rebecca C. Spillmann⁴, Erin Torti⁵, Sruthi Dontu³, Swarnendu Tripathi⁶, Deepa Ajit³, Kathryn M. Harper^{7,8}, Michael C. Stankewich⁹, Richard E. Person⁵, Yue Si⁵, Elizabeth A. Normand⁵, Amy Blevins⁵, Alison S. May¹⁰, Louise Bier¹¹, Vimla Aggarwal^{11,12}, Grazia M. S. Mancini¹³, Marjon A. van Slegtenhorst¹³, Kirsten Cremer¹⁴, Jessica Becker¹⁴, Hartmut Engels¹⁴, Stefan Aretz¹⁴, Jennifer J. MacKenzie¹⁵, Eva Brilstra¹⁶, Koen L. I. van Gassen¹⁶, Richard H. van Jaarsveld¹⁶, Renske Oegema¹⁶, Gretchen M. Parsons¹⁷, Paul Mark¹⁷, Ingo Helbig^{18,19}, Sarah E. McKeown^{18,19}, Robert Stratton²⁰, Benjamin Cogne^{21,22}, Bertrand Isidor^{21,22}, Pilar Cacheiro²³, Damian Smedley²³, Helen V. Firth^{24,25}, Tatjana Bierhals²⁶, Katja Kloth²⁶, Deike Weiss²⁷, Cecilia Fairley^{28,29}, Joseph T. Shieh^{28,29}, Amy Kritzer³⁰, Parul Jayakar³¹, Evangeline Kurtz-Nelson³², Raphael Bernier³², Tianyun Wang³³, Evan E. Eichler^{33,34}, Ingrid M.B.H. van de Laar¹³, Allyn McConkie-Rosell⁴, Marie McDonald⁴, Jennifer Kemppainen^{1,35}, Brendan C. Lanpher^{1,35}, Laura E. Schultz-Rogers^{1,2}, Lauren B. Gunderson^{1,35}, Pavel N. Pichurin¹, Grace Yoon³⁶, Michael Zech^{37,38}, Robert Jech³⁹, Juliane Winkelmann^{37,38,40,41}, Undiagnosed Diseases Network[#], Genomics England Research Consortium[#], Michael T. Zimmermann^{6,42,43}, Brenda Temple⁴⁴, Sheryl S. Moy^{7,8}, Eric W. Klee^{1,2,35}, Queenie K.-G. Tan⁴, Damaris N. Lorenzo^{3,8,45*}

1. Center for Individualized Medicine, Mayo Clinic, Rochester, MN, USA

2. Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA

3. Department of Cell Biology and Physiology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

4. Department of Pediatrics, Duke University Medical Center, Duke University, Durham, NC, USA

5. GeneDx, Gaithersburg, MD, USA

6. Bioinformatics Research and Development Laboratory, Genomic Sciences and Precision Medicine Center, Medical College of Wisconsin, Milwaukee, WI, USA

- 24 7. Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
- 25 8. Carolina Institute for Developmental Disabilities, University of North Carolina at Chapel Hill,
- 26 Chapel Hill, NC, USA
- 27 9. Department of Pathology, Yale University, New Haven, CT, USA
- 28 10. Department of Neurology, Columbia University, NY, USA
- 29 11. Institute for Genomic Medicine, Columbia University, NY, USA
- 30 12. Laboratory of Personalized Genomic Medicine, Department of Pathology and Cell Biology,
- 31 Columbia University, NY, USA
- 32 13. Department of Clinical Genetics, Erasmus MC University Medical Center, 3015GD Rotterdam,
- 33 The Netherlands
- 34 14. Institute of Human Genetics, University of Bonn, School of Medicine & University Hospital Bonn,
- 35 Bonn, Germany
- 36 15. McMaster University, Hamilton, ON, Canada
- 37 16. Department of Genetics, University Medical Center Utrecht, Utrecht, The Netherlands
- 38 17. Spectrum Health Medical Genetics, USA
- 39 18. Division of Neurology, Departments of Neurology and Pediatrics, The Children's Hospital of
- 40 Philadelphia and the Perelman School of Medicine at the University of Pennsylvania,
- 41 Philadelphia, PA, USA
- 42 19. The Epilepsy NeuroGenetics Initiative, Children's Hospital of Philadelphia, Philadelphia, PA, USA
- 43 20. Genetics, Driscoll Children's Hospital, Corpus Christi, TX, USA
- 44 21. Service de Génétique Médicale, CHU Nantes, Nantes, France

- 45 22. Université de Nantes, CNRS, INSERM, L'Institut du Thorax, Nantes, France
- 46 23. William Harvey Research Institute, School of Medicine and Dentistry, Queen Mary University of
- 47 London, London, UK.
- 48 24. Cambridge University Hospitals, UK
- 49 25. Wellcome Sanger Institute, Hinxton, UK
- 50 26. Institute of Human Genetics, University Medical Center Hamburg-Eppendorf, Hamburg,
- 51 Germany
- 52 27. Department of Pediatrics, Neuropediatrics, University Medical Center Hamburg-Eppendorf,
- 53 Hamburg, Germany
- 54 28. Institute for Human Genetics, University of California San Francisco, San Francisco, CA, USA.
- 55 29. Division of Medical Genetics, Department of Pediatrics, University of California San Francisco,
- 56 San Francisco, CA, USA.
- 57 30. Boston Children's Hospital, Division of Genetics and Genomics, MA, USA
- 58 31. Nicklaus Children's Hospital, Miami, FL, USA
- 59 32. Department of Psychiatry and Behavioral Sciences, University of Washington, Seattle, WA, USA
- 60 33. Department of Genome Sciences, University of Washington School of Medicine, Seattle, WA,
- 61 USA
- 62 34. Howard Hughes Medical Institute, Seattle, WA, USA
- 63 35. Department of Clinical Genomics, Mayo Clinic, Rochester, MN, USA
- 64 36. Divisions of Clinical/Metabolic Genetics and Neurology, The Hospital for Sick Children, University
- 65 of Toronto, Toronto, Canada

37. Institute of Neurogenomics, Helmholtz Zentrum München, Munich, Germany
38. Institute of Human Genetics, Technical University of Munich, Munich, Germany
39. Department of Neurology, Charles University, 1st Faculty of Medicine and General University Hospital in Prague, Prague, Czech Republic
40. Lehrstuhl für Neurogenetik, Technische Universität München, Munich, Germany
41. Munich Cluster for Systems Neurology, SyNergy, Munich, Germany
42. Department of Biochemistry, Medical College of Wisconsin, Milwaukee, WI, USA
43. Clinical and Translational Sciences Institute, Medical College of Wisconsin, Milwaukee, WI, USA
44. Department of Biochemistry, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
45. Neuroscience Center, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

*Corresponding author

#A complete list of consortia members is provided in the Supplementary Note

Abstract

SPTBN1 encodes β II-spectrin, the ubiquitously expressed member of the β -spectrin family that forms micrometer-scale networks associated with plasma membranes. β II-spectrin is abundantly expressed in the brain, where it is essential for neuronal development and connectivity. Mice deficient in neuronal β II-spectrin expression have defects in cortical organization, global developmental delay, dysmorphisms, and behavioral deficiencies of corresponding severity. These phenotypes, while less severe, are observed in haploinsufficient animals, suggesting that individuals carrying heterozygous variants in this gene may also present with measurable compromise of neural development and function. Here we report the identification of heterozygous *SPTBN1* variants in 29 individuals who present with global developmental, language and motor delays, mild to severe intellectual disability, autistic features, seizures, behavioral and movement abnormalities, hypotonia, and variable dysmorphic facial features. We show that these *SPTBN1* variants lead to loss-of-function, gain-of-function, and dominant negative effects that affect protein stability, disrupt binding to key protein partners, and disturb cytoskeleton organization and dynamics. Our studies define the genetic basis of this new neurodevelopmental syndrome, expand the set of spectrinopathies affecting the brain and neural development, and underscore the critical role of β II-spectrin in the central nervous system.

Introduction

Spectrins are ubiquitously expressed, elongated polypeptides that bind membrane lipids and ankyrins to line the plasma membrane^{1,2}. The spectrin meshwork is formed by heterodimeric units of α - and β -spectrin assembled side-to-side in antiparallel fashion, which then form head-to-head tetramers that crosslink F-actin to form spectrin-actin arrays^{1,2}. Mammalian neurons express the most diverse repertoire of spectrins (α II- and β I-V spectrins) of any cell type³. Together with ankyrins, spectrins self-assemble with both remarkable long-range regularity and micro- and nanoscale specificity to precisely position and stabilize cell adhesion molecules, membrane transporters, ion channels, and other cytoskeletal proteins³. Some spectrins also enable intracellular organelle transport³. Unsurprisingly, deficits in spectrins underlie several neurodevelopmental and neurodegenerative disorders⁴⁻⁶. For example, inherited autosomal dominant variants in β III-spectrin (encoded by *SPTBN2*) cause late onset spinocerebellar ataxia type 5 (SCA5)⁵, while pathogenic *de novo* variants have been associated with early childhood ataxia, intellectual disability (ID), and developmental delay (DD)⁷⁻¹². Similarly, autosomal recessive *SPTBN2* variants¹³⁻¹⁵ are associated with childhood ataxia, which may also present with ID and DD¹³, collectively referred to as autosomal recessive spinocerebellar ataxia 14 (SCAR14). *De novo* pathogenic variants in *SPTAN1*, which encodes α II-spectrin, cause West syndrome, an early-infantile epileptic encephalopathy (EIEE) characterized by frequent severe seizures and persistent abnormality of cortical function⁵, and other childhood onset epileptic syndromes¹⁶⁻²⁰. Some patients co-present with spastic quadriplegia, DD, and various brain defects⁵. In addition, dominantly inherited *SPTAN1* nonsense variants were recently linked to juvenile onset hereditary motor neuropathy²¹. Biallelic alterations in β IV-spectrin (encoded by *SPTBN4*) result in congenital hypotonia, neuropathy, and deafness, with and without ID^{6,22,23}.

Neuronal β II-spectrin, encoded by *SPTBN1*, is the most abundant β -spectrin in the brain and forms tetramers with α II-spectrin, which intercalate F-actin rings to build a sub-membranous periodic skeleton (MPS)²⁴. A cytosolic pool of β II-spectrin promotes bidirectional axonal organelle transport^{25,26}. We

previously reported that mice lacking β II-spectrin in all neural progenitors (*Sptbn1*^{flox/flox};Nestin-Cre; referred to as β II^{Sp}-KO) show early postnatal lethality, reduced long-range cortical and cerebellar connectivity, spontaneous seizures, and motor deficits²⁶. However, the impact of human genetic variation in *SPTBN1* on β II-spectrin function and its association with disease has not been studied. Here we describe a cohort of 29 individuals carrying rare, mostly *de novo* variants in *SPTBN1* affected by a novel autosomal dominant neurologic syndrome presenting with global developmental, language and motor delays, mild to severe ID, autistic features, seizures, behavioral abnormalities, hypotonia, and variable dysmorphisms. This suggests conserved roles for β II-spectrin in neuronal development and function. The most damaging variants clustered within the actin-binding calponin homology domain (CH) and led to aberrant neuronal morphology, decreased neurite outgrowth, and deficient axonal organelle transport in primary neurons. Consistent with these deficiencies, our biochemistry, microscopy, and molecular modeling studies indicate that *SPTBN1* variants lead to loss-of-function (LOF), gain-of-function (GOF), and dominant negative effects that affect protein stability, disrupt binding to key protein partners, and affect cytoskeleton organization and dynamics. Consequently, histology and behavioral studies in brain β II-spectrin-deficient mice showed neuron-autonomous brain connectivity defects and recapitulated developmental and behavioral phenotypes observed in patients with *SPTBN1* variants. Collectively, our data strongly support pathogenic mechanisms of *SPTBN1* variants as the genetic cause of a novel neurodevelopmental syndrome and underscores the multifaceted role of β II-spectrin in the nervous system.

Results

Patients with *SPTBN1* variants present with a novel neurodevelopmental syndrome

A cohort of 29 individuals from 28 families (one pair of siblings) who carry heterozygous variants in *SPTBN1* was identified through whole genome (WGS) or exome (WES) sequencing. These probands presented with

neurodevelopmental delay and variable neurologic, behavioral, and dysmorphic features (Fig. 1, Table 1, Supplementary Note). Twenty-four of the 29 affected individuals carry *de novo* variants, while the remaining have unknown inheritance due to lack of parental samples for testing (Supplementary Table 1). In proband P17 the *SPTBN1* variant inheritance was unknown but sequencing revealed mosaicism at 13.3% of reads suggesting the variant occurred *de novo*. Twenty-eight unique variants are described (P10 has two *de novo* variants in cis) of which 22 are missense, three are nonsense, and three are canonical splice-site variants, with two predicted by SpliceAI²⁷ to lead to in-frame deletions and one predicted to result in a frameshift that introduces a premature stop codon (Fig. 1a, Supplementary Table 1). Approximately half of the variants cluster in the CH domain, predominantly in the second CH domain (CH2), with the rest distributed in various spectrin repeats (SR) (Fig. 1a).

The phenotypic findings are summarized in Table 1 and detailed clinical and family histories are included in the Supplementary Note. The cohort included 17 male and 12 female probands with the age at last evaluation spanning from 6 months to 26 years of age. All had early onset of symptoms with primarily developmental delay (DD) presentations with the exception of two individuals. For 28 individuals with phenotypic data, 26 reported some level of delays. Of those with DD, 21 reported both speech (SD) and motor (MD) delays, three reported only SD (P16, P26 and P27), and one reported only MD (P29). Twenty individuals reported ID, with seven of those being severe or moderate to severe, and six being mild or low to normal. Developmental regression was noted in three individuals. Although mild delays were noted in the clinical evaluation of proband P29 (c.5961+2T>C, p.(I1988Afs*90)), the primary symptom, dystonia, was observed at age 13. Similarly, while delay in speaking was noted for proband P23 (p.A1086T), the primary phenotype in this individual was liver-related. Only partial phenotype information was obtained for previously reported probands P4 and P25²⁸, and no specific phenotypic information is available for P3 from the Deciphering Developmental Disorders (DECIPHER) database²⁹, although this individual was presumed to have some form of DD (Supplementary Note).

Nine individuals have a history of seizures, with two being febrile only, and six with electroencephalogram (EEG) correlates. Seizure episodes in four of the cases were resolved later in childhood or were successfully managed through medication. Four of these patients were diagnosed with frontal lobe or generalized epilepsy. Seven individuals had abnormal brain MRI findings, including three patients with thinning of the corpus callosum (CC), two with ventriculomegaly, two with delayed myelination, and two showing some volume loss (P1: diffuse cerebral parenchymal; P10: mild cerebellar and vermian). The remaining brain MRI findings were unique (Table 1, Supplementary Note). Behavioral concerns were common within the cohort. Six individuals displayed autistic features or had an autism spectrum disorder (ASD) diagnosis, including two (P19 and P24) previously reported as part of a WES study of a 2,500 ASD patient cohort³⁰. One other proband (P25) was originally identified as part of a WES study of over 500 trios from Tourette syndrome cohorts²⁸. Thirteen individuals presented with other behavioral concerns, including attention deficit and hyperactivity disorder (ADD/ADHD) (n=11), anxiety (n=3), emotional lability including tantrums and depression (n=7), and aggressive or self-injurious behaviors (n=6). Seven individuals experienced sleep disturbances, in some cases co-occurring with seizure episodes. Additional phenotypic findings include changes in muscle tone and movement abnormalities. Ten had hyper- (n=4) or hypotonia (n=8). Seven individuals reported movement abnormalities including dystonia (n=3), ataxic or unsteady gait (n=5), spasticity (n=1), and tremor (n=2). Less common features included hearing impairment, reported for five individuals, generally mild to moderate, with one individual suspected to have conductive hearing loss due to recurrent ear infections. Dysmorphic features were noted in 18 individuals with a subset of individuals shown in Fig. 1c, but no consistent findings were observed. Macrocephaly (n=5) and microcephaly (n=5) were both observed as well as other head shape anomalies (n=6).

SPTBN1 is intolerant to both missense and loss-of-function variants (gnomAD v2.1.1)³¹, and protein sequence alignment of human β II-spectrin and its orthologues across several species shows a high degree

of evolutionary conservation of the residues impacted by these putative pathogenic variants (Fig. 1b). Consistent with their implied functional relevance, the majority of the variants are predicted to be likely damaging to protein function by multiple prediction tools (PolyPhen-2, Mutation Taster, SIFT, PROVEAN, M-CAP, PredictSNP2, and CADD) (Supplementary Table 1). Additionally, all variants are absent or extremely rare in the population (gnomAD v2.1.1)³¹ (Supplementary Table 1). Missense variants in codons G205, T268, R411 and R1003 were identified in more than one individual (Fig. 1a). The p.R1003W variant was identified in two maternal half-siblings (P21 and P22) inherited from their unaffected mother, who was found to be mosaic for the variant at a low level (found in 1.8% of next generation sequencing reads). Unrelated individual P20 also carries the *de novo* p.R1003W variant, and has the common DD features, but also presented with some non-overlapping clinical features. Similarly, variants in unrelated duos P4 and P5, and P15 and P16 affect the same p.G205 and p.R411 residues, respectively, but result in different amino acid substitutions. All of these individuals have DD, both P4 and P5 had an abnormal EEG, and both P15 and P16 had hypotonia but each also has some distinct features consistent with the variability in the cohort. Likewise, unrelated patients P10, P11, and P12 carry different amino acid substitutions in residue p.T268, and present with overlapping phenotypes. Notably, P10 has two β II-spectrin variants *in cis* (p.F344L and p.T268A), which may contribute to the more severe phenotype observed. The partial clinical divergence within these patients likely stems from differences in sex, age, and genomic background, which in turn may determine their corresponding penetrance and physiological consequences.

In sum, the above clinical presentations suggest that *SPTBN1* variants converge to impair cellular and physiological mechanisms that lead to delays in motor and language development and cognitive skills. Additionally, the results of these evaluations suggest that several of these variants also result in additional neurological and behavioral phenotypes. These observations are consistent with pleiotropic functions of β II-spectrin including its diverse and critical roles in brain development and function²⁶.

Human β II-spectrin mutations affect protein cellular distribution and alter cell morphology

To begin to assess the pathogenic mechanisms of *SPTBN1* variants, we introduced a subset of the mutations in the coding sequence of human β II-spectrin cloned into the peGFP-C3 plasmid, transfected the constructs into HEK293 cells, either alone or together with pmCherry-C1, and monitored their effects on GFP- β II-spectrin (GFP- β IIsp) levels, localization, size, and stability by confocal microscopy and western blot. Of the 22 mutations tested, protein levels, of 10 were unchanged relative to control (Fig. 2a, Extended Data Fig. 1a). Variants p.H275R and p.A850G resulted in 25% and 50% β II-spectrin overexpression, respectively. Expression of nonsense p.C183*, p.E892* and p.W1787* variants yielded truncated proteins of the expected size. However, while p.W1787* was expressed normally, the levels of the p.C183* and p.E892* mutants were significantly reduced. While neither of the two CH domain in-frame deletions (I159_Q160del and Y190_R216del) affected protein levels, several mutations in this region showed lower expression. Both p.G205D and p.G205S mutations reduced protein levels, with p.G205D also impacting β II-spectrin solubility, which largely precipitated into the Triton-X100 insoluble fraction (Fig. 2b, Extended Data Fig. 1b). This indicates that these amino acid substitutions at p.G205 likely affect the structural conformation of β II-spectrin, and result in unfolded and unstable polypeptides. Notably, wildtype (WT) GFP- β IIsp localized through the cytosol and at the cell membrane whereas GFP- β IIsp bearing p.I159_Q160del at the end of CH1, or the p.C183*, p.Y190_R216del, p.G205D, and p.G205S mutations in the proximal CH2 domain accumulated in large cytosolic aggregates (white arrowheads Fig. 2c and Extended Data Fig. 1c). Expression of the CH2 mutations p.T268A, T268N, T268S, V271M, and H275R resulted in normal GFP- β IIsp distribution, but induced noticeable changes in cell morphology. Overall, cells were enlarged and had increased membrane protrusions (Fig. 2c and Extended Data Fig. 1c, asterisks). These changes are indicative of modified cytoskeleton arrangements and dynamics, and likely reflect altered F-actin binding.

Cell morphology phenotypes induced by SR mutations varied. For example, cells expressing the p.R411W mutation, located in SR1, which is required for dimerization with α II-spectrin³² and thought to contribute

to actin binding³³, were enlarged and with increased membrane protrusions. Among the five mutations clustered within SR6-8, p.A850G and p.R1003W induced the most noticeable morphological changes (Fig. 2c, Extended Data Fig. 1c). Interestingly, p.A850G, which results in β II-spectrin overexpression, also causes a striking increase in membrane protrusions, which suggests a GOF effect (Fig. 2c and Extended Data Fig. 1c). Since SR3-S14 have no assigned functional specificity nor contain known binding sites for partners, a molecular rationale for how mutations in this region affect β II-spectrin function is lacking. SR15 binds ankyrins, with amino acid p.Y1874 known to be critical for this interaction³⁴. However, p.E1886Q GFP- β II_{Sp} does not show apparent cellular changes, likely because it does not disrupt binding to ankyrins³⁴, or this complex is not required for cytoskeleton organization or dynamics in HEK293 cells. Expression of E892* and W1787* GFP- β II_{Sp}, lacking the polypeptide portions from SR6 to C-terminus and SR14 to C-terminus respectively, did not cause apparent cellular phenotypes. This is surprising because in addition to loss of ankyrin binding, these truncated proteins also lack the tetramerization³⁵ and pleckstrin homology (PH) domains³⁶, the latter being important for binding lipids in the cell and organelle membranes. Together, these data indicate that human β II-spectrin mutations can lead to cellular phenotypes through LOF and GOF mechanisms that likely involve changes in cytoskeleton architecture and dynamics.

Human β II-spectrin mutations affect its interaction with submembrane cytoskeleton partners

A subcortical network of F-actin- and ankyrin-bound β II-/ α II-spectrin tetramers promotes membrane stability and helps organizing membrane proteins within specialized microdomains¹⁻³. Thus, pathogenic *SPTBN1* variants could impair neuronal development and/or function by altering β II-spectrin interaction with F-actin and other cytoskeletal partners or their submembrane availability. Consistent with the latter prediction, we found that both actin and mCherry- α II_{Sp} were sequestered in GFP- β II_{Sp} aggregates caused by expression of various CH domain variants in HEK293 cells (arrowheads, Fig. 3a, Extended Data Fig. 2a). These mutations in the CH domain (F-actin binding region) also resulted in GFP- β II_{Sp} aggregation and in

clustering of endogenous actin and α II-spectrin within GFP aggregates when expressed in cortical neurons from β II-SpKO mice (arrowheads, Fig. 3b). To further evaluate the ability of mutant β II-spectrin to associate with molecular partners, we conducted binding assays and co-immunoprecipitation (co-IP) experiments. We prioritized the evaluation of a subset of variants based on the likelihood that they would have an effect on the interaction tested, given their position on the specific domains known to be critical for binding to that specific partner. We first assessed the effect of the mutations on the formation of β II-spectrin/ α II-spectrin complexes by incubating GFP beads coupled to WT or mutant GFP- β II-Sp with cell lysates expressing mCherry- α II-Sp and measuring the amount of mCherry- α II-Sp in eluates from GFP pulldowns by western blot. As expected, mutant C183* GFP- β II-Sp neither associated with mCherry- α II-Sp in pulldown assays nor sequestered mCherry- α II-Sp or endogenous α II-spectrin into GFP- β II-Sp aggregates in HEK293 or neurons because it lacks the heterodimer nucleation SR1-SR2 region³⁰ (Fig. 3a-c and Extended Data Fig. 2b). Similarly, the pulldown of α II-spectrin with G205D and G205S GFP- β II-Sp baits yielded less α II-/ β II-spectrin complexes, partly due to the lower expression of these mutant polypeptides, but it also indicates a lower affinity for α II-spectrin (Fig. 3a-c and Extended Data Fig. 2b). That a single substitution in CH2 reduces α II-spectrin affinity is surprising because this domain has not been linked to α II-spectrin binding. With the exception of the p.R1003W mutant located in SR7, which reduces association with α II-spectrin by 40%, none of the other variants tested affects α II-spectrin binding (Fig. 3c and Extended Data Fig. 2a, b). The lower α II-spectrin binding to R1003W GFP- β II-Sp might result from local or long-range conformational changes that weakens interactions along the dimer.

Next, we evaluated whether mutations in the CH domains, the known actin-binding domain in β II-spectrin, affect its binding to F-actin using a co-sedimentation assay. β II-spectrin proteins containing a PreScission protease (PP) recognition site between GFP and the initiation codon of β II-spectrin (GFP-PP- β II-Sp) were produced in HEK293 cells and captured on GFP beads. Purified WT and mutant β II-spectrin were recovered from beads upon PP cleavage and mixed with purified F-actin. The partition of β II-spectrin between the

soluble (S) and actin-containing pellet (P) fractions was used to estimate the relative binding proclivity between both proteins. We found that CH1 mutation p.T59I and deletion p.I159_Q160del in the CH1-CH2 linker led to approximately 40% and 70% less β II-spectrin association with F-actin, respectively (Fig. 3d and Extended Data Fig. 2c). Similarly, p.Y190_R216del and p.D255E CH2 variants also resulted in reduced F-actin binding. In contrast, the p.V271M and p.H275R mutations increased F-actin binding to β II-spectrin by 60-90%, while all three p.T268S/N/A mutants bound F-actin at similar levels as the WT protein (Fig. 3d and Extended Data Fig. 2c). This range in binding affinity is likely caused by the balance of individual or combined effects of both local and CH domain-wide conformational changes caused by modified intramolecular interactions, which in turn results in modified intermolecular contacts at the β II-spectrin/F-actin interface. Given that the p.A850G mutant causes a cell morphology phenotype similar to the ones induced by some of the CH domain mutants (Fig. 2c and Extended Data Fig. 1c), we also tested its binding to F-actin. Surprisingly, this mutant resulted in approximately 50% higher F-actin binding (Fig. 3d and Extended Data Fig. 2c). While this higher propensity for actin binding is likely to underlie the abnormal cell morphology, it is not clear how this substitution, several SR away from the CH domain, can modify this interaction.

Finally, we evaluated the impact of β II-spectrin mutations on its interaction with ankyrins. In this experiment, HA-tagged 220kDa ankyrin-B was expressed in HEK293 cells together with WT or mutant GFP- β IIsp proteins. The presence of HA signal in eluates from GFP- β IIsp complexes was detected by western blot. Consistent with previous reports³⁴, expression of the Y1874 mutation in SR15 (the known ankyrin-binding domain) almost entirely abrogated binding to ankyrin-B (Fig. 3e and Extended Data Fig. 2d). As expected, truncated β II-spectrin polypeptides that lack SR15 caused by p.E892* and p.W1787* variants also disrupted binding between these partners (Fig. 3e and Extended Data Fig. 2d). Interestingly, the SR15 p.E1886Q variant did not affect binding to ankyrin-B, despite its spatial proximity to the Y1874 binding site³⁴.

Molecular modeling predicts effects of β II-spectrin variants on protein stability and F-actin binding

We further assessed the impact of *SPTNB1* variants through molecular modelling. We first modeled the 10 missense variants involving seven residues in the β II-spectrin CH1-CH2 domain. The CH domain is a protein module of around 100 residues composed of four alpha helices³⁷ found in cytoskeletal and signal transduction actin-binding proteins (ABP)³⁸. Multiple biochemical studies using ABP containing CH1-CH2 domains, such as spectrin superfamily members α -actinin-4 (*ACTN4*) and utrophin (*UTRN*), suggest dynamic transitions between “closed” and “open” configurations of the tandem domains, whereas the open state is thought to expose CH1 residues to enable its predominant role of binding actin, with CH2 regulating the conformational state through autoinhibition³⁸. The electrostatic surface profile of β II-spectrin CH1 and CH2 domains modeled using an available crystal structure of utrophin³⁹ indicates that they each have one electrically active side complementary to each other and one neutral side, consistent with an energetically balanced closed conformation (Fig. 4a-c). This model also indicates that six of the eight mutated CH domain residues reside at the CH1-CH2 dimer interface, potentially impacting interdomain helix-helix interactions, thereby dysregulating the natural autoinhibition (Fig. 4b,c).

To refine our prediction of the closed conformation of the β II-spectrin CH1-CH2 domain and to identify interactions at the interface, we docked the CH2 domain (residues 173-278) of β II-spectrin⁴⁰ onto the available modeled structure of the CH1 domain (residues 55-158) of β III-spectrin (95% homologous with β II-spectrin)⁴¹ and chose the top docking pose (Fig. 4d). This pose was also the same pose compared to the actinin-4 (PDB ID 6oa6) (unpublished) and utrophin (PDB ID 1qag)³⁹ closed conformations (Extended Data Fig. 3a-c). Figure 4i summarizes the key predicted interacting residues at the CH1-CH2 interface and the structural consequences of the variants in those domains. Four of the eight residues affected by *SPTNB1* missense variants (T59 in CH1, and L250, T268, and H275 in CH2) are predicted to be involved in interdomain interactions (Fig. 4d,i). The *SPTNB1* missense variants in these and the two other interface residues (D255 and V271) are predicted to introduce destabilizing effects (Fig. 4i). For example,

substitutions of T268 by Ala (smaller and more hydrophobic), Ser (loss of methyl group), and Asn (larger and more hydrophilic) likely have different degrees of alteration of the original T268 hydrophobic interaction with L155 in CH1 and I159 in the CH1-CH2 linker. However, if these amino acid changes in T268 affect the CH1-CH2 domain conformation, they do not result in appreciable changes in F-actin binding (Fig. 3d and Extended Data Fig. 2c), despite their marked cellular (Fig. 2c, Extended Data Fig. 1c) and disease-linked effects. Similarly, substitution of glutamic for aspartic acid in D255E is a relatively small change that does not result in changes in F-actin affinity. On the other hand, both the V271M (larger and hydrophobic) and the H275R (longer and significantly more hydrophilic) substitutions may impair CH1 binding to cause a shift toward the open CH1-CH2 conformation and higher F-actin affinity. This is also likely the case for the L250R mutation, which is expected to cause significant steric hindrance by the clashing of the large, charged residue with a hydrophobic CH1 pocket (Fig. 4e). The molecular and cellular consequences of this variant remain to be assessed. Conversely, the T59I mutation introduces a slightly longer, but more hydrophobic group that might promote a stronger interaction with L250 in CH2, potentially shifting the equilibrium to a CH1-CH2 closed configuration consistent with less F-actin binding (Fig. 3d and Extended Data Fig. 2c).

The amino acid substitutions in the two CH2 sites in the interior of the domain (G205S/D and L247H) are predicted to cause significant steric hindrance, which likely results in CH2 domain instability (Fig. 4f-h). G205S/D introduces destabilization by positioning a negative charge on the interior and steric hindrance against the neighbor N233 and L234 sidechains (Fig. 4g,h), which likely underlies the aggregation of the mutant protein in cells (Figs. 2c and 3b). The Y190_R216del mutation, which eliminates 27 amino acids in the CH2 domain, also results in β II-spectrin aggregation and diminished F-actin binding (Figs. 2c and 3b,d and Extended Data Fig. 2c). In these cases, the autoinhibitory interactions will also be lost if the structure of the CH2 domain is compromised, but through a different mechanism than the genetic variants that alter the CH dimer interface. Besides the open/closed CH1-CH2 domain conformational shifts, some of

these mutants might be directly involved in binding F-actin. To explore this possibility, we independently docked the CH1 and CH2 domains onto an F-actin model built from chains A-F of 6anu⁴⁰ using ClusPro^{42,43} (Extended Data Fig. 3d-f). The top eight CH1 docking poses predicted by the balanced and electrostatic scoring algorithms almost all correspond to the location and orientation of CH1 molecules on F-actin as defined by the cryo-EM structure 6anu (Extended Data Fig. 3d, dark blue). For CH2 docking onto F-actin, the top eight docking poses predicted by the balanced and electrostatic scoring algorithms almost all correspond to symmetry-related locations and poses on F-actin (Extended Data Fig. 3e). In addition, the predicted orientation of CH2 molecules on F-actin is consistent with the known binding site of the CH1 domain, as judged by the length of the linker that would be required to join the C-terminus of the docked CH1 domain to the N-terminus of the docked CH2 domain (Extended Data Fig. 3f). Our model predicts that neither the T59 residue nor its mutated version are directly involved in F-actin binding (Extended Data Fig. 3d). On the CH2 domain, the H275R mutant may result in a stronger interaction with negatively charged D51 in F-actin (Extended Data Fig. 3e), which may further contribute to its higher actin binding propensity (Fig. 3d and Extended Data Fig. 2c). We also modeled the missense mutations in the SR (Extended Data Fig. 3g,h). Except for F344L, all SR variants face outwards, to the solvent, indicating that they could be involved in protein binding at the interface. Interestingly, all mutations within the second and third helices of the spectrin fold are neutral or more hydrophobic for the variants, and those in the first helix of the SR are more hydrophilic. Given the consistency of this trend, we suspect that it may underlie a conserved functional role important for heterodimerization and larger order assemblies.

In sum, our modeling results provide a strong molecular rationale for the biochemical and cellular observations described above, which implicate protein stability, abnormal assembly and dynamics of the β II-spectrin-F-actin skeleton, and potential disruptions of β II-spectrin binding to other molecular partners, consistent with similar LOF and GOF changes observed in other members of the spectrin superfamily^{37,40,43}.

βII-spectrin mutations disrupt neuron architecture and function

Individual with *SPTBN1* variants display developmental deficits and a wide range of neurological phenotypes, which implicate βII-spectrin in neuronal development and cerebral cortex function. These clinical presentations are consistent with phenotypes of neural progenitor-specific βII-spectrin null mice that lack βII-spectrin throughout brain development²⁶. In addition, cortical and hippocampal neurons from these mice show disruption of the spectrin-actin membrane periodic structure²⁴, impaired axonal formation and growth, and reduced axonal organelle transport, all deficits that can be rescued by expression of WT βII-spectrin^{26,44}. These reports, together with our initial cellular and molecular observations shown above, suggest that mutant βII-spectrin may result in defects in the organization and the dynamics of the neuronal submembrane skeleton, and the morphology and function of neurons. Thus, we next investigated the neuronal effects of human βII-spectrin mutations using a structure-function rescue approach in βII-SpKO cortical neurons.

First, we expressed WT and mutant GFP-βII-Sp together with mCherry in day in vitro (DIV) 3 βII-SpKO cortical neurons²⁶ and evaluated their neuronal growth at DIV8. We also evaluated mCherry-expressing WT (*Sptbn1*^{flox/flox} /+) and heterozygous (*Sptbn1*^{flox/+}; Nestin-Cre; henceforth abbreviated as βII-SpHet) neurons grown in parallel. As previously observed²⁶, neuronal growth, quantified through axonal length, was severely impaired in βII-SpKO neurons, but restored upon expression of WT GFP-βII-Sp (Fig. 5a,b and Extended Data Fig. 4a). βII-SpHet neurons grew to roughly only half the length of WT neurons, but their axons were at least twice as long as βII-SpKO neurons (Fig. 5a,b). Most of the βII-spectrin mutants failed to rescue axonal length except for p.G1398S and p.E1886Q, which restored growth to WT levels, while p.A1086T and p.E1110D restored length to heterozygous levels (Fig. 5a,b and Extended Data Fig. 4a). Remarkably, some of the aberrant morphological features observed in HEK293 cells were also present and often more markedly displayed in neurons expressing mutant GFP-βII-Sp. As shown above, p.C183*, p.Y190_R216del, p.G205S, and p.G205D GFP-βII-Sp mutants were almost exclusively distributed in large

protein aggregates localized to the neuronal cell bodies and in some processes (Fig. 3b and Extended Data Fig. 5a). All other mutants within the CH domain invariably produced extensive aberrant membrane features in the form of lamellipodia and filopodia around the cell body and along the neuronal processes (Fig. 6a and Extended Data Fig. 5a). Similarly, the p.A850G mutant resulted in cell bodies and neuronal processes with expanded membranes extensively decorated with filopodia-like protrusions, while the p.R411W mutant led to a milder phenotype (Extended Data Fig. 5a). Neuronal membrane expansion was accompanied by a shift in the boundaries of actin and α II-spectrin distribution (Fig. 6a). These results confirm that clinically relevant β II-spectrin mutations can cause marked disruptions in cell morphology, likely driven by disruptions in the submembrane cytoskeleton organization and dynamics, which may be a pathogenic factor in *SPTBN1* syndrome.

Organelle transport is essential for the maintenance of neuronal processes and viability of neurons and defects in transport can contribute to the pathology of several neurological diseases⁴⁵. We previously showed that β II-spectrin promotes normal organelle axonal transport independently of its role assembling the MPS²⁶. Expression of WT β II-spectrin in cultured β II-spectrin null cortical neurons rescues the processivity, motility, and flux of synaptic vesicles and lysosomes²⁶. To evaluate the effects of selected β II-spectrin mutations on axonal transport, we tracked the dynamics of red fluorescent protein (RFP)-tagged LAMP1 (an endosome/lysosome vesicles marker) in control, β II-SpKO, and β II-SpHet cortical neurons using time-lapse video microscopy. As previously observed²⁶, loss of β II-spectrin impairs the bidirectional motility of LAMP1-RFP-positive vesicles and causes significant deficits in their run length and retrograde velocity (Fig. 5c,d and Extended Data Fig. 4b,c). Remarkably, β II-spectrin haploinsufficiency causes similar deficits (Fig. 5c,d and Extended Data Fig. 4b,c), indicating that 50% reduction of β II-spectrin levels is not sufficient to maintain normal organelle transport. As expected²⁶, deficits in transport of lysosomes in β II-SpKO neurons are rescued by expression of WT GFP β II-Sp. However, selected mutants that do not rescue axonal length also fail to restore normal organelle dynamics (Fig. 5c,d and Extended Data Fig. 4b,c). Within

the mutants tested, p.E892* and p.W1787* lack the PH domain, which is required for β II-spectrin coupling to organelle membranes and normal organelle transport²⁵. It is possible that the abnormal binding to molecular partners observed in other mutants unable to rescue organelle dynamics interfere with the formation of complexes between β II-spectrin and molecular motors, its coupling to organelle membranes, or its cytosol to MPS partitioning.

Collectively, our results suggest that human β II-spectrin mutations we report likely cause *SPTBN1* syndrome through molecular and cellular mechanisms that include the individual or combined effects of toxic protein aggregation, disruption of intracellular organelle transport, insufficient axonal growth, and aberrant cytoskeletal organization and dynamics, which in turn may affect neuronal connectivity and function.

***SPTBN1* variant classification**

The *SPTBN1* variants described in this study were classified using the 2015 ACMG Guidelines⁴⁶ and interpretation recommendations⁴⁷⁻⁴⁹ and are listed in Supplementary Table 2 with a summary of functional evidence herein. Of the 28 unique variants in the cohort, 14 were classified as pathogenic, 12 as likely pathogenic, and two as variants of uncertain significance (VUS). Importantly, proband P10 has two *de novo* variants in *cis* in *SPTBN1*, p.T268A and p.F344L. The p.T268A variant has two allelic variants p.T268N and p.T268S and functional studies suggesting a damaging effect support a pathogenic classification. The p.F344L variant is classified as a VUS since it is in *cis* with a pathogenic variant and showed no significant differences from wild type in functional studies, thus its contribution to the phenotype of this individual is unclear.

β II-spectrin haploinsufficiency causes cell-autonomous deficits in neuronal connectivity

β II-spectrin is widely expressed in both neurons and in brain non-neuronal cells⁵⁰. β II-spectrin loss in both neurons and glial cells of β II-SpKO mice results in significant reduction of long-range axonal tracts in the

cerebellum and of those tracts connecting cerebral hemispheres, including the CC²⁶. These white matter connectivity deficits are likely caused by the impaired axonal growth of neurons lacking β II-spectrin²⁶. Since β II-spectrin haploinsufficiency affects axonal growth *in vitro* (Fig. 5a, b), we next assessed cortical axonal connectivity in β II-SpHet mice. Consistent with a diminished axonal growth, PND25 β II-SpHet mice exhibit callosal hypoplasia (Fig. 6b,c). CC thinning is also detected by MRI in three of the probands in this cohort (P2, P10, and P28) (Fig. 1d and Table 1 and Supplementary Note), which further implicates β II-spectrin in regulating brain cytoarchitecture. Deficits in connectivity of long axonal tracts can also result from defects in neuronal migration and axonal pathfinding, which in turn can be affected by non-neuronal cells⁵¹. To determine the neuron-specific effects of β II-spectrin depletion on cortical wiring, we generated mice lacking β II-spectrin only in cortical and hippocampus projection neurons by crossing *Sptbn1*^{flox/flox} to Nex-Cre⁵² animals (*Sptbn1*^{flox/flox};Nex-Cre; henceforth referred to as β II-Sp-Nex KO) (Extended Data Fig. 5b). β II-spectrin loss or haploinsufficiency only in projection neurons is sufficient to induce CC hypoplasia (Fig. 6c,d). These results suggest that partial β II-spectrin LOF can produce neuronal miswiring in the cortex and those defects are at least in part neuron-autonomous.

β II-spectrin deficiency causes developmental and behavioral deficits in mice

Individuals bearing *SPTBN1* variants exhibit a wide range of facial dysmorphisms, brain growth defects, including microcephaly and macrocephaly, and DD (Table 1, Supplementary Note). We found that embryonic day 19 (E19) β II-SpKO mice have enlarged head circumference, and both E19 β II-SpKO and β II-SpHet animals exhibit increased distance between the eyes (Fig. 7a-c), consistent with the observed hypertelorism in some of the patients (Fig. 1c and Supplementary Note). In line with reported DD in patients, β II-SpKO mice show arrested development (Fig. 7d,e)²⁶. In addition, β II-spectrin haploinsufficiency is sufficient to yield animals of an intermediate body size and weight (Fig. 7d-f). The global DD changes observed in mice with β II-spectrin deficits arise in part due to neuronal-autonomous

effects, given that they are also observed in β II-Sp-NexKO mice that only lack the protein in cortical and hippocampal projection neurons (Extended Data Fig. 6a).

Since individuals carrying *SPTBN1* variants present with various behavioral phenotypes, including ASD, ADHD as well as learning and mild motor deficits (Table 1, Supplementary Note), we assessed behavioral effects of brain β II-spectrin deficiency in mice. First, we evaluated the effects of complete LOF using β II-SpKO mice. Because these animals do not survive longer than five weeks²⁶, they were only challenged with open field and acoustic startle tests at PND30. β II-SpKO mice had overt hyperactivity at every time point during the open field test (Fig. 7g) and profound deficits in rearing, a response requiring good hind limb function and balance (Fig. 7h). β II-SpKO mice also showed decreases in startle response amplitudes in the acoustic startle test, but normal levels of prepulse inhibition (PPI) (Extended Data Fig. 6b,c), suggesting that reduced startle responses were due to motor deficits, rather than alterations in auditory function or sensorimotor gating. This is consistent with impaired motor abilities likely due to the severe loss of cerebellar connectivity in these mice²⁶.

Our clinical, cellular, and animal data indicates that β II-spectrin haploinsufficiency, or the altered function of only one copy of *SPTBN1* due to GOF or dominant-negative effects, is sufficient to cause a neurodevelopmental disorder. Thus, we next characterized behavioral phenotypes of β II-SpHet mice, whose normal lifespan allowed for an expanded battery of tests. In contrast to β II-SpKO mice, β II-SpHet animals had normal activity during an open field test (Fig. 7i,l). Further, β II-SpHet and control mice had comparable performance in an acoustic startle test for PPI and in the rotarod test (Extended Data Fig. 6d-f), indicating that expression of half levels of β II-spectrin is sufficient to rescue motor problems. β II-SpHet mice also exhibited normal spatial and reversal learning in the Morris water maze test (Extended Data Fig. 6g,h). On the other hand, in the 3-chamber choice test, β II-SpHet demonstrated no preference for spending more time in proximity to a stranger mouse versus an empty cage and made significantly fewer entries into the side containing the stranger mouse (Fig. 7k,l). These genotype differences were not

observed in the subsequent test for social novelty preference, in which β II-SpHet and β II-SpWT littermates demonstrated the typical shift in preference to the newly introduced stranger 2 (Extended Data Fig. 6k,l). Notably, there was a non-significant trend for the β II-SpHet mice to make fewer entries than the littermate controls in the social novelty test. The lack of sociability in the β II-SpHet mice was not associated with changes in anxiety-like behavior or olfactory function (Extended Data Fig. 7m). Overall, these results suggest that β II-spectrin LOF impairs global development and has a selective impact on social motivation and reward that may contribute to the autistic features and social behavior impairments manifested in some affected individuals.

Discussion

In this study, we report for the first time the identification of *de novo* *SPTBN1* variants in individuals as a cause of a neurodevelopmental disorder most commonly characterized by motor and speech delays, ID, and various neurologic and behavioral comorbidities. In addition to DD and ID, eleven individuals in our cohort have been diagnosed with ADD/ADHD and six with ASD, with three having co-occurrence. This observation is consistent with a recent WES study of a Danish cohort of approximately 8,000 children with ASD and/or ADHD and 5,000 controls that identified *SPTBN1* as a top hit among genes with rare truncating variants co-occurring in these disorders at a significantly higher rate than in controls⁵³. *SPTBN1* variants had previously been reported in probands with ASD²⁹, Tourette²⁸, and DD (all included in our study). Noteworthy, β II-spectrin's canonical partner ankyrin-B is encoded by high confidence ASD gene *ANK2*²⁷ and some ASD patients with *ANK2* variants also exhibit ID⁵⁴. Loss of ankyrin-B isoforms in mice result in axonal transport deficits⁵⁵ and developmentally regulated defects in brain connectivity^{54,55}, two overlapping phenotypes we observed in our β II-spectrin mouse models. Although ankyrin-B and β II-spectrin are independent modulators of axonal transport²⁶, *SPTBN1* and *ANK2* may otherwise converge through mechanisms that affect other neuronal functions. For example, loss of ankyrin-B affects the polarized distribution of β II-spectrin in neurites, which gives rise to its more even portioning between

axons and dendrites causing a higher than normal prevalence of the MPS in dendrites⁵⁶. Conversely, disruption of the MPS due to loss of β II-spectrin^{24,26} may disrupt the periodic distribution of ankyrin-B and its membrane partners in axons⁵⁴, which together may be essential for critical signal transduction events⁵⁷. In addition to their strong correlation with DD, our results together with these observations support the association of *SPTBN1* pathogenic variants with ASD and ADHD.

Seizures and epilepsy were other noticeable re-occurring phenotypes in our cohort. That *SPTBN1* variants may have epileptogenic effects is not surprising, given the strong association of *de novo* and inherited variants in the partner gene *SPTAN1* (α II-spectrin) with epileptic syndromes^{5,16-21}. Although the precise pathogenic mechanism of *SPTAN1* in epilepsy is unknown, α II-spectrin protein aggregation has been reported for several of the putative pathogenic variants^{16,20}. As we show above, α II-spectrin cellular distribution can be disrupted by mutant β II-spectrin to cause these partners to co-aggregate, or otherwise continue to associate in aberrant cellular patterns. Since β II- and α II-spectrin are critically involved in localizing and stabilizing ion channels¹⁻³, going forward it will be critical to elucidate whether these tightly intertwined partners share pathways disrupted in channelopathies underlying seizures and epilepsy.

Besides the widely shared DD phenotype in our cohort, further supporting evidence of the pathogenicity of *SPTBN1* variants is the re-occurrence of *de novo* variants in the same amino acid position in unrelated individuals which are not found in the general population. These individuals share other co-occurring clinical manifestations, but also diverge in some of the clinical presentations, which may originate in part by differences in the identity of the amino acid substitution, sex, age, and genetic background. Another striking indicator of convergence in the pathogenic mechanism of the β II-spectrin mutations we report is their partial clustering (14 of 28) within the CH domains. The region of *SPTBN1* encoding the CH domains has a higher degree of missense variant constraint in the population (ExAC v.10)⁵⁸, indicating the importance of the CH domains for protein function and supporting the pathogenicity of the variants within. Our cellular and biochemical findings suggest that CH domain mutants generally affect β II-

spectrin's interaction with F-actin and α II-spectrin and result in modified spectrin/actin cytoskeleton dynamics and cellular morphology. The aberrant accumulation of mutant β II-spectrin within cytosolic aggregates suggests that a subset of the CH mutations introduce destabilizing effects on the protein structure, which is supported by our structural modeling. These changes in β II-spectrin distribution, as well as in binding to submembrane cytoskeleton partners, likely underlie GOF effects, such as aberrant neuronal membrane morphology, and contribute to LOF deficits, such as impaired organelle transport and reduced axonal growth. In turn, these cellular defects likely result in the deficient or aberrant brain connectivity and function observed in β II-spectrin-deficient mice and in patients. Interestingly, pathogenic CH domain variants have been reported in β I-spectrin⁵⁹, which cause spherocytosis, and β III-spectrin^{4,13}, which leads to cerebellar ataxia, DD, and ID, and have been shown to affect F-actin binding⁴¹. Together with our results, this evidence indicates that the abnormal modulation of actin binding by CH domain variants likely constitute a conserved pathogenic mechanism in spectrinopathies.

Like in other spectrinopathies⁴⁻²³, missense mutations affecting SR are likely to be disease-causing in the *SPTBN1* syndrome, although the molecular mechanisms are not fully understood. For example, it is not clear how p.A850G phenocopies the cellular phenotypes caused by some of the CH domain mutants. It is possible that this mutant affects β II-spectrin/F-actin dynamics through allosteric mechanisms or dominant negative effects due to overexpression. Alternatively, this and the other SR mutants may disrupt β II-spectrin association with undefined binding partners or its coupling to organelles and motor proteins, which may explain their detrimental effect on axonal growth²⁶. Our cellular assays failed to identify a potential pathogenic mechanism for a small subset of *SPTBN1* variants. However, it is possible that these variants affect other less explored neuronal β II-spectrin roles, such as dendritic and postsynaptic development and function, which are associated with ASD²⁹. Additionally, given the wide expression of β II-spectrin in non-neuronal brain cells, it will be of interest to assess if their function is affected by *SPTBN1* variants. It is likely that the clinical variability is at least partly rooted in the multifunctionality and

ubiquitous expression of β II-spectrin, although we cannot rule out that some clinical manifestations unique to affected individuals in the cohort may be caused by an alternate etiology. For example, a few individuals in the cohort have additional genetic variants that might be contributing to their clinical phenotype. Proband P19 has a pathogenic *NF1* variant (NM_000267.3:c.3449C>G; p.S1150*) and has neurofibromatosis, which could also have associated learning disabilities, but likely would not explain the behavioral challenges and autism seen in this individual. Proband P27 has a variant in *GNB1* (NM_001282539.1:c.700-1G>T) inherited from her mother also affected with delays. However, the *SPTBN1* variant was not present in the mother, and could be *de novo* or paternal as her father has moderate ID, suggesting both variants could be contributory. Finally, given the critical roles β II-spectrin plays in other organs^{60,61} and its association with other non-neurological disorders, including clinical presentations beyond the nervous system in patients in our cohort, the *SPTBN1* syndrome warrants thorough clinical assessment and further studies beyond the brain.

Materials and methods

Identification of Pathogenic *SPTBN1* Variants

Pathogenic variants in *SPTBN1* were identified by whole exome or genome sequencing performed on whole blood DNA from probands identified through diagnostic clinical practice or Institutional Review Board approved research studies. Affected individuals were identified through professional communication, connections through GeneMatcher⁶², and by searching the Undiagnosed Diseases Network (UDN) and the Deciphering Developmental Disorders (DDD) Research Study²⁹ repositories. Variants were reported according to standardized nomenclature defined by the reference human genome GRCh37 (hg19) and *SPTBN1* transcript GenBank: NM_003128.2. The minor-allele frequency of each variant was determined from genomic sequencing data derived from the gnomAD.

Patient consent

Patient consent for participation and phenotyping was obtained through the referring clinical teams. Referring clinicians were requested to complete a comprehensive questionnaire that was based upon our current understanding of the phenotypic associations of *SPTBN1*. They included sections related to neurodevelopmental screening, behavior, dysmorphology, muscular, cardiac, and other systemic phenotypic features. Consent and collection of information conformed to the recognized standards of the Declaration of Helsinki and approved by local Institutional Review Boards.

Variant interpretation and classification

SPTBN1 variants were interpreted using the NM_003128.2 transcript and splice variants were evaluated using SpliceAI²⁷ to predict the most likely mRNA splicing outcome. The *SPTBN1* variants identified in this study were classified according to the ACMG 2015 Guidelines⁴⁶. Based on the recommendations of PVS1 loss-of-function criterion under the ACMG/AMP specifications⁴⁷, PVS1_strong was used as a maximum weight of evidence. This is appropriate for this criterion as we have shown moderate clinical validity⁴⁸, unrelated probands with a consistent phenotype, and robust functional evidence showing that these nonsense variants remove downstream portions of the protein known to be essential for protein function, and that both null and haploinsufficient mouse models recapitulate disease phenotypes. The maximum weight of functional evidence (PS3) used was moderate under the ACMG/ACMP specifications⁴⁹.

Mouse lines and animal care

Experiments were performed in accordance with the guidelines for animal care of the Institutional Animal Care and Use Committee of the University of North Carolina at Chapel Hill. To generate neural progenitor-specific β II-spectrin null (*Sptbn1*^{flox/flox}/Nestin-Cre, β II-Sp-KO) and haploinsufficient (*Sptbn1*^{flox/+}/Nestin-Cre, β II-Sp-Het) mice, *Sptbn1*^{flox/flox} animals, a gift from Dr. Mathew Rasband⁵⁰, were crossed with the Nestin-Cre mouse line [B6.Cg-Tg(Nes-cre)1Kln/J, stock number 003771; The Jackson Laboratory]. *Sptbn1*^{flox/flox} animals negative for the Cre transgene were used as littermate controls in all experiments. Mice lacking

β II-spectrin in cortical projection neurons (*Sptbn1*^{flox/flox}/Nex-Cre, β II^{Sp}-Nex KO) were generated by crossing *Sptbn1*^{flox/flox} and Nex-Cre, a gift from Dr. Klaus-Armin Nave⁵², animals for multiple generations. All mice were housed at 22°C ± 2°C on a 12-hour-light/12-hour-dark cycle and fed ad libitum regular chow and water.

Generation of human β II-spectrin mutations

The human β II-spectrin cDNA was subcloned into peGFP-C3 vector (Clontech) using HindIII and SacI sites to generate the peGFP- β II^{Sp} plasmid. For purification of full-length β II-spectrin proteins, both a prescission protease site (LEVLFQGP) and a 6x histidine tag were respectively introduced between the GFP and start codon and before the C-terminal stop codon of peGFP- β II-spectrin using site-directed mutagenesis to generate the peGFP-PP- β II-^{Sp}-6xHis construct. peGFP- β II^{Sp} and peGFP-PP- β II-^{Sp}-6xHis plasmids bearing the human mutations included in the study were generated using the In-Fusion HD Cloning Plus system (Takara) and primers specific for each mutation site (Supplementary Table 3). All plasmids were verified by full-length sequencing.

Plasmids

Plasmid used in transfection experiments included: pLAMP1-RFP (Addgene plasmid #1817, gift from Walther Mothes), pmCherry-C1 (Clontech) and peGFP-C3 vector (Clontech). To generate mCherry-tagged α II-spectrin (pmCherry- α II^{Sp}), the cDNA sequence of human α II-spectrin (NM_001130438.3) was amplified by PCR as a BsrGI/XhoI fragment and cloned into the corresponding sites of pmCherry-C1 (Clontech). peGFP-C3-Y1874A- β II-spectrin and HA-tagged 220 kDa ankyrin-B (pAnkB-3X HA) plasmids were previously reported.²⁶ All plasmids were verified by full-length sequencing prior to transfection.

Antibodies

Affinity-purified rabbit antibodies against GFP and β II-spectrin, used at a 1:500 dilution for immunohistochemistry and 1:5000 for western blot, were generated by Dr. Vann Bennett laboratory and

have been previously described.^{26,51} Other antibodies used for western blot analysis and immunoprecipitation included mouse anti-GFP (1:1000, #66002-1-Ig), rabbit anti-GFP (1:1000, #50430-2-AP), rabbit anti-HA tag (1:1000, #51064-2-AP), and mouse anti-6*His tag (1:1000, # 66005-1-Ig) all from Proteintech, and rabbit anti-mCherry (1:2000, #ab24345) from Abcam. Commercial antibodies used for immunofluorescence included mouse anti-neurofilament (1:200, # 837801) from BioLegend and chicken anti-GFP (1:1000, #GFP-1020) from Aves. Secondary antibodies purchased from Life Technologies were used at 1:400 dilution for fluorescence-based detection by confocal microscopy, and included donkey anti-rabbit IgG conjugated to Alexa Fluor 568 (#A10042), donkey anti-mouse IgG conjugated to Alexa Fluor 488 (#A21202), goat anti-chicken conjugated to Alexa Fluor 488 (#A11039), and donkey anti-rat IgG conjugated to Alexa Fluor 647 (#A21247). Fluorescent signals in western blot analysis were detected using goat anti-rabbit 800CW (1:15000, #926-32211) and goat anti-mouse 680RD (1:15000, #926-68070) from LICOR.

Neuronal culture

Primary cortical neuronal cultures were established from E17 mice. Cortices were dissected in Hibernate E (Life Technologies) and digested with 0.25% trypsin in HBSS (Life Technologies) for 20 min at 37°C. Tissue was washed three times with HBSS and dissociated in DMEM (Life Technologies) supplemented with 5% fetal bovine serum (FBS, Genesee), and gently triturated through a glass pipette with a fire-polished tip. Dissociated cells were filtered through a 70 µm cell strainer to remove any residual non-dissociated tissue and plated onto poly-D-lysine-coated 1.5 mm coverglasses or dishes (MatTek) for transfection and time-lapse microscopy imaging. For all cultures, media was replaced 3 hours after plating with serum-free Neurobasal-A medium containing B27 supplement (Life Technologies), 2 mM Glutamax (Life Technologies), and penicillin/streptomycin (Life Technologies). 5 µM cytosine-D-arabinofuranoside (Sigma) was added to the culture medium to inhibit the growth of glial cells three days after plating. Neurons were fed twice a week with freshly made culture medium until use.

Plasmid transfection for time-lapse live imaging and immunofluorescence analysis

For time-lapse imaging experiments DIV5 cortical neurons were co-transfected with 1µg of each pLAMP1-RFP and peGFP-βIIISp plasmids using lipofectamine 2000 (Life Technologies) and imaged 48-96 hours after transfection. For experiments that evaluate axonal length, DIV3 control and βIIISp-Het neurons were transfected with 500 ng of pmCherry-C1 and 1µg of peGFP-C3. βII-SpKO neurons were transfected with 500 ng of pmCherry-C1 and 1µg of peGFP-βIIISp rescue plasmids bearing full-length wildtype of mutant β2-spectrin. Neurons were processed for immunofluorescence 5 days after transfection. Immunofluorescence evaluations of βII-spectrin distribution in HEK293 cells was conducted in cells transfected with 100 ng of peGFP-βIIISp plasmids, or co-transfected with 100 ng of each peGFP-βIIISp and pmCherry-αIIISp plasmids 48 hours post-transfection.

Plasmid transfection for biochemistry analysis

All transfections were conducted in HEK293 cells grown in 10 cm culture plates using the calcium phosphate transfection kit (Takara). To purify full-length βII-spectrin proteins, cells were transfected with 8 µg of peGFP-PP-βII-Sp-6xHis plasmids. To determine levels and stability of βII-spectrin proteins, HEK293T cells were co-transfected with 8 µg of eGFP-PP-βII-Sp-6xHis and 4 µg of pmCherry-C1 plasmids. To determine interaction between ankyrin-B and βII-spectrin, cells were co-transfected with 8µg of each peGFP-PP-βII-Sp-6xHis and pAnkB-3X HA plasmids. For assessment of binding between βII-spectrin and αII-spectrin, cells were separately transfected with 8 µg of peGFP-PP-βII-Sp-6xHis or 4 µg peGFP-C3 and 8 µg of pmCherry-αIIISp.

Histology and immunohistochemistry

Brains from mice two-weeks and older were fixed by transcardial perfusion with phosphate-buffered saline (PBS) and 4% paraformaldehyde (PFA) followed by overnight immersion in the same fixative. Brains from PND0-PND14 mice were fixed by direct immersion in 4% PFA for 36 hours. After fixation, brains were

rinsed with PBS, transferred to 70% ethanol for at least 24 hours, and paraffin-embedded. 7- μ m coronal and sagittal brain sections were cut using a Leica RM2155 microtome and mounted on glass slides. Sections were analyzed by hematoxylin and eosin (H&E) staining or immunostaining. For antibody staining, sections were deparaffinized and rehydrated using a standard protocol of washes: 3 $\times$ 3-min Xylene washes, 3 $\times$ 2-min 100% ethanol washes, and 1 $\times$ 2-min 95%, 80%, and 70% ethanol washes followed by at least 5 min in PBS. Sections were then processed for antigen retrieval using 10 mM sodium citrate, pH 6 in the microwave for 20 min. Sections were allowed to cool, washed in PBS, and blocked using antibody buffer (2% bovine serum albumin (BSA), 1% fish oil gelatin, 5% donkey serum, and 0.02% Tween 20 in PBS) for 1 hour at room temperature. Tissue sections were then subsequently incubated overnight with primary antibodies at 4°C and with secondary antisera for 1.5 hours at 4°C, washed with PBS, and mounted with Prolong Gold Antifade reagent (Life Technologies). Neuronal cultures and HEK293 cells were washed with cold PBS, fixed with 4% PFA for 15 min, and permeabilized with 0.2% Triton-X100 in PBS for 10 min at room temperature. Neurons and HEK293 cells were blocked in antibody buffer for 1 hour at room temperature and processed for fluorescent staining as tissue sections. For actin labeling, Alexa Fluor 568- or Alexa Fluor 633-conjugated phalloidin (1:100) was added to the secondary antibody mix. DAPI was added to the last PBS rinse for nuclei staining.

Immunoblots

Protein homogenates from mouse brains or transfected cells were prepared in 1:9 (wt/vol) ratio of homogenization buffer (8M urea, 5% SDS (wt/vol), 50mM Tris pH 7.4, 5mM EDTA, 5mM N-ethylmeimide, protease and phosphatase inhibitors) and heated at 65°C for 15 min to produce a clear homogenate. Total protein lysates were mixed at a 1:1 ratio with 5x PAGE buffer (5% SDS (wt/vol), 25% sucrose (wt/vol), 50mM Tris pH 8, 5mM EDTA, bromophenol blue) and heated for 15 min at 65°C. Samples were resolved by SDS-PAGE on 3.5-17.5% acrylamide gradient gels in Fairbanks Running Buffer (40mM Tris pH 7.4, 20mM NaAc, 2mM EDTA, 0.2%SDS (wt/vol)). Proteins were transferred overnight onto 0.45 μ m nitrocellulose

membranes (#1620115, BioRad) at 4°C. Transfer efficiency was determined by Ponceau-S stain. Membranes were blocked in TBS containing 5% non-fat milk for 1 hour at room temperature and incubated overnight with primary antibodies diluted in antibody buffer (TBS, 5% BSA, 0.1% Tween-20). After 3 washes in TBST (TBS, 0.1% Tween-20), membranes were incubated with secondary antibodies diluted in antibody buffer for two hours at room temperature. Membranes were washed 3x for 10 minutes with TBST and 2x for 5 minutes in TBS. Protein-antibody complexes were detected using the Odyssey® CLx Imaging system (LI-COR).

Immunoprecipitation

For immunoprecipitation experiments, total protein homogenates from transfected HEK293 cells were prepared in TBS containing 150 mM NaCl, 0.32 M sucrose, 2 mM EDTA, 1% Triton X-100, 0.5% NP40, 0.1% SDS, and complete protease inhibitor cocktail (Sigma). Cell lysates were incubated with rotation for 1 hour at 4°C and centrifuged at 100,000 x g for 30 min. Soluble fractions were collected and precleared by incubation with Protein-G magnetic beads (#1614023, Bio-Rad) for 1 hour in the cold. Samples were subjected to immunoprecipitation in the presence of protein-G magnetic beads/antibody or protein-G magnetic beads/isotype control complexes overnight at 4°C. Immunoprecipitation samples were resolved by SDS-PAGE and western blot and signal detected using the Odyssey® CLx imaging system.

Purification of full-length β II-spectrin proteins

Ten 10-cm plates of HEK293 cells expressing each peGFP-PP- β II-Sp-6xHis construct were used per purification. Total protein homogenates from transfected HEK293 cells were prepared in TBS containing 150 mM NaCl, 0.32 M sucrose, 2 mM EDTA, 1% Triton X-100, 0.5% NP40, 0.1% SDS, and complete protease inhibitor cocktail (Sigma) (IP buffer). Cell lysates were incubated with rotation for 1 hour at 4°C and centrifuged at 100,000 x g for 30 min. Soluble fractions were incubated overnight with Protein A/G magnetic beads (#88802, Life Technologies) coupled to GFP antibodies with rotation at 4°C. Beads were

extensively washed with IP buffer, followed by washes in TBS containing 300 mM NaCl, and TBS. Full-length β II-spectrin proteins were eluted from GFP-protein A/G magnetic beads by incubation with HRV-3C protease, which cleaves between GFP and the start codon of β II-spectrin in prescission protease buffer (25 mM HEPES, 150 mM NaCl, 1 mM EDTA, 1 mM DTT) for 36 hours at 4°C. The efficiency of cleavage and purity of the eluates was analyzed by western blot using validated antibodies specific for β II-spectrin and GFP and 6*His tags, and by Coomassie blue stain. Eluates were concentrated using Pierce™ Protein Concentrators PES.

Pulldown assays

For detection of β II-spectrin/ α II-spectrin complexes, control and mutant GFP- β IIsp proteins were coupled to GFP-bound Protein-A/G magnetic beads and incubated with lysates from HEK293 cells expressing mCherry- α IIsp in IP buffer overnight at 4°C. Beads complexes were washed sequentially with IP buffer, followed by washes in TBS containing 400 mM NaCl, and TBS. Proteins were eluted in 5x PAGE loading buffer and analyzed by SDS-PAGE and western blot.

Actin co-sedimentation assay

Interaction between purified full-length β II-spectrin proteins and actin was evaluated using the Actin Binding Protein Spin-Down Biochem Kit (#BK001, Cytoskeleton) following the manufacturer's recommendations. In brief, full-length β II-spectrin (1 mg/ml) and α -actinin (20 mg/ml, positive control) were prepared in general actin buffer (5 mM Tris-HCl pH 8.0 and 0.2 mM CaCl_2) and centrifuged at 150,000 x g for 1 h at 4°C. F-actin (1 mg/ml) was prepared by incubation of purified actin in general actin buffer for 30 min on ice followed by the actin polymerization step in actin polymerization buffer (50 mM KCl, 2 mM MgCl_2 , 1 mM ATP) for 1 hour at 24°C. F-actin (21 μ M) was incubated with either β II-spectrin (10 μ M), α -actinin (2 μ M), or BSA (2 μ M, negative control) for 30 min at 24°C. F-actin-protein complexes were pelleted

by ultracentrifugation at 150,000 x g for 1.5 h at 24°C. The presence of F-actin together with interacting proteins was assessed in the supernatant and pellet fractions by SDS-PAGE and Coomassie blue stain.

Fluorescence image acquisition and image analysis

Confocal microscope images were taken using a Zeiss LSM780 using 405-, 488-, 561-, and 633-nm lasers. Single images and Z-stacks with optical sections of 1 µm intervals and tile scans were collected using the ×10 (0.4 NA) and ×40 oil (1.3 NA) objective lens. Images were processed, and measurements taken and analyzed, using Zeiss Zen, Volocity (Perkin Elmer), or NIH ImageJ software. Three-dimensional rendering of confocal Z-stacks was performed using Imaris (Bitplane).

Time-lapse video microscopy and movie analyses

Live microscopy of neuronal cultures was carried out using a Zeiss 780 laser scanning confocal microscope (Zeiss) equipped with a GaAsP detector and a temperature- and CO₂-controlled incubation chamber. Movies were taken in the mid-axon and captured at a rate of 1 frame/second for time intervals ranging from 60-300 seconds with a 40x oil objective (1.4NA) using the zoom and definite focus functions. Movies were processed and analyzed using ImageJ (<http://rsb.info.nih.gov/ij>). Kymographs were obtained using the KymoToolBox plugin for ImageJ (https://github.com/fabricecordelieres/IJ_KymoToolBox). In details, space (x axis in µm) and time (y axis in sec) calibrated kymographs were generated from video files. In addition, the KymoToolBox plugin was used to manually follow a subset of particles from each kymograph and report the tracked particles on the original kymograph and video files using a color code for movement directionality (red for anterograde, green for retrograde and blue for stationary particles). Quantitative analyses were performed manually by following the trajectories of individual particles to calculate dynamic parameters including, net and directional velocities and net and directional run length, as well as time of pause or movement in a direction of transport. Anterograde and retrograde motile vesicles were defined as

particles showing a net displacement >3 μm in one direction. Stationary vesicles were defined as particles with a net displacement <2 μm .

Statistical analysis

GraphPad Prism (GraphPad Software) was used for statistical analysis. Two groups of measurements were compared by unpaired Student's t test. Multiple groups were compared by one-way ANOVA followed by a Dunnett's multiple comparisons test.

Molecular modeling of *SPTBN1* Variants

We used the closed conformation of utrophin CH1-CH2 closed dimer (PDB 1qag)³⁹ as a template for the analogous β II-spectrin conformation to estimate its electrostatic surface profile. Molecular structures from the 6.9 Å cryo-EM structure of the CH1 actin-binding domain of β III-spectrin bound to F-actin (PDB ID 6anu)⁴¹ and the structure of the CH2 domain of β II-spectrin (PDB ID 1bkr)⁴⁰ were used for protein-protein docking predictions. The ClusPro protein-protein docking webserver^{42,43} was used to 1) dock the CH1 domain of spectrin onto F-actin, 2) dock the CH2 domain of spectrin onto F-actin, and 3) dock the CH2 domain of spectrin onto the CH1 domain of spectrin. The CH1 structure used for the dockings reported here was the model of the CH1 domain of β III-spectrin from 6anu (chain a)⁴¹. This CH1 model was built based on the crystal structure of plectin (PDB ID 1mb8)⁶³ by I-TASSER⁶⁴. The CH1 domain of β III-spectrin shares 95% sequence identity with the CH1 domain of β II-spectrin. The actin model corresponded to chains A-F of 6anu, which in turn was generated from the cryo-EM structure of actin (PDB ID 5jlh)⁶⁵. The molecular structure of the CH2 domain of β II-spectrin from 1bkr was of a 1.1 Å crystal structure⁴⁰.

To identify the inactive closed conformation of the tandem domain (CH1-CH2) of β II-spectrin, the CH2 domain of β II-spectrin was docked onto the CH1 domain of β II-spectrin using the ClusPro webserver. The top 15 docking poses for each of the four scoring algorithms were evaluated for the placement of β II-spectrin residue L250 from the CH2 domain at the interface of the CH2/CH1 closed conformation. The top

docking pose in the electrostatic scoring algorithm corresponded to a pose with a deeply buried L250 at the interface of the CH1/CH2 complex. The mutation of the equivalent residue in β III-spectrin (L253P) might disrupt the closed structure and drive the spectrin ensemble to a more open state suitable for binding to actin⁴⁰. This same docking pose was also a top docking pose (pose 4) within the set of poses calculated by the balanced scoring algorithm. This pose was used for evaluation of the β II-spectrin mutants. It was also the same pose compared to the actinin-4 (PDB ID 6oa6) (unpublished) and utrophin (PDB ID 1qag)³⁹ closed conformations.

For each of the three ClusPro protein docking analyses, the webserver provided up to 30 docking poses for each of four scoring algorithms (balanced; electrostatic-favored; hydrophobic-favored; VdW+Elec). The top 15 poses from each of the four scoring algorithms were included in the final analysis. For the dockings of the CH1 and CH2 domains from β II-spectrin onto F-actin, several of the top docking poses were to the ends of the actin segment defined as the receptor. These docking poses were immediately rejected as other actin molecules would be binding at those locations in F-actin and these sites would not be available for binding to spectrin. For CH1 docking onto F-actin, the remaining poses within the top eight docking poses predicted by the balanced and electrostatic scoring algorithms almost all corresponded to the location and orientation of CH1 molecules on actin as defined by the cryo-EM structure 6anu. For CH2 docking onto F-actin, the remaining poses within the top 8 docking poses predicted by the balanced and electrostatic scoring algorithms almost all corresponded to symmetry-related locations and poses on the F-actin. In addition, the predicted orientation of the CH2 molecules on F-actin was consistent with the known binding site of the CH1 domains, as judged by the length of the linker that would be required to join the C-terminus of the docked CH1 domain to the N-terminus of the docked CH2 domain.

β II-spectrin is a large multi-domain protein that requires a different approach for each type of domain. The SR have relatively low sequence identity to each other, and only a few have been experimentally solved, requiring independent models to be generated for each. We used RaptorX⁶⁶ homology modeling

to generate each model and assembled them into a linear conformation using Discovery Studio [Dassault Systèmes BIOVIA, Discovery Studio Modeling Environment, Release 2019, San Diego: Dassault Systèmes. 2019]. We calculated protein electrostatics using APBS⁶⁷ and visualized structures using PyMOL [The PyMOL Molecular Graphics System, Version 2.0.7 Schrödinger, LLC.]. Individual spectrin repeats were also superimposed onto each other using a geometric algorithm⁶⁸ as implemented in PyMOL, to investigate patterns across the fold.

Behavioral assessment

Animals. Because the *Sptbn1*^{flox/flox}/Nestin-Cre (βII-SpKO) mice have early mortality (typically between PND30 and PND40), testing in these mice was conducted late in the juvenile period. Subjects were 15 wildtype (*Sptbn1*^{flox/flox}/+, βII-SpWT) and 5 βII-SpKO mice, taken from 5 litters. βII-SpKO mice were evaluated in two tests: open field (at PD 28-31) and acoustic startle (at PD 29-32). *Sptbn1*^{flox/+}/Nestin-Cre (βII-SpHet), which have normal survival rates, were subjected to a more expansive battery of tests. βII-SpHet mice (n=12 per genotype, all males) underwent the following tests, with order planned so that more stressful procedures occurred closer to the end of the study.

Age (wk)	Procedure
5	Elevated plus maze test for anxiety-like behavior
6	Locomotor activity and exploration in a 1-hr open field test
7	Rotarod test for motor coordination and motor learning
8-9	Social approach in a three-chamber choice test
9-10	Marble-bury assay for anxiety-like behavior and perseverative responses
	Prepulse inhibition of acoustic startle responses
10-11	Buried food test for olfactory ability

862 **11-12** Morris water maze; visible platform test for vision and swimming ability

863 **12-13** Water maze, hidden platform test for spatial learning

864 **13-14** Reversal learning in water maze

865 **Elevated plus maze.** A five-min test for anxiety-like behavior was carried out on the plus maze (elevation,
866 50 cm H; open arms, 30 cm L; closed arms, 30 cm L, walls, 20 cm H). Mice were placed in the center (8 cm
867 x 8 cm) at the beginning of the test. Measures were taken of percent open arm time and open arm entries,
868 and total number of arm entries.

869 **Open field.** Exploratory activity was evaluated by a 1-hr test (30-min for β II-SpKO mice) in a novel open
870 field chamber (41 cm x 41 cm x 30 cm) crossed by a grid of photobeams (VersaMax system, AccuScan
871 Instruments). Counts were taken of photobeam breaks in 5-min intervals, with separate measures for
872 locomotor activity (total distance traveled) and vertical rearing movements. Anxiety-like behavior was
873 assessed by measures of time spent in the center region.

874 **Accelerating rotarod.** Mice were first given three trials on the rotarod (Ugo Basile, Stoelting Co.), with 45
875 seconds between each trial. Two additional trials were conducted 48 hr later, to evaluate consolidation
876 of motor learning. Rpm (revolutions per minute) progressively increased from 3 to a maximum of 30 rpm.
877 across five minutes (the maximum trial length), and latency to fall from the top of the rotating barrel was
878 recorded.

879 **Social approach in a three-chamber choice test.** Mice were evaluated for the effects of *Sptbn1* deficiency
880 on social preference. The procedure had three 10-minute phases: habituation, sociability, and social
881 novelty preference. In the sociability phase, mice were presented with a choice between proximity to an
882 unfamiliar C57BL/6J adult male ("stranger 1"), versus an empty cage. In the social novelty phase, mice
883 were presented with the already-investigated stranger 1 and a new unfamiliar mouse ("stranger 2"). The
884 test was carried out in a rectangular, three-chambered Plexiglas box (60 cm L, 41.5 cm W, 20 cm H). An

automated image tracking system (Noldus Ethovision) provided measures of time in spent within 5 cm proximity to each cage and entries into each side of the social test box.

Marble-burying. Mice were tested for exploratory digging in a Plexiglas cage, placed inside a sound-attenuating chamber with ceiling light and fan. The cage floor had 5 cm of corncob bedding, with 20 black glass marbles (14 mm diameter) set up in a 5 X 4 grid on top of the bedding. Measures were taken of the number of marbles buried by the end of the 30-min test.

Buried food test. Mice were presented with an unfamiliar food (Froot Loops, Kellogg Co.) in the home cage several days before the test. All home cage food was removed 16-24 hr before the test. The assay was conducted in a tub cage (46 cm L, 23.5 cm W, 20 cm H), containing paper chip bedding (3 cm deep). One Froot Loop was buried in the cage bedding, and mice were given 15 min to locate the buried food. Latency to find the food was recorded.

Acoustic startle. This procedure was used to assess auditory function, reactivity to environmental stimuli, and sensorimotor gating. The test was based on the reflexive whole-body flinch, or startle response, that follows exposure to a sudden noise. Mice were evaluated for startle magnitude and prepulse inhibition, which occurs when a weak prestimulus leads to a reduced startle in response to a subsequent louder noise. Startle amplitudes were measured by force displacement of a piezoelectric transducer (SR-Lab, San Diego Instruments). The test had 42 trials (7 of each type): no-stimulus trials, trials with the acoustic startle stimulus (40 msec; 120 dB) alone, and trials in which a prepulse stimulus (20 msec; either 74, 78, 82, 86, or 90 dB) occurred 100 msec before the onset of the startle stimulus. Levels of prepulse inhibition at each prepulse sound level were calculated as $100 - [(response\ amplitude\ for\ prepulse\ stimulus\ and\ startle\ stimulus\ together / response\ amplitude\ for\ startle\ stimulus\ alone) \times 100]$.

Morris water maze. The water maze (diameter = 122 cm) was used to assess spatial and reversal learning, swimming ability, and vision. The procedure had three phases: visible platform, acquisition in the hidden

platform task, and reversal learning (with the platform moved to a new location). For each phase, mice were given 4 60-sec trials per day. Measures were taken of time to find the escape platform (diameter = 12 cm) and swimming velocity by an automated tracking system (Noldus Ethovision). Criterion for learning was an average group latency of 15 sec or less to locate the platform. At the end of the acquisition and reversal phases, mice were given a one-min probe trial in the maze without the platform. Selective quadrant search was evaluated by measuring number of crosses over the location where the platform (the target) had been placed during training, versus the corresponding areas in the other three quadrants.

Statistical Analyses for behavioral tests

All testing was conducted by experimenters blinded to mouse genotype. Statview (SAS, Cary, NC) was used for data analyses. One-way or repeated measures analysis of variance (ANOVA) were used to determine effects of genotype. Post-hoc analyses were conducted using Fisher's Protected Least Significant Difference (PLSD) tests only when a significant F value was found in the ANOVA. For all comparisons, significance was set at $p < 0.05$.

Web Resources

Genome Aggregation Database (GnomAD), <https://gnomad.broadinstitute.org/>

ClinVar, <https://www.ncbi.nlm.nih.gov/clinvar/>

Combined Annotation Dependent Depletion (CADD), <https://cadd.gs.washington.edu/>

Mutation Taster, <http://www.mutationtaster.org/>

PolyPhen2, <http://genetics.bwh.harvard.edu/pph2/>

Protein Variation Effect Analyzer (PROVEAN), <http://provean.jcvi.org/index.php>

Sorting Intolerant from Tolerant (SIFT), <https://sift.bii.a-star.edu.sg/>

PredictSNP2, <http://loschmidt.chemi.muni.cz/predictsnp2/>

M-CAP, <http://bejerano.stanford.edu/mcap/>

Acknowledgments

We thank all the families who participated in this study. We thank Drs. Matthew Rasband and Klaus-Armin Nave for the gift of the β II-spectrin conditional null and the Nex-cre mice, respectively. We thank Dr. Natallia V. Riddick for her assistance with the behavioral studies. We thank Drs. Beverly Koller and Karen Mohlke for their insightful comments on this manuscript and Dr. James Bear for helpful discussions. M.A.C, L.E.S-R., and E.W.K. were supported by the Center for Individualized Medicine at Mayo Clinic. D.N.L. was supported by the University of North Carolina at Chapel Hill (UNC-CH) School of Medicine as a Simmons Scholar, by the National Ataxia Foundation, and by the US National Institutes of Health (NIH) grant R01NS110810. Microscopy was performed at the UNC-CH Neuroscience Microscopy Core Facility, supported, in part, by funding from the NIH-NINDS Neuroscience Center Grant P30 NS045892 and the NIH-NICHD Intellectual and Developmental Disabilities Research Center Support Grant U54 HD079124, which also supported, in part, the behavioral studies. Research reported in this manuscript was supported by the NIH Common Fund, through the Office of Strategic Coordination/Office of the NIH Director under Award Number U01HG007672 (Duke University to Dr. Vandana Shashi). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH. This research was made possible through access to the data and findings generated by the 100,000 Genomes Project. The 100,000 Genomes Project is managed by Genomics England Limited (a wholly owned company of the Department of Health and Social Care). The 100,000 Genomes Project is funded by the National Institute for Health Research and NHS England. The Wellcome Trust, Cancer Research UK and the Medical Research Council have also funded research infrastructure. The 100,000 Genomes Project uses data provided by patients and collected by the National Health Service as part of their care and support.

Author Contributions

M.A.C. and D.N.L. conceived and planned the study with input from Q.K.T and R.C.S. M.A.C. managed the collection, analysis, and interpretation of patient clinical data with Q.K.T., R.C.S., and D.N.L. D.N.L. designed the cell biology, histology, and biochemistry studies, performed these with K. A. B., B. A. C., D. A., and S. D., and analyzed the data. S.T., M.T.Z., B.T. and D.N.L. performed the structural modeling. K.M.H. and S.M. performed the mouse behavioral studies. M.C.S. contributed reagents. M.A.C. and D.N.L. wrote the manuscript with contributions from R.C.S., S.M, M.T.Z, and B.T. E.W.K. and D.N.L. supervised the study. All other authors including Q.K.T. and R.C.S. contributed clinical data. All authors approved the final manuscript.

Competing interests

E.T., R. E.P., Y.S., E.A.N., and A.B. are employees of GeneDx, Inc. E.E.E. The authors declare no other competing interests.

References

1. Bennett, V. & Lorenzo, D. N. Spectrin- and ankyrin-based membrane domains and the evolution of vertebrates. *Curr. Top. Membr.* **72**, 1-37 (2013).
2. Bennett, V. & Lorenzo, D. N. An adaptable spectrin/ankyrin-based mechanism for long-range organization of plasma membranes in vertebrate tissues. *Curr. Top. Membr.* **77**, 143-184 (2016).
3. Lorenzo, D. N. Cargo hold and delivery: Ankyrins, spectrins, and their functional patterning of neurons. *Cytoskeleton (Hoboken)*, **77**, 129-148 (2020).
4. Ikeda, Y., et al. Spectrin mutations cause spinocerebellar ataxia type 5. *Nat. Genet.* **38**, 184–190 (2006).

- 973 5. Saitsu, H., et al. Dominant-negative mutations in alpha-II spectrin cause West syndrome with severe
974 cerebral hypomyelination, spastic quadriplegia, and developmental delay. *Am. J. Hum. Genet.* **86**,
975 881–891 (2010).
- 976 6. Wang, C. C., et al. β IV spectrinopathies cause profound intellectual disability, congenital hypotonia,
977 and motor axonal neuropathy. *Am. J. Hum. Genet.* **102**, 1158–1168 (2018).
- 978 7. Jacob, F. D., Ho, E. S., Martinez-Ojeda, M., Darras, B. T. & Khwaja, O. S. Case of infantile onset
979 spinocerebellar ataxia type 5. *J. Child Neurol.* **28**, 1292–1295 (2013).
- 980 8. Parolin Schnekenberg, R., et al. De novo point mutations in patients diagnosed with ataxic cerebral
981 palsy. *Brain*, **138**, 1817–1832 (2015).
- 982 9. Nuovo, S., et al. Between SCA5 and SCAR14: delineation of the SPTBN2 p.R480W-associated
983 phenotype. *Eur. J. Hum. Genet.* **26**, 928–929 (2018).
- 984 10. Nicita, F., et al. Heterozygous missense variants of SPTBN2 are a frequent cause of congenital
985 cerebellar ataxia. *Clin. Genet.* **96**, 169-175 (2019).
- 986 11. Mizuno, T., et al. Infantile-onset spinocerebellar ataxia type 5 associated with a novel SPTBN2
987 mutation: A case report. *Brain Development*, **41**, 630-633 (2019).
- 988 12. Accogli, A., et al. Heterozygous missense pathogenic variants within the second spectrin repeat of
989 SPTBN2 lead to infantile-onset cerebellar ataxia. *J. Child Neurol.* **35**, 106-110 (2019).
- 990 13. Lise, S., et al. Recessive mutations in SPTBN2 implicate β -III spectrin in both cognitive and motor
991 development. *PLoS Genet.* **8**, e1003074 (2012).
- 992 14. Yıldız Bölükbaşı, E., et al. Progressive SCAR14 with unclear speech, developmental delay, tremor, and
993 behavioral problems caused by a homozygous deletion of the SPTBN2 pleckstrin homology domain.
994 *Am. J. Med. Genet. Part A*, **173**, 2494-2499 (2017).

- 995 15. Al-Muhaizea, M., et al. A novel homozygous mutation in SPTBN2 leads to spinocerebellar ataxia in a
996 consanguineous family: Report of a new infantile-onset case and brief review of the literature.
997 *Cerebellum*, **17**, 276-285 (2018).
- 998 16. Writzl, K., et al. Early onset West syndrome with severe hypomyelination and coloboma-like optic
999 discs in a girl with SPTAN1 mutation. *Epilepsia*, **53**, e106–e110 (2012).
- 1000 17. Hamdan, F. F., et al. Identification of a novel in-frame de novo mutation in SPTAN1 in intellectual
1001 disability and pontocerebellar atrophy. *Eur. J. Hum. Genet.* **20**, 796–800 (2012).
- 1002 18. Nonoda, Y. et al. Progressive diffuse brain atrophy in West syndrome with marked hypomyelination
1003 due to SPTAN1 gene mutation. *Brain Development*, **35**, 280–283 (2013).
- 1004 19. Tohyama, J., et al. SPTAN1 encephalopathy: distinct phenotypes and genotypes. *J. Hum. Genet.* **60**,
1005 167-173 (2015).
- 1006 20. Syrbe, S., et al. Delineating SPTAN1 associated phenotypes: from isolated epilepsy to
1007 encephalopathy with progressive brain atrophy. *Brain*, **140**, 2322-2336 (2017).
- 1008 21. Beijer, D., et al. Nonsense mutations in alpha-II spectrin in three families with juvenile onset
1009 hereditary motor neuropathy. *Brain*, **142**, 2605-2616 (2019).
- 1010 22. Knierim, E., et al. A recessive mutation in beta-IV-spectrin (SPTBN4) associates with congenital
1011 myopathy, neuropathy, and central deafness. *Hum. Genet.* **136**, 903-910 (2017).
- 1012 23. Häusler, M. G., et al. A novel homozygous splice-site mutation in the SPTBN4 gene causes axonal
1013 neuropathy without intellectual disability. *Eur. J. Med. Genet.* **63**, 103826 (2020).
- 1014 24. Xu, K., Zhong, G., Zhuang, X. Actin, spectrin, and associated proteins form a periodic cytoskeletal
1015 structure in axons. *Science*, **339**, 452-456 (2013).

- 1016 25. Cheney, R., Hirokawa, N., Levine, J., & Willard, M. Intracellular movement of fodrin. *Cell Motility*, **3**,
1017 649–655 (1983).
- 1018 26. Lorenzo, D.N., et al. β II-spectrin promotes mouse brain connectivity through stabilizing axonal
1019 plasma membranes and enabling axonal organelle transport. *Proc. Natl. Acad. Sci. U S A.* **116**, 15686-
1020 15695 (2019).
- 1021 27. Jaganathan, K., et al. Predicting Splicing from Primary Sequence with Deep Learning. *Cell*, **176**, 535-
1022 548 (2019).
- 1023 28. Willsey, A. J., Fernandez, T. V., Yu, D., et al. De novo coding variants are strongly associated with
1024 Tourette Disorder. *Neuron*, **94**, 486–499 (2017).
- 1025 29. Firth, H.V., et al. DECIPHER: Database of chromosomal imbalance and phenotype in humans using
1026 ensembl resources. *Am. J. Hum. Genet.* **84**, 524-533 (2009).
- 1027 30. Iossifov, I., et al. The contribution of de novo coding mutations to autism spectrum disorder. *Nature*,
1028 **515**, 216–221. (2014).
- 1029 31. Karczewski, K. J., et al. The mutational constraint spectrum quantified from variation in 141,456
1030 humans. *Nature*, **581**, 434-443 (2020).
- 1031 32. Speicher, D. W., Weglarz, L., DeSilva, T. M. Properties of human red cell spectrin heterodimer (side-
1032 to-side) assembly and identification of an essential nucleation site. *J. Biol. Chem.* **267**, 14775-14782
1033 (1992).
- 1034 33. Li, X., Bennett, V. Identification of the spectrin subunit and domains required for formation of
1035 spectrin/adducin/actin complexes. *J. Biol. Chem.* **271**, 15695-15702 (1996).
- 1036 34. Davis, L., Abdi, K., Machius, M., et al. Localization and structure of the ankyrin-binding site on beta2-
1037 spectrin. *J. Biol. Chem.* **284**, 6982-6987 (2009).

- 1038 35. Bignone, P. A., Baines, A. J. Spectrin alpha II and beta II isoforms interact with high affinity at the
1039 tetramerization site. *Biochem. J.* **374**, 613-624 (2003).
- 1040 36. Hyvönen, M., et al. Structure of the binding site for inositol phosphates in a PH domain. *EMBO J.* **14**,
1041 4676-4685 (1995).
- 1042 37. Korenbaum E, Rivero F. Calponin homology domains at a glance. *J. Cell Sci.* **115**, 3543-3545 (2002).
- 1043 38. Yin, L. M., Schnoor, M., Jun, C. D. Structural characteristics, binding partners and related diseases of
1044 the calponin homology (CH) domain. *Front. Cell. Dev. Biol.* **8**, 342 (2020).
- 1045 39. Keep NH, Winder SJ, Moores CA, Walke S, Norwood FL, Kendrick-Jones J. Crystal structure of the
1046 actin-binding region of utrophin reveals a head-to-tail dimer. *Structure*, **7**, 1539-1546 (1999).
- 1047 40. Bañuelos S, Saraste M, Djinić Carugo K. Structural comparisons of calponin homology domains:
1048 implications for actin binding. *Structure*, **6**, 1419-1431 (1998).
- 1049 41. Avery, A.W., et al. Structural basis for high-affinity actin binding revealed by a β -III-spectrin SCA5
1050 missense mutation. *Nat. Commun.* **8**, 1350 (2017).
- 1051 42. Vajda, S., et al. New additions to the ClusPro server motivated by CAPRI. *Proteins: Structure*,
1052 *Function, and Bioinformatics.* **85**, 435-444 (2017).
- 1053 43. Kozakov, D., et al. The ClusPro web server for protein-protein docking. *Nat. Prot.* **12**, 255-278 (2017).
- 1054 44. Galiano, M. R., et al. A distal axonal cytoskeleton forms an intra-axonal boundary that controls axon
1055 initial segment assembly. *Cell*, **149**, 1125-1139 (2012).
- 1056 45. Sleight, J. N., Rossor, A. M., Fellows, A. D., Tosolini, A. P., Schiavo, G. Axonal transport and
1057 neurological disease. *Nat. Rev. Neurol.* **15**, 691-703 (2019).

- 1058 46. Richards, S., et al. Standards and guidelines for the interpretation of sequence variants: a joint
1059 consensus recommendation of the American College of Medical Genetics and Genomics and the
1060 Association for Molecular Pathology. *Genet. Med.* **17**, 405-424 (2015).
- 1061 47. Abou Tayoun, A. N., et al. Recommendations for interpreting the loss of function PVS1 ACMG/AMP
1062 variant criterion. *Hum. Mutat.* **39**, 1517-1524 (2018).
- 1063 48. Strande, N. T., et al. Evaluating the Clinical Validity of Gene-Disease Associations: An Evidence-Based
1064 framework developed by the Clinical Genome Resource. *Am. J. Hum. Genet.* **100**, 895-906 (2017).
- 1065 49. Brnich, S. E., et al. Recommendations for application of the functional evidence PS3/BS3 criterion
1066 using the ACMG/AMP sequence variant interpretation framework. *Genome Med.* **12**, 3 (2019).
- 1067 50. Susuki, K., et al. Glial β II spectrin contributes to paranode formation and maintenance. *J. Neurosci.*
1068 **38**, 6063-6075 (2018).
- 1069 51. Gobius, I. et al. Astroglial-mediated remodeling of the interhemispheric midline is required for the
1070 formation of the corpus callosum. *Cell Rep.* **17**, 735-747 (2016).
- 1071 52. Goebbels, S., et al. Genetic targeting of principal neurons in neocortex and hippocampus of NEX-Cre
1072 mice. *Genesis*, **44**, 611-621 (2006).
- 1073 53. Satterstrom, F. K., et al. Autism spectrum disorder and attention deficit hyperactivity disorder have a
1074 similar burden of rare protein-truncating variants. *Nat. Neurosci.* **22**, 1961-1965 (2019).
- 1075 54. Yang, R. et al. ANK2 autism mutation targeting giant ankyrin-B promotes axon branching and ectopic
1076 connectivity. *Proc. Natl. Acad. Sci. U S A.* **116**, 15262-15271 (2019).
- 1077 55. Lorenzo, D. N., et al. A PIK3C3–ankyrin-B–dynactin pathway promotes axonal growth and
1078 multiorganelle transport. *J. Cell Biol.* **207**, 735-752 (2014).

1079 56. Zhong, G., et al. Developmental mechanism of the periodic membrane skeleton in axons. *Elife*, **3**,
1080 e04581 (2014).

1081 57. Zhou, R., Han, B., Xia, C., Zhuang, X. Membrane-associated periodic skeleton is a signaling platform
1082 for RTK transactivation in neurons. *Science*, **365**, 929-934 (2019).

1083 58. Lek, M., et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature*, **536**, 285-291
1084 (2016).

1085 59. Park, J. et al. Mutational characteristics of ANK1 and SPTB genes in hereditary spherocytosis. *Clin.*
1086 *Genet.* **90**, 69-78 (2016) [published correction appears in *Clin. Genet.* **95**, 341 2019].

1087 60. Baek, H. J. et al. Transforming growth factor- β adaptor, β 2-spectrin, modulates cyclin dependent
1088 kinase 4 to reduce development of hepatocellular cancer. *Hepatology*. **53**, 1676-1684 (2011).

1089 61. Derbala, M. H., Guo, A. S., Mohler, P. J., Smith, S. A. The role of β II spectrin in cardiac health and
1090 disease. *Life Sci.* **192**, 278-285 (2018).

1091 62. Sobreira, N., Schiettecatte, F., Valle, D., Hamosh, A. GeneMatcher: a matching tool for connecting
1092 investigators with an interest in the same gene. *Hum. Mutat.* **36**, 928-930 (2015).

1093 63. García-Alvarez, B., Bobkov A, Sonnenberg A, de Pereda JM. Structural and functional analysis of the
1094 actin binding domain of plectin suggests alternative mechanisms for binding to F-actin and integrin
1095 beta4. *Structure*, **11**, 615-625 (2003).

1096 64. Zhang, Y. I-TASSER server for protein 3D structure prediction. *BMC Bioinformatics*, **9**, 40 (2008).

1097 65. von der Ecken, J., Heissler, S. M., Pathan-Chhatbar, S., Manstein, D. J., Raunser, S. Cryo-EM structure
1098 of a human cytoplasmic actomyosin complex at near-atomic resolution. *Nature*, **534**, 724-728
1099 (2016).

66. Källberg, M., Margaryan, G., Wang, S., Ma, J., Xu, J. RaptorX server: a resource for template-based protein structure modeling. *Methods Mol. Biol.* **1137**, 17-27 (2014).
67. Jurrus, E., et al. Improvements to the APBS biomolecular solvation software suite. *Protein Science: a Publication of the Protein Society.* **27**, 112-128 (2018).
68. Shindyalov, I.N., Bourne, P.E. Protein structure alignment by incremental combinatorial extension (CE) of the optimal path. *Protein Eng.* **11**, 739-747 (1998).

Figure Legends

Fig. 1: *SPTBN1* variants found in individuals with neurodevelopmental disorders. **a**, Schematic representation of functional domains of β II-spectrin. CH1=calponin homology domain 1 (teal), CH2=calponin homology domain 2 (red), SR=spectrin repeat (green), and PH=pleckstrin homology domain (purple). The locations of *SPTBN1* variants are indicated. **b**, Alignment of protein sequences for β II-spectrin and orthologues show that missense variants identified in the patients in this study are located at highly conserved residues across species from humans to *Drosophila*. The position of *SPTBN1* variants analyzed in the sequenced of human β II-spectrin is shown for reference. **c**, Photos of individuals with *SPTBN1* variants. Ages at the time of photograph are: P8: 7y8m, P9:16, P12: 11y, P13: 6y, P21 left: unknown, Right: 11y, P22: 15y, P27: 16y11m, P28: 3y11m. **d**, Examples of brain MRI findings: diffuse cerebral parenchymal volume loss (L>R) and asymmetric appearance of hippocampi (P1, acquired at <1y), white matter disease in the supratentorial and infratentorial regions (P18, acquired at 7y), thinning of the posterior body of the corpus callosum without significant volume loss (P28, acquired at 10m).

Fig. 2: *SPTBN1* variants alter protein expression and subcellular distribution. **a**, Levels of GFP- β II_{Sp} mutant proteins in HEK293T cells co-transfected with GFP- β II_{Sp} and mCherry plasmids relative to expression of WT GFP- β II_{Sp}. **b**, Partition of indicated GFP- β II_{Sp} proteins expressed in HEK293T cells between Triton-X100 soluble and insoluble fractions relative to total WT GFP- β II_{Sp} levels. Data in **b** and **c**

is representative of three independent experiments. All data represent mean $\pm$ SEM. One-way ANOVA with Dunnett's post hoc analysis test for multiple comparisons, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$. **c**, Immunofluorescence images of HEK293T cells transfected with indicated GFP- β IIISp plasmids and stained for actin (phalloidin) and DAPI. Scale bar, 10 μ m. Expression of variants within the distal end of CH1 and the proximal portion of the CH2 domains result in cytosolic GFP-positive aggregates (white arrowheads). Expression of variants within the C-terminal portion of the CH2 domain and a subset of variants in SRs increase the number of membrane protrusions (white asterisks). Data is representative of at least six independent experiments.

Fig. 3: SPTBN1 variants alter interaction with critical cytoskeleton partners. **a**, Immunofluorescence images of HEK293T cells transfected with mCherry- α IIISp and with either WT or mutant GFP- β IIISp plasmids. Cells were stained for actin (phalloidin) and DAPI. Scale bar, 10 μ m. **b**, Immunofluorescence images of DIV8 mouse β IIISp-KO cortical neurons transfected with indicated GFP- β IIISp plasmids and stained for actin (phalloidin) and endogenous α II-spectrin. Scale bar, 5 μ m. In **a** and **b** GFP-positive cytoplasmic aggregates (orange arrowheads) also contain either actin or α II-spectrin proteins, or both. **c**, Quantification of binding of mCherry- α IIISp to GFP- β IIISp proteins relative to the abundance of mCherry- α IIISp/WT GFP- β IIISp complexes. **d**, Binding of purified β II-spectrin proteins to purified F-actin assessed through an actin co-sedimentation assay. **e**, Binding of GFP- β IIISp proteins to 220-kDa AnkB-2HA assessed via co-IP from HEK293T cells. The Y1874A β II-spectrin mutation known to disrupt the formation of AnkB/ β IIISp complexes was used as control. Graphs in **c** and **d** summarize results from three independent experiments. Data in **e** summarizes four independent experiments. All data represent mean $\pm$ SEM. One-way ANOVA with Dunnett's post hoc analysis test for multiple comparisons, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Fig. 4: β II-spectrin CH domain variants likely alter CH1-CH2 dimer stability. **a**, Closed conformation of the β II-spectrin CH1-CH2 dimer modeled after utrophin³⁷ showing the sites of β II-spectrin variants at the

interface and the electrostatic surface of each domain calculated independently. **b, c**, Electrostatic complementarity shows that both CH domains have a polar side, where CH2 is negatively charged (**b**) and CH1 is positively charged(**c**), and both have a neutral side. **d**, Closed conformation of the β II-spectrin CH1-CH2 dimer modeled by docking the CH2 domain of β II-spectrin³⁸ onto the CH1 domain modeled after β III-spectrin³⁹. **e**, The L250R variant introduces a large, positively charged residue that clashes with a hydrophobic CH1 pocket through steric hindrance and electric instability. **f**, L247H introduces a large aromatic amino acid and likely disrupts normal CH2 folding. **g, h** Steric hindrance and negative charge introduced by (**g**) G205D and (**h**) G205S in the interior of CH2 likely disrupts normal CH2 folding. **i**, Key interactions at the CH1-CH2 interface (site of mutations in CH1 (teal) and CH2 (red)) and likely molecular perturbations caused by *STPBN1* variants.

Fig. 5: *SPTBN1* variants affect neuronal axonal growth and organelle transport. **a**, Images of DIV8 β II-SpWT, β II-SpHet, and β II-SpKO neurons transfected at DIV3 with mCherry. A subset of β II-SpKO neurons was co-transfected with GFP- β II-Sp and mCherry plasmids. Scale bar, 15 μ m. **b**, Axonal length of β II-SpWT, β II-SpHet, β II-SpKO, and rescued β II-SpKO DIV8 neurons (n=12-34 neurons/genotype) compiled from three independent experiments. Data represent mean $\pm$ SEM. One-way ANOVA with Dunnett's post hoc analysis test for multiple comparisons, ****p < 0.0001. **c**, Kymographs showing the mobility of RFP-tagged LAMP1-positive cargo in axons from DIV8 β II-SpKO and rescued β II-SpKO neurons. Analyzed trajectories are shown with a color code with green for anterograde, red for retrograde, and blue for static vesicles. Scale bar, 10 μ m and 60 s. **d**, Quantification of percent of motile LAMP1-RFP-positive cargo in axons from β II-SpWT, β II-SpHet, β II-SpKO, and rescued β II-SpKO neurons. The box and whisker plots represent data from minimum to maximum collected in n=9-13 axons per genotype. One-way ANOVA with Dunnett's post hoc analysis test for multiple comparisons, ****p < 0.0001.

Fig. 6: β II-spectrin deficiency disrupt proper neuronal development in cortical neuronal cultures and mouse brains. **a**, Images of DIV8 β II-SpKO cortical neurons rescued with WT GFP- β II-Sp or with GFP- β II-Sp

bearing mutations within the distal portion of the CH2 domain. Neurons were stained for actin (phalloidin) and endogenous α II-spectrin. Yellow dotted lines demark the cell edge. Scale bar, 5 μ m. **b**, Images of PND25 β II-SpWT and β II-SpHet brains stained for neurofilament to label axons and DAPI. Scale bar, 50 μ m. **c**, Quantification of CC thickness at the brain midline. (n=4 mice/genotype). Data represent mean $\pm$ SEM. Unpaired *t*-test, **p* < 0.05. **d**, Images of PND25 β II-SpNexWT, β II-SpNexHet and β II-SpNexKO brains stained for neurofilament to label axons and DAPI. Scale bar, 50 μ m. White dotted lines in **b** and **d** denote the position and boundaries of the corpus callosum (CC). **e**, Quantification of CC thickness at the brain midline. (n=6-7 mice/genotype). Data in **c** and **e** represent mean $\pm$ SEM. One-way ANOVA with Dunnett's post hoc analysis test for multiple comparisons, ****p* < 0.001, *****p* < 0.0001.

Fig. 7: β II-spectrin deficiency causes developmental and behavioral deficits in mice. **a**, Images of male E19 β II-SpWT, β II-SpHet, and β II-SpKO embryos. **b**, **c**, Quantification of head circumference (**b**) and eye distance (**c**) at E19. Data represents mean $\pm$ SEM (n=5-6 embryos/genotype). One-way ANOVA with Dunnett's post hoc analysis test for multiple comparisons, **p* < 0.05, ***p* < 0.01. **d**, Images of male PND25 wildtype (β II-SpWT) mice and mice with partial (β II-SpHet) and complete (β II-SpKO) loss of β II-spectrin in neural progenitors. **e**, Body length at PND25 for indicated genotypes. Data represent mean $\pm$ SEM (n=12 male mice/genotype). One-way ANOVA with Dunnett's post hoc analysis test for multiple comparisons, *****p* < 0.0001. **f**, Growth curve (as body weight) of β II-SpWT and β II-SpHet mice. Data represent mean $\pm$ SEM (n=12 male mice/genotype). Unpaired *t*-test, ***p* < 0.01, ****p* < 0.001. **g**, Locomotor activity and **h**, rearing assessed during a 30-minutes open-field test in PND30 β II-SpWT and β II-SpKO mice. Data in **g** and **h** represent mean $\pm$ SEM (n=15 β II-SpWT and n=5 β II-SpKO male mice). Unpaired *t*-test, **p* < 0.05, *****p* < 0.0001. Statistical comparisons were not conducted for **h** due to zero scores in the β II-SpKO group. **i**, **j**, Locomotor activity measured as distance traveled (**i**) and number of rears (**j**) during a one-hour test in a novel open field. **k**, Lack of social preference in β II-SpHet mice during a three-chamber choice

1194 task. I, Decreased entries by β II-SpHet mice into a chamber with stranger mouse. Data represent
1195 mean $\pm$ SEM (n=12 male mice/genotype). Unpaired t-test, *p< 0.05.

1196 **Table 1. Summary of clinical features observed among *SPTBN1* mutations carriers.** Abbreviations are as
1197 follows: Ma=male, Female=F, P=partial, C=complete, ID=intellectual disability, *Emotional Liability
1198 includes tantrums and depression, Mi=mild, Mo=moderate, S=severe, M-M=mild to moderate, M-
1199 S=moderate to severe, L-N=low to normal, U=unknown, ND=not diagnosed, NA=not assessed, F=febrile
1200 seizures only, +=presence, -=absence, $\downarrow$ = micro or hypo, $\uparrow$ = macro or hyper, **=half-siblings.

b

	59	159	183	205	247	250	255	268	271	275	344
Human SPTBN1	KKTFT	FQIQD	LWCQM	RDGMA	LGLTK	TKLLD	PEDIS	IITYV	YVVTY	YYHYF	QAFNT
Chimp SPTBN1	KKTFT	FQIQD	LWCQM	RDGMA	LGLTK	TKLLD	PEDIS	IITYV	YVVTY	YYHYF	QAFNT
Dog SPTBN1	KKTFT	FQIQD	LWCQM	RDGMA	LGLTK	TKLLD	PEDIS	IITYV	YVVTY	YYHYF	QAFNT
Mouse Sptbn1	KKTFT	FQIQD	LWCQM	RDGMA	LGLTK	TKLLD	PEDIS	IITYV	YVVTY	YYHYF	QAFNT
Worm unc-70	KKTFT	FQIQD	LWCQM	RDGLA	LGLAK	AKFLD	PEDIS	IITYV	YVVTY	YYHYF	TEFND
Fly Beta-spec	KKTFT	FQIQD	LWCQM	RDGLA	LGLAK	AKLLD	PEDIS	IITYV	YVVTY	YYHYF	AQFSN

	411	491	850	892	1003	1086	1110	1398	1674	1787	1886
Human SPTBN1	ALRNE	EAENY	TLALY	RFESL	MERDL	AIASE	KNEID	LHGLE	RQSKV	EAWAD	KRENE
Chimp SPTBN1	ALRNE	EAENY	TLALY	RFESL	MERDL	AIASE	KNEID	LHGLE	RQSKV	EAWAD	KRENE
Dog SPTBN1	ALRNE	EAENY	TLALY	RFESL	MERDL	AIASE	KNEID	LHGLE	RQSKV	EAWAD	KRENE
Mouse Sptbn1	ALRNE	EAENY	TLALY	RFESL	MERDL	AIASE	KNEID	LHGLE	RQSKV	EAWAD	KRENE
Worm unc-70	VLKEE	EAENY	ALSly	RFDTL	MERDL	QVASE	REEID	ATQLE	RQAQI	EAWEN	SRESE
Fly Beta-spec	ALREE	ESERY	ALSly	RFEGF	MDRDL	DVASE	REEID	ITDLE	KQSQL	ESWQD	NREQE

Fig.3

Closest CH1 - CH2 closed conformation interface residues		Mutation	Residues impacted by mutation	Predicted molecular consequences
CH1	CH2	T59I	L250	More hydrophobic side chain may increase CH2 binding
K62	D252	G205D	N233 & L234	Steric hindrance likely disrupts normal CH2 folding
T59, W63, W151, I154	L250	G205S		
L155	T268	L247H	Y273 & Y276	Steric hindrance likely disrupts normal CH2 folding
I159	H275	L250R	T59, W63, W151, & I154	Steric hindrance of large charged residue in hydrophobic CH1 pocket
		D255E	K62	Longer side chain may impair CH1 binding
		T268A	L155 & I159	Methyl group loss may impair hydrophobic interaction with CH1 residues
		T268S		Longer, more hydrophilic side chain may impair CH1 binding
		T268N		
		V271M	I159	Longer side chain may impair CH1 binding or new hydrophobic interaction may increase binding
		H275R	I159	Longer, charged basic side chain may impair CH1 binding and interact with acidic residue D51 on actin, enhancing binding

Table 1

Proband ID	P1	P2	P3	P4	P5	P6	P7	P8	P9	P10	P11	P12	P13	P14	P15	P16	P17	P18	P19	P20	P21**	P22**	P23	P24	P25	P26	P27	P28	P29		
Variant	p.T59I	c.475-1G>A p.(1159_Q160del)?	p.C183*	p.G205S	p.G205D	c.647+1G>T p.(Y190_R216del)?	p.L247H	p.L250R	p.D255E	p.T268A:F344L	p.T268N	p.T268S	p.V271M	p.H275R	p.R411W	p.R411Q	p.E491Q	p.A850G	p.E892*	p.R1003W	p.R1003W	p.R1003W	p.A1086T	p.E1110D	p.G1398S	p.S1674P	p.W1787*	p.E1886Q	c.5961+2T>C p.(11988As>90)?	TOTAL	
Phenotype evaluation	C	C	-	P	C	P	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	C	P	C	C	C	C	C		
Age at last exam (y/m)	11m	13y	U	26y	6y 2m	NA	8y 9m	7y 8m	12y 5m	10y	6y 7m	11y	3y 7m	9y	6y	6m	21m	3y 2m	7y	5y 10m	10y	15y	2y	5y 10m	U	9y	16y	3y 11m	18y		
Sex	Ma	Fe	Ma	Fe	Ma	Fe	Ma	Fe	Ma	Fe	Ma	Ma	Ma	Ma	Ma	Ma	Ma	Fe	Ma	Ma	Fe	Ma	Ma	Fe	Ma	Fe	Fe	Fe	Fe	17 male	
Neurodevelopment																															
Developmental Delay	+	+	U	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	U	+	+	-	+	26/27		
Speech Delay	+	S	U	+	+	U	+	+	S	+	+	+	+	+	+	+	+	+	+	+	+	+	Mi	+	U	+	+	-	-	24/26	
Motor Delay	+	+	U	+	+	U	+	+	+	+	+	+	+	+	+	-	+	+	+	+	+	+	Mi	+	U	-	-	-	Mi	22/26	
ID	Mo	S	U	Mi	Mi	U	+	S	S	S	U	S	Mi	Mo	Mo	U	U	M-S	L-N	M-M	Mo	+	-	-	U	Mi	Mi	M-S	-	20/23	
Regression	-	-	U	U	-	U	-	-	+	+	-	+	-	-	-	-	-	-	-	-	-	-	-	U	-	-	-	-	3/25		
Other Neurological																															
Epilepsy/seizure	+	U	U	U	+	U	-	NA	+	+	-	+	U	-	-	-	-	F	-	-	+	-	F	-	U	-	+	-	-	9/22	
EEG abnormalities	+	U	U	+	+	U	NA	NA	+	-	NA	+	U	-	-	NA	NA	-	NA	NA	+	-	NA	NA	U	-	NA	NA	-	6/13	
Abnormal brain MRI	+	+	U	U	-	U	-	NA	-	+	-	-	-	-	-	+	-	+	NA	NA	-	+	NA	NA	U	NA	-	+	NA	7/18	
Micro-/macrocephaly	↑	↓	U	U	-	U	-	-	↓	↓	↓	↑	↑	-	-	-	U	↑	U	U	-	-	U	↑	U	↓	NA	-	U	↓5, ↑5/19	
Sleep disturbances	-	-	U	U	-	U	+	+	+	-	U	+	U	U	-	U	-	+	-	-	-	-	+	U	U	U	-	+	-	7/20	
Behavior																															
ASD/autistic features	ND	-	U	+	-	U	-	+	+	ND	-	-	U	U	-	U	-	-	+	-	-	-	-	+	U	-	-	+	-	6/21	
ADD/ADHD	U	U	U	+	-	U	+	+	-	ND	+	+	+	+	U	U	-	U	-	+	-	+	-	+	U	-	+	-	-	11/20	
Anxiety	U	U	U	+	-	U	-	-	-	ND	U	+	-	-	+	U	-	-	-	-	-	U	-	-	U	U	U	NA	-	3/17	
Obsessive behavior	U	U	U	+	-	U	-	-	-	ND	-	-	-	-	-	U	-	U	-	-	-	U	-	-	+	U	U	-	-	2/19	
Emotional liability*	U	U	U	U	-	U	-	+	-	U	-	-	+	+	-	U	-	-	-	+	-	+	-	+	U	U	+	-	-	7/20	
Aggressive behavior	U	U	U	U	-	U	-	+	-	U	+	+	-	-	+	U	-	-	-	U	U	U	-	+	U	U	+	-	-	6/17	
Systemic problems																															
Abnormal Movement	-	+	U	U	-	U	+	NA	+	+	-	-	-	+	+	-	-	-	-	-	-	-	-	-	U	U	-	-	+	7/23	
Hypo-/hypertonia	↓	↑	U	U	-	U	-	NA	↓	↑	-	-	-	-	↓	↓	↓	↑	-	-	↓	↓	-	U	U	U	↓	-	↑	↓8, ↑4/22	
EMG abnormality	NA	NA	U	U	NA	U	NA	NA	NA	NA	NA	+	NA	NA	NA	NA	NA	NA	U	NA	NA	U	NA	NA	U	NA	NA	NA	-	1/2	
Hearing impairment	-	U	U	U	-	U	-	+	-	+	-	-	-	-	-	-	-	+	-	-	+	-	-	-	U	+	-	-	-	5/24	
Dysmorphology	+	+	U	+	-	U	+	+	+	NA	+	+	+	+	+	+	+	-	-	+	-	-	-	-	U	+	+	+	+	18/25	

Ma=male, Fe=female, P=partial, C=Complete, ID=intellectual disability, *Emotional Liability includes tantrums and depression, Mi=mild, Mo=moderate, S=severe, M-M=mild to moderate, M-S=moderate to severe, L-N=low to normal, U=unknown, ND=not diagnosed, NA=not assessed, F=febrile seizures only, ↓= micro or hypo, ↑= macro or hyper, **=half-siblings