1 Emergence of the D614A mutation in the spike protein of SARS-CoV-2: Imported cases
2 to the South Korea

3

4 Ae Kyung Park, Jin Sun No, Eunkyung Shin, Yoon-Seok Chung, Il-Hwan Kim, Byung-Hak
 5 Kang, Gi-Eun Rhie, Heui Man Kim, Jeong-Min Kim, Ye Eun Park, Min Joon Kim, Jae Sun
 6 Park, Cheon-Kwon Yoo, Junyoung Kim

7

8 Division of High-risk Pathogens, Center for Laboratory Control of Infectious Disease, Korea
 9 Centers for Disease Control & Prevention, Cheongju-si, Republic of Korea

10

11

12

13

14

15

16

17

18

19

20 Abstract

21 We identified imported cases of a novel D614A mutation in the spike glycoprotein of SARS-
22 CoV-2 in the South Korea. SARS-CoV-2 harboring the novel mutation was isolated from
23 returning travelers from Uzbekistan. These results emphasized the possibility that new
24 mutations have emerged in that area.

25

26 Since a new novel beta-coronavirus, SARS-CoV-2, was first reported in December 2019,
27 there has been ongoing explosive spread of this virus throughout the world, severely
28 threatening human health (1). South Korea was one of the first countries to report a
29 significant Coronavirus Disease-19 (COVID-19) outbreak; however, the epidemic curve has
30 been quickly flattened due to the outstanding response of the Korean government (2).
31 Currently, community spread of Covid-19 has slowed in the South Korea. On the other hand,
32 more than half of confirmed cases were related to travel (2). The Korea Centers for Disease
33 Control and Prevention (KCDC) has begun to conduct whole-genome sequencing (WGS) of
34 COVID-19 isolated from travelers to obtain real-time genomic epidemiology data, rapidly
35 monitor how the virus mutates and investigate sequence variations.

36 On July, 2020, a man who entered Korea from Uzbekistan (Uzbekistan nationality) and had
37 symptoms of cough, fever, and headache was diagnosed with COVID-19 (3). For WGS, total
38 RNA was extracted, and cDNA was amplified using primer pools ([https://artic.network/ncov-](https://artic.network/ncov-2019)
39 2019). Libraries were prepared using the Nextera DNA Flex Library Prep Kit (Illumina,
40 USA), and sequencing was performed on a MiSeq instrument with 2 x 75 base pairs using a
41 MiSeq reagent kit V3 (Illumina, USA). The reads were mapped to the reference genome
42 MN908947.3 using CLC Genomics Workbench version 20.0.3 (CLC Bio, Denmark). A total

43 of 663,666 (98.50%) out of a total of 673,800 reads mapped to the reference genome. We
44 obtained 99.72% genome coverage at the median genome coverage depth of 1554×
45 (Supplementary Table 1). The genomic sequences of the SARS-CoV-2 strain were deposited
46 in GISAID (Accession no. EPI_ISL_516783, EPI_ISL_516784, EPI_ISL_516785). Using the
47 reference genome, single-nucleotide variants (SNVs) were called using the BioNumerics
48 version 7.6 SARS-CoV-2 plugin (Applied Maths, Belgium).

49 We identified 14 polymorphic sites in the genome, which corresponded to all SNVs (Table 1).
50 In total, 13 appeared within the gene-coding regions. Of note, we identified a GAT→GCT
51 (A23403C) nucleotide change, which leads to a nonsynonymous change (D614A) in the
52 spike (S) glycoprotein. Based on 77,577 viral genomes downloaded on August 5, 2020, from
53 the GISAID database (4), the DNA sequences corresponding to the S protein of each genome
54 were extracted and aligned to the mutated S protein (D614A) (Table 1). The alignment results
55 indicated that the D614A mutation of the S protein identified in this study was a novel
56 mutation that has not been identified previously.

57 In addition, two travelers who visited Uzbekistan and returned to Korea were diagnosed with
58 COVID-19 immediately after entering Korea. They arrived in the South Korea on July after
59 short-term travel (Supplementary Table 1). The WGS results of the above two isolates also
60 indicated that they had D614A mutations (Table 1). Due to the lack of WGS data on isolates
61 from Uzbekistan in GISAID, we were unable to compare our mutation results to those from
62 Uzbekistan. However, the identification of D614A in the S protein from three individuals
63 who traveled Uzbekistan in recent days emphasized that new types of mutations might be
64 emerging and spreading in that area.

65 The significance of the D614G mutation of the S protein for SARS-CoV-2 infection was

66 elucidated in a recent report by Korver *et al.* (5). Korver *et al.* suggested that an association
67 exists between the G614 variant and higher levels of viral nucleic acid in the upper
68 respiratory tract in patients, which make it more infectious than D614 and has allowed the
69 rapid spread of G614 throughout the world. Additionally, this higher level of infection might
70 require a higher level of antibody, which is correlated with vaccines or antibody-based
71 therapeutic agents. From a structural perspective, the D614G mutation destabilizes the
72 interaction between the S1 and S2 units, resulting in the promotion of S1 shedding.
73 Additionally, this mutation induces the conformational change of the receptor-binding
74 domain (RBD) of the S protein allosterically to promote the binding of angiotensin-
75 converting enzyme 2 (ACE2) (6). Additionally, the D614G mutation may modify the
76 secondary structure of the furin cleavage site to facilitate the cleavage of S1 and S2 (7).
77 Considering that the D614G mutation is expected to have significant effects on the spread of
78 SARS-CoV-2, it will be necessary to study and characterize the D614A mutation to evaluate
79 its phenotypes.

80 In this study, we identified novel D614A mutations in SARS-CoV-2 isolated from travelers
81 returning to Korea from Uzbekistan in recent days. The results of this study emphasized the
82 need for enhanced surveillance, which will be essential to combat COVID-19, especially
83 where genetic epidemiology data are scarce.

84

85 Dr. Park is a senior researcher at the Korea Centers for Diseases Control & Prevention. Her
86 research interests include the molecular epidemiology of viral and bacterial diseases.

87

88

89 **References**

- 90 1. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak
91 associated with a new coronavirus of probable bat origin. *Nature*. 2020 Mar;579(7798):270-3.
- 92 2. Korea Centers for Disease Control and Prevention. Press Release. 2020.
93 ncov.mohw.go.kr
- 94 3. Kim JM, Chung YS, Jo HJ, Lee NJ, Kim MS, Woo SH, et al. Identification of
95 Coronavirus Isolated from a Patient in Korea with COVID-19. *Osong Public Health Res*
96 *Perspect*. 2020 Feb;11(1):3-7.
- 97 4. Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data - from
98 vision to reality. *Euro surveillance : bulletin European sur les maladies transmissibles =*
99 *European communicable disease bulletin*. 2017 Mar 30;22(13).
- 100 5. Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al.
101 Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the
102 COVID-19 Virus. *Cell*. 2020 Jul 3.
- 103 6. Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al. Spike
104 mutation pipeline reveals the emergence of a more transmissible form of SARS-CoV-2.
105 *BioRxiv*. 2020.
- 106 7. Tang L, Schulkins A, Chen C-N, Deshayes K, Kenney JS. The SARS-CoV-2 Spike
107 Protein D614G Mutation Shows Increasing Dominance and May Confer a Structural
108 Advantage to the Furin Cleavage Domain. *Preprints*. 2020.

109

110 Table 1. Single-nucleotide variants (SNVs) of SARS-CoV-2 isolates from travelers returning to Korea from Uzbekistan with respect to the
111 reference genome

Gene	Nucleotide position	Reference (MN908947.3)	#1	#2	#3	Mutation
ORF1ab	4579	T	A	-	-	Synonymous
	8782	C	T	T	T	Synonymous
	11230	G	T	T	T	Nonsynonymous (M3655I)
	11970	A	-	G	G	Nonsynonymous (K3902R)
	20580	G	T	T	T	Synonymous
S	23403	A	C	C	C	Nonsynonymous (D614A)
ORF3a	26028	C	T	T	T	Synonymous
	26111	C	T	-	-	Nonsynonymous (P240L)
ORF8	28144	T	C	C	C	Nonsynonymous (L84S)
	28167	G	A	A	A	Nonsynonymous (E91K)
N	28878	G	A	A	A	Nonsynonymous (S202N)
	28887	C	-	T	T	Nonsynonymous (T205I)
	29224	G	-	-	A	Nonsynonymous (M317I)
Noncoding region (3' UTR)	29742	G	A	-	-	Noncoding region