Protocol

Efficacy and safety of *Boswellia* containing formulations for the treatment of osteoarthritis: protocol for a systematic review and meta-analysis of randomised controlled trials

Zhiqiang Wang^a, Ambrish Singh^a, Salman Hussain^b, Pablo Molina García^{a,c}, Benny Antony^{a*}

^aMenzies Institute for Medical Research, University of Tasmania, Hobart, Tasmania, Australia

^bCzech National Centre for Evidence-Based Healthcare and Knowledge Translation (Cochrane Czech Republic, Czech EBHC: JBI Centre of Excellence, Masaryk University GRADE Centre), Institute of Biostatistics and Analyses, Faculty of Medicine, Masaryk University, Brno, Czech Republic

^cDepartment of Physical Education and Sport, Faculty of Sport Sciences, University of Granada, Spain

Abstract

Osteoarthritis (OA) is a common chronic joint disease with limited pharmacological management options. *Boswellia* extracts (BE) or formulations containing BE are generally considered as safe and offers a modest efficacy for the treatment of OA. Although previous systematic reviews have assessed the efficacy of BE in OA, these reviews had excluded trials assessing various formulations containing BE and excluded various combinations of BE. Hence, this study aims to systematically review the evidence from RCTs assessing the efficacy and safety of both BE and formulations containing BE for the treatment of OA. Biomedical databases such as PubMed, Embase, and Google Scholar will be searched to identify the RCTs of BE or formulations containing BE in patients with OA (hand, knee, hip, or any other OA). Cochrane risk of bias tool will be used to assess the quality of included studies. Review Manager 5 (Rev Man) and STATA Version 16 will be used to conduct the statistical analysis.

* Corresponding author: Dr Benny Antony.

E-mail: Benny.EathakkattuAntony@utas.edu.au

Menzies Institute for Medical Research

University of Tasmania

17 Liverpool St, Hobart TAS 7000

Tasmania, Australia

Background and rationale

Osteoarthritis (OA) is a common chronic joint disease that imparts a substantial and ever-increasing health burden, especially in older people^[1]. Current treatment guidelines of OA often recommend non-pharmacological treatment as first-line, such as education and self-management, exercise, weight loss if overweight or obese^[2, 3]. The options for pharmacological management of OA is limited to analgesics, intra-articular corticosteroids, and NSAIDs^[4]. These medications only have a mild to moderate effect on pain^[5], often resulting in patient dissatisfaction and hastening joint replacement^[6]. There remains unmet healthcare need for chronic pain management^[7, 8]; as a result, patients are increasingly turning to complementary and alternative medicine^[9] such as *Boswellia* extracts (BE) or various formulations containing BE^[10, 11].

Frankincense, or BE, has a long history of use (such as religious ceremonies and perfume production), and its medicinal properties have been appreciated for ages^[12]. Preclinical and clinical evidence suggests that BEs are an effective and safe option for the treatment of OA^[13]. BEs were often used for rheumatoid arthritis, Crohn's disease, osteoarthritis, and collagenous colitis^[14]. Various formulations containing BE often combine BE with other natural products such as curcumin^[15], ginger [16], methylsulfonylmethane [17]. A recent systematic review included the randomised clinical trials (RCTs) evaluating the efficacy and safety of BE for the treatment of OA^[18]. However, studies evaluating formulations containing BE, such as Acujoint™, Lanconone® (E-OA-07), were not included in the review. We hypothesise that BE formulations exhibit a similar or better effect on OA as these combinations often help in increased bio-absorption of the active ingredients of BE such as Acetyl-11-keto-β-boswellic Acid (AKBA).

Objectives

We aim to systematically review and meta-analyse the evidence from RCTs on the efficacy and safety of BE or formulations containing BE for OA, in terms of pain and physical function improvement, change in biomarkers, and adverse events.

Methodology

Eligibility criteria

We will select the eligible RCTs following the specific criteria of PICOS items: Participants, Interventions, Comparators, Outcomes, and Study design. The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement will be employed for the conduction and reporting of this systematic review^[19]. We will include studies published both in English and Chinese.

Characteristics of participants (P)

We will include adult patients, at least 40 years old, of any sex, and diagnosed with OA according to the criteria proposed by the American College of Rheumatology or similar^[20].

Characteristics of interventions (I)

This study will focus on BE obtained from trees of the genus *Boswellia* (*Boswellia sacra*, *B. carterii*, *B. frereana*, *B. serrata*, and *B. papyrifera*) and any formulations containing these BE. RCTs evaluating combination therapy of BE as adjuvant or complementary to conventional drugs (eg. Paracetamol or NSAIDs) will also be included. We will include trials with multi-herbal formulations containing BE and any bioavailability enhancing combination.

Characteristics of comparators (C)

We will include the studies comparing BE and formulations containing BE with active control (e.g. but not limited to NSAIDs) or placebo for the treatment of OA.

Characteristics of outcomes (O)

The studies reporting data for at least one of the following outcomes: pain, physical function, imaging biomarkers (x-ray joint space narrowing, MRI structural measures), biochemical markers, medication change, and adverse events will be included in the analysis.

For each outcome, data on baseline and follow-up values and/or mean change from baseline will be extracted. If the data are expressed in the graphical information, the numerical data will be extracted from graphs using the procedure (adapted) suggested by Guyot et al.^[21]. If the studies do not provide complete data, authors of primary studies will be contacted through email to request missing or additional data.

Characteristics of study design (S)

Studies will be included if they used a randomised, quasi-randomised, controlled, blinded or non-blinded design. Observational and non-randomised studies will be excluded.

Primary and secondary outcomes

Our primary outcome will be to evaluate the efficacy of BE and BE formulations on improvement in pain. Furthermore, we will consider improvement in physical function, change in biochemical markers, and imaging biomarkers as secondary outcomes.

Information sources and search procedure

We will implement the search procedure consistently with the following criteria.

Electronic source and search strategy

We will retrieve the currently available RCTs on the use of BE and BE formulations for OA through a systematic search of the biomedical database. We will search online databases such as PubMed, Embase, Web of Science, and Cochrane Central Register of Controlled Trial, Google scholar, etc. from inception to September 2020. Search terms for interventions will include: *Boswellia*, *Boswellia serrata*, *Boswellia carteri*, *Boswellia frereana*, *Boswellia papyrifera*, *Boswellia sacra*, ru xiang (乳香, Chinese name for *Boswellia serrata*), boswellic acid, Frankincense, Shallaki, Salai, aflapin, 5-loxin and other formulations such as Acujoint, Lanconone. Search terms for participants will include: Osteoarthritis, Osteoarthritis, Osteoarthritis, Osteoarthritis, Degenerative Arthritis, Arthritis. Other search terms will be included: randomized controlled trial, random*, random allocation, single-blind, double-blind.

Both published and unpublished trials will be included, with the later including e.g. abstracts, conference proceedings, and posters with available data. Studies that have compared interventions of interest and reported extractable data for at least one measure of pain, physical function, imaging biomarkers, biochemical markers, medication change, and adverse events will be included.

Hand-searching

Abstract booklet from conference proceedings, abstracts, and poster sessions will be hand searched using the online sources of major international association involved in OA research: European League Against Rheumatism (EULAR), Osteoarthritis Research Society International (OARSI), American Academy of Orthopaedic Surgeons (AAOS) and American College of Rheumatology (ACR). Moreover, references retrieved from relevant meta-analysis and review articles will be hand-searched and analysed for inclusion as per the eligibility criteria.

Study selection

An Inclusion/Exclusion Form was adapted and used for data extraction (Appendix 1). Study selection will be performed by two reviewers independently, and each potential discrepancy will be discussed and solved through consensus with other authors and independent expert consultation.

Assessment of risk of bias

The methodological quality of the selected studies will be evaluated using the Cochrane risk of bias tool^[22]. A Risk-of-Bias form was adapted and used for data extraction (Appendix 2). Each article will be evaluated independently by two researchers using the Cochrane risk of bias tool. The Cochrane risk of bias tool

considers characteristics of the following items: sequence generation, allocation concealment, blinding of participants, study personnel and outcome assessors, incomplete outcome data, selective outcome reporting and other potential sources of bias. At the end of quality assessment, a consensus on final evaluation will be reached; any disagreements will be resolved by the discussion with senior authors.

Data extraction

A data extraction form was adapted and used for data extraction (Appendix 3). Two reviewers will extract data independently from the included studies for the following information: study design, characteristics of the population (age, sex, and BMI), sample size, intervention details and dosage, duration of follow-up, type of placebo or other control, outcome measurements, mean change values of the relevant outcome, and the number of adverse event report and medication change. Intention-to-treat data will be used whenever available. The missing data in the literature will be obtained by emailing the corresponding author; otherwise, it will be estimated by the appropriate method according to Cochrane Handbook 5.1.0^[23].

Statistical analysis

The fixed-effect model will be used if included studies are homogeneous; otherwise, the random-effect model with a restricted maximum-likelihood will be employed for the meta-analysis of both continuous and binary outcomes. The heterogeneity of the effect size across the trials will be tested using the Q statistics ($P < 0.05$ was considered heterogeneous) and I^2 statistic ($I^2 > 50\%$ will be considered heterogeneous)^[24]. Publication bias will be assessed visually with funnel plots^[25], and the trim-and-fill method was used to estimate the effect of publication bias (if any)^[26].

Due to different outcome measures, the change from baseline to follow-up scores will be translated into standard mean differences (SMD) using Hedges' g effect sizes, as per the data availability. Subgroup analysis will be conducted based on characteristics of RCTs, such as different formulation types, and comparator types. To further explore the potential heterogeneity among the trials, we will perform the meta-regression of the SMDs on covariates^[27], such as baseline characteristics of participants (age, gender, BMI), duration, formulation types (BSE or BSE formulations), type of pain measures (VAS vs. WOMAC/KOOS), dosage, study quality (assessed by the RoB), and types of funding (investigator-initiated vs. industry). Statistical analysis will be performed using STATA version 16 (STATA Corp., Texas, USA) and Review Manager 5 (RevMan 5.3) (Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration, 2014).

References

1. Cross M, Smith E, Hoy D, Nolte S, Ackerman I, Fransen M, et al. The global burden of hip and knee osteoarthritis: estimates from the Global Burden of Disease 2010 study. *Annals of the Rheumatic Diseases*. 2014; 73(7):1323-1330.
2. Bannuru RR, Osani MC, Vaysbrot EE, Arden NK, Bennell K, Bierma-Zeinstra SMA, et al. OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis and Cartilage*. 2019.
3. Kolasinski SL, Neogi T, Hochberg MC, Oatis C, Guyatt G, Block J, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Management of Osteoarthritis of the Hand, Hip, and Knee. *Arthritis Rheumatol*. 2020 Feb; 72(2):220-233.
4. Hunter DJ, Bierma-Zeinstra S. Osteoarthritis. *The Lancet*. 2019; 393(10182):1745-1759.
5. Gregori D, Giacobelli G, Minto C, Barbetta B, Gualtieri F, Azzolina D, et al. Association of Pharmacological Treatments With Long-term Pain Control in Patients With Knee Osteoarthritis. *JAMA*. 2018; 320(24):2564.
6. Roos EM, Arden NK. Strategies for the prevention of knee osteoarthritis. *Nat Rev Rheumatol*. 2016 Feb; 12(2):92-101.
7. LaChance J, Booth R, Befus D. Exploring unmet healthcare needs, healthcare access, and the use of practitioner based complementary and alternative medicine in adults with chronic pain. *Contemp Nurse*. 2020 Mar 25:1-15.
8. Zhang W, Ouyang H, Dass CR, Xu J. Current research on pharmacologic and regenerative therapies for osteoarthritis. *Bone Research*. 2016; 4(1):15040.
9. Chen L, Michalsen A. Management of chronic pain using complementary and integrative medicine. *BMJ*. 2017 Apr 24; 357:j1284.
10. Liu X, Machado GC, Eyles JP, Ravi V, Hunter DJ. Dietary supplements for treating osteoarthritis: a systematic review and meta-analysis. *British Journal of Sports Medicine*. 2018; 52(3):167-175.
11. Henrotin Y, Mobasheri A. Natural Products for Promoting Joint Health and Managing Osteoarthritis. *Current Rheumatology Reports*. 2018; 20(11).
12. Hughes K. *The Incense Bible: Plant Scents that Transcend World Culture, Medicine, and Spirituality*: Haworth Press New York, 2007.
13. Krishnaraju AV, Sundararaju D, Vamsikrishna U, Suryachandra R, Machiraju G, Sengupta K, et al. Safety and toxicological

- hr/>
- evaluation of Aflapin: a novel Boswellia-derived anti-inflammatory product. *Toxicol Mech Methods*. 2010 Nov; 20(9):556-563.
14. Ernst E. Frankincense: systematic review. *BMJ*. 2008 Dec 17; 337:a2813.
 15. Haroyan A, Mukuchyan V, Mkrtchyan N, Minasyan N, Gasparyan S, Sargsyan A, et al. Efficacy and safety of curcumin and its combination with boswellic acid in osteoarthritis: a comparative, randomized, double-blind, placebo-controlled study. *BMC Complementary and Alternative Medicine*. 2018; 18(1).
 16. Srivastava S, Chaudhary JA, Girandola RN. Effect of E-OA-07 on improving joint health and mobility in individuals with knee osteoarthritis: A randomized, double-blind, placebo-controlled, parallel group study. *Journal of Pain Research*. 2019; 12(pp 3365-3379).
 17. Notarnicola A, Tafuri S, Fusaro L, Moretti L, Pesce V, Moretti B. The "MESACA" study: methylsulfonylmethane and boswellic acids in the treatment of gonarthrosis. *Adv Ther*. 2011 Oct; 28(10):894-906.
 18. Yu G, Xiang W, Zhang T, Zeng L, Yang K, Li J. Effectiveness of Boswellia and Boswellia extract for osteoarthritis patients: a systematic review and meta-analysis. *BMC Complement Med Ther*. 2020 Jul 17; 20(1):225.
 19. Hutton B, Salanti G, Caldwell DM, Chaimani A, Schmid CH, Cameron C, et al. The PRISMA Extension Statement for Reporting of Systematic Reviews Incorporating Network Meta-analyses of Health Care Interventions: Checklist and Explanations. *Annals of Internal Medicine*. 2015; 162(11):777.
 20. Altman R, Asch E, Bloch D, Bole G, Borenstein D, Brandt K, et al. Development of criteria for the classification and reporting of osteoarthritis: Classification of osteoarthritis of the knee. *Arthritis & Rheumatism*. 1986; 29(8):1039-1049.
 21. Guyot P, Ades A, Ouwers MJ, Welton NJ. Enhanced secondary analysis of survival data: reconstructing the data from published Kaplan-Meier survival curves. *BMC Medical Research Methodology*. 2012; 12(1):9.
 22. Higgins JPT, Altman DG, Gotzsche PC, Juni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ*. 2011; 343(oct18 2):d5928-d5928.
 23. Higgins JPT. *Cochrane handbook for systematic reviews of interventions*: Wiley-Blackwell, 2020.
 24. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Bmj*. 2003; 327(7414):557-560.
 25. Stuck AE, Rubenstein LZ, Wieland D. Bias in meta-analysis detected by a simple, graphical test. Asymmetry detected in funnel plot was probably due to true heterogeneity. *BMJ: British Medical Journal*. 1998; 316(7129):469.
 26. Duval S, Tweedie R. A Nonparametric "Trim and Fill" Method of Accounting for Publication Bias in Meta-Analysis. *Journal of the American Statistical Association*. 2000; 95(449):89-98.
 27. Higgins JP, Thompson SG, Spiegelhalter DJ. A re-evaluation of random-effects meta-analysis. *Journal of the Royal Statistical Society: Series A (Statistics in Society)*. 2009; 172(1):137-159.

Appendix 1

Inclusion/Exclusion Form for Primary Studies

Study ID:

Reviewer:

Date:

Identification Details

Author	Year	Journal/ Conference	Source

On Endnote database Yes / No Full

Text availability Yes / No

Study Eligibility

Study design is one of the following:

Randomised clinical trial..... Yes/ No

The study concerns osteoarthritis..... Yes/ No

The study concerns Boswellia or Boswellia containing formulations..... Yes/ No

The study concerns pain or physical function measurements..... Yes/ No

The study is a human study, not animal/laboratory experiment..... Yes/ No

Please Tick Only One Box Below

Included	Excluded	Pending*

* Issue relates to selective reporting – when authors may have taken pain or physical function measurements, but not reported these within the paper. Reviewers should contact the corresponding author for information on possible non-reported pain or physical function & reasons for exclusion from publication. Study should be listed as ‘Pending’ until clarified. If no clarification is received after three attempts, study should then be excluded.

References to Other Trials

Did this report include any references to published reports of potentially eligible trials not already identified for this review?		
First author	Journal / Conference	Year of publication

Did this report include any references to **unpublished data** from potentially eligible trials not already identified for this review? If yes, give list contact name and details

Appendix 2

Risk-of-Bias Form

Risk of bias	Author's judgement	Support for judgement
Random sequence generation (selection bias) Was the allocation sequence adequately generated?	Low risk of bias Unclear risk of bias High risk of bias	Describe selection bias (biased allocation to interventions) due to inadequate generation of a randomised sequence
Allocation concealment (selection bias) Was allocation adequately concealed?	Low risk of bias Unclear risk of bias High risk of bias	Describe selection bias (biased allocation to interventions) due to inadequate concealment of allocations prior to assignment
Blinding (performance bias and detection bias) Was knowledge of the allocated interventions adequately prevented during the study?	Low risk of bias Unclear risk of bias High risk of bias	Performance bias or detection bias due to knowledge of the allocated interventions after assignment
Incomplete outcome data (attrition bias) Were incomplete outcome data adequately addressed?	Low risk of bias Unclear risk of bias High risk of bias	Evaluate attrition bias due to amount, nature or handling of incomplete outcome data
Selective reporting (reporting bias) Are reports of the study free of suggestion of selective outcome reporting?	Low risk of bias Unclear risk of bias High risk of bias	Evaluate reporting bias due to selective outcome reporting
Other bias Was the study apparently free of other problems that could put it at a high risk of bias?	Low risk of bias Unclear risk of bias High risk of bias	Bias due to problems not covered elsewhere in the table

Appendix 3

Data Extraction Form

Study and Participants characteristics

Study characteristics	
	Further details
Country / Countries	
Single centre / multicentre	
Study registration	
Funding	
How was participant eligibility defined? (e.g. How osteoarthritis was defined?)	
How pain was measured?	
How physical function was measured?	
How many people were recruited?	
Duration of study (randomised clinical trial)	
Other	
Participant characteristics	
	Further details
Number of participants (cases versus controls)	
Age (mean, median, range, etc)	
BMI (mean, median, range, etc)	
Female of OA participants (numbers / %, etc)	Female of OA participants (numbers / %, etc)
Type of OA (hand, knee, hip, etc)	
Other	

Measures relevant to the review					
Pain		VAS / WOMAC/ KOOS			
Physical Function		WOMAC/ KOOS			
Adverse events		YES/ NO			
Others					
For dichotomous data					
OR/ RR (95% CI, p value)	Treatment group (n) n = number of participants, not number of events		Control group (n) n = number of participants, not number of events		
Other Information (eg. adjustment, cut-off, etc):					
For continuous data					
Unit	Treatment group		Control group		Details
	n	Mean (SD)	n	Mean (SD)	
Other relevant data					