A Comparative Study to Find a Suitable Model for an Improved Real-Time Monitoring of The Interventions to Contain COVID-19 Outbreak in The High Incidence States of India

Running Title: *Monitoring the interventions to control COVID19*

Amrutha G.S(0000-0002-7480-7857)¹, Abhibhav Sharma (0000-0001-5513-6210)², Anudeepti Sharma (0000-0003-1542-3818)³

1. M.Phil. Population Studies; MPS (Master of Population Studies); M.Sc. Statistics.
Affiliation: International Institute for Population Sciences (IIPS), Mumbai, Maharashtra, 400088, India (Email: amruthags12@gmail.com)
2. MTech Statistical Computing; M.Sc. Mathematics
Affiliation: School of Computer and Systems Sciences (SC&SS), Jawaharlal Nehru University (JNU), New Delhi, 110067, India (Email: mail.abhibhav@gmail.com)
3. M.Sc. Computer Science
Central University of Rajasthan, Ajmer, Rajasthan, 305801, India (Email: anudeepti.sharma04@gmail.com)

Corresponding Author:

Abhibhav Sharma ²

School of Computer and Systems Sciences (SC&SS),
Jawaharlal Nehru University (JNU),
New Delhi, India, 110067

Email: mail.abhibhav@gmail.com

Contact No: +91 9354595608

Abstract

Background

On March 11, 2020, The World Health Organization (WHO) declared coronavirus disease (COVID-19) as a global pandemic. There emerged a need for reliable models to estimate the imminent incidence and overall assessment of the outbreak, in order to develop effective interventions and control strategies. One such vital metrics for monitoring the transmission trends over time is the time-dependent effective reproduction number (R_t). R_t is an estimate of secondary cases caused by an infected individual at a time t during the outbreak, given that a certain population proportion is already infected. Misestimated R_t is particularly concerning when probing the association between the changes in transmission rate and the changes in the implemented policies. In this paper, we substantiate the implementation of the instantaneous reproduction number (R_{ins}) method over the conventional method to estimate R_t viz case reproduction number (R_{case}), by unmasking the real-time estimation ability of both methodologies using credible datasets.

Materials & Methods

We employed the daily incidence dataset of COVID-19 for India and high incidence states to estimate R_{ins} and R_{case} . We compared the real-time projection obtained through these methods by corroborating those states that are containing high number of COVID-19 cases and are conducting high and efficient COVID-19 testing. The R_{ins} and R_{case} were estimated using R0 and EpiEstim packages respectively in R software 4.0.0.

Results

Although, both the R_{ins} and R_{case} for the selected states were higher during the lockdown phases (March 25 - June 1, 2020) and subsequently stabilizes co-equally during the unlock phase (June 1- August 23, 2020), R_{ins} demonstrated variations in accordance with the interventions while R_{case} remained generalized and under- & overestimated. A larger difference in R_{ins} and R_{case} estimates was also observed for states that are conducting high testing.

Conclusion

Of the two methods, R_{ins} elucidated a better real-time progression of the COVID-19 outbreak conceptually and empirically, than that of R_{case} . However, we also suggest considering the assumptions corroborated in the implementations which may result in misleading conclusions in the real world.

Keywords

COVID-19; Instantaneous Reproduction Number; Effective Reproduction Number; Estimation; India

1. Introduction

Coronavirus disease (COVID-19) is a severe acute respiratory syndrome caused by a novel Coronavirus (SARS CoV 2). As of March 11, around 53,370 and 4,627 COVID-19 cases and deaths were reported respectively, including 62 cases and 1 death in India.^[1] In the light of its rapid spreading in India and an alarming number of daily deaths caused by it, the government of India along with state governments took many strong preventive measures and non-pharmaceutical interventions to flatten the daily incidence and daily death curve. A varying trend of the infection spread have been reported for different locations, which remained ambiguous due to the novel nature of the virus. To unmask this underlying pattern behind the anomalous behavior of the outbreak in response to an intervention, studies to ascertain a suitable epidemiological model are now in the center stage. Accurate modeling of disease transmission is crucial for optimizing control interventions and often-limited healthcare facilities. One of the vital metrics for the assessment of transmission rate over time is the time-dependent effective reproduction number (R_t).^[3] R_t is an estimate of secondary cases caused by an infected individual at a time t during the outbreak, given that a certain population proportion is already infected. The $R_t < 1$ indicates the mitigation of the pandemic while a higher R_t i.e. $R_t > 1$, indicates the outbreak progression. A faulty estimate of R_t can systematically under- or overestimate the true transmission rate. Biases are particularly concerning if they occur near the critical threshold viz $R_t = 1$. The most prevailing and extensively used methodology for the estimation of the effective reproduction number is proposed by Wallinga and Teunis (2004), often called the case reproduction number (R_{case}).^[4] R_{case} is basically a probabilistic reconstruction of the transmission network, counting the average secondary cases produced by an individual. However, this method amalgamates the incidence data from time later than t , which when applied on a data aggregated on a shorter time step, generates an erroneous R_t estimate producing significantly negative autocorrelation when considered the true incidence.^[5] Also, Implementation of R_{case} sometimes requires sound statistical expertise to produce credible estimates, which is often absent in non-modelers. Previous studies assert that correct modeling is essential in the assessment of policy, population susceptibility and other transmission affecting factors.^[6] A fallible R_{case} estimate of the time-dependent reproduction number could be potentially pernicious. To overcome these shortcomings, Cori et al. proposed the instantaneous reproduction number (R_{ins}) for real-time estimation of R_t by corroborating incidences that occurred prior to time t , thus indicating a near real-time estimation of disease transmission.^[7]

Both R_{ins} and R_{case} incorporates the generation interval for the estimation of the reproduction number. Precisely, the generation interval is the time interval between the exposure of the infector and the exposure of the infectee. In the case of COVID-19, the accurate time of exposure is extremely difficult to report.^[8] The standard proxy for generation time is serial interval viz the time duration between the onset of the symptom in infector and infectee.^{[8][9]} In this paper, we incorporated the serial interval to estimate R_{ins} and R_{case} for India and high incidence states. The study aims to unfold the varying projections of R_t obtained by R_{ins} and R_{case} and to probe their credibility by employing the incidence dataset of the states that contain high incidence numbers and has performed robust testing.

2. Methods

R_{ins} and R_{case} requires the incidence dataset and the serial interval distribution to estimate R_{ins} . Conceptually different from Cori et al. (2013) and Wallinga et al. (2014), Bettencourt et al. (2008) proposed a methodology to estimate R_t based on the SIR (susceptible-infected-removed) model, that incorporates the growth rate of SIR and estimates the number of infections between the time intervals.^[10] Although Bettencourt et al. method has the capacity for a credible real-time R_t estimation, the results could be highly misleading as (1) the model requires a great deal of structural assumptions; (2) SIR based models do not account for the latency period.^{[5][11]} In this work, we focus on R_{ins} and R_{case} methods and probe their behavior over the benchmark dataset. The R_{ins} and R_{case} estimations were carried out under the standard assumptions: (1) The serial interval of COVID-19 follows a gamma distribution with mean 4 and standard deviation 2. Previous studies investigating the serial interval of COVID-19 validate the same.^{[5][12][13]} (2) The outbreak within the individual state is entirely driven by local transmission. In order to corroborate this assumption, we estimated R_t s (i.e. R_{ins} and R_{case}) from the date of the imposition of the first national lockdown viz March 25, 2020. The lockdown as a non-pharmaceutical intervention, resulted in a transmission arrest, ideally confining the outbreak within the state. (3) R_t estimate tends to be misleading in the situation where there is underreporting, however, if the rate of underreporting is constant, then it has no significant impact over the R_t estimate.^[14] Sequentially, the rate of underreporting in this study is assumed to be constant.

Here we also analyze the credibility of R_{ins} and R_{case} by juxtaposing their respective estimations over two different time periods viz (1) March 25, 2020 to August 23, 2020, and (2) June 1, 2020 to August 23, 2020. The former interval provides an overview of the R_{ins} and R_{case} behavior throughout the entire lockdown interventions (Lockdown 1: March 25; Lockdown 2: April 15; Lockdown 3: May 4; Lockdown 4: May 18) and the unlock phases (Unlock 1: June 1; Unlock 2: July 1; Unlock 3: August 23) (detailed description of the initiatives and strategies administered during each phase is given in **Table S2**). The later interval is crucial for a comparative analysis of R_{ins} and R_{case} as unlock phases invite abruptions in the infection transmission and we assert that an efficient model should be able to capture these variations. In order to investigate the credibility of the model, the later interval is an ideal window to probe the estimates by the methods.

2.1 Dataset

In relevance to the aim of this study, a benchmark dataset is essential to project a true comparison of the reproduction number estimates obtained by the methods. The study corroborates the dataset for the state: (1) If the total incidence reported in a state is $\geq 50,000$; (2) If total cases per million in the state are ≥ 1000 ; (3) If the state has conducted more than 9000 testing per million as on August 23; and, (4) if the percent (%) of positive cases detected among the sample tested (test positive rate TPR) is greater than 5.0 %. This resulted in a sample of eight states namely *Andhra Pradesh, Delhi, Gujarat, Karnataka, Maharashtra, Tamil Nadu, Telangana* and *West Bengal* (**Table 1**). The benchmark dataset of state-wise daily new cases and the daily test counts were obtained from *covid19india.org*. The data of the test sampling efficiency for a state were provided by ICMR (Health departments of State governments).

Link to the repository containing all the datasets involved in this study is made available in the additional material.

2.2 Case Reproduction Number (R_{case})

Wallinga et al. proposed a likelihood-based estimation method to estimate R_t , also called case or cohort reproduction number because it measures the average number of infections that an infected cohort may cause, given that the cohort was infected at time t . Precisely, the methods infer “who infected whom” through an observed serial interval distribution while reducing the computational burden by incorporating infector-infectee pairs rather than the complete infection network to measure the likelihood of transmission. Let t_i denotes the time of the onset of symptoms in the individual i . Given a serial interval distribution $w(\tau)$, τ being the generation interval, the relative likelihood (P_{ab}) that individual a has been infected by the individual b is given as (eq 1):

$$P_{ab} = \frac{w(t_a - t_b)}{\sum_{a \neq k}^n w(t_a - t_k)} \dots \dots \dots (\text{eq 1})$$

Where n is the total incidence reported for each reported case i , and $w(t)$ is the probability of a t -day interval i.e. the duration between the onset of symptoms in the infector-infectee pair. The quantity P_{ab} is a relative likelihood, normalized by the likelihood of the event where a is infected by the individual k . Sequentially, the Wallinga et al. method obtains the effective reproduction number R_b as (eq 2):

$$R_b = \sum_{a=1}^n p_{ab} \dots \dots \dots (\text{eq 2})$$

A preeminent overview of the underlying mathematical derivation is described in Wallinga et al.^[4]

2.3 Instantaneous Reproduction Number (R_{ins})

Based on the work of Fraser C. (2007), Cori et al. estimated instantaneous reproduction number R_{ins} through the ratio of new incidence (I_t) produced at time step t to the total infectiousness of infected individuals at time t .^{[16][7]} The latter is given as eq 3.

$$\sum_{s=1}^t I_{t-s} w_s \dots \dots \dots (\text{eq 3})$$

Where w_s is the probability distribution of the infectiousness profile of an individual at s units of time from the time of the infection induced in the individual, independent of calendar time t . The probability distribution of infectiousness profile depends on biological factors such as immunity or genetics for an individual. Serial interval may be employed as a proxy for w_s . Subsequently, the average incidence at t is (eq 4),

$$E[I_t] = R_t \sum_{s=1}^t I_{t-s} w_s \dots\dots\dots (eq\ 4)$$

Where I_{t-s} is the incidence reported at time $t-s$. Based on this transmission model and assuming gamma distribution as the prior distribution for R_t , the underlying posterior distribution for R_t is derived through Bayesian statistical inference. The underlying mathematical modelling is elaborated in additional material. An elaborative overview of R_{ins} is also provided in the work of Cori et al. The R_{case} and R_{ins} were estimated using the package R0 and EpiEstim in the R software (Version 4.0.0).^{[17][18]} The simulations were performed under the aforementioned assumptions within a confidence interval (CI) of 95%.

3. Results and Discussion

The average R_{case} and R_{ins} for India during the imposition of Lockdown-1 (March 25, 2020) to the later phase of Unlock-3 (August 23, 2020) was 1.22 (1.19, 1.26; 95% CI) and 1.20 (1.18, 1.21; 95% CI) respectively. Both the metrics were indicating high infection transmission during the early time in the Lockdown-1, consistently declining with two consecutive minor yet significant peaks reported around the end of Lockdown-2 (~ April 30, 2020) and around the beginning of Lockdown-4 (~ May 18, 2020); and eventually continues to stabilize from Unlock-1 and remain steady thereafter (Fig1. A, B). We then compared R_{case} and R_{ins} estimates during the unlock phase 1-3 as the state border of the containment zone were sealed but the inter-state movement was unrestricted. The average R_{case} and R_{ins} for the post unlock period (i.e. June 1, 2020 to August 23, 2020) was 1.15 (1.13, 1.16; 95% CI) and 1.12 (1.11, 1.12; 95% CI) respectively (Fig1. C, D). The R_{case} consistently remained higher than average R_{ins} and also elucidated a noisy projection of R_t . R_{case} remained high (4.05 (3.77, 4.32; 95% CI) & 3.33 (3.31, 3.36; 95% CI)) early in the time series for both the interval which is very unpragmatic. For entire incidence across India, the two methods almost resembled each other however when estimated over the unlock phase, R_{case} indicated a significantly higher R_t value of 3.33 (3.31, 3.36; 95% CI) in the beginning of the time interval, which contradicts with its own R_t estimates of 1.14 (1.11, 1.16; 95% CI) when estimated over the entire Lockdown-Unlock phase (i.e. March 25 to August 23) (Fig1. A, C). On the other hand, R_{ins} estimates for unlock phases remained in line with the estimated R_{ins} for the entire lockdown-unlock period (Fig 1.B, D).

On the same note, these irregularities in R_{case} estimates were observed while probing the transmission of COVID-19 in the high incidence states (Fig S1, S2). For each state, the average R_{case} remained overestimated than average R_{ins} for both the time intervals viz entire lockdown-unlock duration and the Unlock 1-3. Average estimates of R_{case} and R_{ins} for the high incidence states including India is given in Table 2. We found that R_{case} consistently estimated a high R_t in the initial days of the time series independent of the intervals, which implicates misleading conclusions regarding the infection transmission during the unlock phases (Fig S2). While the real-time estimate of R_t by R_{ins} was found superior to R_{case} for all the sample states, this was best illustrated in the cases for Delhi and West Bengal (Fig2). As on August 23, 2020, Delhi contained the highest incidence per million (7996.14) in India and

has conducted the highest number of COVID-19 testing per million (70,870.8). The daily new cases in Delhi started to fall significantly from around June 23, 2020, and continued to decline with irregular local minimums (Fig2. A). This consistent fall was best captured by R_{ins} while R_{case} remained sensitive for the local peaks and overestimated the incidence during unlock 2-3 (Fig2. B). More importantly, the R_{ins} detected the variations at earliest while the R_{case} estimations remained delayed. The underlying mathematical reason behind this anomalistic estimation by R_{case} is that the R_{case} amalgamates a weighted sum across transmission events observed after time t independent of the trend of transmission before t , and this leads to misleading estimations in the latter portion of the time series. While 138,871 total cases of COVID-19 were recorded for West Bengal as of August 23, 2020, the growth of the incidence remained slow during the Lockdown Phase 1-2 viz March 25, 2020 to April 15, 2020, possibly due to underreporting. This anomaly in data was highly underestimated by R_{case} , indicating a complete end of the transmission ($R_{case}=0$) (Fig2. C), which is very contradictory. On the other hand, R_{ins} reflected a subtle decline in the estimate but remained in line with the overall true incidence projection by remaining immune to such data inconsistency (Fig2. C). Similar contradictory over- & underestimations by R_{case} were observed for the cases of other states such as Maharashtra, Gujrat and Telangana (Fig S1, S2).

These deflections of R_{case} from R_{ins} estimates for a state were found significantly associated with the amount of testing the state has conducted (Fig3). Higher deflection was observed for the states that have conducted a high number of testing (R square 0.801, $p=0.003$; CI 95%) (Fig3. B). This association remained intact for the Unlock 1-3 interval as well (Fig3. C). We assert that (1) Because of high testing, the incidence data represents close to real-time incidence occurring in the state, which subsequently facilitates a better comparison of the models. (2) The deflection of R_{case} from R_{ins} estimate could also serve as an indicator for assessment of the sufficiency of testing against the infection transmission.

4. Conclusion

Although many studies in the past have employed R_{case} for the modelling of effective reproduction number R_t in order to probe COVID-19 outbreak in India, this study retrospectively investigated the contrasting estimates of R_t by R_{case} and R_{ins} for the high incidence states of India, during lockdown and unlock phases.^{[12][19]} We found that, (1) while R_{case} remained deflected from the true incidence, R_{ins} captured the variations in line with the real-time trend of incidence recorded in the states during the phases. (2) R_{case} estimates were lagging from the R_{ins} values and delayed in the measuring transmission events that occurred days ago. (3) R_{case} misestimated the transmission rate at beginning of the times while R_{ins} estimations remained credible near the threshold value which is a crucial region to assess the implemented policies.

Since a reliable model is vital for the assessment of the effectiveness of intervention to contain an outbreak, through this work we highly recommend for the implementation of a fine tuned R_{ins} estimate over R_{case} to retrospectively monitor the effectiveness of the interventions. However, we also emphasize not to overlook the underlying assumptions made which could invalidate the R_{ins} estimation and deliver

misleading conclusions if not corroborated properly. For instance, in COVID-19 infection, the serial interval distribution might vary from the generation time distribution as the infector can infect the infectee prior to the symptom's onset, thus indicating a negative value for the serial interval [8]. We hope this work would facilitate an improved assessment modelling for the COVID-19 outbreak for the optimization and designing of effective interventions to combat the COVID-19 pandemic.

5. Author Contributions

Amrutha G.S: Conceptualization, Investigation, Formal Analysis, Writing- Reviewing and Editing. **Abhibhav Sharma:** Conceptualization, Methodology, Data curation, Software, Writing- Original draft preparation. **Anudeepti Sharma:** Formal analysis, Visualization, Writing- Reviewing and Editing.

6. References

[2][3][4][5] [6][7][8][9][10][11][12][13][14][15][16][17][18][19]

1. Worldometers.info. Coronavirus Update [Internet]. Available from: <https://www.worldometers.info/coronavirus/#countries>
2. Zhao S, Stone L, Gao D, He D. Modelling the large-scale yellow fever outbreak in Luanda, Angola, and the impact of vaccination. *PLoS Negl Trop Dis* 2018; 12(1)
3. Van Den Driessche P, Watmough J. Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Math Biosci* 2002; 180(1–2), 29–48
4. Wallinga J, Teunis P. Different epidemic curves for severe acute respiratory syndrome reveal similar impacts of control measures. *Am J Epidemiol* 2004; 160(6) 509–516
5. Gostic KM, McGough L, Baskerville E, Abbott S, Joshi K, Tedijanto C, et al. Practical considerations for measuring the effective reproductive number, Rt. *medRxiv Prepr Serv Heal Sci* 2020;
6. Panovska-Griffiths J. Can mathematical modelling solve the current Covid-19 crisis? *BMC Public Health* 2020; 20(1).
7. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-varying reproduction numbers during epidemics. *Am J Epidemiol* 2013; 178(9) 1505–1512
8. Nishiura H, Linton NM, Akhmetzhanov AR. Serial interval of novel coronavirus (COVID-19) infections. *Int J Infect Dis* 2020; 93 284–286
9. Ganyani T, Kremer C, Chen D, Torneri A, Faes C, Wallinga J, et al. Estimating the generation interval for coronavirus disease (COVID-19) based on symptom onset data, March 2020. *Eurosurveillance* 2020; 25(17)
10. Bettencourt LMA, Ribeiro RM. Real time bayesian estimation of the epidemic potential of emerging infectious diseases. *PLoS One* 2008; 3(5)
11. Cintrón-Arias A, Castillo-Chavez C, Bettencourt LMA, Lloyd AL, Banks HT. The estimation of the effective reproductive number from disease outbreak data. *Math Biosci Eng* 2009; 6(2) 261–

282

12. Marimuthu S, Joy M, Malavika B, Nadaraj A, Asirvatham ES, Jeyaseelan L. Modelling of reproduction number for COVID-19 in India and high incidence states. Clin Epidemiol Glob Heal 2020;
13. Kim S, Jeong YD, Byun JH, Cho G, Park A, Jung JH, et al. Evaluation of COVID-19 epidemic outbreak caused by temporal contact-increase in South Korea. Int J Infect Dis 2020; 96 454-457
14. Saberi M, Hamedmoghadam H, Madani K, Dolk HM, Morgan AS, Morris JK, et al. Accounting for Underreporting in Mathematical Modeling of Transmission and Control of COVID-19 in Iran. Front Phys 2020; 8
15. National commission on population ministry of health & family welfare nirman bhawan, new delhi – 110011. 2019;Available from: https://nhm.gov.in/New_Updates_2018/Report_Population_Projection_2019.pdf
16. Fraser C. Estimating individual and household reproduction numbers in an emerging epidemic. PLoS One 2007; 2(8)
17. Boelle P-Y, Obadia T. R0: Estimation of R0 and Real-Time Reproduction Number from Epidemics [Internet]. 2015;Available from: <https://cran.r-project.org/package=R0>
18. Cori A. EpiEstim: Estimate Time Varying Reproduction Numbers from Epidemic Curves [Internet]. 2020;Available from: <https://cran.r-project.org/package=EpiEstim>
19. Basu D, Salvatore M, Ray D, Kleinsasser M, Purkayastha S, Bhattacharyya R, et al. A Comprehensive Public Health Evaluation of Lockdown as a Non-pharmaceutical Intervention on COVID-19 Spread in India: National Trends Masking State Level Variations. medRxiv Prepr Serv Heal Sci 2020;

Table 1

State	Population*	Total cases as of August 23, 2020	Total cases per million as of August 23, 2020	Total test per million as of August 23, 2020	Test Positive Rate (TPR)
India	1381446532	3104558	2247.32	25537	
Andhra Pradesh	52504000	353111	6725.411	61672.22	10.2%
Delhi	20193000	161466	7996.137	70870.8	11.6%
Gujarat	68862000	86779	1260.187	25502.21	5.9%
Karnataka	66322000	277814	4188.866	36397.44	11.2%
Maharashtra	1.23E+08	682383	5534.555	29371.19	19.2%
Tamil Nadu	76049000	379385	4988.692	55314.56	9.2%
Telangana	37473000	105272	2809.276	24866.94	12.2%
West Bengal	97516000	138871	1424.084	16010.82	8.9%

Table 1: Metrics of India and Indian states containing high incidence of COVID-19. * Projected Total Population as on 1st March – 2020.^[15]

Table 2

State	March 25, 2020 to August 23, 2020				June 01, 2020 to August 23, 2020			
	Average R_{case}	R_{case} 95% CI	Average R_{ins}	R_{ins} 95% CI	Average R_{case}	R_{case} 95% CI	Average R_{ins}	R_{ins} 95% CI
India	1.22	(1.19, 1.26)	1.20	(1.18, 1.21)	1.15	(1.13, 1.16)	1.12	(1.11, 1.12)
Andra Pradesh	1.42	(1.13, 1.74)	1.28	(1.20, 1.36)	1.34	(1.26, 1.42)	1.26	(1.23, 1.28)
Delhi	1.49	(1.24, 1.73)	1.19	(1.14, 1.23)	1.37	(1.32, 1.42)	1.02	(1.00, 1.03)
Gujrat	1.22	(1.05, 1.42)	1.19	(1.13, 1.26)	1.09	(1.02, 1.16)	1.06	(1.03, 1.09)
Karnataka	1.27	(1.02, 1.54)	1.23	(1.14, 1.33)	1.27	(1.20, 1.34)	1.20	(1.18, 1.23)
Maharashtra	1.24	(1.16, 1.33)	1.21	(1.18, 1.24)	1.13	(1.11, 1.16)	1.10	(1.09, 1.11)
Tamil Nadu	1.37	(1.20, 1.55)	1.23	(1.19, 1.28)	1.19	(1.15, 1.22)	1.10	(1.08, 1.11)
Telangana	1.28	(1.03, 1.56)	1.21	(1.12, 1.31)	1.23	(1.15, 1.31)	1.17	(1.14, 1.20)
West Bengal	1.26	(1.04, 1.51)	1.23	(1.14, 1.33)	1.16	(1.10, 1.23)	1.13	(1.11, 1.16)

Table 2: Average estimation of case reproduction number (R_{case}) and instantaneous reproduction number (R_{ins}) for India and high incidence states for Lockdown-Unlock interval (March 25, 2020 to August 23, 2020) and Unlock 1-2 (June 01, 2020 to August 23, 2020) intervals with 95% CI.

It is made available under a [CC-BY 4.0 International license](#).

Figure_2(C)

Case reproduction number (R_{case}) and Instantaneous reproduction number (R_{ins}) estimation for West Bengal from the time of Lockdown-1 till late Unlock-3 (March 25, 2020 to August 23, 2020).

It is made available under a [CC-BY 4.0 International license](#).

Figure_2(B)

Case reproduction number (R_{case}) and Instantaneous reproduction number (R_{ins}) estimation for Delhi from the time of Lockdown-1 till late Unlock-3 (March 25, 2020 to August 23, 2020).

Figure_3(C)

Plots showing the relationship between the difference of and (i.e. $R_{\text{case}} - R_{\text{ins}}$) measured from the time of Unlock-1 till late Unlock-3 (June 1, 2020 to August 23, 2020) for the state and the number of testing conducted per million by the state as on August 23, 2020.

Figure_3(B)

Plots showing the relationship between the difference of and (i.e. $R_{\text{case}} - R_{\text{ins}}$) measured from the time of Lockdown-1 till late Unlock-3 (March 25, 2020 to August 23, 2020) for the state and the number of testing conducted per million by the state as on August 23, 2020.

It is made available under a [CC-BY 4.0 International license](#).

Figure_1(D)

Instantaneous reproduction number (R_{ins}) estimate for India from the time of Unlock-1 till late Unlock-3 (June 1, 2020 to August 23, 2020).

It is made available under a [CC-BY 4.0 International license](#).

Figure_1(C)

Case reproduction number (R_{case}) estimate for India from the time of Unlock-1 till late Unlock-3 (June 1, 2020 to August 23, 2020).

It is made available under a [CC-BY 4.0 International license](#).

Figure_1(B)

Instantaneous reproduction number (R_{ins}) estimate for India from the time of Lockdown-1 till late Unlock-3 (March 25, 2020 to August 23, 2020).

It is made available under a [CC-BY 4.0 International license](#).

Figure_1(A)

Case reproduction number (R_{case}) estimate for India from the time of Lockdown-1 till late Unlock-3 (March 25, 2020 to August 23, 2020).

It is made available under a [CC-BY 4.0 International license](#).

Figure_3(A)

Illustrating daily tests conducted per million in high incidence states of India from the time of Lockdown-1 till late Unlock-3 (March 25, 2020 to August 23, 2020).

It is made available under a [CC-BY 4.0 International license](#).

Figure_2(A)

Illustrating daily new cases per million in high incidence states of India from the time of Lockdown-1 till late Unlock-3 (March 25, 2020 to August 23, 2020).