Keen relationship between COVID_19 and food supply suggest some animal origin foods as excellent vehicle of SARS-Cov-2

Nazli Saeedi¹, Seyed Abbas Rafat^{2*}

¹ Research Center for Pharmaceutical Nanotechnology, Biomedicine Institute, Tabriz University of Medical Sciences, Tabriz, Iran Email : saeedi@hotmail.fr, ² Department of Animal Science, Faculty of Agriculture, University of Tabriz, Tabriz, Iran, Email; abbasrafat@hotmail.com *Corresponding Author

Abstract

In recent months, nearly all countries tried to decrease human-to-human contact as the principal mode of transmission of SARS-CoV-2. However, other modes of transmission also need to be clarified in more depth, especially, the foodborne transmission. We assessed the effect of animal origin foods consumption on the pandemic of COVID-19. For this purpose, we studied the relationship among 20 food supply as independent variables, and the parameter of Total Cases as dependent variable. Here we show a relationship between a group of animal origin foods and total cases. Regression, Bayes, and Lasso results showed that eggs and fresh water fish have positive coefficient. So, among the transmission ways of COVID_19, the role of foodborne transmission should be more significant than previously thought. The possibility of animal origin foodborne transmission should be taken into more consideration. The perspective is to expand the surveillance of SARS-Cov-2 during the food production chain. In conclusion, the results of the present study indicate that one important vehicle for SARS-Cov-2 may be some of animal origin

foods. It is recommended that virologists examine the possibility of freshwater fish and chicken's eggs being as excellent vehicles/preservatives for SARS-Cov-2.

Introduction

At the time of writing this manuscript, the number of total cases of COVID-19 past 28 mark. During the pandemic, most countries around the world tried to minimize human-to-human contact and travels(Merler et al., 2020). However, the prevalence of COVID-19 continued. Despite efforts of many countries to reduce the transmission by person-to-person way the rapid outbreak shows the necessity of research on other possible transmission routes. During early pandemics, in Italy and the USA, for example, the chances of being surprised by pandemics were low, but both countries faced epidemic dramatically. Among the various routes of transmission, FBT is most likely. We suppose the role of COVID-19's transmission by other ways should be more significant than previously thought. Transmission of MERS, a similar pathogenic virus to SARS-Cov-2, through food has strengthened our hypothesis.

Furthermore, some researchers cited the possibility of FBT for SARS-Cov-2(Hi, 2020; Kingsbury et al., 2020; Newell et al., 2010; Panel and Biohaz, 2011; Todd et al., 2020). Also, FAO/WHO experts mentioned the data gap in the relationship between viruses and food borne disease (FAO/WHO, 2008) . The motivation for the present study is to find the effects of human food consumptions (independent variables) on COVID_19 (As dependent variable).

Material and Methods

Table 1 shows descriptive statistics of studied variables. We assessed the possible relationship between the consumption of main foods and total cases od covid_19. Accordingly, we studied

total cases as a dependent variable and 20 food supply data as independent variables. The primary purpose of the study was to find effect of animal origin food on total cases, however, we considered six plant origin foods for comparison reasons. The statistical methods included correlation, stepwise and LASSO regression, and Bayes analysis. Food supply (kg/capita/yr.) was quantity in each country based on the latest available data of FAO. Adjusted R-square of fitted models are presented in Table 2, respectively. Detail material and methods are included in supplementary materials.

Table 1

Descriptive statistics of studied foods (n=20) besides total cases.

#	Variable (kg/capita/yr)	Abbreviation	N	Mean	Std Dev	Minimum	Maximum
1	Pig meat	pig	169	14.66	15.52	0.00	71.48
2	Bovine Meat	bov	174	11.20	9.00	0.78	55.48
3	Cheese	che	173	4.97	6.55	0.00	30.82
4	Cream	cre	159	1.21	2.76	0.00	18.45
5	Eggs	Egg	174	7.08	5.09	0.16	19.15
6	Freshwater Fish	freshf	174	3.38	4.54	0.00	34.00
7	Honey	hon	168	0.36	0.46	0.00	3.51
8	Whole milk	milk	174	59.17	48.93	0.96	258.44
9	Poultry Meat	poul	174	20.56	16.46	0.42	72.87
10	Demersal Fish	demer	173	3.40	4.93	0.00	33.45
11	Cephalopods	ceph	171	0.44	0.96	0.00	6.70
12	Butter, Ghee	butg	174	1.25	1.64	0.00	9.25
13	Marine Fish, Other	marf	172	2.74	4.87	0.00	28.19
14	Aquatic Plants	aqp	162	0.20	1.91	0.00	22.41
15	Beverages, Alcoholic	bev	171	3.20	3.87	0.00	26.90
16	Rice(Milled Equivalent)	rice	174	30.95	36.20	0.95	171.73
17	Potatoes and products	potato	174	34.89	33.65	0.05	183.16
18	Maize and products	maiz	170	21.71	28.52	0.00	158.40
19	Wheat and products	wheat	174	70.92	46.85	1.89	222.40
20	wine	wine	169	6.27	9.49	0.00	51.41
	Total Cases (Per Country)	TC	215	130744.74	619823.34	3	6558095

Table 2

R-Square, Adjusted R-square, AIC and number of data in the fitted models for food consumption.

	Stepwise	LASSO	BAYES
R-Square	0.33	0.47	-
Adj-R Square	0.31	0.42	-
AIC	-	313	562
n	131	131	130

We estimated the relationship between 20 food supply and Total Cases of covid19. The methods include Stepwise Regression with Proc Reg, LASSO regression with GLMSELECT Proc, and Bayes analysis with Proc Genmod (SAS). Food supply was quantity (kg/capita/yr.). Fourteen foods of animal origin and six foods of plant origin were selected. The primary purpose of the study was to find AOF relationship with total cases, but some plant origin foods were chosen for comparison reason.

We started our analyses first time on March 2020. Then we repeated it several times with updated COVID_19 data on April, May, June, and September. Finally we presented the latest results.

Data

We downloaded the data the following websites:

COVID_19: <https://www.worldometers.info/coronavirus/> 10 September 2020

Crops Primary Equivalent: <http://www.fao.org/faostat/en/#data/CC>

Food Supply - Livestock and Fish: <http://www.fao.org/faostat/en/#data/CL>

We used base-*e* logs, also known as natural logs for the transformation of total cases and independent variables. In all analyses, the accuracy of the model with non-transformed data was very low. So, all variables has been transformed to

normalize the distributions. With transformation of data, the accuracy of the model increased to an acceptable level. In comparison of models, Lasso's results are more reliable than stepwise regression and Bayes, because it's AIC was smaller and also it had higher accuracy as R-square (Table 2). Visual examination of the trace plot displays proper mixing in Bayesian analysis.

We provide the SAS code for an application of the used statistical methods (Rafat, 2020). We note that this work was posted as preprint on medrxiv (Rafat, seyed abbas, 2020) .

Results and Discussion

Stepwise Regression, Bayes, and Lasso's results confirmed that eggs and fresh water fish have high positive relevance with total cases. With our results in Table 3, implying the need for tracing of SARS-COV-2 in main AOF. Bayes results, e.g., show there is a 0.99 probability of a positive correlation between eggs and total cases, adjusted for the other covariates. The number of AOF and their role in our results varies depending on the type of statistical method used, but all results are consistent. The results of all three statistical methods confirm each other in the case of eggs. Negative high correlation between marine fish and total cases is curious observation needs more study.

118

119 **Table 3. The results of multivariate analyses.** All variables are logarithmic transformed data.

120

Variable	Stepwise	Lasso	Bayes
Pig meat		-0.26	0.36
Bovine Meat		0.12	0.75
Cheese			0.35
Cream		-0.18	0.19
Eggs	0.85	0.98	0.99
Freshwater fish		0.72	0.99
Honey		-1.18	0.03
Whole milk			0.34
Poultry meat			0.38
Demersal fish			0.47
Cephalopods			0.36
Butter/ghee		0.19	0.84
Marine fish(other)	-1.49	-1.32	-1.00
Aquatic plants			0.38
Beverages(Alcoholic)			0.18
Rice(Milled Equivalent)		0.30	0.95
Potatoes and products		0.37	0.96
Maize and products		0.39	0.99
Wheat and products		0.34	0.91
Wine			0.64

121

122 The effect of a group of AOF on total cases was significant in our analysis. So, we
 123 suppose a noticeable effect of AOF as FBT/vehicle. At present, we are not able to
 124 interpret all significant effects and their coefficients. Interpretation of the factors
 125 could be the subject of future researches. Potential transmission ways of COVID-
 126 19 based on the results of our analysis and bibliography is summarized as Fig 1.
 127 Lake and Kingsbury (2020) provided a review about potential for FBT of COVID-
 128 19, but they could not guarantee the application of the data found on other
 129 coronaviruses to SARS-CoV-2(Kingsbury et al., 2020). The persistence of viruses
 130 by refrigeration temperature(FAO/WHO, 2008) might be an explanation of the
 131 relationship between COVID-19 and AOF. The viral persistence of coronavirus in
 132 soft cheese (Panon et al., 1988) and raw milk(Hamdi, 2013) has been reported. We
 133 don't know what's going on from “Farm to Fork” about AOF. For example, SARS-
 134 CoV-2 may be more resistant to heat when cooking, as it is observed for poliovirus
 135 in beef(Pirtle, 1991). It is recommended not to consume raw or undercooked
 136 animal products(Moy, 2020). Kitchen utensils can transmit the virus among
 137 foods(Sarah C. P. Williams, n.d.), accordingly, AOF may be a suitable vehicle, and
 138 the virus inside them will survive better and longer.

139 Furthermore, the methods of diagnosing the virus in food are not complete, and
 140 there are many limitations(Bosch et al., 2018). In food science technology, the

primary attention paid to bivalve moullouses. Nowadays, it is necessary to expand the detection of viruses to all ANIMAL ORIGIN FOOD.

The high-fat tissue in the body has effect on the pathogenesis of COVID-19(Maffetone and Laursen, 2020). Although the probability of transmitting the virus through wine is zero, we deliberately kept it in the initial model of our analysis. Results showed wine is eliminated among significant variables of the suited model (Table 3). So we concluded that just the energy content of food should not keep a variable in the fitted model. So, in addition to energy content, there must be something else in food-related COVID-19. We concentrate on the transmission of COVID-19 by AOF, rather than the immunity response.

The paper of (Katy Askew, 2020) guided us to construct our hypothesis about the essential relationship between COVID-19 and AOF include.

WHO's recommendations state that COVID_19 have not FBT, and suitable food safety practices could prevent the probability of FBT. However, the realization of proper food safety practices for all AOF is questionable. It is reported that acquisition of SARS coronavirus by consumption of contaminated food is possible (Yepiz 2013). Furthurmore, oral infection of SARS-Cov-2 could not be excluded among transmission ways (FAO/WHO 2008). Promkuntod 2006 recommends

taking precautions to prevent the spread of the avian flu by consumption of quail eggs(PROMKUNTOD et al., 2006). One explanation for our observation is the trapping and accumulation of sars-cov-2 by AOF. Thereafter, along food processing or preparation, AOF conserve SARS-Cov-2 infectively.

Conclusion

Despite the awareness and readiness of many countries for COVID-19 during early 2020, why did it spread so quickly? It seems FBT has received less attention relative to other routes. The perspective is to expand the surveillance of SARS-Cov-2 during the food production chain. It is recommended that virologists examine the possibility of freshwater fish and chicken's eggs being as an excellent vehicle/preservator for SARS-Cov-2. Some animal origin food may have been the first cause of transmission of the SARS-Cov-2 from reservoirs populations to humans.

ACKNOWLEDGEMENTS:

Author contributions: Conceptualization and Visualisation: N. S. K., Methodology: S. A. R. Writing: Both authors. The authors have declared no competing interest. Data and materials availability: All data are available in the manuscript or the supplementary materials.

References

- Bosch, A., Gkogka, E., Le, F.S., Loisy-hamon, F., Lee, A., Lieshout, L. Van, Marthi, B., Myrmel, M., Sansom, A., Charlotte, A., Winkler, A., Zuber, S., Phister, T., 2018. International Journal of Food Microbiology Foodborne viruses: Detection , risk assessment , and control options in food processing. International Journal of Food Microbiology 285, 110–128. <https://doi.org/10.1016/j.ijfoodmicro.2018.06.001>
- Bourgarel, M., Pfukenyi, D., Becquart, P., Morand, S., 2019. Next-Generation Sequencing on Insectivorous Bat Guano: An Accurate Tool to Identify Arthropod Viruses of Potential Agricultural Concern. <https://doi.org/10.3390/v11121102>
- Cavanagh, D., Cavanagh, D., 2007. Coronaviruses in poultry and other birds Coronaviruses in poultry and other birds 9457. <https://doi.org/10.1080/03079450500367682>
- FAO/WHO, 2008. [Food and Agriculture Organization of the United Nations/World Health Organization]. Microbiological Risk Assessment Series 14, 151.
- Geisbert, T.W., Mire, C.E., Geisbert, J.B., Chan, Y.-P., Agans, K.N., Feldmann, F., Fenton, K.A., Zhu, Z., Dimitrov, D.S., Scott, D.P., Bossart, K.N., Feldmann, H., Broder, C.C., 2014. Therapeutic treatment of Nipah virus infection in nonhuman primates with a neutralizing human monoclonal antibody. Science translational medicine 6, 242ra82-242ra82. <https://doi.org/10.1126/scitranslmed.3008929>
- Gibbins, J.D., Hale, C., Silver, M., Fischer, M., Steinberg, J., Basler, C.A., Jacobs, J.R., Kennedy, E.D., Tomasi, S., Trout, D., Hornsby-myers, J., 2020. COVID-19 Among Workers in Meat and Poultry Processing Facilities — 69, 557–561.

202 Guan, Y., Zheng, B.J., He, Y.Q., Liu, X.L., Zhuang, Z.X., Cheung, C.L., Luo, S.W., Li, P.H.,
203 Zhang, L.J., Guan, Y.J., Butt, K.M., Wong, K.L., Chan, K.W., Lim, W., Shortridge, K.F.,
204 Yuen, K.Y., Peiris, J.S.M., Poon, L.L.M., 2003. Isolation and characterization of viruses
205 related to the SARS coronavirus from animals in southern China. Science (New York, NY)
206 302, 276–278. <https://doi.org/10.1126/science.1087139>

207 Hamdi, A., 2013. DETECTION OF ROTA AND CORONA VIRUSES IN RAW MILK AND
208 MILK PRODUCTS. Benha veterinary medical journal 24, 79–85.

209 Hi, Z.S., 2020. REVIEW ARTICLE A brief review of foodborne zoonoses in China 1497–1504.
210 <https://doi.org/10.1017/S0950268811000872>

211 Kai Kupferschmidt, 2013. New Middle Eastern Virus Found in Camels.

212 Katy Askew, 2020. 2020 German slaughterhouses under fire over “insufficient precautionary
213 measures” to block COVID-19 spread.

214 Kingsbury, J., Lake, R., Hewitt, J., Smit, E., King, N., 2020. POTENTIAL FOR FOODBORNE
215 TRANSMISSION OF COVID-19□: 1–51.

216 Maffetone, P.B., Laursen, P.B., 2020. The Perfect Storm□: Coronavirus (Covid-19) Pandemic
217 Meets Overfat Pandemic 8, 1–6. <https://doi.org/10.3389/fpubh.2020.00135>

218 Maori, E., Trust, W., Reports, C., 2019. Discovery of RNA transfer through royal jelly could aid
219 development of honey bee vaccines 3–5. <https://doi.org/10.1016/j.celrep.2019.04.073>

220 McMahon, D.P., Wilfert, L., Paxton, R., Brown, M., Holloway, R., 2018. Emerging Viruses in
221 Bees□: From. <https://doi.org/10.1016/bs.aivir.2018.02.008>

222 Merler, S., Pastore, A., Mu, K., Rossi, L., Sun, K., 2020. The effect of travel restrictions on the

223 spread of the 2019 novel coronavirus (COVID-19) outbreak 400, 395–400.

224 Miller, R.S., Sweeney, S.J., Sloomaker, C., Gear, D.A., Salvo, P.A. Di, Kiser, D., Shwiff, S.A.,
 225 2017. Cross-species transmission potential between wild pigs , livestock , poultry , wildlife ,
 226 and humans□: implications for disease risk management in North America. Scientific
 227 Reports 1–14. <https://doi.org/10.1038/s41598-017-07336-z>

228 Moy, G.G., 2020. IUFoST / CIFST hold an Extraordinary Scientific Roundtable on COVID-19
 229 and Food Safety. npj Science of Food 1–2. <https://doi.org/10.1038/s41538-020-0068-2>

230 Newell, D.G., Koopmans, M., Verhoef, L., Duizer, E., Aidara-kane, A., Sprong, H., Opsteegh,
 231 M., Langelaar, M., Threlfall, J., Scheutz, F., Giessen, J. Van Der, Kruse, H., 2010.
 232 International Journal of Food Microbiology Food-borne diseases — The challenges of 20
 233 years ago still persist while new ones continue to emerge. International Journal of Food
 234 Microbiology 139, S3–S15. <https://doi.org/10.1016/j.ijfoodmicro.2010.01.021>

235 Panel, E., Biohaz, H., 2011. Scientific Opinion on an update on the present knowledge on the
 236 occurrence and control of foodborne viruses 19, 1–96.
 237 <https://doi.org/10.2903/j.efsa.2011.2190>.

238 Panon, G., Tache, S., Labie, C., 1988. Respective stability of rotavirus and coronavirus in bovine
 239 milk 68, 49–64.

240 Papineau, A., Berhane, Y., Wylie, T.N., Wylie, K.M., Sharpe, S., Lung, O., 2019. Genome
 241 Organization of Canada Goose Coronavirus , A Novel Species Identified in a Mass Die-off
 242 of Canada Geese. Scientific Reports 1–8. <https://doi.org/10.1038/s41598-019-42355-y>

243 Pirtle, E.C., 1991. Virus survival in the environment 10, 733–748.

244 PROMKUNTOD, N., ANTARASENA, C., PROMMUANG, PORNTIP, PROMMUANG,
245 PRAISON, 2006. Isolation of Avian Influenza Virus A Subtype H5N1 from Internal
246 Contents (Albumen and Allantoic Fluid) of Japanese Quail (*Coturnix coturnix japonica*)
247 Eggs and Oviduct during a Natural Outbreak. *Annals of the New York Academy of*
248 *Sciences* 1081, 171–173. <https://doi.org/doi:10.1196/annals.1373.020>

249 Rafat, S.A. , saeedi, N. 2020. Human food consumption patterns concerning COVID-19
250 pandemic. [https://www.medrxiv.org/content/10.1101/2020.06.16.20132464v4.article-](https://www.medrxiv.org/content/10.1101/2020.06.16.20132464v4.article-metrics)
251 [metrics](https://www.medrxiv.org/content/10.1101/2020.06.16.20132464v4.article-metrics)

252 Rafat, S.A., 2020. DATA of Preprint: Human food consumption patterns concerning COVID-19
253 pandemic. <https://doi.org/10.13140/RG.2.2.30851.12321>

254 Rizzo, F., Edenborough, K.M., Toffoli, R., Culasso, P., Zoppi, S., Dondo, A., Robetto, S.,
255 Rosati, S., Lander, A., Kurth, A., Orusa, R., Bertolotti, L., Mandola, M.L., 2017.
256 Coronavirus and paramyxovirus in bats from Northwest Italy 1–11.
257 <https://doi.org/10.1186/s12917-017-1307-x>

258 Sarah C. P. Williams, n.d. Kitchen Utensils Transfer Viruses.

259 Tao, Z., Tian, J., Pei, Y., Yuan, M., Zhang, Y., Dai, F., 2020. A new coronavirus associated with
260 human respiratory disease in China 579. <https://doi.org/10.1038/s41586-020-2008-3>

261 Todd, E.C.D., Greig, J.D., Todd, E., Llc, C., 2020. Viruses of foodborne origin□: a review 25–
262 46.

263

264

Figure captions

Fig. 1 Potential transmission ways of COVID-19 based on the results of our analysis and bibliography. It is necessary to study the exchange of human pathogen viruses in wildlife and livestock. Where and how suspicious foods become contaminated must be investigated. The potential spillover of pathogenic viruses from arthropods hosts to humans should be investigated(Tao et al., 2020). The cross-species transmission is reported(Miller et al., 2017). There may be a type of contact among flowers, bees, and bats. Infected humans could have transmitted MERS to the camels (Kai Kupferschmidt, 2013), similar situation is imaginable for SARS-Cov-2. SARS-CoV-2 and MERS-Co are among the most notable examples of spillover into humans(Guan et al., 2003). References for arrows: 1 (Bourgarel et al., 2019; Cavanagh and Cavanagh, 2007; Maori et al., 2019; McMahon et al., 2018; Moy, 2020; Rizzo et al., 2017) 2: (Miller et al., 2017; Moy, 2020) 3: (Miller et al., 2017; Papineau et al., 2019) 4:(Geisbert et al., 2014; Tao et al., 2020) 5: (Gibbins et al., 2020; Newell et al., 2010; Panon et al., 1988) 6 : (Gibbins et al., 2020; Hi, 2020; Katy Askew, 2020)