Article

Correlation between serum 25(OH) vitamin D level and sleep quality in elderly patients with type 2 diabetes

Hongyan Ma^{1,*†}, Danni Fan², Xiaofei Li³, Ying Qi⁴, Yang Zhao⁵, Lin Zhang^{5,6,7†}

1. Department of Geriatrics, The First Affiliated Hospital of University of Science and Technology of China, Hefei 230001, China; (slyyhongyan816@126.com)
2. East Kent Hospitals University NHS Foundation Trust, Ethelbert Road, Canterbury Kent, CT1 3NG, England (Danni.fan@nhs.net)
3. The Second Clinical College of Guangzhou University of Chinese Medicine, Guangzhou, China (lxf13751726381@163.com)
4. School of Health Policy and Management, Chinese Academy of Medical Sciences & Peking Union Medical College, Beijing, China (yingqi@vip.126.com)
5. Melbourne School of Population and Global Health, The University of Melbourne, Victoria 3100, Australia. (xiyunjushi@163.com)
6. School of Population Medicine and Public Health, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing, 100021, China (Tony1982110@gmail.com)
7. Centre of Cancer Research, Victorian Comprehensive Cancer Centre, Melbourne, Victoria, Australia

*Correspondence: Dr. Hongyan Ma, Email: slyyhongyan816@126.com

† These authors contributed equally to this work and should be considered co-first authors.

Abstract:

Background: Untreated sleep disorders have been linked with type 2 diabetes in previous evidence. This study was to investigate the correlation between serum 25(OH) vitamin D level and sleep quality in elderly patients with type 2 diabetes and reveal the therapeutic potential of 25(OH) vitamin D-based treatments to improve sleeping quality and its effect on diabetes management.

Methods: The sleep quality of 186 elderly patients with type 2 diabetes were assessed by questionnaires of Pittsburgh sleep index (PSQI), with a cutoff point of PSQI • 7 defined as poor sleep. The measurement of serum 25(OH) vitamin D, fasting blood glucose, glycosylated hemoglobin, serum creatinine, total cholesterol, triglyceride, low-density lipoprotein cholesterol, and urinary albumin/creatinine ratio were collected. History of diabetic nephropathy, diabetic retinopathy, peripheral neuropathy, blood pressure, height, weight, hypoglycemic drug use, and the duration of diabetes also were recorded. Both univariate and multivariate logistic regression analysis was used to investigate the relationship of serum 25(OH) vitamin D and sleep quality in elderly patients with type 2 diabetes.

Results: The age of elderly patients with type 2 diabetes was 73.6±4.3 years., and duration of diabetes was 8.8 ± 3.0 years. In the study, inadequate quality of sleep was reported in 63% elderly patients with type 2 diabetes. This study revealed that lower level of 25(OH) vitamin D was significantly associated with poor sleep quality in type 2 diabetes patients(P < 0.05). Multivariate logistic regression analysis showed that 25(OH) vitamin D deficiency and the use of insulin were risk factors for of inadequate sleep quality in elderly patients with type 2 diabetes.

Conclusions: Elderly type 2 diabetic patients with 25(OH) vitamin D deficiency showed a high incidence of poor sleep quality, low-level of 25(OH) vitamin D was a risk factor for inadequate quality of sleep in elderly patients with type 2 diabetes.

Keywords: type 2 diabetes; sleep quality; 25(OH) vitamin D; elderly

1. Introduction

Type 2 diabetes is a chronic endocrine ailment, its prevalence rate continues to rise because of the change of lifestyle and is as high as 10% China and India¹. Type 2 diabetes is estimated that around 60 million in Europe had type 2 diabetes by 2017 and would cause serious cardiovascular effects². Type 2 diabetes and sleep disorders both are common health problems and affect each other. Persons with diabetes suffered from sleep disorders and took more sleeping medications^{3,4}. According to latest statistics, more than 30% of adults suffer from sleep disorders, and the incidence of sleep disorders in the elderly over 60 years old is up to 40%^{5,6}. Type 2 diabetes has shown to be associated with higher incidence of sleep disorders. Sleep disorders might be caused by the disease itself or by complications of the disease, such as peripheral neuropathy and polyuria.

25(OH) vitamin D is essential for good bone health and deficiency or insufficiency have also been linked to other health concerns, such as diabetes, cardiovascular disease, hypertension, tumor and autoimmune diseases⁷. The prevalence of Vitamin D deficiency has increased in recent decades.^{8,9} In a recent population-based study of Asian adults, approximately 75% had suboptimal Vitamin D concentrations.¹⁰ The Endocrine Society Clinical Practice Guideline defines Vitamin D deficiency as 25(OH)D level <20 ng/mL and insufficiency as 21 to 29 ng/mL.¹¹ Some studies have linked deficient 25(OH) vitamin D to neuropsychological diseases such as sleep disorders¹²⁻¹⁴, which showed that 25(OH) vitamin D deficiency was associated with a higher risk of inadequate quality of sleep. Massa, Jennifer et al. found a significant trend that lower total serum 25(OH) vitamin D was associated with shorter sleep duration, and that men with the lowest total serum 25(OH) vitamin D (< 20.3 ng/mL) were twice as likely to have a shorter sleep duration than those with the highest total serum 25(OH) vitamin D (•40.06 ng/mL). Some researchers believed that the neural systems that control the opposing states of arousal and sleep are predominantly confined to the brainstem, hypothalamus and the thalamus. It has been shown that 25(OH) vitamin D target neurons are thought to be involved in sleep regulation¹⁵.

There was a statistical difference between 25(OH) vitamin D concentration in diabetic patients and the healthy subjects in the study by Mohammad Ali Bayani. et al. The results showed that the mean concentration of healthy women was higher than that 25(OH) vitamin D in diabetic women (27.03±10.28 ng/dl, 19.3±11.9 ng/dl, respectively) (P =0.0001)¹⁶ Therefore, we may believe that 25(OH) vitamin D is important for diabetic women. In addition, a study was been shown that 25(OH) vitamin D affects sleep in people between the ages of 20 and 50. 25(OH) vitamin D supplementation can improve the sleep quality, reduce the sleep latency, prolong the sleep time, and improve the subjective sleep quality of patients with sleep disorders. However, there are still few studies on the effect of 25(OH) vitamin D on sleep quality in elderly patients with type 2 diabetes¹⁷.

This study was designed to investigate the relationship between serum 25(OH) vitamin D levels and sleep quality in elderly patients with type 2 diabetes in China and to fill in the literature gap on this topic.

84 2. Materials and Methods

85 2.1 Patients and definitions

86 A total of 186 inpatients were selected, who were admitted in the Department of
87 Endocrinology and Geriatric Medicine of the First Affiliated Hospital of the University of Science
88 and Technology of China (USTC) from January 2019 to December 2019. All patients were over 60
89 years old and diagnosed with type 2 diabetes according to the 1999 WHO Diabetes Diagnostic
90 Criteria. Exclusion criteria : Patients with type 1 diabetes and other special types of diabetes; with
91 type 2 diabetes-related acute complications; with thyroid and parathyroid function disorders; with
92 cardiac insufficiency, hepatic and renal dysfunction, respiratory failure, malignant tumors, and
93 other severe illness; with incorporative psychiatric illnesses, mental retardation and cognitive
94 deficit; being treated with glucocorticoids, 25(OH) vitamin D, calcium and other drugs that may
95 affect 25(OH) vitamin D levels in the past 3 months. Patient informed consents were obtained. All
96 the data received Institutional Review Board (IRB) approval by the Ethics Committee.

98 2.2 Study design and Method

99 2.2.1 Clinical data collection

100 The age, blood pressure, height, body weight, the duration of type 2 diabetes patients,
101 medications (including oral hypoglycemic drugs and insulin) of all patients were recorded. Body
102 mass index (BMI) was calculated according in kg/m^2 . All patients were subjected for fundus
103 examination to identify diabetic retinopathy after mydriasis, excluding other causes of retinopathy
104 and physical examination and nerve conduction velocity (NCV) examination to identify diabetic
105 peripheral neuropathy, excluding other neurological diseases and other causes of peripheral
106 neuropathy. Venous fasting blood was sampled including fasting blood glucose (FBG), glycosylated
107 hemoglobin (HbA1c), serum creatinine (Cr), total cholesterol (TC), triacylglycerol (TG) and
108 low-density lipoprotein (LDL-C) were measured. Urine albumin was assayed by
109 immunoturbidimetry, and urine creatinine concentration was determined by benedict-behre
110 creatinine colorimetry urine albumin/creatinine ratio (UACR) was calculated, and diabetic
111 nephropathy was defined by an UACR of $\geq 30\text{mg/g}$.

112 2.2.2 Sleep quality assessment

113 The Pittsburgh Sleep Quality Index (PSQI) was originally developed in 1989 to measure sleep
114 quality for psychiatric research as well as clinical practice¹⁸. The index has since been used to assess
115 sleep quality in patients suffering T2DM¹⁹. The PSQI scale was used to evaluate the sleep status of
116 all subjects in the past 4 weeks. The questionnaire included 19 self-assessments and 5 other
117 assessment items. Higher scores indicated worse sleep quality. The investigator made use of
118 questionnaire adjustment method. The adjustment tables were independently completed by the
119 study subjects. Those who could not be accomplished independently due to visual impairment, low
120 cultural level, etc., were asked by the investigator according to the content of the scale and the
121 answer results were truthfully recorded. Poor sleep quality was defined by a total PSQI score²⁰ of
122 >7 .

123

124 2.3 Statistical analyses

125 Normal distribution was determined by the Kolmogorov-Smirnov test, and the data were
 126 expressed as mean $\pm$ standard deviation. Group comparison was performed by t test. The data that
 127 did not meet the normal distribution was compared by the Mann-Whitney U test. Count data were
 128 reported in rates, and comparisons between groups were performed using the χ^2 test. Univariate
 129 logistic regression analysis was used to screen variables for selected research factors. Univariate
 130 analysis was performed with the presence of sleep disturbance as the dependent variable and
 131 gender, age, body mass index (BMI), duration of diabetes, systolic blood pressure, diastolic blood
 132 pressure, FBG, HbA1c, TG, TC, LDL-C, Cr, 25(OH) vitamin D, diabetic nephropathy, diabetic
 133 peripheral neuropathy, diabetic retinopathy, and insulin use as the independent variables.
 134 Multivariate logistic regression analysis was used to explore the influencing factors of sleep
 135 disorder in elderly patients with type 2 diabetes. All tests were two-tailed test. A P value of <0.05
 136 indicated statistically significant. SPSS software version 22.0 (SPSS Inc. Chicago, IL, USA,) was used
 137 for statistical analysis of the data.

138

139 3. Results

140 3.1 Baseline characteristics of two groups according to sleep quality in the elderly patients with type 2 141 diabetes

142 186 type 2 diabetes elderly patients were recruited in the study with average age (73.6 ± 4.3)
 143 years and the average duration of diabetes (8.8 ± 3.0) years. There were 68 cases with diabetic
 144 nephropathy, 62 cases with diabetic retinopathy, and 50 cases with diabetic peripheral neuropathy;
 145 100 cases were treated with insulin. According to the total PSQI score, 186 elderly patients with
 146 type 2 diabetes were divided into a normal sleep group (PSQI ≤ 7) and an inadequate sleep quality
 147 group (total PSQI score > 7). 117 cases presented with poor sleep quality, accounting for 63% of all
 148 subjects under study. Compared with the normal sleep group, the inadequate sleep quality group
 149 had lower 25(OH) vitamin D level, higher HbA1c level, and a higher incidence of diabetic
 150 nephropathy and more insulin use ($P < 0.05$). The differences in gender, age, BMI, duration of
 151 diabetes, systolic blood pressure, diastolic blood pressure, FBG, TG, TC, LDL-C, and Cr levels in the
 152 two groups were not statistically significant (all $P > 0.05$). The results were showed in **Table 1**.

Table 1 Baseline characteristics of two groups according to sleep quality in the elderly patients with type 2 diabetes

	Normal sleep group (n=69)	Inadequate sleep group (n=117)	t (• ²)	P
Sex (male/female)	46/23	76/41	0.056	0.813
Age (years)	73.0±4.1	73.9±4.3	-1.295	0.197
BMI (kg/m ²)	21.5±2.9	21.3±3.4	0.425	0.672
Duration of diabetes (years)	8.6±2.8	8.9±3.1	-0.623	0.534
Systolic pressure (mmHg)	129.5±8.2	131.9±8.2	-1.931	0.055
Diastolic pressure (mmHg)	74.1±7.8	76.0±7.9	-1.514	0.132
FBG (mmol/L)	7.24±1.9	7.52±2.35	-0.855	0.393
HbA1c (%)	6.43±2.6	7.35±2.69	-2.274	0.024
TG (mmol/L)	1.53±0.7	1.67±0.67	-1.308	0.192
TC (mmol/L)	4.54±1.3	4.92±1.68	-1.645	0.102
LDL-C (mmol/L)	3.87±0.9	4.14±0.96	-1.824	0.07
Cr (•mol/L)	88.77±23.1	92.42±29.2	-0.888	0.376
25(OH) vitamin D (ng/mL)	20.674±2.6	13.131±2.9	18.004	<0.01
Diabetic nephropathy (%)	18 (26.1)	50 (42.7)	5.187	0.023
Diabetic peripheral neuropathy (%)	16 (23.2)	34 (29.1)	0.761	0.383
Diabetic retinopathy (%)	20 (29.0)	42 (35.9)	0.933	0.334
Insulin use (%)	25 (36.2)	75 (64.1)	13.563	<0.01

3.2 Comparison of sleep quality in different 25(OH) vitamin D level groups in elderly patients with type 2 diabetes

There were 141 cases presented with 25(OH) vitamin D deficiency, accounting for 76% of the 186 elderly patients with type 2 diabetes recruited in this study. The average PSQI total score of the 25(OH) vitamin D-deficient group was higher than that of the 25(OH) vitamin D -nondeficient group, and the difference was statistically significant (P <0.05); The scores of 25(OH) vitamin D deficient group were higher than those of normal 25(OH) vitamin D group (P <0.05) in 7 dimensions of subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleeping medication and daytime dysfunction (**Table 2**).

Table 2 Comparison of sleep quality in two groups, which were divided based on vitamin D level in the elderly patients with type 2 diabetes

	25(OH) vitamin D -nondeficient group (n=45)	25(OH) vitamin D -deficient group (n=141)	Z (•2)	P
Sex (male/female)	31/14	91/50	0.286	0.593
PSQI total score	6.69±1.4	10.6±1.9	-9.190	<0.01
Subjective sleep quality	1.47±0.84	1.81±0.88	-2.329	0.02
Sleep latency	0.96±0.77	1.54±0.94	-3.664	<0.01
Sleep duration	0.98±0.15	1.28±0.83	-2.565	<0.01
Habitual sleep efficiency	0.91±0.56	1.46±0.64	-4.903	<0.01
Sleep disturbances	0.89±0.32	1.65±0.49	-7.856	<0.01
Use of sleeping medication	0.38±0.49	0.70±0.46	-3.900	<0.01
Daytime dysfunction	0.86±0.34	2.15±0.90	-7.995	<0.01

3.3 Logistic regression analysis of risk factors for sleep disorders in elderly patients with type 2 diabetes

The results showed diabetic nephropathy, HbA1c, 25(OH) vitamin D, use of insulin were related to the occurrence of inadequate quality of sleep in patients. Multivariate logistic regression analysis of the above indicators suggested that 25(OH) vitamin D deficiency and the use of insulin were risk factors for of inadequate sleep quality in elderly patients with type 2 diabetes. (Table 3)

Table 3. logistic analysis of sleep quality risk factors in elderly patients with type 2 diabetes

Risk factors	Exp(B)	95% CI	P
25(OH) vitamin D	0.21	0.11- 0.41	<0.01
Use of Insulin *	28.68	2.54-323.54	<0.01

* Use of insulin , no = 0 , yes = 1

4. Discussion

In this study, we revealed the correlation between serum 25(OH) vitamin D deficiency and sleep quality in elderly patients with type 2 diabetes.

Sleep plays an important role in human health, and sleep disorders is closely related to diabetes²¹. In recent years, sleep disorders have been taken into account as an important risk factor for the development of type 2 diabetes in the elderly. Sleep disorders and depression are more prevalent in people with type 2 diabetes. Van Dijk et al. reported in the study that the probability of increased nocturnal enuresis, pain, dythermesthesia, and habitual snoring in diabetic populations were significantly increased²². Epidemiological studies found that the incidence of sleep disorders in people with type 2 diabetes was 45% to 55%^{23 24}. This study showed the prevalence of inadequate sleep quality, which is only a subset of all various sleep disorders, was high up to 63% in elderly patients with type 2 diabetes. This indicated the incidence of sleep disorders would be significantly higher in elderly patients among all type 2 diabetic patients⁵.

Inadequate sleep quality could cause sympathetic nerve excitement, increase the secretion of antagonist hormone of insulin such as nighttime growth hormone, glucagon, catecholamine, and cortisol, disturb glucose homeostasis; and increase insulin resistance²⁵. One has shown One study also showed that long-term sleep deprivation could reduce insulin secretion in the body leading to increased blood glucose levels in patients with diabetes²⁶. This study also found that HbA1c levels in diabetic patients with sleep disorders were higher than those in the normal sleep group, indicating that blood glucose in the sleep disorder group was more difficult to control.

Since serum 25(OH) vitamin D concentration is affected by many factors, including sun exposure, lifestyle, and skin tone, etc. 25(OH) vitamin D deficiency has become a worldwide problem and the prevalence is rising²⁷. Bayani et al. found that compared with healthy controls, serum 25(OH) vitamin D levels were significantly lower in patients with type 2 diabetes, and the decline was more pronounced in women²⁸. This study also showed that 25(OH) vitamin D deficiency was prevalent in elderly people with type 2 diabetes, accounting for 76% of the study population. The potential causes of low 25(OH) vitamin D in elderly patients could be less outdoor activities, shorter sunshine hours, and reduced gastrointestinal digestion and absorption in elderly patients.

The data in this study show that in elderly patients with type 2 diabetes, the serum 25(OH) vitamin D level in the sleep disorders group was lower than that in the normal sleep group, and the insulin use rate was higher than in the normal sleep group. The average total PSQI score of the 25(OH) vitamin D -deficient group was higher than those of the normal vitamin D group, and the 25(OH) vitamin D -deficient group scored higher in 7 dimensions of subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleeping medication and daytime dysfunction. Further logistic regression analysis showed that low level of 25(OH) vitamin D and more insulin use were associated with sleep disturbance in elderly patients with type 2 diabetes. Logistic regression analysis of risk factors for sleep disorders in elderly patients with type 2 diabetes demonstrated that the low-level serum 25(OH) vitamin D was an important risk factor for inadequate quality of sleep in elderly patients with type 2 diabetes.

Knhanes result showed that lower serum 25(OH) vitamin D levels were linked to a significantly higher risk of short sleep duration^{29,30}. McCarty et al. found that patients with 25(OH) vitamin D deficiency scored lower on the Epworth sleepiness scale (ESSs)³¹. Another study measured objective sleep duration and sleep effectiveness of men over the age of 65 and found that lower serum 25(OH) vitamin D levels were associated with shorter sleep duration and poorer sleep effectiveness³². These results mentioned above were consistent with our data.

So far, the mechanisms for a link between sleep duration and 25(OH) vitamin D were not clearly understood. Vitamin D receptors (VDR) were shown to be present in nearly all tissues of the body, including many parts of the human brain, such as the hypothalamus, prefrontal cortex, midbrain central gray, and substantia nigra, etc., which all play an important role in sleep regulation³²⁻³⁴.

The classical clinical presentation of Vitamin D deficiency is bone pain. Increased pain could be associated with sleep disturbances³⁵. Epidemiologic evidence for impact of 25(OH) vitamin D supplements on sleep quality was emerging. One study showed that people's pain levels, sleep quality, and various aspects of Quality of Life (QoL) can be significantly improved after standardized vitamin D supplementation³⁶. Another double-blind clinical trial showed improvements and increases in sleep quality and duration in patients with sleep disorders after eight weeks of continuous supplementation with 25(OH) vitamin D (50,000 IU/ 2 weeks)¹⁷.

As far as we know, this was the first study to analyze the relationship between serum 25(OH) vitamin D concentration and sleep quality in elderly patients with type 2 diabetes. The significance of this study is that the concentration of 25(OH) vitamin D can be used to identify sleep disorders in elderly patients with type 2 diabetes. These findings have positive implications for the prevention and treatment of inadequate sleep quality in elderly patients with type 2 diabetes. One limitation of

this study was that the dietary habits and psychological state of patients were not assessed, which, may also be related to sleep quality. Further, because of the character of the cross-sectional study, we could not conclude the causal relationship among 25(OH) vitamin D and sleep quality. Therefore, further studies with larger scale cohort studies and well-designed randomized controlled trials would be needed to verify this relationship and reveal possible mechanisms.

5 Conclusion

In conclusion, this study confirms that 25(OH) vitamin D deficiency was more common in elderly type 2 diabetic patients. There was a clear correlation between vitamin D deficiency and sleep quality in this population. This initial work warrants further larger scale studies of ethnic 25(OH) vitamin D variations and potential causal interrelationship between sleep quality, insulin usage and 25(OH) vitamin D deficiency state among elderly type 2 diabetic patients. Further studies are required to investigate possible biological mechanisms of the association between 25(OH) vitamin D deficiency, insulin usage and sleep quality, and to discover whether effective interventions for 25(OH) vitamin D supplementation could improve sleeping quality and effectiveness of insulin treatment in decrease the risk of geriatric syndromes in elders with type 2 diabetes.

Author Contributions: Conceptualization, Hongyan Ma, Lin Zhang; Methodology, Hongyan Ma, Lin Zhang; Writing – original draft, Hongyan Ma, Dani Fan, Lin Zhang; Writing – review & editing, Hongyan Ma, Dani Fan, Xiaofei Li, Yang Zhao, Lin Zhang. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: This study was supported by the First Affiliated Hospital of University of Science and Technology of China for data access.

Conflicts of Interest: The authors declare no conflict of interest

Ethical approval: Informed consents of patients were obtained for diagnosis and treatment, and the study protocol was approved by the First Affiliated Hospital of University of Science and Technology of China. All the data received Institutional Review Board (IRB) approval by the Ethics Committee.

References

1. Gwatidzo SD, Stewart Williams J. Diabetes mellitus medication use and catastrophic healthcare expenditure among adults aged 50+ years in China and India: results from the WHO study on global AGEing and adult health (SAGE). *BMC Geriatrics* 2017;17(1) doi: 10.1186/s12877-016-0408-x
2. Cho NH, Shaw JE, Karuranga S, et al. IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Research & Clinical Practice* 2018;271.
3. Skomro RP, Ludwig S, Salamon E, et al. Sleep complaints and restless legs syndrome in adult type 2 diabetics. *Sleep Medicine* 2001;2(5):417-22.

- 277 4. Association AD. 3. Comprehensive Medical Evaluation and Assessment of Comorbidities: *Standards of*
278 *Medical Care in Diabetes—2018*. *Diabetes Care* 2018;41(Suppl 1):S28.
- 279 5. Vittoria E, Sisti D, Pascucci P, et al. [Prevalence of Obstructive Sleep Apnea Syndrome among holders of a
280 category B driver's license and among professionals in the Province of Pesaro-Urbino (Italy)]. *Igiene E*
281 *Sanita Pubblica* 2017;73(6):605.
- 282 6. Owens, J. Insufficient Sleep in Adolescents and Young Adults: An Update on Causes and Consequences.
283 *Pediatrics* 2014;134(3):e921.
- 284 7. Mayte M, Estrella CC, Isabel M, et al. Vitamin D: Effect on Haematopoiesis and Immune System and Clinical
285 Applications. *International Journal of Molecular Sciences* 2018;19(9):2663-.
- 286 8. Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *The American*
287 *journal of clinical nutrition* 2008;87(4):1080S-6S. doi: 10.1093/ajcn/87.4.1080S [published Online First:
288 2008/04/11]
- 289 9. Hilger J, Friedel A, Herr R, et al. A systematic review of vitamin D status in populations worldwide. *Br J*
290 *Nutr* 2014;111(1):23-45. doi: 10.1017/S0007114513001840 [published Online First: 2013/08/13]
- 291 10. Man RE, Li LJ, Cheng CY, et al. Prevalence and Determinants of Suboptimal Vitamin D Levels in a
292 Multiethnic Asian Population. *Nutrients* 2017;9(3) doi: 10.3390/nu9030313 [published Online First:
293 2017/03/23]
- 294 11. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention of vitamin D
295 deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2011;96(7):1911-30.
296 doi: 10.1210/jc.2011-0385 [published Online First: 2011/06/08]
- 297 12. Ewelina, ?ukaszzyk, Katarzyna, et al. Cognitive Functioning of Geriatric Patients: Is Hypovitaminosis D the
298 Next Marker of Cognitive Dysfunction and Dementia? *Nutrients* 2018
- 299 13. Hiller AL, Murchison CF, Lobb BM, et al. A randomized, controlled pilot study of the effects of vitamin D
300 supplementation on balance in Parkinson's disease: Does age matter? *PLoS ONE* 2018;13(9)
- 301 14. Pierrot-Deseilligny C, Souberbielle JC. Vitamin D and multiple sclerosis: An update. *Multiple Sclerosis &*
302 *Related Disorders* 2017;14:35.
- 303 15. Gominak SC, Stumpf WE. The world epidemic of sleep disorders is linked to vitamin D deficiency. *Medical*
304 *Hypotheses* 2012;79(2):132-35. doi: 10.1016/j.mehy.2012.03.031
- 305 16. Bayani. MA, . RA, . BB, et al. Status of Vitamin-D in diabetic patients. 2014;5(1):3. [published Online First:
306 42]
- 307 17. Majid MS, Ahmad HS, Bizhan H, et al. The effect of vitamin D supplement on the score and quality of sleep
308 in 20-50 year-old people with sleep disorders compared with control group. *Nutr Neurosci*
309 2018;21(7):511-19. doi: 10.1080/1028415X.2017.1317395
- 310 18. Buysse DJ, Reynolds CF, 3rd, Monk TH, et al. The Pittsburgh Sleep Quality Index: a new instrument for
311 psychiatric practice and research. *Psychiatry research* 1989;28(2):193-213. doi:
312 10.1016/0165-1781(89)90047-4 [published Online First: 1989/05/01]
- 313 19. Telford O, Diamantidis CJ, Bosworth HB, et al. The relationship between Pittsburgh Sleep Quality Index
314 subscales and diabetes control. *Chronic Illn* 2019;15(3):210-19. doi: 10.1177/1742395318759587
315 [published Online First: 2018/02/23]
- 316 20. Liu HX, Lin J, Lin XH, et al. Quality of sleep and health-related quality of life in renal transplant recipients.
317 *International Journal of Clinical & Experimental Medicine* 2015;8(9):16191-98.

- 318 21. Buxton OM, Marcelli E. Short and long sleep are positively associated with obesity, diabetes, hypertension,
319 and cardiovascular disease among adults in the United States. *Social ence & Medicine*
320 2010;71(5):1027-36.
- 321 22. Dijk MV, Donga E, Dijk JGV, et al. Disturbed subjective sleep characteristics in adult patients with
322 long-standing Type 1 diabetes mellitus. *Diabetologia* 2011;54(8):1967-76.
- 323 23. Luyster FS, Dunbar-Jacob J. Sleep Quality and Quality of Life in Adults With Type 2 Diabetes. *Diabetes*
324 *Educator* 2011;37(3):347-55.
- 325 24. Amarabalan, Rajendran, Shruthi, et al. Prevalence and correlates of disordered sleep in southeast asian
326 indians with type 2 diabetes. *Diabetes & metabolism journal* 2012
- 327 25. P CF, Lanfranco DE, Pasquale S, et al. Quantity and quality of sleep and incidence of type 2 diabetes: a
328 systematic review and meta-analysis.[J]. *Diabetes care*,2010,33(2). *Diabetes care* 2010;33(2) doi:
329 10.2337/dc09-1124
- 330 26. Darukhanavala A, Booth JN, Bromley L, et al. Changes in Insulin Secretion and Action in Adults With
331 Familial Risk for Type 2 Diabetes Who Curtail Their Sleep. *Diabetes Care* 2011;34(10):2259-64.
- 332 27. Norman AW, Bouillon R. Vitamin D nutritional policy needs a vision for the future. *Experimental Biology &*
333 *Medicine* 2010;235(9):1034-45.
- 334 28. Bayani MA, Akbari R, Banasaz B, et al. Status of Vitamin-D in diabetic patients. *Caspian J Intern Med*
335 2014;5(1):3. [published Online First: 42]
- 336 29. Lee HJ, Lee HD, Hwang HS, et al. Associations between Serum Vitamin D Level and Mean Hours of Sleep:
337 2011Korea National Health and Nutrition Examination Survey. *Metabolism-Clinical and Experimental*
338 2014;7(8):6346-61.
- 339 30. Kim JH, Chang JH, Kim DY, et al. Association between self-reported sleep duration and serum vitamin D
340 level in elderly Korean adults. *Journal of the American Geriatrics Society* 2014;62(12):2327-32.
- 341 31. Mccarty DE, Reddy A, Keigley Q, et al. Vitamin D, Race, and Excessive Daytime Sleepiness. *Journal of*
342 *Clinical Sleep Medicine Jcsm Official Publication of the American Academy of Sleep Medicine* 2012;8(6):693-7.
- 343 32. Massa J, Stone KL, Wei EK, et al. Vitamin D and Actigraphic Sleep Outcomes in Older
344 Community-Dwelling Men: The MrOS Sleep Study. *Sleep* 2015;38(2):251-57. doi: 10.5665/sleep.4408
- 345 33. A EG, B WB, A MM, et al. New clues about vitamin D functions in the nervous system. *Trends in*
346 *Endocrinology & Metabolism* 2002;13(3):100-05.
- 347 34. Musiol IM, Stumpf WE, Bidmon HJ, et al. Vitamin D nuclear binding to neurons of the septal, substriatal
348 and amygdaloid area in the Siberian hamster (*Phodopus sungorus*) brain. *Neuroence* 1992;48(4):841-48.
- 349 35. Monika H, Elsa S, Mullington JM. Elevated inflammatory markers in response to prolonged sleep
350 restriction are associated with increased pain experience in healthy volunteers. *Sleep* (9):1145.
- 351 36. Huang W, Shah S, Long Q, et al. Improvement of Pain, Sleep, and Quality of Life in Chronic Pain Patients
352 With Vitamin D Supplementation. *Clinical Journal of Pain* 2013;29(4):341-47.
- 353