

The impact of host resistance on cumulative mortality and the threshold of herd immunity for SARS-CoV-2

José Lourenço ^a, Francesco Pinotti ^a, Craig Thompson ^a & Sunetra Gupta ^{a*}

Affiliations: ^a Department of Zoology, University of Oxford, Oxford, United Kingdom;

Correspondence: * sunetra.gupta@zoo.ox.ac.uk

Supplementary Figures & Tables

	Proportionate mixing ($\rho = \delta$)					Assortative mixing ($\delta = 1$)				
	$R_0 = 1.25$	$R_0 = 1.5$	$R_0 = 2$	$R_0 = 2.5$	$R_0 = 3$	$R_0 = 1.25$	$R_0 = 1.5$	$R_0 = 2$	$R_0 = 2.5$	$R_0 = 3$
$\rho = 0$	0.2	0.33	0.5	0.6	0.66	0.2	0.33	0.5	0.6	0.66
$\rho = 0.1$	0.1	0.23	0.4	0.5	0.56	0.18	0.3	0.45	0.54	0.6
$\rho = 0.2$	0	0.13	0.3	0.4	0.46	0.16	0.26	0.4	0.48	0.53
$\rho = 0.3$	0	0.03	0.2	0.3	0.36	0.14	0.23	0.35	0.42	0.46
$\rho = 0.4$	0	0	0.1	0.2	0.26	0.12	0.2	0.3	0.36	0.4
$\rho = 0.5$	0	0	0	0.1	0.16	0.1	0.16	0.25	0.3	0.33

Table S1 - Herd-immunity threshold (HIT). Model output for proportionate mixing ($\rho = \delta$, left, light grey) and assortative mixing ($\delta = 1$, right, white) for varying basic reproduction number R_0 , proportion resistant ρ , and $R_{0_1} = 0$, $R_{0_2} = R_0$.




Figure S1 - Herd-immunity threshold under proportionate mixing and incomplete resistance. (left) Herd-immunity threshold (z^*) under proportionate mixing ($\delta = \rho$) and incomplete resistance $F = R_{0_1}/R_{0_2}$ with $R_0 = 1.5$. (right) Same output as in the left panel but for with $R_0 = 2.5$. The scenario of $F = 0$ is fully explored in the main text. Simulations ran for 365 days with an infectious period ($1/\sigma$) of 5 days, $\gamma = 0$, $\delta = \rho$.




Figure S2 - Effect of duration of immunity. Time series of proportion infected (y) for varying proportion of resistant (ρ) and duration of immunity ($1/\gamma$, color scale) under proportionate mixing ($\delta = \rho$). Proportion of resistant (ρ) for each panel is presented on the title. Infectious period $1/\sigma = 5$ days.