A systematic review of the knowledge, attitudes, and practices of physicians, health workers, and the general population about Coronavirus disease 2019 (COVID-19)

Saeede Saadatjoo¹, ORCID: 0000-0003-4163-4299, ss.saadatjoo1360@gmail.com

Maryam Miri¹, ORCID: 0000-0002-8125-5093, mmiri1062@gmail.com

Soheil Hassanipour², ORCID: 0000-0002-6661-4908, soheil.epid@gmail.com

Hosein Ameri³, ORCID: 0000-0002-9914-034X, hamery7@yahoo.com

Morteza Arab-Zozani¹, ORCID: 0000-0001-7223-6707, arab.hta@gmail.com

1. Social Determinants of Health Research Center, Birjand University of Medical sciences, Birjand, Iran.
2. Gastrointestinal and Liver Diseases Research Center, Guilan University of Medical Sciences, Rasht, Iran.
3. Health Policy and Management Research Center, School of Public Health, Shahid Sadoughi University of Medical Sciences, Yazd, Iran

Corresponding author: Morteza Arab-Zozani

Email: arab.hta@gmail.com

ORCID: 0000-0001-7223-6707

Mobile: +989153317843

Fax: +985631631651

Tel: +985632381272

Address: South Khorasan, Birjand, Moallem Street, Birjand University of Medical Sciences, School of Health, Social Determinants of Health Research Center (SDHRC), P.Box: 32430076.

Data availability statement: All relevant data are with the paper and the attached supplementary information.

Funding: The author(s) received no specific funding for this work.

Competing Interest: The authors have declared that no competing interests exist.

A systematic review of the knowledge, attitudes, and practices of physicians, health workers, and the general population about Coronavirus disease 2019 (COVID-19)

Abstract

Background: Understanding people, physicians, and healthcare workers' knowledge, attitude, and practices (KAPs) can help to achieve the outcomes of planned behavior. The aim of this study was to investigate and synthesize the current evidence on KAPs regarding COVID-19.

Methods: We conducted a systematic search on PubMed/LitCovid, Scopus, and Web of Sciences databases for papers in the English language only, up to 1 Jul 2020. We used the Joanna Briggs Institute (JBI) checklist developed for cross-sectional studies to appraise the quality of the included studies. All stages of the review conducted by two independent reviewers and potential discrepancies solved with a consultation with a third reviewer. We reported the result as number and percentage. PROSPERO registration code: (CRD42020186755).

Results: Fifty-two studies encompassing 49786 participants were included in this review. 45.76% of the participants were male. The mean age of the participants was 32.6 years. 44.2% of the included studies were scored as good quality, 46.2% as fair quality, and remaining (9.6%) as low quality. 30.76% examined all three components of the KAPs model. The knowledge component was reported as good, fair, and poor in 59%, 34%, and 7%, respectively. Of the studies that examined the attitude component, 82% reported a positive attitude, 11% a fairly positive attitude, and 7% a negative attitude. For the practice component, 52% reported good practice, 44% fair practice, and 4% poor practice.

Conclusion: This systematic review showed that the overall KAP components in the included studies were at an acceptable level. In general, knowledge was at a good level, the attitude was

positive and practice was at a fairly good level. Using an integrated international system can help better evaluate these components and compare them between countries.

Keywords

COVID-19; Knowledge; Attitude; Practice; Systematic review

Introduction

Coronavirus disease 2019 (COVID-19) was reported on 31st December 2019 from Wuhan, China, and announced by the World Health Organization (WHO) as a pandemic on 11th March 2020 [1, 2]. To date, it was estimated that about 30 million people were infected with COVID-19 worldwide, of which about one million have died [3].

COVID-19 is characterized by a number of flu-like symptoms including fever, respiratory problems (dry cough, shortness of breath or difficulty breathing, sore throat), chills, headache, and loss of taste. Also, this disease is much more severe with men, higher age groups, and patients with other pre-existing conditions, such as cardiovascular disease, chronic respiratory disease, diabetes, and hypertension [4, 5]. Based on existing evidence, about 81% of COVID-19 cases are mild, 14% are severe and 5% is critical. The median time from symptoms onset to clinical recovery is approximately two weeks for mild cases and three to six weeks for severe or critical cases [6]. The incubation period for this disease was reported as 2-14 days based on WHO reports. The mortality rate for this disease is different among countries and was reported between two% and 5% [7, 8]. The most important ways to prevent this disease are to use a mask and maintain social distance [9-11]. So far, there have been several cases of infection in the general public, especially doctors and medical staff, some of which have led to death [12-14].

Considering the extent and progress of COVID-19 disease and its major effects on economic, social, political, and cultural dimensions of all countries [15, 16], it is essential that people with COVID-19 are provoked, informed, and engaged in all aspects of the disease. From the onset of the disease until now, various studies conducted worldwide have investigated on this disease and some of these studies have examined the knowledge, attitudes, and practices (KAPs) of people

with COVID-19. Having enough knowledge about a disease can always affect people's attitudes and practices, and on the other hand, improper attitudes and practices can increase the risk of disease and death. Therefore, understanding people, physicians, and healthcare workers' KAPs and knowing potential risk factors can help to achieve the outcomes of planned behavior [17, 18].

Given the importance of the issue, conducting a review of studies that have examined the KAPs of individuals and summarizing the results can provide solid evidence for decision-makers in all countries to better manage the disease. Thus, this study aimed at conducting a systematic review to synthesize current evidence on KAPs of people with COVID-19 worldwide.

Materials and Methods

Protocol and registration

We conducted a systematic review of the existing evidence related to KAPs of COVID-19 patients worldwide following Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) statements (Appendix Supplementary file 1) [19]. We also registered a protocol for this systematic review in the International Prospective Register of Systematic Reviews [20] (CRD42020186755).

Eligibility criteria

We included all studies which met the following inclusion criteria: 1) cross-sectional survey; 2) investigate at least one component of the KAPs model regarding COVID-19 patients worldwide; 3) published, in-press or preprint original paper; 4) in English; 5) with a sample of the general population, physicians or other healthcare workers. No restrictions were applied to the setting, time, or quality of the study.

Information sources, search and study selection

We search the PubMed/LitCovid, Scopus, and Web of Sciences for papers in the English language only, up to 1 Jul 2020 (MA-Z). We also conducted a search in Google Scholar for retrieving studies that were not cited in the above-mentioned databases. In addition, the reference lists to final articles were hand-searched. The keywords used in the search were: attitude, knowledge, practice, awareness, perception, action, COVID-19, coronavirus disease, SARS-CoV-2, and severe acute respiratory syndrome coronavirus 2. The full search strategy for the Scopus database is provided in Appendix Supplementary file 2. When the search was complete, all records were transferred to the Endnote software (V. X8; Clarivate Analytics, Philadelphia, PA) and duplicates were removed. Then, studies based on the title, abstract, and full text were screened by two researchers independently by considering the pre-specified eligibility criteria (MM and SS). Disagreements were solved through consultation with a third researcher (MA-Z).

Data collection process and data item

Two researchers independently engaged in the data collection process and extracted data including author, year, journal name, location, study design, data collection tools, sample size, focusing group, mean age or range, gender percent, and result related to KAPs model components (MM and SS). Potential disagreements were solved through consultation with a third researcher (MA-Z).

Quality appraisal

Included studies were critically appraised by two researchers independently (MM and SS). We used the Joanna Briggs Institute (JBI) checklist developed for cross-sectional studies to appraise the quality of the included studies [21]. This checklist contains eight simple and clear questions

that cover topics such as inclusion criteria for sample; details about study subjects and setting; validity and reliability; criteria for measurement of the condition; confounding variables; and statistical analysis [22]. The answer to each questions is yes, no, unclear, and not applicable. Potential discrepancies were resolved by consultation with a third researcher (SH).

Synthesis of results

Due to potential heterogeneity between included studies the meta-analysis was not conducted. Therefore we carried out the descriptive analysis in most sections and report the pooled data as number or percentage for similar data items. We used Microsoft Excel software to design the charts. Also, we report the result of the included studies in a narrative manner.

Results

Study selection

A total of 4829 records were retrieved from our database search. After removing duplicate, 3085 records were screened by title, abstract, and full-text based on eligibility criteria, of which fifty-two studies were included in the final review [23-74]. The PRISMA flow diagram for the complete study selection process is presented in Fig 1.

Figure 1: PRISMA flow diagram

Study characteristics

Fifty-two studies encompassing 49786 participants were included. Also, 45.76% of the participants were male. The mean age of the participants was 32.6 years. Most studies were from Asia, Africa, South America, Europe, and multinational, respectively (Fig 2A). The most important method of data collection was online questionnaires (Fig 2B). Focusing groups were

diverse between the studies and included health care workers, students, general population, adult population, patients groups, residents, and pregnant women (Fig 2C). Most studies examined all three components of the KAPs model, but some studies examined two components or one component. More details about the characteristics of included studies are presented in Table 1.

Figure 2: The percentage of the included studies based on location (A), data collection methods (B), and focusing group (C).

Table 1: Summary characteristics of the included studies

Reference (Author, Year)	Journal	Location	Study Design	Data Collection tool	Sample Size	Focusing Group	Male (%)	Mean Age or range
Abdelhafiz et al, 2020 [23]	Journal of community health	Egypt	Cross-Sectional	Online questionnaire/Google Form	384	Egyptian Adults	37.7	18-60+
Al Nsour et al, 2020 [24]	JMIR Medical Education/preprints	Jordan, Sudan, and Yemen	Cross-Sectional	Online Questionnaire/Email	57	Epidemiology Training Program graduates	70	39.2
Alhaj et al, 2020 [25]	World neurosurgery	Multinational	Cross-Sectional	Questionnaire/on site	52	Neurosurgery Residents	73.1	NR*
Khan et al, 2020 [46]	Ophthalmology retina	USA	Cross-Sectional	Online questionnaire/SurveyMonkey	87	Ophthalmology Trainees	NR	NR
Alzoubi et al, 2020 [26]	Journal of pure and applied microbiology	Jordan	Cross-Sectional	Online self-administered questionnaire/Facebook	592	Undergraduate Students	34.5	NR
Escalera-Antezana et al, 2020 [36]	Travel medicine and infectious disease	Bolivia and Colombia	Cross-Sectional	Questionnaire/On site, before lockdowns; and online, thereafter	1165	Healthcare workers and students	61.9	33.5
Apaijitt and Wiwanitkit 2020 [27]	Infection control & hospital epidemiology	Thailand	NR	Questionnaire/on site	124	Medical Personnel	33.87	36.7
Azlan et al, 2020 [28]	PLOS one	Malaysia	Cross-Sectional	Online questionnaire/Survey Monkey	4850	General population	42.1	34
Basheti et al, 2020 [29]	Research in social and administrative pharmacy	Jordan	Cross-Sectional	Online questionnaire / Facebook and WhatsApp	726	Pharmacists and Pharmacy Students	28.1	26.9
Bhagavathula et al, 2020 [30]	JIMIR public health and surveillance	Multinational	Cross-Sectional	Web-Based questionnaire/Telegram	529	Health Care Workers	51.6	25-34

Bragazzi et al, 2020 [31]	Dermatologic therapy	Italy	Cross-Sectional	Questionnaire/on site	98	Dermatological Patients	52	44.36
Cai et al, 2020 [32]	Laryngoscope	USA	Cross-Sectional Survey	Online questionnaire/Qualtrics survey platform	82	Otolaryngology Residents	NR	NR
Chesser et al, 2020 [33]	Health education & behavior	USA	Survey	Questionnaire/E mail	1136	University Students	30	27
Clements, 2020 [34]	JMIR public health and surveillance	USA	Cross-Sectional	Online questionnaire/MTurk platform	1034	U.S. Residents	58.2	37.11
Galle et al, 2020 [37]	International journal of environmental research and public health	Italy	Survey	Web-based questionnaire	2125	Undergraduate Students	37.2	22.5
Geldsetzer et al, 2020 [38]	Journal of medical internet research	USA and UK	Cross-Sectional	Online questionnaire / Prolific platform	5974	Adults	US=49.1 UK=48.8	18-58+
Giao et al, 2020 [39]	Asian pacific journal of tropical medicine	Vietnam	Cross-Sectional	Self-Administered Questionnaire/on site	327	Health Care Workers	26	30.1
Liu et al, 2020 [50]	Brain, behavior, and immunity	China	Cross-Sectional	Self-Administered Questionnaire/on site	118	Patients with COVID-19	52	41
Kamate et al, 2020 [42]	Dental and medical problems	Multinational	Survey	Online Questionnaire/Google forms	860	Dental Practitioners	NR	NR
Karasneh et al, 2020 [43]	Research in social and administrative pharmacy	Jordan	Cross-Sectional	Web-Based self-reported questionnaire/social media	486	Pharmacists	21.4	28.1
Kebede et al, 2020 [44]	Plos one	Ethiopia	Cross-Sectional	Questionnaire/on site	247	Medical Center Visitors	76.5	30.5
Erbas et al, 2020 [35]	Sao Paulo medical journal	Turkey	Cross-Sectional	Online questionnaire/Survey Monkey	248	Intensive care physicians	49.1	37.2
Lau et al, 2020 [49]	Journal of global health	Philippines	Cross-Sectional	Online questionnaire/SurveyCTO platform	2224	Income-poor households	7.3	41.3
Abdel Wahed et al, 2020 [70]	Journal of Community Health	Egypt	Cross-Sectional	Online questionnaire/Google Forms /Facebook and WhatsApp	407	Health Care Workers	50.6	34.9
Sari et al, 2020 [65]	Journal of Community Health	Indonesia	Cross-Sectional	Online questionnaire/Google Forms /WhatsApp	201	General population	46.3	35.5
Khasawneh et	Frontiers in	Jordan	Cross-	Online	1404	Medical	40.5	NR

al, 2020 [47]	public health		Sectional	questionnaire/ Google Forms		students		
Khader et al, 2020 [45]	JMIR public health and surveillance	Jordan	Cross-Sectional	Online questionnaire/ Google Forms	368	Dentists	33.4	32.9
Kumar et al, 2020 [48]	Cureus	Pakistan	Cross-Sectional	Semi-structured questionnaire/ on site	392	Health Care Workers	86.98	42.37
McFadde et al, 2020 [51]	PLOS ONE	US	Survey	Online questionnaire/ Qualtrics	718	Adult population	49	18-55+
Modi et al, 2020 [52]	Cureus	India	Survey	Online questionnaire	1562	Health and Students, Allied Health Professionals	24.1	18-45+
Moro et al, 2020 [53]	Acta Biomedica	Italy	Survey	Online questionnaire/ SurveyMonkey	2046	Hospital staff	NR	NR
Olapegba et al, 2020 [55]	Data in brief	Nigeria	Survey	Online questionnaire	1357	General population	58	26.85
Olum et al, 2020 [56]	Frontiers in Public Health	Uganda	Cross sectional	Online questionnaire/ Google Forms	581	Health Care Workers	87	34
Parikh et al, 2020 [57]	Cureus	India	Cross-sectional	Online questionnaire/ social media platforms	1246	General public, healthcare professionals	52.40	30.60
Ricco et al, 2020 [60]	Acta Biomedica	Italy	cross-sectional	Online questionnaire/ Google Forms	561	Medical workforce	42.6	NR
Roy et al, 2020 [61]	Asian Journal of Psychiatry	India	Cross-sectional	Online questionnaire/ Google Forms	662	General population	48.6	29.9
Sahraian et al, 2020 [62]	Multiple Sclerosis and Related Disorders	Iran	Cross-sectional	Online questionnaire/ Google Forms	233	Multiple Sclerosis patients	22.7	34.2
Salman et al, 2020 [63]	Drugs & Therapy Perspectives	Pakistan	Cross-sectional	Self-administered questionnaire/ on site	417	Students and employees at two higher educational institutions	41.2	22.6
Saqlain et al, 2020 [64]	The journal of Hospital Infection	Pakistan	Cross-sectional	Online questionnaire/ Google Forms	414	Healthcare professionals	50.5	NR
Shi et al, 2020 [66]	Brain, Behavior and Immunity	China	Cross-sectional	Online questionnaire	311	Medical staff in psychiatric hospitals	48.2	33.7
Singh et al, 2020 [67]	Journal of Community Health	Nepal	cross-sectional	Online questionnaire	871	Nepalese adults	40.4	26.4
Siniscalchi et al, 2020 [68]	Digestive and Liver Disease	Italy	cross-sectional	E-mail	276	Celiac adult patients	24.3	39
Taghrir et al, 2020 [69]	IRANIAN MEDICINE	Iran	cross-sectional	Online	240	Medical Students	40.8	23.6

				questionnaire				
Nwafor et al, 2020 [54]	Gynecology Obstetrics	Nigeria	cross-sectional	Self-administered questionnaire	284	Pregnant women	NR	<30->40
Wolf et al, 2020 [71]	Annals of Internal Medicine	USA	Cross-sectional	Telephone	630	Adults aged	40.3	23 - 88
Yassa et al, 2020 [72]	Maternal-fetal & neonatal medicine	Turkey	cross-sectional	Telephone	172	Pregnant women	NR	27.5
Zhang et al, 2020 [73]	Journal of Hospital Infection	China	cross-sectional	NR	1357	Healthcare workers	53.4	NR
Zhong et al, 2020 [74]	International Journal of Biological Sciences	China	cross-sectional	Online questionnaire	6910	Residents	34.3	16-50≤
Reuben et al, 2020 [59]	Journal of Community Health	Nigeria	Cross-sectional	Online questionnaire	589	Residents	59.6	18-50
Jemal et al, 2020 [40]	Research Square	Ethiopia	Cross-sectional	Structured questionnaire/on site	422	Healthcare workers	56.7	29.28
Raza et al, 2020 [58]	International Journal of Homeopathy Complementary and Alternative Medicine	Pakistan	Cross-sectional	Questionnaire/on site	150	Patients	36	12-75
Kakemam et al, 2020 [41]	Pre-print MedRxiv	Iran	Cross-sectional	Online questionnaire/Portfolio/Social media	1480	General population	42.8	31.2

160 *NR: not reported

161 Quality appraisal

162 The overall mean quality score of the included studies was 4.90. Of the included studies, 23
163 studies (44.2%) were scored as good quality (score ≥ 6), 24 (46.2%) as fair quality (score 3-5),
164 and remaining (9.6%) as low quality (score < 3) (Fig 3). The lowest and highest quality scores in
165 the studies were two and six, respectively. None of the studies scored on questions 5 and 6,
166 which were related to identification and deal with confounding variables in the studies (for more
167 details about items see Appendix Supplementary file 3.

168 Figure 3: The percentage of included studies based on quality score

Synthesis of results

Among the included studies, 30.76% examined all three components of the KAPs model simultaneously. The most studied component in the studies was the knowledge component with about 84.61%, followed by attitude and practice with 53.84% and 44.23%, respectively. Also, 34.61% and 17.3% of the articles have examined other components including perception and awareness, respectively (Table 2, Fig 4).

Figure 4: The number of investigated components in the included studies

Of the studies that examined the knowledge component, 59% reported good knowledge, 34% fair knowledge, and 7% poor knowledge. As well as, of the studies that examined the attitude component, 82% reported a positive attitude, 11% a fairly positive attitude, and 7% a negative attitude. For the practice component, 44% reported good practice, 52% fair practice, and 4% poor practice. Also, 39% of the perception component reported a high level of perception and for the awareness component, 44% reported good awareness (Table 2, Fig 5).

Figure 5: The percentage of studies based on the knowledge (K), attitudes (A), and practices (P)

Table 2: Results related to Coronavirus- related KAPs components of the included studies

Reference (Author, Year)	KAPs components			Other components	
	Level of Knowledge	Level of Attitudes	Level of Practices	Level of Perception	Level of Awareness
Abdelhafiz et al, 2020 [23]	Good	Positive	NA*	Moderate	NA
Al Nsour et al, 2020 [24]	NA	NA	NA	NA	Good
Alhaj et al, 2020 [25]	Good	NA	NA	NA	NA
Khan et al, 2020 [46]	NA	NA	NA	High	NA
Alzoubi et al, 2020 [26]	Good	Positive	Good	NA	NA
Escalera-Antezana et al, 2020 [36]	Good	NA	NA	NA	NA
Apaijitt and Wiwanitkit 2020 [27]	Fair	NA	NA	NA	NA
Azlan et al, 2020 [28]	Good	Positive	Fair	NA	NA
Basheti et al, 2020 [29]	NA	NA	NA	Moderate	Fair
Bhagavathula et al, 2020 [30]	Poor	NA	NA	High	NA
Bragazzi et al, 2020 [31]	Fair	Fairly positive	NA	NA	NA

Cai et al, 2020 [32]	NA	NA	Good	High	NA
Chesser et al, 2020 [33]	Fair	NA	NA	NA	NA
Clements, 2020 [34]	Good	NA	Fair	NA	NA
Galle et al, 2020 [37]	Good	NA	Fair	NA	Good
Geldsetzer et al, 2020 [38]	Good	NA	NA	Low	NA
Giao et al, 2020 [39]	Good	Positive	NA	NA	NA
Liu et al, 2020 [50]	Poor	NA	NA	Low	NA
Kamate et al, 2020 [42]	Good	Positive	Good	NA	NA
Karasneh et al, 2020 [43]	Fair	NA	NA	High	Fair
Kebede et al, 2020 [44]	Fair	NA	Fair	Moderate	NA
Erbas et al, 2020 [35]	Fair	Fairly positive	NA	NA	NA
Lau et al, 2020 [49]	Good	Positive	Good	NA	NA
Abdel Wahed et al, 2020 [70]	Good	Positive	NA	High	NA
Sari et al, 2020 [65]	Good	Positive	NA	NA	NA
Khasawneh et al, 2020 [47]	Good	Positive	Good	NA	NA
Khader et al, 2020 [45]	NA	Good	NA	Moderate	Good
Kumar et al, 2020 [48]	Poor	Positive	Fair	NA	NA
McFadde et al, 2020 [51]	NA	NA	NA	Low	NA
Modi et al, 2020 [52]	NA	NA	NA	NA	Fair
Moro et al, 2020 [53]	Fair	Positive	NA	NA	NA
Olapegba et al, 2020 [55]	Fair	NA	Fair	Moderate	NA
Olum et al, 2020 [56]	Fair	Negative	Good	NA	NA
Parikh et al, 2020 [57]	Good	NA	NA	High	NA
Ricco et al, 2020 [60]	Fair	Negative	NA	NA	Fair
Roy et al, 2020 [61]	Fair	Positive	NA	NA	Good
Sahraian et al, 2020 [62]	Good	NA	NA	NA	NA
Salman et al, 2020 [63]	Fair	Positive	Fair	NA	NA
Saqlain et al, 2020 [64]	Good	Positive	Good	Moderate	NA
Shi et al, 2020 [66]	Good	Positive	NA	NA	NA
Singh et al, 2020 [67]	Good	NA	NA	High	NA
Siniscalchi et al, 2020 [68]	NA	NA	NA	Moderate	NA
Taghrir et al, 2020 [69]	Good	NA	Good	Moderate	NA
Nwafor et al, 2020 [54]	Fair	NA	Poor	NA	NA
Wolf et al, 2020 [71]	Fair	Fairly positive	Fair	NA	Fair
Yassa et al, 2020 [72]	Fair	Positive	NA	NA	NA
Zhang et al, 2020 [73]	Good	Positive	Fair	NA	NA
Zhong et al, 2020 [74]	Good	Positive	Good	NA	NA
Reuben et al, 2020 [59]	Good	Positive	Good	NA	NA
Jemal et al, 2020 [40]	Good	Positive	Fair	NA	NA
Raza et al, 2020 [58]	Good	Positive	Fair	NA	NA
Kakemam et al, 2020 [41]	Good	Positive	Fair	NA	NA

*NA: not assessed

Discussion

COVID-19 has had serious, long-term, and sometimes irreparable effects on all aspects of the daily lives of individuals and society [75, 76]. Getting informed from the knowledge, attitude,

and practice of different people can play a vital role in preventing diseases [77, 78], so the study of these components in different communities and between different groups seems necessary.

Strength and weakness

Meta-analysis was not possible due to the different questionnaires and their dimensions, the target group, and study setting, so we reported the results descriptively in the form of tables and figures. Also, a large number of the included studies did not report the validity and reliability of the questionnaires, and therefore, the overall quality of the included studies was moderate. The main reason for this is the rush to publish articles related to coronavirus disease. Also, due to the high volume of articles published in this field, some new articles that may not have been included in this article may have been published during the writing and reviewing process. One of the most important strengths of this study was that all stages of the study were conducted with two researchers and in all stages, in cases of disagreement, the third person and consensus were used. Also, registering the protocol of this study and reviewing and modifying it in the PROSPERO platform is the strength of this study.

Summary of study findings

We found that about 60% of the samples had good knowledge of COVID-19. Also, 82% of the samples were reported positive attitudes regarding COVID-19 and slightly more than 50% of samples performed good practices. About 80% of the studies used an online questionnaire to collect data, and the most used platforms included Google form, SurveyMonkey, and Qualtrics. The most important social media through which the questionnaires were distributed were Facebook, WhatsApp, and Telegram. The most important sources for learning and staying up to

date about COVID-19 mentioned in the studies were television, social media, internet, radio, and friend and relatives.

The finding of our systematic review demonstrated fairly good knowledge about COVID-19. In most studies, more than 70% of the participants had a good knowledge of issues such as causes, symptoms, ways of transmission, and ways of prevention. Also, the majority of participants had a high level of knowledge about symptoms such as high fever and dry cough, breathing difficulty and a small number had sufficient knowledge about other symptoms such as chills, headache, muscle pain, sore throat, and loss of taste or smell [23, 25, 26, 34, 36-38, 41, 42, 64, 66]. More than 90% of the participants considered air droplets as a way to spread. This good level of knowledge can be due to widespread information through various means such as public media (television and radio), social media, and government announcements. In addition, preparing several guidelines and reports by WHO, CDC and local government in times of outbreak and easy access to them have increased the level of information and knowledge of individuals regarding COVID-19 [79, 80]. On the other hand, factors such as low literacy level, older age, and the presence of the rural population in the samples were among the factors that have reduced the level of knowledge in the studies [23, 27, 31, 33, 71].

In this review, participants showed a positive attitude regarding COVID-19. Almost all participants believed in the importance of handwashing, disinfecting surfaces, using masks to prevent the spread of infection, resting at home in the event of symptoms, and maintaining social distance and limited contact. Of course, in some cases, there was a negative belief that it could be due to differences in instructions and guidelines by different institutions, such as what was about wearing a face mask at the beginning of the pandemic, and then it was recommended that the whole population should use a mask [56, 60]. Such cases show the importance of integrated

guidelines and the focus of decision-making in times of crisis [81, 82]. Although having a responsible organization can help make better and faster decisions, in such cases, political pressure is exerted by governments that such organizations should put the health of the people at the top and not refuse to make the right decisions due to political pressures [11, 83, 84].

In general, the level of practice of the participants in the studies was fair. However, despite the high knowledge and positive attitude of the participants, the level of practices was still sometimes lower than expected. Numerous reasons for a low level of practice have been cited in studies. Lack of availability (for example, masks and disinfectants), imposing financial costs on participants, ambiguity in instructions, not getting used to new conditions such as staying home and wearing a mask, exhaustion from existing conditions, and anxiety and stress of disease were among the causes mentioned in the studies [28, 34, 40, 44, 54, 58, 71]. In this regard, some countries have imposed strict laws and penalties on people who do not follow the guidelines to improve their performance, but in many countries under study, such laws do not exist and have not been applied [85, 86]. Another factor that affects the performance of individuals was the presence of decision-makers in public and social media. Seeing a person without a mask at the height of a pandemic can have a negative impact on a person's good practices.

Given the diversity of settings and questionnaires, the authors of this paper recommend that there be a need to design an integrated online system to assess the knowledge, attitudes, and practices of the population about health-related crises. Designing such an integrated system can help better compare countries because integrated items are used for comparison. On the other hand, designing such a system and disseminating its results can accelerate integrated decision-making and improve crisis management. On the other hand, the existence of such an integrated system

can lead to an increase in solidarity, which was emphasized by the World Health Organization during the corona pandemic [87, 88].

Limitations

The included studies were from both high and low-income countries and therefore generalization of results to all countries should be done with caution. Also, many of the questionnaires in the studies did not have sufficient validity and reliability or did not report it. Our review study also had some limitations. Due to differences in studies and the use of different questionnaires, conducting a meta-analysis was not possible in this study. We have only reviewed studies published in English. On the other hand, due to the high speed of publication of articles in this field, some other studies may be published at the time of writing the article and the review process, which has been missed. Of course, due to the high speed of publishing articles, this limitation is inevitable.

Conclusion

This systematic review showed that the KAP components in the participants are at an acceptable level. In general, knowledge was at a good level, the attitude was positive and practice was at a fairly good level. Providing accurate and up-to-date information in times of crises and disseminating them through responsible institutions and through the mass media and holding online training courses can help increase people's knowledge, attitudes, and practices.

Supporting Information

Supplementary file 1; PubMed search strategy

Supplementary file 2; quality appraisal of the included studies

Acknowledgments

We thank the PROSPERO institute for accelerating the review process in the time of Coronavirus. We also thank Birjand University of Medical Sciences for approving our proposal and giving it a code of ethics (IR.BUMS.REC.1399.099).

Author Contributions

Conception and design: MA-Z. Screen the records, data extraction, and quality appraisal: MM, SS and SH. Data analysis: MA-Z and HA. Draft manuscript: MA-Z. Critical review: SH and HA. All authors approved the final version of the manuscript for publication.

References

1. Cao Z, Li T, Liang L, Wang H, Wei F, Meng S, et al. Clinical characteristics of coronavirus disease 2019 patients in Beijing, China. PloS one. 2020;15(6):e0234764.
2. Arab-Zozani M, Hassanipour S. Features and limitations of LitCovid hub for quick access to literature about COVID-19. Balkan Med J. 2020.
3. <https://www.worldometers.info/coronavirus/#countries>.
4. Arab-Zozani M, Hassanipour S, Ghoddoosi-Nejad D. Favipiravir for treating patients with novel coronavirus (COVID-19): protocol for a systematic review and meta-analysis of randomised clinical trials. BMJ open. 2020;10(7):e039730.
5. Struyf T, Deeks JJ, Dinnes J, Takwoingi Y, Davenport C, Leeftang MM, et al. Signs and symptoms to determine if a patient presenting in primary care or hospital outpatient settings has COVID-19 disease. Cochrane Database of Systematic Reviews. 2020;(7).
6. Gaythorpe K, Imai N, Cuomo-Dannenburg G, Baguelin M, Bhatia S, Boonyasiri A, et al. Report 8: Symptom progression of COVID-19. 2020.

- 297 7. Organization WH. Novel Coronavirus (2019-nCoV) situation report-7 (2020). URL <https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200127-sitrep-7-2019--ncov.pdf>. 2020.
- 298
- 299 8. Wang C, Horby PW, Hayden FG, Gao GF. A novel coronavirus outbreak of global health concern.
- 300 The Lancet. 2020;395(10223):470-3.
- 301 9. Feng S, Shen C, Xia N, Song W, Fan M, Cowling BJ. Rational use of face masks in the COVID-19
- 302 pandemic. The Lancet Respiratory Medicine. 2020;8(5):434-6.
- 303 10. Siedner MJ, Harling G, Reynolds Z, Gilbert RF, Venkataramani A, Tsai AC. Social distancing to
- 304 slow the US COVID-19 epidemic: an interrupted time-series analysis. MedRxiv. 2020.
- 305 11. Arab-Zozani M, Ghoddooosi-Nejad D. Covid-19 in Iran: the good, the bad and the ugly strategies
- 306 for preparedness—A report from the field. Disaster Medicine and Public Health Preparedness. 2020:1-6.
- 307 12. Dewey C, Hingle S, Goelz E, Linzer M. Supporting clinicians during the COVID-19 pandemic.
- 308 American College of Physicians; 2020.
- 309 13. Zhan M, Qin Y, Xue X, Zhu S. Death from Covid-19 of 23 health care workers in China. New
- 310 England Journal of Medicine. 2020.
- 311 14. Ing E, Xu Q, Salimi A, Torun N. Physician deaths from corona virus (COVID-19) disease.
- 312 Occupational Medicine. 2020;70(5):370-4.
- 313 15. Yoosefi Lebni J, Abbas J, Moradi F, Salahshoor MR, Chaboksavar F, Irandoost SF, et al. How the
- 314 COVID-19 pandemic effected economic, social, political, and cultural factors: A lesson from Iran.
- 315 International Journal of Social Psychiatry. 2020:0020764020939984.
- 316 16. Yezli S, Khan A. COVID-19 social distancing in the Kingdom of Saudi Arabia: Bold measures in the
- 317 face of political, economic, social and religious challenges. Travel Medicine and Infectious Disease.
- 318 2020:101692.

17. Wu ZMcGoogan J. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese Center for Disease Control and Prevention. JAMA. 2020;323:1239-42.
18. McEachan R, Taylor N, Harrison R, Lawton R, Gardner P, Conner M. Meta-Analysis of the Reasoned Action Approach (RAA) to Understanding Health Behaviors. Annals of behavioral medicine : a publication of the Society of Behavioral Medicine. 2016;50(4):592-612. Epub 2016/05/14. doi: 10.1007/s12160-016-9798-4. PubMed PMID: 27169555; PubMed Central PMCID: PMC4933736.
19. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med. 2009;6(7):e1000097.
20. Booth A, Clarke M, Dooley G, Ghera D, Moher D, Petticrew M, et al. The nuts and bolts of PROSPERO: an international prospective register of systematic reviews. Systematic reviews. 2012;1(1):1-9.
21. Porritt K, Gomersall J, Lockwood C. JBI's systematic reviews: study selection and critical appraisal. AJN The American Journal of Nursing. 2014;114(6):47-52.
22. Arab-Zozani M, Pezeshki MZ, Khodayari-Zarnaq R, Janati A. Inappropriate Rate of Admission and Hospitalization in the Iranian Hospitals: A Systematic Review and Meta-Analysis. Value in Health Regional Issues. 2020;21:105-12. doi: <https://doi.org/10.1016/j.vhri.2019.07.011>.
23. Abdelhafiz AS, Mohammed Z, Ibrahim ME, Ziady HH, Alorabi M, Ayyad M, et al. Knowledge, Perceptions, and Attitude of Egyptians Towards the Novel Coronavirus Disease (COVID-19). Journal of Community Health. 2020. doi: 10.1007/s10900-020-00827-7.
24. Al Nsour M, Khader Y, Al Serouri A, Bashier H, Osman S. Awareness and preparedness of Field Epidemiology Training Program graduates in the Eastern Mediterranean Region to respond to COVID-19: A cross-section study. JMIR Med Educ. 2020. doi: 10.2196/19047. PubMed PMID: 32406852.

25. Alhaj AK, Al-Saadi T, Mohammad F, Alabri S. Neurosurgery Residents Perspective on the COVID-19: Knowledge, Readiness, and Impact of this Pandemic. *World Neurosurg.* 2020. doi: 10.1016/j.wneu.2020.05.087. PubMed PMID: 32426064.
26. Alzoubi H, Alnawaiseh N, Al-Mnayyis A, Abu-Lubad M, Aqel A, Al-Shagahin H. Covid-19 - Knowledge, attitude and practice among medical and non-medical university students in Jordan. *Journal of Pure and Applied Microbiology.* 2020;14(1):17-24. doi: 10.22207/JPAM.14.1.04.
27. Apaijitt P, Wiwanitkit V. Knowledge of Coronavirus disease 2019 (COVID-19) by medical personnel in a rural area of Thailand: An observation. *Infection Control and Hospital Epidemiology.* 2020. doi: 10.1017/ice.2020.159.
28. Azlan AA, Hamzah MR, Sern TJ, Ayub SH, Mohamad E. Public knowledge, attitudes and practices towards COVID-19: A cross-sectional study in Malaysia. *PloS one.* 2020;15(5):e0233668. doi: 10.1371/journal.pone.0233668.
29. Bashedi IA, Nassar R, Barakat M, Alqudah R, Abufarha R, Muqatash T, et al. Pharmacists' readiness to deal with the coronavirus pandemic: Assessing awareness and perception of roles. *Research in Social and Administrative Pharmacy.* 2020. doi: 10.1016/j.sapharm.2020.04.020.
30. Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK. Knowledge and Perceptions of COVID-19 Among Health Care Workers: Cross-Sectional Study. *JMIR Public Health Surveill.* 2020. doi: 10.2196/19160. PubMed PMID: 32320381.
31. Bragazzi NL, Ricco M, Pacifico A, Malagoli P, Kridin K, Pigatto P, et al. COVID-19 knowledge prevents biologics discontinuation: data from an Italian multi-center survey during RED-ZONE declaration. *Dermatol Ther.* 2020. doi: 10.1111/dth.13508. PubMed PMID: 32415727.
32. Cai Y, Jiam NT, Wai KC, Shuman EA, Roland LT, Chang JL. Otolaryngology Resident Practices and Perceptions in the Initial Phase of the U.S. COVID-19 Pandemic. *Laryngoscope.* 2020. doi: 10.1002/lary.28733.

33. Chesser A, Drassen Ham A, Keene Woods N. Assessment of COVID-19 Knowledge Among University Students: Implications for Future Risk Communication Strategies. *Health Educ Behav.* 2020. doi: 10.1177/1090198120931420. PubMed PMID: 32460566.
34. Clements JM. Knowledge and behaviors toward COVID-19 among us residents during the early days of the pandemic: Cross-sectional online questionnaire. *Journal of Medical Internet Research.* 2020;22(5). doi: 10.2196/19161.
35. Erbas M, Dost B. Evaluation of knowledge and attitudes among intensive care physicians during the COVID-19 pandemic: a cross-sectional survey. *Sao Paulo Medical Journal.* 2020;(AHEAD).
36. Escalera-Antezana JP, Cerruto-Zelaya PE, Apaza-Huasco M, Miranda-Rojas SH, Flores-Cárdenas CA, Rivera-Zabala L, et al. Healthcare workers' and students' knowledge regarding the transmission, epidemiology and symptoms of COVID-19 in 41 cities of Bolivia and Colombia. *Travel Medicine and Infectious Disease.* 2020. doi: 10.1016/j.tmaid.2020.101702.
37. Galle F, Sabella EA, Molin GD, Giglio OD, Caggiano G, Onofrio VD, et al. Understanding Knowledge and Behaviors Related to CoViD-19 Epidemic in Italian Undergraduate Students: The EPICO Study. *Int J Environ Res Public Health.* 2020. doi: 10.3390/ijerph17103481. PubMed PMID: 32429432.
38. Geldsetzer P. Knowledge and Perceptions of COVID-19 Among the General Public in the United States and the United Kingdom: A Cross-sectional Online Survey. *Annals of internal medicine.* 2020. doi: 10.7326/M20-0912.
39. Giao H, Le An P, Khanh T.V, Ngan V.K, Tam V.V, Le An P. Knowledge and attitude toward COVID-19 among healthcare workers at District 2 Hospital, Ho Chi Minh City. *Asian Pacific Journal of Tropical Medicine.* 2020;13.
40. Jemal B, Ferede ZA, Mola S, Hailu S, Abiy S, Wolde GD, et al. Knowledge, attitude and practice of healthcare workers towards COVID-19 and its prevention in Ethiopia: a multicenter study. 2020.

41. Kakemam E, Ghoddoosi-Nejad D, Chegini Z, Momeni K, Salehinia H, Hassanipour S, et al. Knowledge, attitudes, and practices among the general population during COVID-19 outbreak in Iran: A national cross-sectional survey. medRxiv. 2020.
42. Kamate SK, Sharma S, Thakar S, Srivastava D, Sengupta K, Hadi AJ, et al. Assessing Knowledge, Attitudes and Practices of dental practitioners regarding the COVID-19 pandemic: A multinational study. Dent Med Probl 2020;57(1):11-7.
43. Karasneh R, Al-Azzam S, Muflih S, Soudah O, Hawamdeh S, Khader Y. Media's effect on shaping knowledge, awareness risk perceptions and communication practices of pandemic COVID-19 among pharmacists. Research in Social and Administrative Pharmacy. 2020. doi: 10.1016/j.sapharm.2020.04.027.
44. Kebede Y, Yitayih Y, Birhanu Z, Mekonen S, Ambelu A. Knowledge, perceptions and preventive practices towards COVID-19 early in the outbreak among Jimma university medical center visitors, Southwest Ethiopia. PloS one. 2020;15(5):e0233744. doi: 10.1371/journal.pone.0233744.
45. Khader Y, Al Nsour M, Al-Batayneh O.B, Saadeh R, Bashier H, Alfaqih M, et al. Dentists' Awareness, Perception, and Attitude Regarding COVID-19 and Infection Control: Cross-Sectional Study Among Jordanian Dentists. JMIR Public Health Surveill. 2020;6(2).
46. Khan MA, Sivalingam A, Haller JA. Perceptions of Occupational Risk and Changes in Clinical Practice of U.S. Vitreoretinal Surgery Fellows during the COVID-19 Pandemic. Ophthalmol Retina. 2020. doi: 10.1016/j.oret.2020.05.011. PubMed PMID: 32450265.
47. Khasawneh AI, Abu Humeidan A, Alsulaiman JW, Bloukh S, Ramadan M, Al-Shatanawi TN, et al. Medical Students and COVID-19: Knowledge, Attitudes and Precautionary Measures. A Descriptive Study from Jordan. Frontiers in Public Health. 2020;8:253.
48. Kumar J, Katto MS, Siddiqui AA, Sahito B, Jamil M, Rasheed N, et al. Knowledge, Attitude, and Practices of Healthcare Workers Regarding the Use of Face Mask to Limit the Spread of the New

413 Coronavirus Disease (COVID-19). Cureus. 2020;12(4). doi: 10.7759/cureus.7737. PubMed PMID:
414 WOS:000526848800001.

415 49. Lau LL, Hung N, Go DJ, Ferma J, Choi M, Dodd W, et al. Knowledge, attitudes and practices of
416 COVID-19 among income-poor households in the Philippines: A cross-sectional study. Journal of global
417 health. 2020;10(1).

418 50. Liu H, Li X, Chen Q, Li Y, Xie C, Ye M, et al. Illness perception, mood state and disease-related
419 knowledge level of COVID-19 family clusters, Hunan, China. Brain Behav Immun. 2020. doi:
420 10.1016/j.bbi.2020.05.045. PubMed PMID: 32425333.

421 51. McFadden SM, Malik AA, Aguolu OG, Willebrand KS, Omer SB. Perceptions of the adult US
422 population regarding the novel coronavirus outbreak. PLoS ONE. 2020;15(4). doi:
423 10.1371/journal.pone.0231808.

424 52. Modi PD, Nair G, Uppe A, Modi J, Tuppekar B, Gharpure AS, et al. COVID-19 Awareness Among
425 Healthcare Students and Professionals in Mumbai Metropolitan Region: A Questionnaire-Based Survey.
426 Cureus. 2020;12(4). doi: 10.7759/cureus.7514. PubMed PMID: WOS:000523461700009.

427 53. Moro M, Vigezzi GP, Capraro M, Biancardi A, Nizzero P, Signorelli C, et al. 2019-novel
428 coronavirus survey: Knowledge and attitudes of hospital staff of a large Italian teaching hospital. Acta
429 Biomedica. 2020;91:29-34. doi: 10.23750/abm.v91i3-S.9419.

430 54. Nwafor JI, Aniukwu JK, Anozie BO, Ikeotuonye AC, Okedo-Alex IN. Pregnant women's knowledge
431 and practice of preventive measures against COVID-19 in a low-resource African setting. International
432 Journal of Gynecology and Obstetrics. 2020. doi: 10.1002/ijgo.13186.

433 55. Olapegba PO, Iorfa SK, Kolawole SO, Oguntayo R, Gandhi JC, Ottu IFA, et al. Survey data of COVID-
434 19-related Knowledge, Risk Perceptions and Precautionary Behavior among Nigerians. Data in Brief.
435 2020;30. doi: 10.1016/j.dib.2020.105685.

436 56. Olum R, Chekwech G, Wekha G, D.A N, Bongomin F. Coronavirus Disease-2019: Knowledge,
437 Attitude, and Practices of Health Care Workers at Makerere University Teaching Hospitals,Uganda. Front
438 Public Health. 2020;8.

439 57. Parikh PA, Shah BV, Phatak AG, Vadnerkar AC, Uttekar S, Thacker N, et al. COVID-19 Pandemic:
440 Knowledge and Perceptions of the Public and Healthcare Professionals. Cureus. 2020;12(5). doi:
441 10.7759/cureus.8144. PubMed PMID: WOS:000533196100017.

442 58. Raza A. Public Awareness Towards COVID-19: A Cross Sectional Survey Among Patients of
443 Pakistan Homeopathic Trust Hospital Lahore, Pakistan. International Journal of Homeopathy,
444 Complementary and Alternative Medicine. 2020;7-.

445 59. Reuben RC, Danladi MM, Saleh DA, Ejembi PE. Knowledge, Attitudes and Practices Towards
446 COVID-19: An Epidemiological Survey in North-Central Nigeria. Journal of Community Health. 2020;1-14.

447 60. Riccò M, Ferraro P, Gualerzi G, Ranzieri S, Bragazzi NL, Balzarini F, et al. Point-of-Care diagnostic
448 of SARS-CoV-2: knowledge, attitudes, and perceptions (KAP) of medical workforce in Italy. Acta bio-
449 medica : Atenei Parmensis. 2020;91(2):57-67. doi: 10.23750/abm.v91i2.9573.

450 61. Roy D, Tripathy S, Kar SK, Sharma N, Verma SK, Kaushal V. Study of knowledge, attitude, anxiety
451 & perceived mental healthcare need in Indian population during COVID-19 pandemic. Asian Journal of
452 Psychiatry. 2020;51. doi: 10.1016/j.ajp.2020.102083.

453 62. Sahraian MA, Gheini MR, Rezaeimanesh N, Ghajarzadeh M, Naser Moghadasi A. Knowledge
454 regarding COVID-19 pandemic in patients with multiple sclerosis (MS): A report from Iran. Multiple
455 Sclerosis and Related Disorders. 2020;42. doi: 10.1016/j.msard.2020.102193.

456 63. Salman M, Mustafa ZU, Asif N, Zaidi HA, Hussain K, Shehzadi N, et al. Knowledge, attitude and
457 preventive practices related to COVID-19: a cross-sectional study in two Pakistani university populations.
458 Drugs and Therapy Perspectives. 2020. doi: 10.1007/s40267-020-00737-7.

64. Saqlain M, Munir MM, Rehman SU, Gulzar A, Naz S, Ahmed Z, et al. Knowledge, attitude, practice and perceived barriers among healthcare professionals regarding COVID-19: A Cross-sectional survey from Pakistan. J Hosp Infect. 2020. doi: 10.1016/j.jhin.2020.05.007. PubMed PMID: 32437822.
65. Sari DK, Amelia R, Dharmajaya R, Sari LM, Fitri NK. Positive Correlation Between General Public Knowledge and Attitudes Regarding COVID-19 Outbreak 1 Month After First Cases Reported in Indonesia. Journal of community health. 2020:1-8.
66. Shi Y, Wang J, Yang Y, Wang Z, Wang G, Hashimoto K, et al. Knowledge and attitudes of medical staff in Chinese psychiatric hospitals regarding COVID-19. Brain Behav Immun Health. 2020. doi: 10.1016/j.bbih.2020.100064. PubMed PMID: 32289123.
67. Singh DR, Sunuwar DR, Karki K, Ghimire S, Shrestha N. Knowledge and Perception Towards Universal Safety Precautions During Early Phase of the COVID-19 Outbreak in Nepal. Journal of Community Health. 2020. doi: 10.1007/s10900-020-00839-3.
68. Siniscalchi M, Zingone F, Savarino EV, D'Odorico A, Ciacci C. COVID-19 pandemic perception in adults with celiac disease: an impulse to implement the use of telemedicine: COVID-19 and CeD. Dig Liver Dis. 2020. doi: 10.1016/j.dld.2020.05.014. PubMed PMID: 32425731.
69. Taghrir MH, Borazjani R, Shiraly R. COVID-19 and iranian medical students; A survey on their related-knowledge, preventive behaviors and risk perception. Archives of Iranian Medicine. 2020;23(4):249-54. doi: 10.34172/aim.2020.06.
70. Wahed WYA, Hefzy EM, Ahmed MI, Hamed NS. Assessment of knowledge, attitudes, and perception of health care workers regarding COVID-19, a cross-sectional study from Egypt. Journal of community health. 2020:1-10.
71. Wolf MS, Serper M, Opsasnick L, O'Connor RM, Curtis LM, Benavente JY, et al. Awareness, Attitudes, and Actions Related to COVID-19 Among Adults With Chronic Conditions at the Onset of the

482 U.S. Outbreak: A Cross-sectional Survey. *Ann Intern Med.* 2020. doi: 10.7326/M20-1239. PubMed PMID:
483 32271861.

484 72. Yassa M, Birol P, Yirmibes C, Usta C, Haydar A, Yassa A, et al. Near-term pregnant women's
485 attitude toward, concern about and knowledge of the COVID-19 pandemic. *J Matern Fetal Neonatal*
486 *Med.* 2020. doi: 10.1080/14767058.2020.1763947. PubMed PMID: 32429780.

487 73. Zhang M, Zhou M, Tang F, Wang Y, Nie H, Zhang L, et al. Knowledge, attitude, and practice
488 regarding COVID-19 among healthcare workers in Henan, China. *Journal of Hospital Infection.*
489 2020;105(2):183-7. doi: 10.1016/j.jhin.2020.04.012.

490 74. Zhong B.L, Luo W, Li H.M, Zhang Q.Q, Liu X.G, Li W.T, et al. Knowledge, attitudes, and practices
491 towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a
492 quick online cross-sectional survey. *Int J Biol Sci.* 2020;16(10).

493 75. Tull MT, Edmonds KA, Scamaldo K, Richmond JR, Rose JP, Gratz KL. Psychological Outcomes
494 Associated with Stay-at-Home Orders and the Perceived Impact of COVID-19 on Daily Life. *Psychiatry*
495 *research.* 2020:113098.

496 76. Haleem A, Javaid M, Vaishya R. Effects of COVID 19 pandemic in daily life. *Current Medicine*
497 *Research and Practice.* 2020.

498 77. Ghazanfari Z, Azattalab F. Knowledge, Attitude and Practice (Kap) Study Regarding Prevention of
499 Breast Cancer in Working Women of Chaloos City. *SSU_Journals.* 2006;14(2):51-5.

500 78. Namikawa K, Kikuchi H, Kato S, Takizawa Y, Konta A, Iida T, et al. Knowledge, attitudes, and
501 practices of Japanese travelers towards malaria prevention during overseas travel. *Travel Medicine and*
502 *Infectious Disease.* 2008;6(3):137-41.

503 79. Organization WH. Coronavirus disease 2019 (COVID-19): situation report, 82. 2020.

504 80. Organization WH. Coronavirus disease 2019 (COVID-19): situation report, 72. 2020.

81. Hong KH, Lee SW, Kim TS, Huh HJ, Lee J, Kim SY, et al. Guidelines for laboratory diagnosis of coronavirus disease 2019 (COVID-19) in Korea. *Ann Lab Med*. 2020;40(5):351-60.
82. Jin Y-H, Cai L, Cheng Z-S, Cheng H, Deng T, Fan Y-P, et al. A rapid advice guideline for the diagnosis and treatment of 2019 novel coronavirus (2019-nCoV) infected pneumonia (standard version). *Military Medical Research*. 2020;7(1):4.
83. Flinders M. Democracy and the Politics of Coronavirus: Trust, Blame and Understanding. *Parliamentary Affairs*. 2020.
84. Davies S. The politics of staying behind the frontline of coronavirus. *Wellcome Open Research*. 2020;5(131):131.
85. Parmet WE, Sinha MS. Covid-19—the law and limits of quarantine. *New England Journal of Medicine*. 2020;382(15):e28.
86. Griffith R. Using public health law to contain the spread of COVID-19. *British Journal of Nursing*. 2020;29(5):326-7.
87. Arab-Zozani M, Hassanipour S. Sharing Solidarity Experiences to Overcome COVID-19. *Annals of Global Health*. 2020;86(1).
88. Arora G, Kroumpouzou G, Kassir M, Jafferany M, Lotti T, Sadoughifar R, et al. Solidarity and transparency against the COVID-19 pandemic. *Dermatologic Therapy*. 2020.

Supporting information captions

Supplementary file 1: PRISMA 2009 checklist

Supplementary file 2: Complete search strategy

Supplementary file 3: Quality appraisal of the included studies

Identification

Records identified through
database searching
(n = 4829)

Additional records identified
through other sources
(n = 12)

Screening

Records after duplicates removed
(n = 3085)

Eligibility

Records screened
(n = 3085)

Records excluded
(n = 3012)

Full-text articles assessed
for eligibility
(n = 73)

Full-text articles excluded,
with reasons
(n = 21)
Non original articles= 14
Not reported KAPs
components= 7

Included

Final included studies
(n = 52)

A

B

C

Quality appraisal of included studies

low fair good

K

A

P