

Supplement Tables

Supplement Table 1 Clinical symptoms of whole cohort

	Total (n=386)	Vitamin D deficient (n = 61)^{a,b}	Non-Vitamin D deficient (n = 325)^{a,b}	P value
Cough no. (%)				
Yes	117 (30%)	15 (25%)	102 (31%)	0.362
No	269 (70%)	46 (75%)	223 (69%)	
Fever no. (%)				
Yes	235 (61%)	44 (72%)	191 (59%)	0.063
No	151 (39%)	17 (28%)	134 (41%)	
Breathlessness no. (%)				
Yes	186 (48%)	28 (46%)	158 (49%)	0.780
No	200 (52%)	33 (54%)	167 (51%)	
Cough + fever + breathlessness no. (%)				
Yes	87 (23%)	18 (30%)	69 (21%)	0.181
No	299 (77%)	43 (70%)	256 (79%)	
Loss of smell or taste no. (%)				
Yes	169 (44%)	32 (52%)	137 (42%)	0.160
No	217 (56%)	29 (48%)	188 (58%)	
Body aches and pains no. (%)				
Yes	274 (71%)	50 (82%)	224 (69%)	0.045
No	112 (29%)	11 (18%)	101 (31%)	
Fatigue no. (%)				
Yes	339 (88%)	53 (87%)	286 (88%)	0.831
No	47 (12%)	8 (13%)	39 (12%)	
Diarrhoea no. (%)				
Yes	115 (30%)	23 (38%)	92 (28%)	0.169
No	271 (70%)	38 (62%)	233 (72%)	
Sore throat (%)				
Yes	197 (51%)	37 (61%)	160 (49%)	0.124
No	189 (49%)	24 (39%)	165 (51%)	

^a Vitamin D deficient is Serum 25(OH)D₃ < 30 nmol/l while not deficient is ≥ 30 nmol/l; ^b Where proportions are shown, they were calculated using the n numbers shown in columns as denominator; p values were calculated using Fisher's exact test for data showing proportions, p value <0.05 is considered significant.