1 **Current Practice and Challenges in Screening for Visual Perception Deficits after**

2 **Stroke: a Qualitative Study**

3 Kathleen Vancleef, (corresponding author)

4 Department of Experimental Psychology, University of Oxford, Anna Watts Building,

5 Radcliffe Observatory Quarter, Woodstock Road, University of Oxford, OX2 6GG,

6 United Kingdom

7 Email: kathleen.vancleef@psy.ox.ac.uk

8 Twitter: @katvancleef

9 ORCID: 0000-0002-9943-9341

10 Michael J Colwell,

11 Department of Experimental Psychology, University of Oxford

12 Twitter: @MichaelJColwell

13 Olivia Hewitt,

14 The Oxford Institute of Clinical Psychology Training and Research, University of

15 Oxford

16 Nele Demeyere,

17 Department of Experimental Psychology, University of Oxford

18 Twitter: @NeleDemeyere

19 ORCID: 0000-0003-0416-5147

20

21 Abstract

22 PURPOSE: We document current clinical practice and needs in screening for visual
23 perception problems after stroke to inform development of new screening tools.

24 MATERIALS AND METHODS: We interviewed 12 occupational therapists and 13
25 orthoptists. Interviews were thematically analysed using the Value Proposition Canvas, a
26 model which establishes challenges and facilitators in what people want to achieve.

27 RESULTS: Participants' understanding of visual perception varied and often included
28 sensory and cognitive deficits. Occupational therapists commonly screened for visual field
29 deficits and hemispatial neglect, while other aspects of visual cognition were rarely assessed.
30 A positive screening result triggered an orthoptic referral. Screening generally occurred
31 during functional assessments and/or with in-house developed tools. Challenges to practice
32 were: lack of time, lack of training, environmental and stroke survivor factors (e.g. aphasia),
33 insufficient continuation of care, and test characteristics (e.g. not evidence-based).
34 Facilitators were: quick and practical tools, experienced staff or tools with minimal training
35 requirements, a streamlined care pathway.

36 CONCLUSIONS: Screening employs non-standardised assessments and rarely covers higher
37 visual perceptual deficits. We demonstrates the need for an evidence-based visual perception
38 screen, which should ideally be 15 minutes or less, be portable, and require minimal
39 equipment. The screen should be suitable for bedside testing and aphasia-friendly.

40 Keywords

41 Visual perception, screening, stroke, assessment, clinical needs, facilitators, barriers

42

43

44 Introduction

45 Visual perception is the dynamic process of perceiving the environment through sensory
 46 inputs and translating the sensory input into meaningful concepts associated with visual
 47 knowledge of the environment.(1) Visual perception problems are therefore distinct from
 48 sensory visual impairments such as reduced visual acuity, visual field and eye movements.(2)
 49 Where sensory visual impairments result from damage to the eye or early visual pathways
 50 from the eye to the primary visual cortex, visual perception deficits are attributed to impaired
 51 function in later visual processing areas in the occipital, parietal and temporal cortex.(3)
 52 Examples of visual perceptual deficits include apperceptive and associative agnosia (object
 53 recognition difficulties), prosopagnosia (face recognition difficulties), akinetopsia
 54 (difficulties in perceiving motion), achromatopsia (difficulties in perceiving motion),
 55 problems in visual memory (remembering what you have seen before), and in visuospatial
 56 abilities (e.g. judging distances or spatial relations between objects).(3) Visual inattention or
 57 hemispatial neglect is sometimes considered to be part of visual perception,(1) (4) though
 58 neuropsychology research attributes this to an attentional deficit.(5) In particular, the
 59 presence of preserved perception when attention is stretched to focus on the stimuli, the
 60 existence of cross-modal neglect and manipulations of stimulus density on the extent of
 61 neglect support the classification of hemispatial neglect as a disorder of attention. (6,7)
 62 Rowe and colleagues reported that 20.5% of stroke survivors with a suspected visual
 63 difficulty have visual perception deficits such as visual inattention (14%), visual
 64 hallucinations (2.5%), and object agnosia (2.2.%(8) This study made use of reports by stroke
 65 survivors and carers rather than formal assessment (except for visual inattention). With
 66 systematic screening with the Rivermead Perceptual Assessment Battery, Edmans and
 67 Lincoln identified visual perception problems in 76% of hemiplegic stroke survivors.(9) The
 68 discrepancy between prevalence with self-reports compare to neuropsychological assessment

69 suggests that not all visual perceptual problems are picked up based on self-report. This
70 means many stroke survivors are discharged without the appropriate rehabilitation or
71 adjustments in their home environment or care packages.

72 Under-diagnosis of visual perception problems can severely impact stroke survivors' quality
73 of life, functional outcome, participation in the community, independence and pose
74 substantial risk. (10–14) For instance, participation in traffic with visual perception problems
75 can be dangerous and even life-threatening. Risks are also heightened indoors when
76 preparing a meal or in perceiving trip hazards. Better diagnosis of visual perception
77 difficulties after stroke will allow better care planning and substantially impact stroke
78 survivors' life.

79 The first step toward better diagnosis is an in-depth understanding of the clinical reality and
80 the reasons behind under-diagnosis. We need to understand what the challenges are in
81 screening for visual perceptual difficulties after stroke. This can then inform the development
82 of solutions in the future.

83 In the current study, we aimed to conduct an in-depth exploration of current clinical practice,
84 challenges and facilitators of screening for visual perception problems after stroke. To
85 achieve a rich understanding of these issues, we performed semi-structured interviews with
86 orthoptists and occupational therapists. These professionals are most commonly involved in
87 visual perception screening after stroke in the United Kingdom's National Health Service.

88 **Materials and methods**

89 ***Participants***

90 All participants were recruited via opportunistic and snowball sampling. Invitations for the
 91 interviews were sent out through (i) email to the British and Irish Orthoptic Society Stroke
 92 and Neuro Rehab Clinical Advisory Group, (ii) Twitter, (iii) a sign-up sheet at a conference
 93 poster and (iv) informal face-to-face conversations at the United Kingdom Stroke Forum
 94 2018. Participants met inclusion criteria if they were working in the National Health Service
 95 as an occupational therapist or an orthoptist, and were involved in the assessment of visual
 96 perceptual problems after stroke. The participants were informed about the aims of the study,
 97 the organisations running the study (University of Oxford and North Bristol NHS Trust), and
 98 the planned outcomes and dissemination. The project was reviewed and approved by the
 99 Patient Safety Assurance & Audit Service at NHS North Bristol Trust as a Clinical
 100 Effectiveness study (CE45999). All participants gave verbal consent.

101 ***Interviewer's training***

102 All interviews were conducted over the phone by the first author, KV, who holds an MSc in
 103 Clinical Psychology and a PhD in Experimental Psychology. At the time of the research
 104 study KV was working as a postdoctoral research fellow at the University of Oxford. KV had
 105 nine years' experience as a researcher, including one year as a qualitative researcher. Her
 106 formal training in qualitative research included an undergraduate course in qualitative
 107 research methods at the University of Leuven which included a research project. In
 108 preparation for the current project, she was trained and supervised by senior author, OH. OH
 109 is a clinical psychologist who has worked predominantly as a qualitative researcher and
 110 health care professional within the field of intellectual disabilities. OH has numerous
 111 publications within this field, teaches qualitative research methods at the University of
 112 Oxford, and has supervised a number of doctoral projects using various qualitative

methodologies. KV's training included reading and reflecting on reference works in qualitative research,(15,16) a question and answer session with an experienced moderator, and guidance during coding and analyses via regular meetings with OH.

Model

The interview process and qualitative analysis was guided by the Value Proposition Canvas. This model was originally designed to guide product development, but can be applied to different contexts.(17) Central to this model is the principle that development of products or services are best informed by the users. A good product or service takes into account the goals of the user, it reduces the pains a user experiences while trying to achieve their goals, and it provides gains in achieving their goals. For instance a good electric car helps a person to get to work (goal to achieve) without having to fear that the battery will run out (reduce pains) while at the same time provides an attractive design to impress colleagues and family (increase gains). We can apply this model in trying to understand current clinical practice in visual perception screening after stroke (goal), the challenges (pains) it brings, and what facilitates the process (gains).

Data collection

A semi-structured interview guide was developed by KV in collaboration with an occupational therapist and an orthoptist (table 1). The interview guide was reviewed by an experienced moderator and piloted with one orthoptist and three occupational therapists. The interviews consisted of open-ended questions about current practice in visual perception screening after stroke, and the challenges and facilitators of practice. The interviewer conducted interviews at the participant's workplace or over the phone in a quiet office with no other people present. The participants took the call in a location of their choice. The location and presence of others at the participant's side was not recorded. After obtaining participant consent, the interview was audio recorded. All participants were interviewed once.

138 Recruitment was ongoing during data collection and interviews were conducted until data
139 saturation was reached.

140 Table 1. Interview structure

Section	Interview route
Introduction	<i>Q.</i> Which duties are central to your job role?
Definition of visual perception	<i>Q.</i> What comes to your mind when I say visual perception problems in stroke survivors?
Current practice	<i>Q.</i> Do you do visual perception screening as a routine on your stroke survivors? (Orthoptists only)
	<i>Q.</i> What did you do the last time you suspected visual perception problems in a stroke survivor?
	<i>Cues: standardised tests, unstandardized tests, observations, conversation with stroke survivor, family members, other staff, medical notes</i>
	<i>Q.</i> What is the care pathway for these stroke survivor? <i>Cues: Which health care professional (occupational therapist,, neuropsychologist, orthoptist, consultant, physiotherapist) does what (clinical judgement, systematic screen, in depth assessment) and where (A&E, Acute stroke ward, rehab ward, care home, at home, outpatient appointment at eye clinic), referral process</i>
Facilitators to practice	<i>Q.</i> What does an initial visual perception screen need to be like for you? <i>Cues: time it takes, format, what to include, what result, who should do it</i>
Challenges to practice	<i>Q.</i> Can you name three things that frustrate you about visual perception screening on the stroke ward? <i>Cues: assessment methods, time, training, other staff, care pathway,</i>
	<i>Q.</i> Which of the three challenges you have mentioned is the most important?
Ending	<i>Q.</i> If you could give one piece of advice to someone making a visual perception screen, what would it be? <i>Q.</i> My goal was to have an idea of how visual perception screening is done and what the clinical needs are. Do you think we've talked about everything or is there anything we've missed?

Qualitative Analyses

As a method of data analysis we used thematic analysis.(18) Interviews were transcribed verbatim by a professional typist. The transcripts were not returned to the participants for feedback. Two researchers (KV and MC) were involved in data analysis. KV was familiar with the data from conducting the interviews (Phase 1 in thematic analysis). The first 15 interviews were checked and coded by KV in NVivo Software (Version 12.0). She developed an initial coding tree (Phase 2 in thematic analysis). Overall themes were set in advance in alignment with the Value Proposition Model: Jobs, Challenges, and Facilitators. The coding tree was reviewed and substantially adapted after coding of five interviews. MC familiarised himself with the data by checking transcripts and reading through interviews coded by KV. Two of these interview transcripts were checked, coded and discussed by both KV and MC to align their coding. MC checked and coded the remaining 10 interview transcripts. The frequency of text extracts coded under each node was visualised with a hierarchical chart of nodes in NVivo to explore overall patterns in the data describing current practice. The use of frequencies in qualitative analysis can be a valuable strategy when used as a complement to an overall process orientation to the research. They can provide evidence for generalizability of the findings between participants, highlight diversity, reveal larger patterns beyond a participant's immediate experience, minimise cherry picking of the data.(19) This chart shows the frequency of references for each node when participants were talking about current practice in routine visual and visual perception screening. If a participant referred to a node multiple times during one interview, each reference was counted. Only nodes with at least ten references were included in the hierarchical chart of nodes. Overall themes and subthemes were derived from all data by KV (Phase 3 of thematic analysis). This was done in two stages: (1) we performed search queries of coded text for challenges and facilitators of visual perception screening in current practice and (2) read through the resulting text. The themes

were subsequently reviewed by reading all answers to the questions under the headings Challenges and Facilitators (see table 1). The information was then synthesised by KV (Phase 4 of thematic analysis). Lastly, themes and subthemes were reviewed, refined and defined by KV (Phase 5 of thematic analysis) and a report was prepared (Phase 6 of thematic analysis). This report was sent out to the participants for validation. Ten out of 25 participants replied and nine agreed with the thematic analysis. One participant pointed out that participants' understanding of visual perception was underexplored. We added a results section on this topic posthoc.

Results

12 occupational therapists and 13 orthoptists from 16 health care organisations in England took part in the study. Characteristics such as gender, years of experience and main clinical setting of each participant are reported in table 2. All participants worked in England, and were based across the East Midlands (n=1), East of England (n=1), Greater London (n=2), North East (n=3), North West (n=4), South East (n=6), South West (n=4), and Yorkshire and the Humber (n=4). The interviewer had no personal relationship with any of the participants. Two orthoptists received referrals from the same acute stroke unit where the first author recruits stroke survivors as participants for other studies (pseudonyms cannot be disclosed to ensure anonymity). One of the participants was also involved in the development of the interview guide. All but one participant (Jessica) agreed to be audio recorded. For Jessica, notes were made during and immediately after the interview. Interviews lasted between 16 and 46 minutes with an average of 27 minutes. For Christopher, the interview was split over two consecutive days.

190 Table 2. Participant characteristics

Pseudonym	Gender	Profession	Clinical setting [†]	Years of experience in stroke care
Amanda	Female	Orthoptist	Eye clinic	10+ years
Amber	Female	Occupational therapist	Rehabilitation service	10+ years
Amy	Female	Orthoptist	Eye clinic	unknown
Brittany	Female	Occupational therapist	Rehabilitation service	10+ years
Christopher	Male	Occupational therapist	Acute stroke service	unknown
Crystal ^a	Female	Occupational therapist	Rehabilitation service	5-10 years
Danielle	Female	Occupational therapist	Acute stroke service	unknown
Elizabeth	Female	Orthoptist	Eye clinic	10+ years
Emily ^a	Female	Occupational therapist	Rehabilitation service	unknown
Erin	Female	Occupational therapist	Acute stroke service	5-10 years
Heather ^b	Female	Orthoptist	Eye clinic	10+ years
Jamie ^b	Female	Orthoptist	Eye clinic	2-5 years
Jessica ^c	Female	Orthoptist	Eye clinic	10+ years
Joshua	Male	Orthoptist	Eye clinic	2-5 years
Kimberley	Female	Occupational therapist	Acute stroke service	2-5 years
Matthew	Male	Orthoptist	Eye clinic	5-10 years
Megan	Female	Occupational therapist	Rehabilitation service	10+ years
Melissa	Female	Occupational therapist	Acute stroke service	10+ years
Michelle	Female	Occupational therapist	Rehabilitation service	2-5 years
Nicole	Female	Orthoptist	Eye clinic	unknown
Rachel	Female	Orthoptist	Eye clinic	unknown
Rebecca	Female	Occupational therapist	Acute stroke service	5-10 years
Sarah ^c	Female	Orthoptist	Eye clinic	10+ years
Stephanie	Female	Orthoptist	Eye clinic	unknown
Tiffany	Female	Orthoptist	Eye clinic	2-5 years

191 ^{a,b,c} Participants working in the same clinical teams A, B or C. [†] All orthoptists were based in eye clinics but
192 could offer inpatient appointments on the stroke unit, inpatient appointments in the eye clinic and outpatient
193 appointments in the eye clinic.

194

195 ***Lack of clarity around visual perception***

196 The interviews highlighted that participants' understanding of visual perception differed from
197 textbook definitions of visual perception. Visual perception is defined as a dynamic process
198 of translating sensory visual information into meaning percepts(1). These percepts can
199 subsequently be linked to higher cognitive functions like visual memory for recognition
200 (ventral or what stream) and praxis for visually guided actions (dorsal or where stream)(3).
201 Visual perception must be differentiated from low level sensory visual functions like acuity,

visual fields, and ocular movements, and from higher cognitive processes like (visual) memory or (visual) inattention, although some authors consider visual inattention to be part of visual perception. Our participants seemed to have a broader understanding of visual perception and often did not differentiate between sensory and higher cognitive functions. Participants typically mentioned the following examples when asked about what they consider visual perceptual problems (in descending order of frequency): visual inattention, agnosia's for objects, faces, letters, or shapes, hallucinations or Charles Bonnet syndrome, difficulties in depth perception, visual field deficits like hemianopia, and spatial vision (e.g. navigating and perceiving space around them).

Participants' understanding seemed to vary widely but we observed general trends in the different professions. Orthoptists seemed to differentiate visual perception difficulties from low level sensory visual impairments, but considered visual inattention the prime (and only) example of visual perception difficulties after stroke.

“[Interviewer: What comes to your mind, when I say visual perception problems in stroke patients?] So I would probably say visual neglect, ignoring one side of their world when the other side is presented to them as well, is the main definition of it. That it's decreased awareness of that side of their vision and their external world as well as themselves as well, on that side. [Interviewer: Anything else, that falls under that term for you?] No, not that I can think of, off the top of my head, no.” (Tiffany, Orthoptist in an eye clinic)

In addition to visual inattention, occupational therapists mentioned impairments that are typically (but not exclusively) related to low level sensory vision like contrast sensitivity, hemianopia, and depth perception. They seemed to interpret visual perception more broadly.

224 “A range really. So we would look at visual perception as things like depth perception, ...
 225 recognition of objects, ... and any kind of visual inattention or neglect, that sort of thing...
 226 But yes, that’s it basically.” (Danielle, Occupation Therapist in an acute stroke service)

227 These observations are relevant in interpreting the subsequent results. Although the interview
 228 questions asked for experiences with visual perception assessment after stroke, the results
 229 below should be interpreted as potentially referring to all or any of the following: sensory
 230 visual impairments (for occupational therapist participants), visual perception impairments
 231 (for all participants), and visual inattention (for all participants).

232 ***Current Practice***

233 A hierarchical chart of nodes is presented in figure 1. The figure illustrates that participants
 234 most often mentioned occupational therapists working in acute stroke or rehabilitation
 235 services when discussing screening for visual neglect, visual field deficits, and ocular
 236 movements. For screening they referred to in-house developed screening tools, standardised
 237 tests or observations in functional tasks like dressing and making a cup of tea.

238 “In terms of the visual field, we do the standard, you know, wiggle your fingers
 239 test to see how people respond. We do the tracking tests. Otherwise it’s mainly
 240 through observing people in functional tasks what they’re doing. ... Through that
 241 you can often realise ... if they’re having problems with their visual perception
 242 because they’re missing things or not seeing things.” (Crystal, occupational
 243 therapist in a stroke rehabilitation service)

244 “I’d rather do it in function. It’s more meaningful especially with stroke patients
 245 because it makes more sense. Rather than getting them to balance cubes on each
 246 other or, you know, I can say ‘Can you find the toothpaste?’ ” (Megan,
 247 occupational therapist in an acute stroke service)

Figure 1. Hierarchical chart of nodes showing the frequency of references for each node. The size of a box

reflects the frequency of references for each node with larger boxes demonstrating more frequency mentioned nodes. Parent nodes are shown in lightly shaded boxes, child nodes are shown in darker shaded boxes within their parent's box. The colours are added for easy distinction between different parent boxes but have no further meaning. Text is filtered for extracts where participants were talking about current practice in routine visual and visual perceptions screening. Only nodes with at least ten references are shown.

Most participants mentioned that the service they work within has a referral pathway in place: if problems are suspected following screening by the occupational therapist, a referral is made to orthoptics. A stroke orthoptist will then see the stroke survivor as an inpatient on the unit or as an outpatient after discharge from the hospital.

“It's really as and how other health professionals have identified them, then an orthoptist will go up to try and assert a diagnosis and then give them the rehabilitation options.” (Amy, orthoptist in an eye clinic)

263 *Challenges*

264 Themes around challenges in visual perception screening are presented in table 3. We
265 identified five major themes and 13 subthemes.

266 *Lack of Time*

267 Occupational therapists and orthoptists both reported that time pressure makes visual
268 perception screening difficult. They mentioned that because of understaffing they spend little
269 time with stroke survivors or experience delays in seeing them following referral. In addition,
270 occupational therapists report that stroke survivors are discharged from an acute stroke unit
271 as soon as medically safe and a suitable discharge location has been organised. For survivors
272 with milder strokes, discharge is reported to happen within 1-2 days, giving very little time
273 for therapists to complete their assessments.

274 “Well one [*challenge*] is just logistically and organisationally, just in terms of
275 caseload and time to hand. We are having to prioritise more direct functional
276 assessments in terms of establishing readiness for discharge.” (Christopher,
277 occupational therapist in an acute stroke service)

278 “There’s not a lot of OTs who are actually for [*delivering*] the therapy, so time-
279 wise we haven’t got a lot of capacity at times.” (Danielle, occupational therapist in
280 an acute stroke service)

281 “Lack of time is always going to be frustrating knowing that you’ve got a clinic
282 here and you can’t get up to the ward and there’s an urgent stroke survivor.”
283 (Heather, orthoptist in an eye clinic)

284 “Erm... it sounds awful doesn’t it when you’re constantly having to slim down
285 what you do.” (Melissa, occupational therapist in an acute stroke service)

Lack of Training

Training is another theme that frequently emerged across both professions. Visual perception difficulties following stroke appear to be only briefly covered in the undergraduate education of orthoptists and occupational therapists, and is mostly learned about post-qualification through on-the-job-training. Many occupational therapists mentioned that they rotate between services, leaving little time for them to be trained in many aspects of stroke care. In particular, they said that this lack of experience made them feel uncertain when it comes to vision and visual perception deficits following stroke. Inexperience and limited training seemed to be an important hurdle for occupational therapists in visual perception screening. Because vision and visual perception was not well understood by the staff (see section ‘Lack of clarity around visual perception’), participants reported that mistakes in assessments are commonly made, leading to a potential under-diagnosis of visual perception deficits.

“*[It]* can be a bit frustrating ... lots of the occupational therapists are on a rotational job. So getting them *[occupational therapists]* trained up *[is difficult]*, I think they *[occupational therapists]* do find vision scary.... Vision, it’s the scariest thing about stroke.” (Stephanie, orthoptists in an eye clinic, using ‘vision’ as a general term to cover a wide range of sensory and perceptual visual functions)

“*[Visual perception is]* one of those things that is perhaps not widely understood.... think unless you’ve got a unit with a special interest in stroke and you’ve taken the time to look into that... then it’s perhaps under-diagnosed as well, so yeah I think it’s recognition that it’s a potential problem.” (Amanda, orthoptist in an eye clinic)

“The key is really, somebody in the hospital to be able to deliver the training and the therapy staff to actually take on board the training and to actually deliver it.” (Sarah, orthoptist in an eye clinic)

“I think it probably is underreported; the amount of incidents of perceptual problems post stroke.” (Rachel, orthoptist in an eye clinic)

Environmental and Stroke Survivor Factors

Participants reported that visual perception screening is limited by the environment in which they take place. They said that in acute stroke care, most assessments are done in a hospital bay with several beds. Even if curtains can be drawn around the stroke survivor’s bed, distractions were reported to be numerous: sounds from neighbouring beds, interruptions for timed caring needs, distractions from medical equipment around the bed and presence of personal objects on a small table in front of the stroke survivor (spectacles, drinks, food, tissues, personal care items, newspaper, get well cards, crossword book, pen, etc.) can be visually distracting and interfere with visual perceptual assessment.

“Frustrating for us is ... that we .. depending on which unit we are *[on]* or where we are, it can be quite difficult to do *[our assessment at]* the bedside if there is a lot *[of]* distractions going on; if there is no quiet space to take the patient. Obviously, yes, we can draw the curtains but it’s sometimes more the noise levels than everything. They get easily distracted. That makes it quite hard actually to do an assessment.” (Amber, occupational therapist in a stroke rehabilitation service)

“It’s *[the]* environment. Often *[it]* doesn’t lend itself to a nice quiet space where patients can attend.” (Danielle, occupational therapist in an acute stroke service)

Participants also mentioned the condition of the stroke survivors as a limiting factor to visual perception screening. Both professions reported that tests and tools are often not suitable for stroke survivors with poor alertness, poor cognition, aphasia or severe weakness in their upper limbs. For instance, stroke survivors may not be able to answer a question on whether their vision has changed because they developed aphasia following stroke. Similarly, copying

and cancellation tasks to assess visual neglect were reported to be unsuitable for a stroke survivor who can no longer hold a pen.

“Another component would be around the client or the patient themselves in terms of factors that might work against screening: so attention, concentration, fatigue, the ability to deliver a verbal response in terms of aphasia or dysarthria difficulties. ... That has significant impact on our ability to undertake formal testing of whatever nature.” (Christopher, occupational therapist in an acute stroke service)

“Our patients who either are fatigued quite quickly, or are ... medically not well enough to do a lot of taking down to the therapy department ..., or their attention span is quite limited, or they’ve got a weakness in their upper limb that makes it more difficult to complete the tests.” (Danielle, occupational therapist in an acute stroke service)

Continuation of Care

A further concern of the participants was the follow-up process after screening for visual perception deficits. They highlighted that quite often the information from screening is not passed on to the next care team. Therefore, even if visual perception problems are picked up in an acute setting, no follow-up assessments are carried out or no rehabilitation is provided in their experience. With respect to rehabilitation, occupational therapists found it frustrating that few treatment options are available. They reported that little information and few guidelines are available on how visual perception problems can be treated or managed. Some noted that the lack of options for rehabilitation reduced their motivation to screen for visual perception problems.

“Yes, and it [*the discharge summary*] is very medical because the occupational therapists or myself don’t like the discharge summaries. ... We do all this

information and then it gets lost Yeah, it's not part of our discharge summary in enough detail, so when they go somewhere new which could be two or three weeks later ... they don't have that information." (Matthew, orthoptist in an eye clinic)

"It might be because it [*visual perception deficits*] is limited in treatment ... that's why it's not the focus. So ... it would be more useful to spend time during my assessment speaking to them about adapting to their field loss and giving them a prism or occlusion for their double vision." (Rachel, orthoptist in an eye clinic)

Test Characteristics

Both professions highlighted limitations in the characteristics of the available tests for visual perception. Occupational therapists more frequently reported to be hindered by practical limitations and the unknown impact on daily life. With respect to the practical limitations, they referred to a large number of test materials like stimulus booklets, stopwatch, cubes, etc. needed for assessment or too many loose pages. With respect to the unknown impact on daily life, they referred to stimuli and tests that were thought to be limited in ecological validity. They considered the tests too abstract with no clear link to the implications for everyday tasks.

"We have our little screening tool. It's a few pages, just to prompt us That's all right but then ... everything else you are ... searching together from different kits. ... That's a little bit frustrating." (Amber, occupational therapist in a stroke rehabilitation service)

"There's an awful lot of bits of paper and a bit of faffing about and preparing to get it all ready before we actually go out to see the patients." (Melissa, occupational therapist in an acute stroke service)

384 “We use Rivermead Perceptual Battery. But to be fair, I don’t tend to use it as
 385 often because I’d rather do it in function. ...It’s more meaningful, especially with
 386 stroke patients because it makes more sense. Rather than getting them to balance
 387 cubes on each other or, you know I can say “Can you find the toothpaste? Can
 388 you take the lid off? Can you brush your own teeth?” and then it just makes more
 389 sense. And then ... it’s easier for them to identify *[a]* goal.” (Megan, occupational
 390 therapist in a stroke rehabilitation service)

391 Orthoptists, on the other hand, seemed to emphasize more the lack of evidence and
 392 consequently the lack of specific clinical guidelines on how to assess visual perception.

393 “If you see the national guidelines for visual problems after stroke, they have a
 394 really small section on vision with orthoptics but we don’t really have anything in
 395 it that’s specific to what we should be screening them with and what we should be
 396 treating them with. So even though we’re using evidence-based practice, the
 397 evidence sometimes isn’t great and I think that’s probably why it’s not in the
 398 guidelines because there’s not really good evidence to support it. But sometimes
 399 you’re seeing a patient and you’re not entirely sure if you’re doing the right thing
 400 because you’re maybe giving them information based on clinical experience, what
 401 other people have been doing but you’re not really sure if that is the best thing to
 402 be doing or could you be doing something else with them.” (Amy, orthoptist in an
 403 eye clinic)

404

Table 3. Challenges in visual perception screening

Major themes	Subthemes	Endorsed by occupational therapists	Endorsed by orthoptists
Lack of time	Limited time with stroke survivor due to discharge pressure	x	
	Limited time with stroke survivor due to staff shortage	x	x
Lack of training	Inexperienced staff with limited training	x	
	Vision and visual perception is not well understood	x	x
	Mistakes made in assessment	x	x
	Vision and visual perception deficits are underdiagnosed	x	x
Environmental and stroke survivor factors	Condition of stroke survivor (e.g. low alertness, poor cognition, aphasia, not mobile)	x	x
	Distractions in test environment (often in hospital bay)	x	x
Continuation of care	Information on vision and visual perception problems gets lost after discharge	x	x
	No information on treatment or management of vision or visual perception problems	x	
Test characteristics	Test materials not readily available or making use clumsy	x	
	Lack of evidence-based screening instruments		x
	No specific recommendations in clinical guidelines		x
	Tests are not meaningful for everyday life	x	

Facilitators

The facilitators to practice that participants reported, largely mirrored the challenges reported above (table 4). We identified three major themes and 12 subthemes.

Practical Tool

On a practical level, participants preferred a tool that can be administered in approximately 10-20 minutes, so it can fit in with busy every-day practice and current staff levels. They also favoured a tool that consists of few test materials and pages to prevent test materials getting lost or mixed up during testing. According to participating orthoptists, the tool should have a 1 to 2 page proforma. They reported that it should include questions to ask the stroke survivor and simple quick tests for a range of functions. They suggested that the proforma could subsequently be used in a referral for orthoptic input. To allow bedside testing or testing in the community, participants preferred portable tools. Lastly, many participants highlighted that a tool should ideally be suitable for all stroke survivors, such as those with aphasia, poor concentration, poor cognition or severe fatigue.

421 “It needs to be something that’s fairly quick and easy to administer.” (Danielle,
 422 occupational therapist in an acute stroke service)

423 “Just pen and paper, quite simple or a few things to hand that will fit in something
 424 that’s quite portable and mobile, I think [*that*] would be handy, yes.” (Brittany,
 425 occupational therapist in a stroke rehabilitation service)

426 “Reasonably short ... for logistical [*reasons*]: ... our time and patient
 427 concentration time, fatigue in the more acute stages. I think really half an hour is
 428 the maximum time.” (Christopher, occupational therapist in an acute stroke
 429 service)

430 “It would have to be a list of standardised questions: Are they able to look
 431 between two points specifically? ... I guess the visual perception question would
 432 be if they’re following a moving object? ... You know are they able to identify
 433 their friends or their family? Can they recognise themselves in a photo?” (Rachel,
 434 orthoptist in an eye clinic)

435 *Training and Experience*

436 Orthoptists and occupational therapists agreed on the importance of training and experience
 437 for robust visual perception screening. The presence of experienced colleagues or training by
 438 other staff members was seen as very advantageous. Given the limited training in visual
 439 perception highlighted above, a tool that is easy to administer and score was preferred by our
 440 participants. Similarly, they reported that results should be easy to interpret, and should
 441 provide clear guidance for management of the deficit (if present).

442 “If it was something that was quite self-explanatory then anybody could do it and
 443 therefore anybody can interpret it, then that would be much clearer.” (Amy,
 444 orthoptist in an eye clinic)

“Just easy to administer and easy to score. ... So making instructions clear and easy to understand would be good.” (Emily, occupational therapist in a stroke rehabilitation service)

Staff Roles and Interactions

Occupational therapists reported that their preferred practice is to screen stroke survivors for vision and visual perception problems during their functional assessments (e.g. washing, making a cup of tea) or via a screening tool. Both professions mention that if a problem is flagged, more in-depth assessment can then take place.

“Whereas since we’ve set up a proper vision screening it just means that things have been a lot more efficient and that’s made a much better service because they’re not having to see a lot of people that don’t need to be seen.” (Nicole, orthoptists in an eye clinic)

“So, I’m happy with the main basic screen to begin with, and if it digs something up then we’ll dig it in some more detail. I think that works quite well.” (Matthew, orthoptists in an eye clinic)

If the problem concerns visual perception difficulties, an occupational therapist is seen by our participants as the most appropriate professional to follow it up. If the problem concerns sensory vision, a referral to an orthoptist was the preferred practice in our sample. Good relationships between stroke occupational therapists and orthoptists in eye clinics were frequently mentioned as invaluable.

“I think the key is having the staff, the occupational therapist and the physio staff actually take that on board.” (Sarah, orthoptist in an eye clinic)

Table 4. Facilitators to visual perception screening

Major themes	Subthemes	Endorsed by occupational therapists	Endorsed by orthoptists
Practical tool	Quick to administer (10-20 min)	x	x
	Paper and pen based without too many test materials or pages	x	x
	Proforma with questions and simple tests to be used in referral		x
Training and experience	Portable	x	
	Suitable for stroke survivors with poor concentration, communication and fatigue	x	
	Easy to administer and score	x	x
	Easy to interpret with management advice included	x	x
Staff roles and interactions	Trained and experienced staff	x	x
	Screening for visual and visual perception problems by occupational therapists during functional assessment or with screening tool	x	x
	In depth assessment with range of complimentary tests	x	x
	Specialised staff for visual (orthoptists) and visual perceptual (occupational therapists) problems		x
	Good relationship between stroke and eye departments		x

Discussion

The aim of this qualitative research was to gain a deeper understanding of current practice in visual perception screening after stroke, the challenges faced by health care professionals. In our participants' stroke care services, occupational therapists most often screened for visual and visual perceptual difficulties. If visual difficulties like hemianopia, double vision or reduced acuity, are found, a referral to an orthoptist is made for an in depth assessment. This referral strategy corresponds to an earlier report of 76.7% of stroke units having access to an orthoptic stroke service.(20) If visual neglect is suspected, occupational therapists will often follow this up themselves with a range of tasks like cancellations, copying, line bisection, etc. The biggest challenges faced by our participants were lack of time and training. Facilitating factors mentioned were simple, easy to use instruments and good relationships between stroke and eye departments.

The qualitative approach of our study allowed us to gather views directly and in depth. Instead of reporting frequency of views (e.g. how many health care professionals report staff shortage), we focus on how central certain views are to a person's experience. Our sample covers the two main professions involved in visual perception screening after stroke.

National Clinical Guidelines recommend that every stroke survivor who appears to have perceptual difficulties should have an assessment with standardised measures.(2) As is evident from our interviews, only a small proportion of stroke survivors receive an assessment with standardised measures for visual perceptions difficulties, and screening is not routinely done. The means of assessment of visual perception deficits varied widely between participants. Orthoptists will often use standardised tools in their assessments, though these are not necessarily validated with stroke survivors.(21,22) Overall, standardised assessments were rarely mentioned by our participants. Occupational therapists expressed preferring assessment during functional tasks like washing and dressing and assessment with in-house developed screening checklists. This may be due to a lack of a validated screening tool for a wide range of visual and visual perceptual deficits that can be used by any health care professional.(21) The diversity in tools that we observed in our group of English participants contrasts with the reported use of standardised tools in an Australian population where the Occupational Therapy Adult Perceptual Screening Test and the Loewenstein Occupational Therapy Cognitive Assessment are most commonly used to evaluate visual perceptual difficulties.(22)

McCluskey and colleagues investigated the barriers and enablers for following recommended practice in stroke care.(23) In line with our findings the occupational therapists they interviewed reported the condition of the stroke survivors as a barrier for standardised assessments. Many tests are not designed and not suitable for stroke survivors with aphasia and/or dysarthria. Occupational therapists in their study also mentioned the lack of training

and knowledge, specifically for visual neglect rehabilitation. Time pressure and fluctuating staff levels meant occupational therapists in their study prioritised assessments and interventions which would produce the best clinical outcome. A lack of time to see stroke survivors was also mentioned by their orthoptists. However, in our study this referred to the lack of time to see all referred stroke survivors before they were discharged, while in the study by McCluskey, the orthoptists referred to the lack of time to treat stroke survivors. The standardised assessments used by Australian occupational therapists, Occupational Therapy Adult Perceptual Screening Test and the Loewenstein Occupational Therapy Cognitive Assessment, require training and take 20-45 minutes to complete. Given the frequently reported lack of training and time, it may appear that the standardised tools recommended by clinical guidelines are not adapted to the clinical reality of the National Health Service in England. Although McCluskey's study was limited to one Australian hospital and only involved 5 occupational therapists and 2 orthoptists, all working in the same service and circumstances, the barriers are remarkably similar to what our 12 occupational therapists and 13 orthoptists reported experiencing in 16 English healthcare services. The stroke survivor's condition, lack of time and staff seem to be universal barriers for providing evidence-based stroke care across domains, not just visual perception screening. Also, the lack of information on treatment or management of vision or visual perception problems that occupational therapists mention as a barrier is not surprising given the very limited evidence for treatment and management options.(24)

The first challenge exposed by our study is the variation between participants in their understanding of constitute visual perception problems after stroke. This led to participants answering interview questions with different reference frameworks. When sharing the challenges of visual perception screening in their clinical practice, some might have been considering challenges in just screening for visual inattention difficulties while others

considered a broad range of sensory and cognitive visual impairments. To maintain an equal relationship between participant and interviewer, and an open non-judgemental environment to freely share their experiences, participants' definition of visual perception was not challenged during the interview. A disadvantage of this approach is that it is unclear whether certain themes apply specifically to one aspect of visual and visual perception screening or to the whole range of vision related difficulties after stroke.

The second challenge lies in our sample size. Our sample, though the largest for a qualitative study in this topic, is not representative for the whole population of health care professionals involved in visual perception screening after stroke. We limited ourselves to occupational therapists and orthoptists as they are most commonly involved in visual perception screening in England. Experiences might be very different in clinical settings without an orthoptic department or with more involvement of neuropsychologists in stroke care. In addition, voluntary participation and asking our participants to commit 30 minutes of their time might have induced a bias for allied health professionals with a keen interest in the topic. We have possibly missed opinions from people with a more negative or neutral attitude towards visual perception screening. In addition, newly qualified health care professionals with limited experience in stroke care might not have felt confident enough to share their opinions on the topic. Lack of confidence on this topic with junior staff was brought up by several of our participants. However, the aim of this qualitative research was not to generalise findings to the population, but rather to provide an in-depth exploration of the topic to generate hypotheses. We are following this research up with a large scale survey for all health care professionals involved in visual perception screening in the United Kingdom and Republic of Ireland. In the recruitment to this survey study, we are emphasizing that we welcome the opinions of health care professionals with all levels of experience.

A third potential limitation is that the interviewer was the principal investigator of the study and was therefore not independent. Unconscious bias of the interviewer might have guided the participants' answers. We have tried to minimize this by using a detailed interview guide that included a list of cues that could be given. In addition, all interviews were recorded and a second coder was involved.

Our study limits itself to health care professionals' perspective on visual perception screening after stroke. We have not explored experiences by stroke survivors, although they can be substantially different. Also, we have not investigated the management of visual perceptual problems after stroke of their impact in daily life as is previously reported for sensory visual problems. (25–27)

The current research highlighted that a clear and consistent definition of visual perception should be provided to orthoptists and occupational therapists. More training is needed on the assessment (and management) of visual perceptual screening. This is preferably provided via professional organisations to ensure consistency and in line with national guidelines(2,28).

This will raise awareness and reduce insecurity experienced by junior staff members.

Establishing good relationships between staff at stroke and eye departments is strongly recommended. We suggest this could be achieved in the form of knowledge exchanges or via a re-evaluation of the existing referral system. Lastly, a quick and portable visual perception screening tool that is easy to administer, score and interpret would highly benefit both staff and stroke survivors if it is inclusive for stroke survivors with aphasia, motor impairments and cognitive problems. The tool should be evidence-based and self-explanatory to use. Such a tool is currently not available. Our research provides test developers with the requirements of occupational therapists and orthoptists for such a screening tool to be clinically useful and thereby increase the likelihood of adoption.

580 **Acknowledgments**

581 We would like to express our sincere gratitude to all the participating allied health
582 professionals who gifted their time. In addition we would like to thank Philip Clatworthy and
583 Claire Murray for facilitating this project. This work was supported by the Stroke Association
584 under TSA PDF 2017/03; and under TSA LECT 2015/02.

585 **Declaration of interest statement**

586 The authors report no conflicts of interest.

587 **Data availability statement**

588 Given the small group of stroke orthoptists and stroke occupational therapists in England,
589 sharing the interview transcripts would potentially disclose participants' identity.

References

1. Bouska MJ, Kauffman NA, Marcus SE. Disorders of the visual perceptual system. In: Umphred DA, editor. Neurological Rehabilitation. 2nd editio. St. Louis, MO: The C. V. Mosby Company; 1990. p. 705–40.
2. Intercollegiate Stroke Working Party. National clinical guideline for stroke. 2016.
3. Kolb B, Whishaw IQ. Fundamentals of human neuropsychology. 7th editio. New York, NY: Worth Publishers; 2003.
4. Rowe FJ, Hepworth LR, Howard C, Hanna KL, Cheyne CP, Currie J. High incidence and prevalence of visual problems after acute stroke: An epidemiology study with implications for service delivery. PLoS One [Internet]. 2019 [cited 2019 Jul 2];14(3):e0213035. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30840662>
5. Làdavias E. The Role of Visual Attention in Neglect: A Dissociation between Perceptual and Directional Motor Neglect. Neuropsychol Rehabil. 1994;4(2):155–9.
6. Mancini F, Bricolo E, Mattioli FC, Vallar G. Visuo-haptic interactions in unilateral spatial neglect: The cross modal Judd illusion. Front Psychol. 2011;2(NOV).
7. Chatterjee A, Thompson KA, Ricci R. Quantitative analysis of cancellation tasks in neglect. Cortex. 1999;35(2):253–62.
8. Rowe FJ, Brand D, Jackson CA, Price A, Walker L, Harrison S, et al. Visual impairment following stroke: Do stroke patients require vision assessment? Age Ageing. 2009;38(2):188–93.
9. Edmans J, Lincoln N. The frequency of perceptual deficits after stroke. Clin Rehabil [Internet]. 1987 Nov 1;1(4):273–81. Available from: <http://cre.sagepub.com/cgi/doi/10.1177/026921558700100403>

- 613 10. Kerkhoff G. Neurovisual rehabilitation: Recent developments and future directions. J
614 Neurol Neurosurg Psychiatry. 2000;68(6):691–706.
- 615 11. Jehkonen M, Ahonen JP, Dastidar P, Koivisto AM, Laippala P, Vilkkilä J, et al. Visual
616 neglect as a predictor of functional outcome one year after stroke. Acta Neurol Scand
617 [Internet]. 2000 Mar [cited 2016 Dec 8];101(3):195–201. Available from:
618 <http://www.ncbi.nlm.nih.gov/pubmed/10705943>
- 619 12. Plante M, Demers L, Swaine B, Desrosiers J. Association Between Daily Activities
620 Following Stroke Rehabilitation and Social Role Functioning Upon Return to the
621 Community. Top Stroke Rehabil. 2010;17(1):47–57.
- 622 13. Mercier L, Audet T, Hébert R, Rochette a, Dubois MF. Impact of motor, cognitive,
623 and perceptual disorders on ability to perform activities of daily living after stroke.
624 Stroke. 2001;32(11):2602–8.
- 625 14. Kalra L, Smith DH, Crome P. Stroke in patients aged over 75 years: outcome and
626 predictors. Postgrad Med J [Internet]. 1993 Jan 1;69(807):33–6. Available from:
627 <http://pmj.bmj.com/cgi/doi/10.1136/pgmj.69.807.33>
- 628 15. Bazeley P. Qualitative data analysis with NVivo. SAGE; 2007. 217 p.
- 629 16. Krueger RA, Casey MA (Mary AW. Focus groups□: a practical guide for applied
630 research. SAGE; 2009. 219 p.
- 631 17. Strategyzer | Value Proposition Canvas [Internet]. [cited 2019 Jul 26]. Available from:
632 <https://www.strategyzer.com/canvas/value-proposition-canvas>
- 633 18. Braun V, Clarke V. Using thematic analysis in psychology. Qual Res Psychol
634 [Internet]. 2006 Jan;3(2):77–101. Available from:
635 <http://www.tandfonline.com/doi/abs/10.1191/1478088706qp063oa>

- 636 19. Maxwell JA. Using Numbers in Qualitative Research. Qual Inq [Internet]. 2010 Jul
637 15;16(6):475–82. Available from:
638 <http://journals.sagepub.com/doi/10.1177/1077800410364740>
- 639 20. Hepworth L, Rowe F. Ten Years On – A Survey of Orthoptic Stroke Services in the
640 UK and Ireland. Br Ir Orthopt J [Internet]. 2019 May 8;15(1):89–95. Available from:
641 <http://www.bioj-online.com/articles/10.22599/bioj.135/>
- 642 21. Hanna KL, Hepworth LR, Rowe FJ. Screening methods for post-stroke visual
643 impairment: a systematic review. Disabil Rehabil [Internet]. 2016 Sep 26;1–13.
644 Available from:
645 <https://www.tandfonline.com/doi/full/10.1080/09638288.2016.1231846>
- 646 22. Kitsos G, Harris D, Pollack M, Hubbard IJ. Assessments in Australian stroke
647 rehabilitation units: a systematic review of the post-stroke validity of the most
648 frequently used. Disabil Rehabil [Internet]. 2011 Jan 9 [cited 2019 Aug 5];33(25–
649 26):2620–32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21554012>
- 650 23. McCluskey A, Vratsistas-Curto A, Schurr K. Barriers and enablers to implementing
651 multiple stroke guideline recommendations: a qualitative study. BMC Health Serv Res
652 [Internet]. 2013 Dec 19 [cited 2019 Aug 5];13(1):323. Available from:
653 <http://www.ncbi.nlm.nih.gov/pubmed/23958136>
- 654 24. Bouwmeester L, Heutink J, Lucas C. The effect of visual training for patients with
655 visual field defects due to brain damage: A systematic review. J Neurol Neurosurg
656 Psychiatry. 2007;78(6):555–64.
- 657 25. Hepworth LR, Rowe FJ. Visual Impairment Following Stroke - The Impact on Quality
658 of Life: A Systematic Review. Ophthalmol Res An Int J [Internet]. 2016;5(2):1–15.
659 Available from: <http://sciencedomain.org/abstract/12864>

- 660 26. Pollock A, Hazelton C, Brady M. Visual Problems After Stroke: A Survey of Current
661 Practice by Occupational Therapists Working in UK Stroke Inpatient Settings. Top
662 Stroke Rehabil. 2011;18(sup1):643–51.
- 663 27. Rowe FJ. Stroke survivors’ views and experiences on impact of visual impairment.
664 Brain Behav [Internet]. 2017 Sep;7(9):e00778. Available from:
665 <http://doi.wiley.com/10.1002/brb3.778>
- 666 28. NATIONAL INSTITUTE FOR HEALTH AND CARE EXCELLENCE (NICE) .
667 Stroke Rehabilitation: Long term rehabilitation after stroke [Internet]. NICE. Londo;
668 2013. Available from: <http://www.nice.org.uk/nicemedia/live/14182/64094/64094.pdf>
669