

Supplements

Figure S1: Sensitivity to severity and duration of stay in ICU. **A.** Projected number of ICU beds according to different severity scenarios: baseline ($p_H = 1.1\%$, red), low ($p_H = 0.6\%$, green), and high ($p_H = 1.8\%$, blue). **B.** Projected number of ICU beds according to different durations of stay in ICU: baseline ($\tau_{ICU} = 11.4$ days, red), short ($\tau_{ICU} = 8.0$ days, brown), and long ($\tau_{ICU} = 15.0$ days, purple). Black dots indicate data used to calibrate the models, while empty circles denote data not available at the time of the analyses.




Figure S2: Model selection. DIC difference (DIC of model M1 - DIC of model M2) for models calibrated from June 19th to June 29th. The dashed lines indicate a DIC difference of 4 units. Model M2 has a change point on June 15th (Figure S3).




Figure S3: Choice of transmission rate change point for model M2. Model DIC for change points ranging from June 6th to June 26th. The lowest DIC is obtained for a change point on June 15th. The dashed lines indicate a DIC difference of 4 units.




Figure S4: Comparison between the final version of M2 and a version of M2 that includes age structure. Solid blue lines indicate model M2 posterior means while color areas indicate 95% credible intervals. The black lines denote trajectories obtained with the model that includes age structure: these were simulated by using the parameters' posterior means obtained by calibrating M2 to data available on August 25th.




Table S1: Key Model Parameters

Parameter	Model M1	Model M2	Model M2 (Final)
<i>Time spent in hospital</i> (τ_H)	12.8 days ^b	11.8 days ^b	11.8 days ^b
<i>Delay hospital to ICU admission</i> ($\tau_H - \tau_{ICU}$)	1.5 days ^a	2.6 days ^b	2.6 days ^b
<i>Time spent in ICU</i> (τ_{ICU})	11.4 days ^b	17.6 days ^b	15.0 [13.1, 17.4] days
<i>Probability of ICU given hospitalization</i> (p_{ICU})	21.5% ^b	11.0% ^b	15.7% [13.9%, 17.6%]
<i>Reproduction number</i>	1.35 [1.26, 1.45] (before 5/20) 1.78 [1.68, 1.88] (after 5/20)	1.40 [1.32, 1.49] (before 5/20) 1.71 [1.65, 1.77] (from 5/20 to 6/15) 1.14 [0.95, 1.31] (after 6/15)	1.44 [1.36, 1.53] (before 5/20) 1.69 [1.65, 1.73] (from 5/20 to 6/15) 1.08 [1.07, 1.10] (after 6/15)

^a Value from [Salje et al. 2020](#), estimated using data from mainland France.

^b Values estimated using a mixture distribution as described in [Lefrancq et al. 2020](#) and using data from French Guiana available on July 2nd 2020.

All other values were estimated with our compartmental model.