1 Successive epidemic waves of cholera in South Sudan, 2014 - 2017

2 Forrest K Jones¹, Joseph F Wamala², John Rumunu³, Pinyi Nyimol Mawien³, Kol Mathew Tut³,
3 Shirlee Wohl¹, Lul Deng³, Lorenzo Pezzoli⁴, Linda Haj Omar⁵, Justin Lessler¹, Marie-Laure
4 Quilici⁶, Francisco J Luquero⁷, Andrew S Azman^{1,8}

5

6 ¹ Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health, Baltimore,
7 USA

8 ² World Health Organization, Juba, South Sudan

9 ³ South Sudan Ministry of Health, Juba, South Sudan

10 ⁴ World Health Organization, Geneva, Switzerland

11 ⁵ World Health Organization, Brazzaville, Republic of Congo

12 ⁶ Institut Pasteur, Paris, France

13 ⁷ Epicentre, Geneva, Switzerland

14 ⁸ Médecins Sans Frontières, Geneva, Switzerland

15

16 Corresponding author: Andrew Azman (azman@jhu.edu)

17

18

19

20

21

22 **Abstract**

23

24 *Background*

25 Between 2014 and 2017, successive cholera epidemics occurred in South Sudan within the
26 context of civil war, population displacement, flooding, and drought. Understanding the
27 determinants of cholera spread in complex settings like this can provide valuable insights for
28 mitigating future cholera risk.

29

30 *Methods*

31 We analyzed cholera linelist and molecular data to describe the spatio-temporal progression of
32 the epidemics. We explored the role of rainfall, population movement and vaccination
33 campaigns in shaping the explaining incidence and the spatial distribution of reported cases.

34

35 *Findings*

36 South Sudan experienced three distinct cholera epidemic waves of cholera ranging from 6-18
37 months with more than 28,000 cases reported and more than 2 million cholera vaccine doses
38 delivered to curb transmission. The 2014 and 2015 epidemics remained spatially limited while
39 the 2016/17 epidemic exploded along the Nile river. Initial cases of each epidemic were
40 reported in or around Juba soon after the start of the rainy season, but we found no evidence
41 that rainfall modulated transmission during each epidemic. All isolates analyzed had similar
42 genotypic and phenotypic characteristics, closely related to sequences from Uganda and
43 Democratic Republic of Congo. The direction of large-scale population movements between
44 counties with cholera outbreaks was consistent with the spatial distribution of outbreaks. As of
45 September 2020, zero cholera cases have been confirmed within South Sudan since 2017.

46

47 *Interpretation*

48 The three epidemic waves were caused by *V. cholerae* of the same clonal origin despite the
 49 periods of no reported cases between waves. While the complex emergency likely shaped
 50 some of the observed spatial and temporal patterns of reported cases, the full scope of
 51 transmission determinants remains unclear. Timely and well targeted use of cholera vaccine
 52 can avert cases and deaths, however, most of the vaccine campaigns occurred after the
 53 epidemic peak highlighting the challenges of delivering vaccines quickly in response to an
 54 outbreak in settings like South Sudan. These analyses provide a multi-faceted template for
 55 examining cholera dynamics through epidemiological, microbiological, climatic, and behavioral
 56 lenses.

57

58 *Funding*

59 *The Bill and Melinda Gates Foundation*

60 **Research in context**

61

62 *Evidence before this study*

63 We searched PubMed for articles published between January 1st, 1970 and September 18th,
64 2020 using "cholera"[title] AND (epidemic OR outbreak) AND (Africa OR emergency OR
65 conflict) AND (molecular OR sequencing OR rain OR precipitation OR mobility OR movement
66 OR vaccine OR WASH). We also downloaded reported cases and deaths from the Global
67 Health Observatory Data repository on all outbreaks in the geographical area of South Sudan.
68 Across the globe, there are an estimated 2.9 million cases and 95,000 deaths annually. Many
69 of these cases occurred in areas with poor infrastructure including peri-urban slums, rural
70 areas dependent on surface water, and areas with complex emergencies .

71

72 Cholera has been introduced into the African continent several times over the past 50 years. In
73 particular, endemic transmission has been maintained in several countries bordering the
74 African Great Lakes, including the Democratic Republic of Congo, Uganda, Tanzania, and
75 Malawi. In contrast, reported cholera in South Sudan appears in cycles, with few reported
76 outbreaks prior to 2005, a period of annual epidemics reported between 2005 and 2009, and
77 then no reported cases between 2010 and 2014. The South Sudanese Civil War began in
78 December 2013, leading to damaged essential infrastructure and displaced hundreds of
79 thousands of individuals and a return of cholera, in successive annual waves starting in 2014.
80 Previous studies have demonstrated associations between epidemic cholera and complex
81 emergencies (e.g., Yemen, Haiti, Democratic Republic of the Congo), mass population
82 movement (e.g., Senegal), and precipitation (e.g., Yemen and Haiti); however it is unclear how
83 these factors interact to shape cholera epidemics. Oral cholera vaccination campaigns have
84 only recently been used both in response to epidemics and preventatively though its impact at
85 the population level has not been well described.

86

87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113

Added value of this study

Our study helps explain the complex story of three epidemic waves of cholera in South Sudan that began soon after the beginning of the civil war. We showed how each wave began in the capital city of Juba at the start of the rainy season and spread to different areas of the country. Using molecular data, isolates from all four years were found to be from the same clonal origin. We presented evidence, including the analyses of previously published whole genome sequences, that cholera was likely introduced in 2013 or 2014 from Uganda or the Democratic Republic of Congo and may have been re-introduced through multiple cross-border transmission events. We identified large-scale population displacement and movement as partially explaining the differences in the number of cases between years and transmission during the dry season of the 2016-2017 wave. Lastly, we describe 36 vaccination campaigns providing over 2 million doses of cholera vaccine: we find that early vaccination may have reduced local outbreak sizes though most vaccination activities occurred after the peak week of cases.

Implications of all the available evidence

Our research highlights the need for regional-level responses to curb outbreaks of cholera in humanitarian settings. In addition to oral cholera vaccine campaigns and interventions to improve water, sanitation, and hygiene in these vulnerable settings, controlling cholera in nearby countries that have the potential to introduce cholera might be an effective additional strategy to reduce cases. Increased sampling of *Vibrio cholerae* from both human cases and environmental samples combined with whole genome sequencing can help us better understand how cholera spreads across borders and maintains itself in the environment. Expanding the use of sequencing and bioinformatics in more countries with *Vibrio cholerae*

114 transmission could lead to improved global representativeness of sequences and integration of
 115 results into local decision-making processes. While increased precipitation has been linked to
 116 increased cholera transmission in many settings, we found that it was associated with the
 117 onset of outbreaks, but not with increases in transmission once an outbreak has begun. While
 118 our study adds to the evidence on the role of population movement in cholera outbreaks,
 119 improved methods for measuring population movement within and between countries during
 120 complex emergencies is needed to better characterize these relationships.

121

122

123

124 Introduction

125

126 The seventh cholera pandemic, first recognized in 1960, continues to plague populations with
127 poor access to safe water and sanitation across the world. Although estimates of the global
128 burden are uncertain, over 140,000 suspected cholera cases are reported annually from Sub-
129 Saharan Africa.¹ Universal access to water and sanitation would likely eliminate cholera
130 transmission but is unlikely given the current pace of progress and financial commitments.²

131

132 In 2017, the World Health Organization (WHO) led Global Task Force on Cholera Control
133 (GTFCC) developed a roadmap to end cholera by 2030 and in 2018 a resolution to end cholera
134 was adopted at the 71st World Health Assembly.³ The roadmap calls for a geographically-
135 targeted approach to use limited resources in areas with a high-risk of cholera, requiring an in-
136 depth understanding of transmission across endemic and epidemic settings. South Sudan is
137 one of the 47 focal countries of this roadmap. Detailed analyses of historical cholera in South
138 Sudan, which has suffered from civil-war, mass population movements, floods and droughts,
139 can provide important lessons for cholera control in similar epidemic-prone settings.

140

141 Three successive epidemic waves of cholera occurred in South Sudan between 2014 and
142 2017, leading to tens of thousands of cases and hundreds of deaths, many among internally
143 displaced persons (IDPs).⁴⁻⁶ Each wave varied greatly by geographic extent, timing, and
144 magnitude and were separated by periods, with few to no reported cases. It is unclear whether
145 each outbreak was the result of a new introduction of *Vibrio cholerae* O1 or whether cholera
146 was present but undetected between outbreaks. Here, using detailed cholera case data
147 collected from 2014 through 2017, we describe the key features of the successive epidemic
148 waves and explore how precipitation, population displacement, and vaccination campaigns
149 may explain the differences between waves.

150

151 **Methods**

152

153 ***Cholera surveillance system***

154 The Ministry of Health Republic of South Sudan (MoH) in collaboration with local health
155 authorities implemented cholera surveillance through both the Integrated Disease Surveillance
156 and Response (IDSR) and Early Warning Alert and Response Network (EWARN) with support
157 from WHO and the Health Cluster. Linelist data were collected from all sites treating cholera
158 patients across the country, including cholera treatment centers and oral rehydration points
159 from June 2014 to December 2017. Prior to the confirmation cholera, a suspected cholera case
160 was defined as any patient aged ≥ 5 years who developed severe dehydration or died from
161 acute watery diarrhoea. After cholera was confirmed within a county, a suspected cholera case
162 was any patient ≥ 2 years who developed acute watery diarrhoea. A confirmed case was a
163 suspected case with culture-confirmed *V. cholerae* O1. We use the term epidemic waves to
164 describe a rise and decline of reported cholera cases at the national-level, which is a different
165 concept than the previously described 'waves' of the 7th pandemic ⁷.

166

167 ***Laboratory analysis***

168 When suspected cases were first reported in a county, stool specimens from select suspected
169 cases were sent to the National Public Health Laboratory (NPHL) for culture confirmation.⁸
170 Additional isolates were collected from suspected cases and tested by culture at the NPHL in
171 2015 in a vaccine effectiveness study.⁹ Overall, 150 isolates from South Sudan (and 10 from
172 Uganda) were sent to the French National Reference Center for Vibrios and Cholera (Institut
173 Pasteur, Paris, France) for serotype, antibiotic resistance pattern testing (disk diffusion
174 following Comité de l'Antibiogramme de la Société Française de Microbiologie 2013 standards
175 for Enterobacteriaceae) and Etests (AB bioMérieux, Solna, Sweden) for the determination of

176 minimum inhibitory concentrations (MICs) of nalidixic acid and ciprofloxacin), genotyping tests
177 (*rstR*, *ctxB*, *tcpA*, *gyrA*, *gyrB*, *parE*, *parC*),¹⁰ and multi-locus variable number tandem repeat
178 analysis (MLVA).^{8,11,12}

179

180 ***Rainfall data***

181 Given that few sites directly measured rainfall in South Sudan during the study period, we used
182 remote sensing-derived estimates. We obtained average daily precipitation estimates from
183 2010-2018 at the country, state, and county level from the Climate Hazards Group InfraRed
184 Precipitation with Station Data dataset (<https://climateserv.servirglobal.net/>).

185

186 ***Population displacement data***

187 We used county-level population estimates from WorldPop
188 (<https://data.humdata.org/organization/worldpop>). To better understand population movements
189 as a result of the conflict, we acquired Mobility Tracking Data from the International Office of
190 Migration (IOM) (<https://www.globaldtm.info/>). During the 2nd Round of Mobility Tracking (Mar-
191 Apr 2018), IOM officers interviewed key informants (e.g. local authorities, community leaders
192 and humanitarian partners) in 46/78 counties to estimate the number of individuals (internally
193 displaced persons and returnees) present at assessment areas. These counties include 18/29
194 cholera-affected counties accounting for 62% of suspected cholera cases across the study
195 period. Key informants also provided information on the year when individuals arrived and from
196 where they came.¹³

197

198 ***Intervention data***

199 WHO collated data from all oral cholera vaccination (OCV) campaigns conducted between
200 2014 and 2018. We attempted to collate data on water, sanitation and hygiene (WASH)
201 interventions but these data were not systematically reported.

202

203 **Data Analysis**

204 We calculated the weekly number of cases, attack rate (AR), case-fatality ratio (CFR), the
205 proportion of cases <5 years old, and proportion male. For most cases, we relied on the self-
206 reported date of symptom onset for each case; when this was not available, we used the date
207 of health facility visit. Administrative vaccination coverage was calculated by dividing the total
208 number of first round doses delivered by the estimated target population size. To explore the
209 potential effect of vaccination campaigns, we compared the attack rates of outbreaks where
210 vaccine was used before the epidemic peak with those where it was used after the peak with
211 Poisson regression models using the population size as an offset.

212
213 We generated a maximum likelihood tree from the 1210 publicly available *V. cholerae* O1
214 whole genomes.^{14–16} This tree closely resembles the phylogeny of 1203 genomes from Weill et
215 al,¹⁴ but includes 7 genomes from Uganda published separately.¹⁶ GenBank accession
216 numbers for all 1210 sequences are available in the supplement.¹⁷ For each reference-based
217 assembly (reference accession:AE003852/AE003853), we masked recombinant sites manually
218 (<https://figshare.com/s/d6c1c6f02eac0c9c871e>) and using Gubbins v2.3.4¹⁸ as previously
219 described.¹⁵ We generated the tree from the `filtered\_polymorphic\_sites` FASTA outputted by
220 Gubbins, which contained only the 10,098 variant sites in our alignment. We used IQ-TREE
221 v1.6.10¹⁹ with a GTR substitution model and 1000 bootstrap iterations.²⁰ We rooted the tree on
222 A6 by first constructing a tree containing an outgroup sequence
223 (M66,accession:CP001233/CP001234) and choosing the most ancestral non-outgroup strain.
224 We visualized phylogenetic trees using FigTree v1.4.4
225 (<http://tree.bio.ed.ac.uk/software/figtree/>). For MLVA analysis, we defined a clonal complex as
226 a group of isolates where each differs (in loci-specific copy number) from at least one other
227 isolate in the complex at no more than one locus.

228

229 We used a distributed lag non-linear model^{21–23} to investigate how accumulated rainfall over the
 230 previous seven days (AR7D) is associated with changes in cholera transmission in Juba
 231 County, as characterized by the instantaneous basic reproductive number, $R_0(t)$. We first jointly
 232 estimated the serial interval and reproductive number for the period of epidemic growth for
 233 each wave using methods described by White and Pagano.²⁴ $R_0(t)$ was calculated using
 234 previously described methods^{25,26} assuming a shifted gamma distributed serial interval with a
 235 mean of 3.9 days, a standard deviation of 5.8 days based on the estimates of the 2015 wave,
 236 which was largely confined to Juba County. We assumed that infection leads to complete
 237 immunity over the study period and that 10% of infections were reported as medically attended
 238 cholera cases²⁷. We fit a quasipoisson model with penalized splines transforming AR7D and
 239 lags up to 10 days. The results from this model provide an estimate of the relative $R_0(t)$ over 10
 240 days following a day with a given value of AR7D compared to an AR7D of 0mm.

241

242 We aggregated population movement data to the county level. The origin of returnees
 243 immigrating to South Sudan is presented at the country level. Data were most often available at
 244 the annual scale (e.g. population movement for 2015/2016 was evenly split between the two
 245 years). We calculated the expected number of cholera cases exported by one county to
 246 another by multiplying the attack rate for each county by the number of people leaving that
 247 area during the same year.

248

249 ***Role of funding sources***

250 The funders of this investigation had no role in study design, data analysis, interpretation, or
 251 report writing.

252

253 **Results**

254

255 ***First Wave – 2014***

256 After three years without cases, the MoH confirmed a cholera case in Juba County (Central
257 Equatoria State) on April 28, 2014 soon after the start of the rainy season. Case reports
258 increased in Juba County and in the neighboring Eastern Equatoria State (Figures 1 and 2),
259 where CFRs reached as high as 5% (Table 1). Cholera was then reported in the northeastern
260 state of Upper Nile (1,044 cases) where most cases came from an IDP camp outside of
261 Malakal (South Sudan's second largest city). In total, over the course of 29 weeks, 13 counties
262 reported 6,389 suspected cholera cases (AR=0.05%) and 139 cholera-related deaths
263 (CFR=2.3%, Table 1). Among suspected cases, 25% were children <5 years old and 52% were
264 male. Overall, 430 suspected cases from five different states underwent laboratory
265 confirmation; 44% were confirmed positive for *V. cholerae* O1 by culture (Table S1). The last
266 confirmed and suspected case reported in 2014 was on October 27 and on November 8 in
267 Eastern Equatoria State.

268

269 **Second Wave – 2015**

270 Following a period of 27 weeks with no suspected cases, the first confirmed case of the second
271 wave was reported in Juba County on May 20, 2015. In 2015, over the course of 21 weeks,
272 1,816 suspected cases (AR=0.02%) and 47 deaths (CFR=2.6%) were reported. Among
273 suspected cases, 18% were <5 years old and 55% were male. Nearly all suspected cases
274 (89%) came from Juba County between May 20 and Oct 15; the remaining 196 cases were
275 reported in two nearby counties (Figure 1B). Out of the 132 samples from suspected cases
276 tested in 2015, 33% were confirmed positive by culture. The last confirmed and suspected
277 cases from the 2015 wave occurred in Juba County on September 11 and October 15,
278 respectively.

279

280 **Third Wave – 2016 & 2017**

281 Following a period of 34 weeks with no suspected cases, South Sudan began to experience its
282 largest reported cholera outbreak in recent history, with 20,438 suspected cases (AR=0.18%)

and 428 deaths (CFR=2.13%). Among suspected cases, 22% were under five and 48% were male. On June 18, 2016, Juba County reported the first confirmed cholera case of the year within the country, later than both of the previous years. In 2016, 139 of 402 (34.6%) suspected cases were confirmed. Confirmed cases occurred in Central Equatoria State, Eastern Equatoria State, Jonglei State, Lakes State, and Unity State (Table S1). As rainfall decreased from October to December 2016, no additional counties reported cholera.

Through the dry season, counties along the Nile River continued to report suspected cases (Figure 2). Beginning in January, outbreaks were observed in new areas of Lakes State, Unity State, and Jonglei State. Ayod County (Jonglei State) experienced a particularly explosive outbreak with 3,855 suspected cases and 39 deaths occurring between March 27 and July 9, 2017 (>250 cases per week) and 3,276 cases over a 3-week period; many of these cases occurred in cattle camps, where nomadic pastoralists have little to no access to safe water and sanitation²⁸. As the rainy season began in 2017, suspected cases were detected in northeastern (Upper Nile State), southeastern (Eastern Equatoria State), and central (Warrap State) South Sudan. In 2017, 231 of 453 (51.0%) suspected cases across seven states were confirmed by culture. The last confirmed and suspected cases were reported on August 30 and November 29 respectively, with no additional cases reported through the time of revising this manuscript (18-Sept-2020).

Vaccination Campaigns

In addition to case management, surveillance, and WASH interventions, over 2 million doses of OCV were administered in South Sudan between 2014 and 2018. There were 36 vaccine campaigns (2014: 6, 2015: 6, 2016: 4, 2017: 17, 2018: 3) across 15 of the 34 cholera-affected counties (10 campaigns occurred in Juba County). The population reached during each campaign's first round (typically out of two) ranged between 1,926 and 206,521. While the administrative coverage (number of doses delivered / estimated population size) of most

310 campaigns in the targeted population was moderate to high (range: 52-119%), the coverage at
311 the county-level was much lower. The median first-dose administrative coverage at the county-
312 level was 18% (IQR: 4-41%) (Figure 2).

313
314 Among 25 campaigns conducted reactively (i.e., in response to cases), only 6 occurred before
315 the county-specific peak week of cases. In 2014 and 2015, 4/6 campaigns occurred before the
316 peak as compared to 2/19 campaigns in 2016 and 2017 (Figures 2 & S1). Five campaigns took
317 place after the last suspected case in a county epidemic, all in 2017. All five counties with
318 vaccination campaigns in 2014 or 2015 reported cases during the 2016/2017 wave, though it is
319 unclear if these were in the same sub-county populations. Counties with campaigns occurring
320 after the county-level epidemic peak had, on average, 2.4 (95% CI: 1.8-3.2) times higher attack
321 rates compared to those where vaccine campaigns occurred before the peak, even after
322 adjusting for vaccine coverage and epidemic wave.

323

324 ***Molecular Analyses***

325 Results from phylogenetic analyses (Figure 3, Figure S2), MLVA, and antibiotic resistance
326 testing (Figure S3) all suggest that the bacteria circulating during the study period were nearly
327 identical *V. cholerae* O1 Inaba strains with the *ctxB1* genotype. Our phylogenetic analysis
328 looked more closely at previously-published T10 lineage sequences from several African
329 countries, including the fourteen sequences from South Sudan within this clade.^{14,15} We added
330 seven recently-published T10 sequences from Uganda in the same phylogenetic tree to better
331 understand the relationship between *V. cholerae* in these neighboring countries.¹⁶ All fourteen
332 whole genome sequences from South Sudan epidemics between 2014 and 2017,¹⁵ from
333 patients in Central Equatoria State and Eastern Equatoria State, had extremely similar
334 sequences. Across these fourteen sequences, there are 21 polymorphic sites (~3% of the
335 diversity observed in the T10 lineage), with no more than 10 SNPs between any two
336 sequences (Figure 3). These sequences all belonged to the T10 lineage, which has circulated

337 in the Democratic Republic of Congo for more than twenty years (Figure S1), was first
338 observed in Uganda in 1998, and is distinct from the T13 lineage recently reported in East and
339 Southern Africa and Yemen.^{14,29} Four sequences from Uganda in 2014–2015 cluster closely
340 with all fourteen sequences from South Sudan.

341
342 MLVA analysis on 150 *V. cholerae* isolates collected from 12 different counties in 7 different
343 states in 2014 (46), 2015 (27), 2016 (42), and 2017 (35) showed that all isolates were from the
344 same clonal complex (Figure S3A). Ten isolates collected in Uganda in 2014 also belonged to
345 the same clonal complex. Nearly all isolates had identical antibiotic resistance profiles,
346 including resistance to colistin, nitrofurantoin, polymyxin B, streptomycin, sulfonamides,
347 trimethoprim-sulfamethoxazole (Figure S3B).

348

349 ***The Role of Precipitation***

350 Over 90% of average annual precipitation during 2010–2018 occurred between March and
351 November, both nationally and in Juba County (Figure S4A). Rainfall in 2014–2017 differed
352 little from that in 2010–2013, when no cholera was reported within the country, despite a 2014–
353 2016 El Niño-event.³⁰ With the exception of 2017, the rainy season preceded the detection of
354 cases every year, with cases in Juba County typically reported in closest proximity (2–10
355 weeks) to the start of the heavy rains (Figure S4B). In Juba County, we found no significant
356 association between accumulated rainfall over 7 days and the instantaneous basic
357 reproductive number in the subsequent 10 days (Figure S5, S6). Though we did not extend this
358 analysis to all counties, it is noteworthy how this finding contrasts with those from analyses of
359 cholera in Yemen using similar methods.²¹

360

361 ***The Role of Population Movement***

362 Between 2014 and 2017, the IOM estimated that over 760,000 individuals moved between
363 counties in South Sudan or immigrated from a neighboring country (Table S2). In 2017,

movement increased nearly two-fold (2014: 164,119, 2015: 154,840, 2016: 166,864, 2017: 274,965). Most movement was between counties (553,689) with substantial international migration from Ethiopia (3,025), Democratic Republic of Congo (4,653), Kenya (16,876), Sudan (87,175), and Uganda (95,370).

The movement of people from counties reporting cholera was generally consistent with the year-to-year geographic extent of each wave (Figure 4, Figure S7). In 2014, movement from Juba County, where the first cases were reported, was primarily eastward with little movement north. That same year, cholera was primarily reported in and around Central Equatoria State and in Eastern Equatoria State. In 2015, there was far less movement from Juba County outwards (66% fewer people leaving Juba County than in 2014), and the epidemic remained geographically limited. In 2016 and 2017, movement up and down the Nile River increased dramatically including an increase in movement from Juba County northwards along the river. During this same period, cholera cases were reported from nearly every county along the Nile. In addition to internal migration, more than 110,000 returnees came from Kenya and Uganda, particularly in 2016/17 when both countries reported cholera (Figure S8).

Discussion

Between 2014–2017, South Sudan experienced three distinct epidemic waves of cholera, all beginning in Juba County. Genotypic and phenotypic analyses indicate that despite periods of no reported cases in between each wave, all were of the same clonal origin and most closely related to *V. cholerae* O1 circulating in Uganda and Democratic Republic of Congo. Each wave occurred soon after the onset of the rainy season though rainfall was not associated with increased transmission in Juba County. Movements of internally displaced people and returnees were consistent with spatial patterns of cholera, although we could not quantify this association due to the resolution of the data.

391

392 Cholera was likely introduced into South Sudan in 2013 or early 2014 through cross-border
393 human population movement.¹¹ It remains unclear whether *V. cholerae* O1 remained in South
394 Sudan in between waves through undetected transmission (e.g., insensitive surveillance or
395 asymptomatic carriers), persistence in the environment, or if it was reintroduced. Given the
396 magnitude of population movement between Uganda and South Sudan, reported cases in
397 Uganda during the lull periods in South Sudan, increased surveillance efforts in South Sudan,
398 and the genetic similarity between Ugandan and South Sudanese isolates, repeated cross-
399 border transmission events may be the most plausible explanation. This highlights the need for
400 a regional approach to fighting cholera: efforts to slow an outbreak in one country could be
401 undone by introductions from elsewhere.

402

403 For each wave, the first suspected cases in Juba County were detected shortly after the onset
404 of the rainy season. Flooded infrastructure or changes in human behavior around the time of
405 the rainy season likely led to conditions favorable to the propagation of *V. cholerae* O1 in Juba
406 County. Interestingly, increases in rainfall once the rainy season had begun were not
407 associated with increases in transmission in Juba, suggesting rain may have played a role in
408 triggering but not amplifying the outbreaks. In each wave, Juba County was the first county
409 (except narrowly in 2016) to report suspected cases of cholera in the country and likely played
410 an important role in maintaining cholera outbreaks in the country due to the size of the
411 population and the significant population movement in and out of the city (Figure 4, S6).²⁸ Juba
412 County had >44% of weeks from 2014-2018 with suspected cholera cases and a mean annual
413 incidence of 2.6 per 1,000, making it a clear 'burden hotspot,' as defined by the GTFCC (Figure
414 S9).

415

416 Both increased human migration and transmission occurring during the dry season might
417 partially explain why the 2016-2017 outbreak spread more broadly across the country. In July

2016, intense fighting began in Juba County, causing mass population displacement towards swampy areas along the Nile river and to Uganda. At the same time, the number of cases in Juba County was skyrocketing and cases began to be reported from a number of counties adjacent to the river.³¹ Cholera persisted during the dry season only in counties along the river (Figure S10). Though the reason for continued transmission remains unclear, most of these communities had not reported cholera cases in recent years and presumably had low population-level immunity. The large geographic extent at the end of the dry season may have then created conditions where cholera transmission could be easily amplified as heavy rains arrived. Similar dynamics were noted in Yemen where cholera expanded geographically during the dry season with cases exploding when rainfall increased.²¹

The massive use of cholera vaccine (both reactively and proactively) in South Sudan highlights some of the challenges in implementing campaigns, especially in humanitarian settings. Cholera control efforts during these outbreaks was continually hampered by conflicts and limited access to areas with ongoing transmission. Of the 2,068,622 doses used reactively, only 6.4% were part of campaigns initiated before the peak of the epidemic in the target county. Due to the global shortage of vaccine supply, campaigns were often limited in size and rarely covered the full population; only those deemed at the highest risk. Given the low coverage at the county-level (though relatively high coverage in the target population), the spatial scale of the most reliable epidemiologic data collected, and the relatively low testing rate among suspected cases, the impact of vaccination on reducing incidence is unclear. Our simple analyses exploring the relative attack rates in places that had OCV campaigns before and after the peak provide hints of impact on the population-level but efforts to collect more spatially resolved incidence data combined with systematic lab testing of suspected cholera cases will allow for more precise quantification of vaccine impact.

444 This study comes with a number of limitations. We relied on reports of suspected cases from
 445 reporting health structures across the country. However, due to the instability in the region,
 446 some health centers may not have reported cases, and some people may not have sought
 447 care for even severe disease. Isolates that underwent whole genome sequencing came
 448 exclusively from the southern portion of the country and were not geographically
 449 representative. However, MLVA analyses and the antibiotic sensitivity testing suggest similarity
 450 of strains across the country. While interventions such as chlorination of public water sources,
 451 hygiene promotion, and enforcement of sanitation standards in public areas did occur, we were
 452 unable to identify datasets to quantify the role of these services. Lastly, the population
 453 displacement data were at an annual temporal scale and did not cover the entire country, thus
 454 limiting our ability to have precise quantitative estimates of the contribution of movement as
 455 shown with cell phone data or theoretical models in other settings.^{32,33} Despite these
 456 limitations, trends of mass migration and spread of cholera were largely consistent.

457
 458 Cholera has not been reported in South Sudan from November 2017 through the time of last
 459 editing this manuscript (19-September-2020), which may be explained by a combination of
 460 natural and vaccine-derived immunity, reduced displacement, and perhaps improved sanitary
 461 conditions. Estimated water and sanitation indicators remain poor in South Sudan with only
 462 36.7% having access to improved water and 9.9% having access to improved sanitation in
 463 2017.² South Sudan will likely continue to remain at risk of introductions of cholera from other
 464 countries in the region followed by large-scale spread. Broad use of oral cholera vaccine
 465 combined with local WASH interventions may provide a stopgap over the coming years to
 466 block transmission, while much needed investments in water and sanitation are made.
 467 However, in South Sudan, peace is likely the first prerequisite to long-lasting cholera
 468 prevention.

469

470 Through better understanding of how cholera spreads and the complex roles of human
471 movement, climate, and cholera-specific interventions, we can potentially plan for quicker and
472 appropriately targeted cholera control measures in future cholera epidemics in South Sudan
473 and beyond. Such advances in humanitarian emergencies may be critical to achieving
474 progress towards the 2030 cholera goals of reducing cholera as a public health threat.
475

476 Acknowledgments

477 We thank Andrea C. Caflisch for his input in interpreting movement data from the IOM. We also
478 thank Daryl Domman for assistance in analyzing sequencing data by sharing alignments and
479 details of masking and alignment methods.

480

481 References

- 482 1 Lessler J, Moore SM, Luquero FJ, *et al.* Mapping the burden of cholera in sub-Saharan
483 Africa and implications for control: an analysis of data across geographical scales. *Lancet*
484 2018; **391**: 1908–15.
- 485 2 Local Burden of Disease WaSH Collaborators. Mapping geographical inequalities in
486 access to drinking water and sanitation facilities in low-income and middle-income
487 countries, 2000–17. *Lancet Glob Health* 2020; **8**: e1162–85.
- 488 3 WHO | Ending Cholera. 2017; published online Oct 31.
489 <http://www.who.int/cholera/publications/global-roadmap/en/> (accessed Nov 1, 2017).
- 490 4 Parker LA, Rumunu J, Jamet C, *et al.* Adapting to the global shortage of cholera vaccines:
491 targeted single dose cholera vaccine in response to an outbreak in South Sudan. *Lancet*
492 *Infect Dis* 2017; **17**: e123–7.
- 493 5 Abubakar A, Azman AS, Rumunu J, *et al.* The First Use of the Global Oral Cholera
494 Vaccine Emergency Stockpile: Lessons from South Sudan. *PLoS Med* 2015; **12**:
495 e1001901.
- 496 6 Burki T. End of a cholera epidemic in South Sudan declared. *Lancet* 2018; **391**: 647.
- 497 7 Mutreja A, Kim DW, Thomson NR, *et al.* Evidence for several waves of global transmission
498 in the seventh cholera pandemic. *Nature* 2011; **477**: 462–5.
- 499 8 Manual for the laboratory identification and antimicrobial susceptibility testing of bacterial
500 pathogens of public health importance in the developing world. World Health Organization,
501 2003
502 https://apps.who.int/iris/bitstream/handle/10665/68554/WHO_CDS_CSR_RMD_2003.6.pdf
503 .
- 504 9 Azman AS, Parker LA, Rumunu J, *et al.* Effectiveness of one dose of oral cholera vaccine
505 in response to an outbreak: a case-cohort study. *Lancet Glob Health* 2016; **4**: e856–63.
- 506 10 Safa A, Nair GB, Kong RYC. Evolution of new variants of *Vibrio cholerae* O1. *Trends*
507 *Microbiol* 2010; **18**: 46–54.
- 508 11 Abubakar A, Bwire G, Azman AS, *et al.* Cholera Epidemic in South Sudan and Uganda
509 and Need for International Collaboration in Cholera Control. *Emerg Infect Dis* 2018; **24**:
510 883–7.
- 511 12 Weill F-X, Domman D, Njamkepo E, *et al.* Genomic insights into the 2016–2017 cholera
512 epidemic in Yemen. *Nature* 2019; **565**: 230–3.

- 513 13 South Sudan — Mobility Tracking Report 5 (19 July 2019) | Displacement.
514 [https://displacement.iom.int/reports/south-sudan-%E2%80%94-mobility-tracking-report-5-](https://displacement.iom.int/reports/south-sudan-%E2%80%94-mobility-tracking-report-5-19-july-2019?close=true)
515 [19-july-2019?close=true](https://displacement.iom.int/reports/south-sudan-%E2%80%94-mobility-tracking-report-5-19-july-2019?close=true) (accessed Jan 10, 2020).
- 516 14 Weill F-X, Domman D, Njamkepo E, *et al.* Genomic insights into the 2016-2017 cholera
517 epidemic in Yemen. *Nature* 2019; **565**: 230–3.
- 518 15 Weill F-X, Domman D, Njamkepo E, *et al.* Genomic history of the seventh pandemic of
519 cholera in Africa. *Science* 2017; **358**: 785–9.
- 520 16 Bwire G, Sack DA, Almeida M, *et al.* Molecular characterization of *Vibrio cholerae*
521 responsible for cholera epidemics in Uganda by PCR, MLVA and WGS. *PLoS Negl Trop*
522 *Dis* 2018; **12**: e0006492.
- 523 17 Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. GenBank. *Nucleic Acids*
524 *Research*. 2016; **44**: D67–72.
- 525 18 Croucher NJ, Page AJ, Connor TR, *et al.* Rapid phylogenetic analysis of large samples of
526 recombinant bacterial whole genome sequences using Gubbins. *Nucleic Acids Res* 2015;
527 **43**: e15.
- 528 19 Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective
529 stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 2015;
530 **32**: 268–74.
- 531 20 Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. UFBoot2: Improving the
532 Ultrafast Bootstrap Approximation. *Mol Biol Evol* 2018; **35**: 518–22.
- 533 21 Camacho A, Bouhenia M, Alyusfi R, *et al.* Cholera epidemic in Yemen, 2016–18: an
534 analysis of surveillance data. *The Lancet Global Health* 2018; **6**: e680–90.
- 535 22 Eisenberg MC, Kujbida G, Tuite AR, Fisman DN, Tien JH. Examining rainfall and cholera
536 dynamics in Haiti using statistical and dynamic modeling approaches. *Epidemics* 2013; **5**:
537 197–207.
- 538 23 Jeandron A, Saidi JM, Kapama A, *et al.* Water supply interruptions and suspected cholera
539 incidence: a time-series regression in the Democratic Republic of the Congo. *PLoS Med*
540 2015; **12**: e1001893.
- 541 24 Forsberg White L, Pagano M. A likelihood-based method for real-time estimation of the
542 serial interval and reproductive number of an epidemic: ESTIMATION OF R 0 AND THE
543 SERIAL INTERVAL. *Stat Med* 2008; **27**: 2999–3016.
- 544 25 Fraser C. Estimating individual and household reproduction numbers in an emerging
545 epidemic. *PLoS One* 2007; **2**: e758.
- 546 26 Cori A. EpiEstim: a package to estimate time varying reproduction numbers from epidemic
547 curves. *R package version* 2013.
- 548 27 King AA, Ionides EL, Pascual M, Bouma MJ. Inapparent infections and cholera dynamics.
549 *Nature* 2008; **454**: 877–80.
- 550 28 Khambira M. Epidemiological study of cholera hotspots and epidemiological basins in East
551 and Southern Africa.
552 [https://plateformecholera.info/index.php/resources/publications/regional-studies/639-](https://plateformecholera.info/index.php/resources/publications/regional-studies/639-epidemiological-study-of-cholera-hotspots-and-epidemiological-basins-in-east-and-)
553 [epidemiological-study-of-cholera-hotspots-and-epidemiological-basins-in-east-and-](https://plateformecholera.info/index.php/resources/publications/regional-studies/639-epidemiological-study-of-cholera-hotspots-and-epidemiological-basins-in-east-and-)

554 southern-africa (accessed Sept 18, 2019).

555 29 Mashe T, Domman D, Tarupiwa A, *et al.* Highly Resistant Cholera Outbreak Strain in
556 Zimbabwe. *N Engl J Med* 2020; **383**: 687–9.

557 30 Exceptionally strong El Niño has passed its peak, but impacts continue. World
558 Meteorological Organization. 2016; published online March 15.
559 [https://public.wmo.int/en/media/press-release/exceptionally-strong-el-ni%C3%B1o-has-](https://public.wmo.int/en/media/press-release/exceptionally-strong-el-ni%C3%B1o-has-passed-its-peak-impacts-continue)
560 [passed-its-peak-impacts-continue](https://public.wmo.int/en/media/press-release/exceptionally-strong-el-ni%C3%B1o-has-passed-its-peak-impacts-continue) (accessed Sept 18, 2019).

561 31 South Sudan president, rival call for cease-fire. AP NEWS. 2016; published online July 11.
562 <https://apnews.com/92b9f2fe32eb4a6fa6b54af1c7076fc2> (accessed Sept 10, 2019).

563 32 Mari L, Bertuzzo E, Righetto L, *et al.* Modelling cholera epidemics: the role of waterways,
564 human mobility and sanitation. *J R Soc Interface* 2012; **9**: 376–88.

565 33 Finger F, Genolet T, Mari L, *et al.* Mobile phone data highlights the role of mass gatherings
566 in the spreading of cholera outbreaks. *Proc Natl Acad Sci U S A* 2016; **113**: 6421–6.

567

568

Figure 1. Successive epidemic curves (A) and geographic spread of suspected cholera cases (B) in South Sudan between Jan 1, 2014, and July 1, 2018. Bar heights represent the weekly number suspected cases in Juba County (red) or in other counties (grey). For each year with cholera cases (2014, 2015, 2016 & 2017), county-level maps indicate the month of the first case was reported. States are labeled and

583 borders are in black. Stars represent the capital city of Juba. Light blue line represents
584 the Nile river.

585

586

587

Figure 2. Relative timing of oral cholera vaccine campaigns and county-level cholera outbreaks in South Sudan. Each circle represents a distinct OCV campaign with its size representing the proportion of the county-level population estimated to be vaccinated during the first round (number of doses delivered / estimated population size). Campaigns are marked on the week of the first round, and cases are marked at the week of reported symptom onset for each county.

Figure 4. County-level suspected case attack rate and population movement in South Sudan, 2014 - 2017. Red fill (A) indicates the percentage of the population which were suspected cholera cases. Blue arrows (B) represent the number of internally displaced persons and returnees moving between counties. Purple arrows (C) represent the expected number of exported cholera cases estimated by multiplying the attack rate of each county by the number of people leaving that county to another location for the same time period. Higher intensity of population movement and exported cases are indicated by darker colors on the $\log_{10}$ scale. Counties in grey were not assessed for arrival of internally displaced persons (IDPs) or returnees. Arrows move to and from county-level centroids.

619

620

State	County	Population	2014				2015				2016 / 2017			
			Cases	AR(%)	Deaths	CFR(%)	Cases	AR(%)	Deaths	CFR(%)	Cases	AR(%)	Deaths	CFR(%)
Central Equatoria	Juba	514 770	2,229	0.43	40	1.79	1,622	0.32	45	2.77	2,928	0.57	37	1.28
	Kajo-keji	268 911	84	0.03	6	7.14	57	0.02	1	1.75	0	0.00	0	0.00
	Terekeke	190 050	0	0.00	0	0.00	0	0.00	0	0.00	15	0.01	5	33.33
	Yei	287 217	48	0.02	2	4.17	0	0.00	0	0.00	0	0.00	0	0.00
Eastern Equatoria	Budi	138 379	0	0.00	0	0.00	0	0.00	0	0.00	805	0.58	81	10.08
	Ikotos	103 380	227	0.29	5	1.88	0	0.00	0	0.00	0	0.00	0	0.00
	Kapoeta	237 797	1	0.00	0	0.00	0	0.00	0	0.00	2,105	0.89	26	1.24
	East Kapoeta	142 072	81	0.08	1	1.23	0	0.00	0	0.00	983	0.70	1	0.10
	North Kapoeta	108 540	14	0.01	0	0.00	0	0.00	0	0.00	1,016	0.95	10	0.90
	South Lafon	84 973	256	0.30	15	5.88	0	0.00	0	0.00	0	0.00	0	0.00
	Magw	238 723	243	0.10	10	4.12	0	0.00	0	0.00	29	0.01	1	3.45
	Turit	148 190	2,067	1.40	36	1.82	0	0.00	0	0.00	0	0.00	0	0.00
	Ayod	199 107	0	0.00	0	0.00	0	0.00	0	0.00	3,855	1.94	39	1.01
Jonglei	Bor South	319 971	0	0.00	0	0.00	139	0.04	1	0.72	307	0.10	8	2.61
	CenallPigi	138 207	0	0.00	0	0.00	0	0.00	0	0.00	206	0.15	5	2.43
	Duk	93,280	0	0.00	0	0.00	0	0.00	0	0.00	542	0.58	36	6.64
	Fangak	154 844	0	0.00	0	0.00	0	0.00	0	0.00	281	0.17	2	0.77
Lakes	Nyirul	164 403	0	0.00	0	0.00	0	0.00	0	0.00	72	0.04	2	2.78
	Awerial	68,058	0	0.00	0	0.00	0	0.00	0	0.00	1,197	1.78	13	1.09
	Yirul East	114 288	0	0.00	0	0.00	0	0.00	0	0.00	1,498	1.31	87	4.48
	Yirul West	97,405	0	0.00	0	0.00	0	0.00	0	0.00	238	0.24	0	0.00
Unity	Leer	77,512	0	0.00	0	0.00	0	0.00	0	0.00	94	0.12	3	3.19
	Mayendit	88,514	0	0.00	0	0.00	0	0.00	0	0.00	226	0.26	5	2.21
	Mayom	179 099	0	0.00	0	0.00	0	0.00	0	0.00	9	0.01	4	44.44
	Panyijar	72,344	0	0.00	0	0.00	0	0.00	0	0.00	012	0.89	30	4.80
	Rubkone	154 075	0	0.00	0	0.00	0	0.00	0	0.00	1,170	0.76	10	0.85
Upper Nile	Fashoda	45,974	0	0.00	0	0.00	0	0.00	0	0.00	584	1.27	9	1.54
	Malakal	178 354	1,035	0.59	22	2.13	0	0.00	0	0.00	19	0.01	0	0.00
	Panyikang	33,303	9	0.03	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00
Warap	Renk	199 008	0	0.00	0	0.00	0	0.00	0	0.00	10	0.01	0	0.00
	Tonj East	173 337	0	0.00	0	0.00	0	0.00	0	0.00	1,611	0.93	42	2.61
	Tonj North	240 681	0	0.00	0	0.00	0	0.00	0	0.00	31	0.01	0	0.00
Western Bahr el Ghazal	Raga	80,910	0	0.00	0	0.00	0	0.00	0	0.00	1	0.00	0	0.00
Western Equatoria	Mundii	83,357	3	0.00	0	0.00	0	0.00	0	0.00	0	0.00	0	0.00
South Sudan	East South Sudan	11,854,000	6,389	0.05	139	2.18	1,818	0.02	47	2.59	20,438	0.18	436	2.13

621

622 **Table 1. Cholera indicators by county and nationally in South Sudan for each of**

623 **the three national epidemic waves. AR = Attack Rate. CFR = Case-fatality risk. A**

624 **nationwide attack rate was calculated for the entire population of South Sudan.**