

S2 Table. Summary of model parameters and their choices for numerical simulations.

Parameter	Definition	Value
n_E	Number of latent phase Erlang states	16
n_P	Number of prodromal Erlang states	16
n_I	Number of fully contagious Erlang states	16
n_L	Number of late infectious Erlang states	16
D_E	Average duration of latency period	3.7 days
D_P	Average duration of prodromal period	1 day
D_I	Average duration of fully contagious period	7 days
D_L	Average duration of late infectious period	7 days
ε	Transition rate of latent states	n_E/D_E
φ	Transition rate of prodromal states	n_P/D_P
γ	Transition rate of fully contagious states	n_I/D_I
δ	Transition rate of late infectious states	n_L/D_L
f_{Sick}	Fraction of symptomatic (sick) infections in Ge and St	58%
$f_{\text{Sick}}^{(\text{Ri})}$	Fraction of symptomatic (sick) infections in Ri	70%
f_{Dead}	Fraction of sick ind. in Ge and St, who die from the disease	4.7%
$f_{\text{Dead}}^{(\text{Ri})}$	Fraction of sick ind. in Ri, who die from the disease	7%
$1/\alpha$	Waiting time for test results in hours	12, 24, 48, 96
$1/\xi$	Average frequency of testing in days	1, 2, 7, 14

General sub-population (Ge), LTCF employees (St), risk group (Ri).