

	A	B	C	D	E	F	G	H	I	J	K	L	M
1	Table 4. Calculation of the Potential Years of Life Lost to COVID-19 (person-years)in the US using the life expectancy (LE) averaged over the 10 year periods indicated												
2													
3	Males												
4	All ages	104,896	Average LE	Person-Years Lost									
5	Under 1 year	13	75.97	987.61									
6	1–4 years	7	73.9875	517.9125									
7	5–14 years	22	67.07	1475.54									
8	15–24 years	223	57.346	12788.158									
9	25–34 years	1,003	48.111	48255.333									
10	35–44 years	2,698	39.017	105267.87									
11	45–54 years	7,063	30.139	212871.76									
12	55–64 years	16,067	22.002	353506.13									
13	65–74 years	25,722	14.769	379888.22									
14	75–84 years	28,239	8.622	243476.66									
15	85 years and over (a)	23,839	4.305	102626.9									
16	Total			1461662.1									
17													
18	Females												
19	All ages	89,191	Average LE	Person-Years Lost									
20	Under 1 year	7	80.96	566.72									
21	1–4 years	8	78.9225	631.38									
22	5–14 years	10	71.993	719.93									
23	15–24 years	139	62.131	8636.21									
24	25–34 years	510	52.456	26752.5									
25	35–44 years	1,258	42.95	54031.1									
26	45–54 years	3,325	33.704	112065.8									
27	55–64 years	8,705	24.998	217607.59									
28	65–74 years	16,143	16.964	273849.85									
29	75–84 years	23,125	10.065	232753.13									
30	85 years and over (a)	35,961	5.084	182825.72									
31	Total			1110439.9									
32													
33													
34	Total person-years Lost	2,572,102 (1,461662 Males and 1,110,440 Females)											
35													
36	Average person years lost per individual			13.25 years									
37				13.93 years per male									
38				12.45 years per female									

Potential sources of error in these calculations

(a) A possible source of error results from the fact that the >85 category presented by the CDC did not break down the number of different years. We took the average life expectancy for the 10-year period 85-94 for the entire category. However, of the population of all individuals 85 and older, only 5.1% on average of men are 95 or older and 7.2% of women with average life

expectancies of 2.2 and 2.5 years, respectively (See Table 7). There are 1200 men and 2600 women in the 95 and greater category who were originally evaluated at 4 and 5 years of life respectively when the real numbers are closer to 2.2 and 2.5 years, respectively, meaning we overestimated men by 1.8 years and women by 2.5 years of life expectancy. Altogether this is an error of less than 10,000 person years assuming the two groups die at an equal rate. If the 95 and greater dies at a 2-fold rate relative to the 85-94 group, that would be adjusted to 20,000 person-years.

- (b) A second source of error derives from the assumption that the individuals in the 10 year groups are evenly distributed for deaths. In the absence of smaller grouping, it is not possible to definitively overcome this at this time.
- (c) A third source of error is the inability to account for comorbidities for which we would need the breakdown of deaths per age group for the subgroups for comorbidities and the average life expectancy tables for each comorbidity without COVID-19.