The interactive effects of ambient air pollutants-meteorological factors on confirmed cases of COVID-19 in 120 Chinese cities

Jianli Zhou ^a, Linyuan Qin ^b, Xiaojing Meng ^{a*}, Nan Liu ^{a, c, d, e*}

^a Department of Occupational Health and Occupational Medicine, Guangdong Provincial Key Laboratory of Tropical Disease Research, School of Public Health, Southern Medical University, Guangzhou, 510515, P. R. China.

^b Department of Epidemiology and Statistics, School of Public Health, Guilin Medical University, Guilin, 541001, P. R. China.

^c Pinghu Hospital, Health Science Center, Shenzhen University, Shenzhen, 518116, P. R. China.

^d Institute of Public Health, School of Nursing, Henan University, Kaifeng, 475004, P. R. China.

^e College of Public Health, Zhengzhou University, Zhengzhou, 540001, P. R. China.

*Corresponding authors:

Xiaojing Meng, Department of Occupational Health and Occupational Medicine, School of Public Health, Southern Medical University, 1838 Guangzhoudadaobei, Guangzhou, Guangdong, China. Tel: +8620-61648471; fax: +8620-61648324. Email: xiaojingmeng@smu.edu.cn.

Nan Liu, Pinghu Hospital, Health Science Center, Shenzhen University, The junction of Hutian Road and No.3 Road, Pinghu Street, Longgang District, Shenzhen, Guangdong, China. Tel: +86-13688869875; fax: 075589222888. Email: nhyy8132@smu.edu.cn.

Abstract

Emerging evidence has confirmed meteorological factors and air pollutants affect novel coronavirus disease 2019 (COVID-19). However, no studies to date have considered the impact of interactions between meteorological factors and air pollutants on COVID-19 transmission. This study explores the association between ambient air pollutants, meteorological factors and their interaction on confirmed case counts of COVID-19 in 120 Chinese cities. We modeled total confirmed cases of COVID-19 with meteorological factors, air pollutants and their interactions. To account for potential migration effects, we included the migration scale index (MSI) from Wuhan to each of the 120 cities included in the model, using data from 15 January to 18 March 2020. As an important confounding factor, MSI was considered in a negative binomial regression analysis. Positive associations were found between the number of confirmed cases of COVID-19 and carbon monoxide, aerodynamic particulate matter with aerodynamic diameter $\leq 2.5 \mu\text{m}$, relative humidity and ozone, with and without MSI-adjustment. Negative associations were also found for sulfur dioxide and wind velocity both with and without controlling for population migration. In addition, air pollutants and meteorological factors had interactive effects on COVID-19 after controlling for MSI. In conclusion, air pollutants, meteorological factors and their interactions all affect COVID-19 cases.

Keywords: air pollutants; COVID-19; interactive effect; meteorological parameters; migration scale index; negative binomial regression

1. Introduction

In Dec. 2019, a novel coronavirus disease (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was reported in Wuhan City, P. R. China (Wu et al. 2020, Zhou et al. 2020). The COVID-19 outbreak rapidly spread across China and to many other countries (Holshue et al. 2020, Huang et al. 2017, Lai et al. 2020, Wang et al. 2020a). Due to its transmissibility and increasing spread, COVID-19 was officially declared a pandemic by the World Health Organization (WHO) on 11 March 2020 (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019>). By 17 August 2020, COVID-19 had become prevalent in 188 countries, with a total of 21,869,433 confirmed cases worldwide (<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/>); and 89906 confirmed cases in China (http://www.nhc.gov.cn/xcs/xxgzbd/gzbd_index.shtml). This has created a worldwide public health crisis.

Air pollutants can significantly affect health and the WHO indicates that 4.6 million individuals die annually from diseases and illnesses directly related to poor air quality (Tobias & Molina 2020). In particular, increasing numbers of studies suggest a relationship between air pollution and infectious diseases. While previous studies report no relationship between air pollution and SARS outbreaks (Cai et al. 2007, Tobias & Molina 2020, Yusuf et al. 2007), a positive association was observed between air pollutants (sulfur dioxide, nitrogen dioxide, carbon monoxide and ground-level ozone) and SARS fatality rates (Cui et al. 2003). Another study in Brisbane suggested that high concentrations of ozone (O₃) and particulate matter <10 microns (PM₁₀) were associated with increased pediatric influenza cases (Xu et al.

2013).

Meteorological factors are another independent factor that can affect the spread of infectious diseases (Cai et al. 2007, Xie &Zhu 2020). Although the association between viruses leading to respiratory disease (e.g., SARS, viral influenza and SARS-CoV-2) and meteorological factors has been explored (Yusuf et al. 2007), results of these studies remain controversial. For example, Tan et al. found a sharp change in ambient temperature was correlated with increased risk of SARS, (Tan et al. 2005), while Yao et al. found ambient temperature had no significant effect on the transmissibility of SARS-CoV-2 (Yao et al. 2020b). Other studies indicate that an increase in average temperature contributes to an increase in average relative risk (RR) but a decrease in daily confirmed case counts of SARS (Cai et al. 2007). These inconsistent results may reflect an interaction occurring between pollutants and meteorological effects on the transmission of these diseases.

The trends of the SARS outbreak may be influenced by air pollutants and meteorological factors that change host susceptibility and survival time of SARS-CoV in vitro (Cai et al. 2007). COVID-19 infections can cause serious respiratory disease. Common symptoms at the beginning of infection include fever, cough, muscle aches or fatigue, etc., similar to SARS-CoV (Liu et al. 2020b, Siordia 2020), and an effect of air pollutants and meteorological factors, similar to that seen in SARS, on confirmed COVID-19 cases has also observed (Jiang et al. 2020, Liu et al. 2020a, Oliveiros et al. 2020, Wang et al. 2020b, Zhu et al. 2020); However, the interactive effects of air pollutants and meteorological factors has not yet been considered. Given the human to human transmission of COVID-19, population migration is also likely to be an important confounding factor when evaluating the effects of meteorological factors and air pollutants on COVID-19, and it has been established that COVID19

cases are almost linearly correlated with the number of people traveled from Wuhan to the destination city (Hu Jianxiong 2020). In this study, we investigate the association between ambient air pollutants, meteorological factors, and their interactive effects on the number of COVID-19 confirmed cases while also controlling for population migration effects, to extend and further our understanding of the relationship between transmission of this disease and environmental factors.

2. Methods

2.1 Data collection

We observed the numbers of the confirmed cases in relation to air pollutants, meteorological parameters and the migration scale index (MSI) in 120 Chinese cities covering the majority of mainland China. The MSI reflects the scale of urban migration into or out of the assigned population, using a horizontal comparison among cities to reflect the trend of population migration between the city pairs. All cities included had no less than five cases from 15 January to 18 March 2020 inclusive. Wuhan was locked down at 10 am on 23 Jan, so we consider only people that traveled from Wuhan to the destination city between 17 and 23 Jan. 2020. All data were taken from the official website of Baidu Migration (<https://qianxi.baidu.com/2020/>). To better fit the model, we do not include the data of Wuhan City since its large number of the confirmed case counts skew the data and thus may reduce the robustness of the model.

Daily confirmed case counts and daily air pollutants were collected from the official website of Harvard University (<https://dataverse.harvard.edu>). Meteorological indexes, including average temperature (AT), diurnal temperature range (DTR), relative humidity (RH), wind velocity (WV), air pressure (AP), precipitation (PRE), and hours of sunshine (HS), were obtained from China's Meteorological Data Sharing

126 Service System (<http://data.cma.cn/site/index.html>). Air pollutant data included
127 particulate matter with aerodynamic diameter ≤ 2.5 μm (PM_{2.5}), nitrogen dioxide
128 (NO₂), sulfur dioxide (SO₂), carbon monoxide (CO), and ozone (O₃). Pollutant
129 concentrations and meteorology parameters were available as daily means in each city;
130 we calculated a single mean for the 63 days of our date range for use in the model.

131 **2.2 Statistical analysis**

132 Descriptive analysis was performed on all data to explore the characteristics of total
133 confirmed case counts, average concentration of O₃, NO₂, SO₂, CO, PM_{2.5}, AT, DTR,
134 RH, WV, AP, PRE, HS and MSI. Meanwhile, to account for the lag effect and the
135 latent period of COVID-19, we evaluated associations with confirmed case counts
136 using lags of 0, 3, 7 and 14 d for air pollutants and meteorological factors (Liu et al.
137 2020a). Lags were incorporated into parameters by shifting the date range of the
138 appropriate number of days. For example, for lag 3, mean parameter values were
139 calculated using daily means from 15 Jan to 15 March for use in the model; confirmed
140 case counts were always for the date range 18 Jan to 18 March. Spearman correlation
141 analysis was employed to explore the correlation between meteorological factors and
142 air pollutants, and among air pollutants, meteorological factors and confirmed case
143 counts. We used generalized linear models with a negative binomial error distribution
144 to analyze associations between air pollutant factors (O₃, NO₂, SO₂, CO and PM_{2.5}),
145 meteorological factors (AT, DTR, RH, WV, AP, PRE and HS), MSI and the total
146 counts of confirmed case since the count data was expected to be clustered and thus
147 over-dispersed. First, we established simple negative binomial regression models
148 (including only one independent variable) to analyze each air pollutant,
149 meteorological factor, and MSI effects on confirmed case counts. We then conducted
150 multiple negative binomial regression models (including multiple independent

variables), with and without MSI, that included all above-mentioned air pollutants and meteorological factors since the independent variables are not strongly correlated (all $|r| < 0.9$). The general negative binomial regression equation is:

$$\text{Log}(\lambda) = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \dots + \beta_m X_m,$$

where λ represents the risk of being a confirmed case counts, $X_1, X_2, \dots, X_m$ are the factors that affect λ ; and $\beta_1, \beta_2, \dots, \beta_m$ are the partial coefficients of X_1, X_2 , and X_m , respectively. To explore the interaction effects, we included $X_i * X_j$ as an interaction term in the model, where X_i and X_j represent an air pollutant and a meteorological factor, respectively. A stratified analysis was done to consider the effect of air pollutants on confirmed COVID-19 cases under different levels of meteorological factors. When a significant interaction effect was present in the negative binomial model, the meteorological factor was stratified into two groups using the median as the dividing point, and a negative binomial model containing all pollutants and remaining meteorological factors was fitted to analyze the effect of air pollutants on the confirmed number of COVID-19 cases was done for each group.

All statistical analyses were performed using SPSS version 22.0 (SPSS Inc., Chicago, IL, USA) and R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/> with a two-sided P-value < 0.05 considered as statistically significant.

3. Results

3.1 Study area

Figure 1(a) displays the location of the 120 selected cities in mainland China and the total confirmed cases in each city collected from 15 January 18 March 2020. The

175 latitude and longitude of these 120 cities range from 20°00' to 47°23' N and from 99°
176 11' to 130°55' E, respectively. Most cases were concentrated in the central and
177 eastern parts of mainland China, with fewer cases in the northwest. The detailed
178 distribution of confirmed COVID-19 cases in the 120 cities is given in Table S1.
179 Hubei province contained the majority of confirmed COVID-19 cases; excluding
180 Wuhan City, Xiangfan, Shiyan, Yichang, Jingzhou, Suizhou, and Xiaogan were the
181 cities with the highest number of cases (>500 per city).

Figure 1

182 3.2 Description of confirmed COVID-19 cases, meteorological factors and air 183 pollutants 184 pollutants

185 **Table1** Summary of total confirmed COVID-19 cases, meteorological data and air
186 pollutants of 120 Chinese cities

Variables	Min	P25	P50	P75	Max
Confirmed case counts					
Confirmed case counts	5	14	36	81	3518
Concentrations of air pollutants					
CO (mg/m ³)	0.42	0.69	0.82	0.98	1.67
NO ₂ (µg/m ³) [□]	5.06	16.65	20.89	28.06	44.01
SO ₂ (µg/m ³)	2.70	5.92	8.23	13.83	41.74
O ₃ (µg/m ³) [□]	29.63	48.42	56.38	62.66	85.94
PM _{2.5} (µg/m ³)	13.62	30.75	43.58	59.54	94.83
Meteorological factors					
RH (%)	35.42	62.43	73.71	79.28	85.50
AT (°C)	-12.48	2.10	8.14	10.77	21.29
PRE (mm)	0.00	0.36	0.95	2.64	6.82
AP (hPa)	758.41	978.83	1008.22	1016.43	1024.43
HS(h)	1.60	3.31	4.48	6.91	9.20
WV (m/s) [□]	1.00	1.78	2.19	2.57	5.05
DTR (°C)	5.13	6.73	8.92	10.90	18.37

MSI	0.01	0.02	0.13	0.37	5.13
-----	------	------	------	------	------

□ The data were normal distribution by normal distribution test.

Descriptive statistics (maximum and minimum values, median, and interquartile range) for the number of confirmed cases, air pollutant measures, and the meteorological parameters of the 120 cities are presented in Table 1. Among these cities, during our study period (15 January to 18 March 2020), the median daily concentrations of CO, NO₂, SO₂, O₃ and PM_{2.5}, were 0.82 mg/m³, 20.89 µg/m³, 8.23 µg/m³, 56.38 µg/m³, and 43.58 µg/m³, respectively. For meteorological parameters, ranges of RH, AT, PRE, AP, HS, WV, DTR were 35.42-85.50%, -12.48 to 21.29 °C, 0.00-6.82 mm, 758.41-1024.43 hPa, 1.60-9.20 h, 1.00-5.05 m/s, and 5.13 to 18.37 °C, respectively. MSI ranged from 0.01 to 5.13.

3.3 Correlation analysis of air pollutants and meteorological factors

Spearman correlation coefficients (r_s) between concentrations of air pollutants and meteorological factors are shown in Figure 1(b). Circle size indicates strength of relationship between two variables. All correlations are less than |0.9|. Therefore, all the variables could be integrated into the multiple negative binomial regression model.

3.4 Effects of air pollutants and meteorological parameters on confirmed case counts

The relationship between pollutants, meteorological parameters and confirmed case counts for the lag of 14 d are presented in a scatter plot in Figure 2. Positive associations were present between the confirmed case counts and the pollutant measure PM_{2.5} ($P = 0.024$), and the meteorological measures RH ($P = 0.001$), PRE ($P = 0.015$) and AP ($P < 0.001$). Negative associations occurred between the confirmed case counts and the pollutant SO₂ ($P < 0.001$) and the meteorological variable HS ($P = 0.018$). Correlations between confirmed case counts and all other variables were not

212 significant.

Figure 2

213

214 3.5 Negative binomial regression model analysis

215 Simple negative binomial regression results are shown in Figure S1(a-d). CO, NO₂,
 216 SO₂, O₃, PM_{2.5}, RH, AT, AP, HS, WV and MSI were significantly associated with
 217 confirmed cases counts (all $P < 0.05$). PRE had no effect on confirmed case counts for
 218 lags of 0, 3, 7 and 14 d, while DTR was significantly associated with confirmed cases
 219 counts at lags of 7 and 14 d. The multiple negative binomial regression model results
 220 (without adjusting for MSI) are shown in Figure 3(a1-d1). CO, PM_{2.5} and RH were
 221 positively associated with confirmed case counts, while SO₂ was negatively
 222 associated with confirmed case counts at lags 0, 3, 7, and 14 d. However, O₃ has a
 223 positive effect on confirmed case counts at lag 0 and WV has a positive effect on
 224 confirmed case counts at lag 14 d. After adjusting for migration using MSI, CO, PM_{2.5},
 225 RH, O₃, SO₂, WV were also statistically significant, as shown in Figure 3(a2-d2).

Figure 3

226

227

228 3.6 Interaction effects between air pollutants and meteorological factors

229 **Table 2** The significant interaction effects of ambient air pollutants and
 230 meteorological factors on COVID-19 confirmed cases after adjusting MSI ($P < 0.05$)

Variables	β	Lower 95% CI	Upper 95% CI	P
CO by HS	-0.811	-1.420	-0.202	0.009
CO by DTR	-0.365	-0.693	-0.037	0.029
CO by RH	0.137	0.058	0.216	0.001
SO ₂ by AP	-0.001	-0.001	0.000	0.014
PM _{2.5} by PRE	0.022	0.008	0.036	0.002

231

232 Significant interactive effects ($P < 0.05$) of ambient air pollutants and meteorological

233 factors on COVID-19 confirmed cases after adjusting with MSI are shown in Table 2.
 234 CO interacted with two meteorological factors, HS and DTR, and SO₂ interacted with
 235 AP to negatively affect the total number of COVID-19 confirmed cases; while CO and
 236 RH, and PM_{2.5} and PRE interacted to positively affect total case count.
 237 **Table 3** Stratified analysis of ambient air pollutants and meteorological factors
 238 interaction effects on COVID-19 confirmed cases

Air pollutant	Meteorological factor	β	Lower 95% CI	Upper 95% CI	P
CO	HS<P50	3.321	-0.230	6.873	0.067
	HS≥P50	1.115	-0.247	2.478	0.108
CO	DTR<P50	5.763	2.402	9.124	0.001
	DTR≥P50	0.931	0.066	1.796	0.035
CO	RH<P50	0.930	-0.574	2.434	0.226
	RH≥P50	6.045	3.247	8.844	<0.001
SO ₂	AP<P50	-0.006	-0.052	0.041	0.812
	AP≥P50	-0.243	-0.307	-0.178	<0.001
PM _{2.5}	PRE<P50	0.004	-0.019	0.027	0.727
	PRE≥P50	0.049	-0.009	0.107	0.100

239 The adjusted interaction effects of CO with HS, DTR, RH, of SO₂ with AP, and of
 240 PM_{2.5} with PRE are listed in Table 3. The remaining adjusted air pollutants,
 241 meteorological factors and MSI and under the levels of HS values below 50th
 242 percentile and HS and above 50th percentile, had no significant effects on COVID-19
 243 case counts by CO. At the low DTR level (<50th percentile), an increase of CO by 1
 244 mg/m³ is expected to increase the log of risk of being confirmed COVID-19 cases by
 245 a factor of >5 ($\beta = 5.763$, $P = 0.001$), while the same increase in CO at the high DTR
 246 level ($\geq 50^{\text{th}}$ percentile) increases the risk by a factor of <1 ($\beta = 0.931$, $P = 0.035$). CO
 247 has no significant effect on confirmed case counts at low RH level, but at a high RH
 248 level, a 1 mg/m³ increase of CO significantly raises the risk of being confirmed case

($\beta = 6.045$, $P < 0.001$). An increase of SO_2 was associated with lower risk of confirmed case counts ($\beta = -0.243$, $P < 0.001$) at high level AP, but was not significantly associated with confirmed case counts at low level AP. $\text{PM}_{2.5}$ variation had no significant effect for either low or high PRE levels. These results indicate that the effects of CO and SO_2 vary depending on the level of meteorological factors.

4. Discussion

In this work, we explored the role of air pollutants, meteorological factors and their interactive effects on confirmed case counts in 120 Chinese cities. We focused on these cities as their meteorological data was available (this is not true for all cities in China). Our results indicate that air pollutants, meteorological factors and their interaction are associated with COVID-19 confirmed case counts, with and without adjusting using MSI.

Previous studies indicate that meteorological factors and air pollutants influence prevalence of infectious diseases (e.g., influenza, SARS, MERS, COVID-19 (Cai et al. 2007, Liu et al. 2020a, Prata et al. 2020, Tan et al. 2005, van Doremalen et al. 2013, Yusuf et al. 2007)). An ecological analysis carried out in China demonstrated that air pollutants increase SARS case fatalities (Cui et al. 2003). As a major component of air pollution, particulate matter is associated with a variety of cardiorespiratory diseases, including acute lower respiratory tract infections. $\text{PM}_{2.5}$ has adverse effects on human health and is associated with many diseases, especially respiratory disease (Horne et al. 2018). $\text{PM}_{2.5}$ may damage bronchial immunity and epithelial cell integrity (Jiang et al. 2020), which, in turn, will reduce the body's ability to fight off viruses, making affected individuals more vulnerable to respiratory disease. Exposure to higher concentrations of $\text{PM}_{2.5}$ results in greater use of healthcare resources for acute lower respiratory infection (Horne et al. 2018). Jiang et al. (Jiang et al. 2020), Suhaimi et al.

(Suhaimi et al. 2020) and Zhu et al. (Zhu et al. 2020) studies suggested that $PM_{2.5}$ was positively associated with confirmed COVID-19 cases, which were consistent with our results (Figure 3). Cai et al. (2007) indicated that no correlation exists between air pollution (including particulate matter, SO_2 , NO_2 , O_3 , and CO) and SARS outbreaks (Cai et al. 2007). However, our results are consistent with Zhu et al. (Zhu et al. 2020), who found that increases in CO were positively associated with the number of confirmed COVID-19 cases, while SO_2 was negatively correlated with COVID-19 cases (Figure 3). The difference is that migration was taken into account in our analysis and it was not in the analysis by Zhu et al. (Zhu et al. 2020). Hu's (Hu Jianxiong 2020) study indicated that the cumulative number of reported COVID-19 cases in each province was positively correlated with the population index of Wuhan moving out to each province, and the correlation coefficient is 0.84, so we believe that it is necessary to control the population migration.

Kesic et al. suggests that exposure to O_3 might disrupt the protease/anti-protease balance of human airways, contributing to an increased influenza A infection rate (Kesic et al. 2012). We similarly found that exposure to $PM_{2.5}$, CO and O_3 increased likelihood of being infected by COVID-19 (Figure 3). Understanding the relationship between air pollutant exposure and health provides clues regarding effective protective measures that can reduce the incidence or risk of diseases. Despite the positive relationship between air pollutants (including $PM_{2.5}$, CO, and O_3) and confirmed COVID-19 cases and the negative relationship between SO_2 and confirmed COVID-19 cases that we found, the mechanism by which pollutants affect COVID-19, whether via changes in the transmission of the virus or in the immunity of the human body, remains unknown. Further experimental studies are required to explore the molecular regulation mechanisms involved.

299 Meteorological factors such as AT, DTR, RH, WV, AP, PRE and HS are important
300 factors influencing the survival of microorganisms in the external environment
301 (Anwar et al. 2019, Bezirtzoglou et al. 2011, Casadevall 2020, Chan et al. 2011, Liu et
302 al. 2020a). For decades, experts have issued warnings that climate change may affect
303 the epidemiology of infectious diseases (Casadevall 2020). Global warming, rising
304 sea levels, and climate change will be accompanied by new and unknown diseases
305 (Bezirtzoglou et al. 2011, Casadevall 2020). Meteorological factors may change the
306 host's behavior, e.g., by affecting time spent indoors or outdoors, affect the host's
307 defense mechanisms (inhalation of dry and cold air will cause damage to the cleaning
308 function of cilia), or change the infectivity and stability of the virus. Although many
309 studies on meteorological factors and respiratory infectious diseases (including
310 COVID-19) exist, research results remain controversial. While temperature is unlikely
311 to act alone in determining the population's health, but it can work synergistically
312 with other weather parameters to do so (Huang et al. 2017).

313 COVID-19 case numbers appear to increase with increasing temperature in the
314 range of -20 to 20 °C (Liu et al. 2020a). Other studies have reported that when AT is
315 below 3 °C, it is positively correlated with confirmed COVID-19 case counts (Xie
316 & Zhu 2020). A Brazilian study suggests a negative linear relationship exists between
317 AT and daily cumulative confirmed cases of COVID-19 in the temperature range of
318 16.8 to 27.4 °C (Prata et al. 2020), while Auler et al.'s research (also conducted in
319 Brazilian) suggests both higher AT and RH favor COVID-19 transmission.(Auler et al.
320 2020). The results of studies in some tropical countries emphasize that COVID-19
321 cases positively correlate with AT (Pani et al. 2020, Suhaimi et al. 2020, Tosepu et al.
322 2020). Another study demonstrated that in DTR increased in range of from 5 to 15 °C,
323 the counts of confirmed COVID-19 cases decreased, and that a non-linear relationship

occurred between COVID-19 cases and absolute humidity (Liu et al. 2020a). A study of 244 cities in China found no significant associations between RH, AT and cumulative incidence rate or R_0 of COVID-19 (Yao et al. 2020a), which is in contrast to our finding that RH and AT are positively associated with COVID-19 case counts (Figure 3 and S1, respectively). Temperature in our study ranged from -12.48 to 21.29 °C, and our results of AT as positively associated with confirmed case counts is consistent with other studies in tropical countries and China indicating that COVID-19 can survive at higher temperatures, this reveals that COVID-19 can survival under higher temperatures. Experimental research on SARS-CoV-2 confirmed that this virus is very stable at 4 °C, but when the temperature increases to 70 °C, inactivation time for the virus is shortened to 5 min (Chin et al. 2020).

Hu et al. reported that low WV might increase the risk of mumps (Hu et al. 2018). Our multiple negative binomial regression models [lag 14 d in Figure 3(d1)] suggest that WV is negatively associated with COVID-19 confirmed case counts, consistent with Hu et al (Hu et al. 2018). Wind can affect the duration of droplets suspended in air. In indoor environments, high wind speed means that rooms are well-ventilated and in outdoor environments, high wind speed enhances dilution and removal of droplets, both of which could shorten droplet suspension time in the air (Cai et al. 2007).

Regarding rainfall, our study showed PRE is positively related to confirmed case counts (Figure 2); however, this effect was not statistically significant in simple and multiple negative binomial regression analysis (Figure S1 and 3). Some epidemiological data generally support a relationship between influenza virus infection and PRE (Pica &Bouvier 2012). Agrawal et al. found a significant association between rainfall and influenza virus infection in India, and fewer reports

of infection in the dry season (Agrawal et al. 2009). The association between PRE and respiratory syncytial virus infection showed similar results in our research and in other studies (Shek & Lee 2003).

To the best of our knowledge, interactive effects of air pollutants and meteorological factors have not previously been identified as affecting COVID-19 cases but have been identified in noninfectious diseases (Guo et al. 2019, Huang et al. 2017) and cause-specific diseases (Chen et al. 2019). In this study, we observed interactive effects on the total number of COVID-19 confirmed cases, especially for CO, which showed interactions with three meteorological factors (HS, DTR and RH; Table 2). The degree of increase in number of COVID-19 confirmed cases with CO depends on the level of certain meteorological factors. For example, effect of CO on confirmed cases is 5.763 ($\beta = 5.763$, $P = 0.001$) at low DTR level (< 50th percentile), but it is 0.931 ($\beta = 0.931$, $P = 0.035$) at high DTR level (Table 3) and the interactive effect was negative ($\beta = -0.365$, $P = 0.029$, Table 2), creating an antagonistic effect between these two factors. Although the interactive effect of CO and HS was statistically significant ($\beta = -0.811$, $P = 0.009$, Table 2), the stratified analysis did not find differences between the two categories of HS (Table 3), and thus the biological meaning of the interaction is unclear. Interactive effects of PM_{2.5} with PRE and SO₂ with AP were also statistically significant. Our findings suggest that environmental pollutants do not influence COVID-19 spread alone, but that they play important roles in combination with meteorological variables. The exact mechanisms of these interactive effects requires for further study.

We have comprehensively evaluated the effect of environmental pollution, meteorological factors and their interaction on the number of confirmed COVID-19 cases with and without controlling for MSI, to increase understanding of these risk

374 factors on the transmission of COVID-19. The limitation of our research is that we
375 only analyzed data from 120 cities in China. Further studies are required in more
376 countries and regions to clarify the effect of environmental air pollutants on
377 COVID-19 infection rates worldwide.

378 **5. Conclusion**

379 Our work verifies that air pollutants, meteorological factors and their interaction
380 affect the number of confirmed cases of COVID-19 even after controlling for
381 migration. Our study suggests that increased RH, CO and PM_{2.5} concentrations are
382 associated with increased numbers of confirmed COVID-19 cases. In addition,
383 pollutant-meteorological factor interaction effects influenced COVID-19 infection
384 levels. Our research provides a theoretical basis for formulating public health policies
385 for the prevention and control of COVID-19 that take into consideration the effects
386 and relationship of air pollutants and meteorological factors.

387

388 **Acknowledgments**

389 This work was supported by National Natural Science Foundation of China (Nos.
390 81872584 and 81273078), National 863 Young Scientist Program (No.
391 2015AA020940), Science and Technology Program of Guangzhou (No.
392 201704020056), and Scientific Research Project for University of Education Bureau
393 of Guangzhou (No. 201831841).

394

395 **Conflict of interest**

396 The authors declare that they have no conflict of interest.

397

398 **Author contributions**

399 Jianli Zhou: conceptualization, methodology, data curation, writing-original draft
400 preparation. Linyuan Qin: methodology, visualization, investigation, writing-original
401 draft preparation. Xiaojing Meng: writing-reviewing and editing. Nan Liu:
402 supervision, validation, writing-reviewing and editing.

403 **Appendix A. Supplementary materials**

404 Supplementary materials to this article can be found online at...

405 **Reference**

- 406 Agrawal AS, Sarkar M, Chakrabarti S, Rajendran K, Kaur H, Mishra AC, Chatterjee MK, Naik
407 TN, Chadha MS, Chawla-Sarkar M (2009) Comparative evaluation of real-time PCR and
408 conventional RT-PCR during a 2 year surveillance for influenza and respiratory syncytial
409 virus among children with acute respiratory infections in Kolkata, India, reveals a distinct
410 seasonality of infection. *J Med Microbiol.* 58: 1616-1622
- 411 Anwar A, Anwar S, Ayub M, Nawaz F, Hyder S, Khan N, Malik I (2019) Climate change and
412 infectious diseases: evidence from highly vulnerable countries. *Iran J Public Health.*
413 2187-2195
- 414 Auler AC, Cássaro FAM, da Silva VO, Pires LF (2020) Evidence that high temperatures and
415 intermediate relative humidity might favor the spread of COVID-19 in tropical climate:
416 A case study for the most affected Brazilian cities. *Sci Total Environ* 729: 139090
- 417 Bezirtzoglou C, Dekas K, Charvalos E (2011) Climate changes, environment and infection: facts,
418 scenarios and growing awareness from the public health community within Europe.
419 *Anaerobe* 17: 337-40
- 420 Cai QC, Lu J, Xu QF, Guo Q, Xu DZ, Sun QW, Yang H, Zhao GM, Jiang QW (2007) Influence
421 of meteorological factors and air pollution on the outbreak of severe acute respiratory
422 syndrome. *Public Health* 121: 258-65
- 423 Casadevall A (2020) Climate change brings the specter of new infectious diseases. *J Clin Invest*
424 130: 553-555
- 425 Chan KH, Peiris JS, Lam SY, Poon LL, Yuen KY, Seto WH (2011) The effects of temperature
426 and relative humidity on the viability of the SARS Coronavirus. *Adv Virol* 2011: 734690
- 427 Chen Y, Zheng M, Lv J, Shi T, Liu P, Wu Y, Feng W, He W, Guo P (2019) Interactions between
428 ambient air pollutants and temperature on emergency department visits: Analysis of
429 varying-coefficient model in Guangzhou, China. *Sci Total Environ* 668: 825-834
- 430 Chin AWH, Chu JTS, Perera MRA, Hui KPY, Yen H-L, Chan MCW, Peiris M, Poon LLM (2020)
431 Stability of SARS-CoV-2 in different environmental conditions. *The Lancet Microbe* 1:
432 e10
- 433 Cui Y, Zhang ZF, Froines J, Zhao J, Wang H, Yu SZ, Detels R (2003) Air pollution and case
434 fatality of SARS in the People's Republic of China: an ecologic study. *Environ Health* 2:
435 15
- 436 Guo Q, Xiong X, Liang F, Tian L, Liu W, Wang Z, Pan X (2019) The interactive effects between

437 air pollution and meteorological factors on the hospital outpatient visits for atopic
438 dermatitis in Beijing, China: a time-series analysis. *J Eur Acad Dermatol Venereol* 33:
439 2362-2370

440 Holshue ML et al. (2020) First case of 2019 novel coronavirus in the United States. *N Engl J*
441 *Med* 382: 929-936

442 Horne BD, Joy EA, Hofmann MG, Gesteland PH, Cannon JB, Lefler JS, Blagev DP, Korgenski
443 EK, Torosyan N, Hansen GI, Kartchner D, Pope CA, 3rd (2018) Short-term elevation
444 of fine particulate matter air pollution and acute lower respiratory infection. *Am J Respir*
445 *Crit Care Med* 198: 759-766

446 Hu Jianxiong HG, Liu Tao, Xiao Jianpeng, Rong Zuhua, Guo Lingchuan, Zeng Weilin, Zhu
447 Zhihua, Gong Dexin, Yin Lihua, Wan Donghua, Zeng Lilian, Ma Wenjun (2020) Risk
448 assessment of exported risk of COVID-19 from Hubei Province *Chinese Journal of*
449 *Preventive Medicine* 54: 632-366

450 Hu W, Li Y, Han W, Xue L, Zhang W, Ma W, Bi P (2018) Meteorological factors and the
451 incidence of mumps in Fujian province, China, 2005-2013: non-linear effects. *Sci Total*
452 *Environ* 619-620: 1286-1298

453 Huang CH, Lin HC, Tsai CD, Huang HK, Lian IB, Chang CC (2017) The interaction effects of
454 meteorological factors and air pollution on the development of acute coronary syndrome.
455 *Sci Rep* 7: 44004

456 Jiang Y, Wu XJ, Guan YJ (2020) Effect of ambient air pollutants and meteorological variables on
457 COVID-19 incidence. *Infect Control Hosp Epidemiol*: 1-11

458 Kesic MJ, Meyer M, Bauer R, Jaspers I (2012) Exposure to ozone modulates human airway
459 protease/antiprotease balance contributing to increased influenza A infection. *PLoS One*
460 7: e35108

461 Lai CC, Shih TP, Ko WC, Tang HJ, Hsueh PR (2020) Severe acute respiratory syndrome
462 coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic
463 and the challenges. *Int J Antimicrob Agents* 55: 105924

464 Liu J, Zhou J, Yao J, Zhang X, Li L, Xu X, He X, Wang B, Fu S, Niu T, Yan J, Shi Y, Ren X, Niu
465 J, Zhu W, Li S, Luo B, Zhang K (2020a) Impact of meteorological factors on the
466 COVID-19 transmission: a multi-city study in China. *Sci Total Environ* 726: 138513

467 Liu K, Chen Y, Lin R, Han K (2020b) Clinical features of COVID-19 in elderly patients: a
468 comparison with young and middle-aged patients. *J Infect* 80: e14-e18

469 Oliveiros B, Caramelo L, Ferreira NC, Caramelo F (2020) Role of temperature and humidity in
470 the modulation of the doubling time of COVID-19 cases. *medRxiv*:
471 2020.03.05.20031872

472 Pani SK, Lin NH, RavindraBabu S (2020) Association of COVID-19 pandemic with
473 meteorological parameters over Singapore. *Sci Total Environ* 740: 140112

474 Pica N, Bouvier NM (2012) Environmental factors affecting the transmission of respiratory
475 viruses. *Curr Opin Virol* 2: 90-5

476 Prata DN, Rodrigues W, Bermejo PH (2020) Temperature significantly changes COVID-19
477 transmission in (sub)tropical cities of Brazil. *Sci Total Environ* 729: 138862

478 Shek LP, Lee BW (2003) Epidemiology and seasonality of respiratory tract virus infections in the
479 tropics. *Paediatr Respir Rev* 4: 105-11

480 Siordia JA, Jr. (2020) Epidemiology and clinical features of COVID-19: A review of current

literature. *J Clin Virol* 127: 104357

Suhaimi NF, Jalaludin J, Latif MT (2020) Demystifying a Possible Relationship between COVID-19, Air Quality and Meteorological Factors: Evidence from Kuala Lumpur, Malaysia. *Aerosol Air Qual. Res* 20: 1520-1529

Tan J, Mu L, Huang J, Yu S, Chen B, Yin J (2005) An initial investigation of the association between the SARS outbreak and weather: with the view of the environmental temperature and its variation. *J Epidemiol Community Health* 59: 186-92

Tobias A, Molina T (2020) Is temperature reducing the transmission of COVID-19 ? *Environ Res* 186: 109553

Tosepu R, Gunawan J, Effendy DS, Ahmad OAI, Lestari H, Bahar H, Asfian P (2020) Correlation between weather and Covid-19 pandemic in Jakarta, Indonesia. *Sci Total Environ* 725: 138436

van Doremalen N, Bushmaker T, Munster VJ (2013) Stability of Middle East respiratory syndrome coronavirus (MERS-CoV) under different environmental conditions. *Euro Surveill* 18: 20590

Wang C, Horby PW, Hayden FG, Gao GF (2020a) A novel coronavirus outbreak of global health concern. *Lancet* 395: 470-473

Wang M, Jiang A, Gong L, Luo L, Guo W, Li C, Zheng J, Li C, Yang B, Zeng J, Chen Y, Zheng K, Li H (2020b) Temperature significant change COVID-19 Transmission in 429 cities. *medRxiv*: 2020.02.22.20025791

Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, Hu YA-O, Tao ZW, Tian JH, Pei YY, Yuan ML, Zhang YL, Dai FH, Liu Y, Wang QM, Zheng JJ, Xu L, Holmes EA-O, Zhang YZ (2020) A new coronavirus associated with human respiratory disease in China. *Nature*. 579: 265-269

Xie J, Zhu Y (2020) Association between ambient temperature and COVID-19 infection in 122 cities from China. *Sci Total Environ* 724: 138201

Xu Z, Hu W, Fau - Williams G, Williams G, Fau - Clements ACA, Clements AC, Fau - Kan H, Kan H, Fau - Tong S, Tong S (2013) Air pollution, temperature and pediatric influenza in Brisbane, Australia. *Environ Int.* 59: 384-388

Yao Y, Pan J, Liu Z, Meng X, Wang W, Kan H (2020a) No association of COVID-19 transmission with temperature or UV radiation in Chinese cities. *Eur Respir J*: 2000517

Yao Y, Pan J, Liu Z, Meng X, Wang W, Kan H (2020b) No association of COVID-19 transmission with temperature or UV radiation in Chinese cities. *Eur Respir J* 55: 2000517

Yusuf S, Piedimonte G, Auais A, Demmler G, Krishnan S, Van Caesele P, Singleton R, Broor S, Parveen S, Avendano L, Parra J, Chavez-Bueno S, Murguia De Sierra T, Simoes EA, Shaha S, Welliver R (2007) The relationship of meteorological conditions to the epidemic activity of respiratory syncytial virus. *Epidemiol Infect* 135: 1077-90

Zhou P et al. (2020) A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579: 270-273

Zhu Y, Xie J, Huang F, Cao L (2020) Association between short-term exposure to air pollution and COVID-19 infection: Evidence from China. *Sci Total Environ* 727: 138704

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567 **Figure Captions**

568 **Figure 1.** Locations of 120 cities and total confirmed cases in each city on 18 Mar.

569 2020, and correlation analysis of meteorological factors and air pollutants.

570 NA: Cities not included in this study.

571

572 **Figure 2.** Correlation analysis of confirmed case counts with air pollution and

573 meteorological parameters.

574 $\text{Log (CCC)} = \log (\text{confirmed case counts})$

575

576 **Figure 3.** The relationship between air pollutants, meteorological factors and

577 confirmed case counts in a multiple negative binomial regression model with and

578 without adjusting for migrations with MSI.

579

580

(a)

(b)

(a1) **(b1)** **(c1)** **(d1)** **(a2)** **(b2)** **(c2)** **(d2)**

Ambient air pollutants

Interaction

Meteorological factors

Adjusting
population
migration

COVID-19
confirmed
cases

Negative
binomial
regression
analysis