

S2 Table. Summary of model parameters.

Parameter	Definition	Value
n_E	No. of latency Erlang states	16
n_P	No. of prodromal Erlang states	16
n_I	No. of fully contagious Erlang states	16
n_L	No. of late-infectious Erlang states	16
D_E	Average duration of latency period	3.7 days
D_P	Average duration of prodromal period	1 day
D_I	Average duration of fully contagious period	7 days
D_L	Average duration of late infectious period	7 days
ε	Transition rate of latent states	n_E/D_E
φ	Transition rate of prodromal states	n_P/D_P
γ	Transition rate of early infectious states	n_I/D_I
δ	Transition rate of late-infectious states	n_L/D_L
α	Transition rate from transient multi infections to multi-infected states	3.2/day
f_{Sick}	Fraction of symptomatic (sick) infections	58%
\tilde{f}_{Sick}	Fraction of sympt. multi infections in fully contagious & late inf. periods	64.4%
f_{Iso}	Fraction of single-infected (sick) who are isolated (or home isolated)	0.5
\tilde{f}_{Iso}	Fraction of multi-infected (sick) who are isolated (or home isolated)	0.55
f_{Dead}	Fraction of single-infected (sick) who die from the disease	3%
\tilde{f}_{Dead}	Fraction of multi-infected (sick) who die from the disease	4%

Summary of parameters describing number of Erlang states, durations, transition rates, morbidity and mortality, and their default parameter choices.