Socio-educational Impact and Psychological Distress of Medical Students amid the COVID-19 Pandemic: A Japanese Cross-Sectional Survey

Yoshito, Nishimura^{1*}, Kanako, Ochi^{1,2}, Kazuki, Tokumasu¹, Mikako, Obika¹, Hideharu, Hagiya¹, Hitomi, Kataoka^{1,3}, Fumio, Otsuka¹

¹Department of General Medicine, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama, 7008558, Japan

²Center for Developing Medical and Health Care Education, Okayama University Medical School, Okayama, 7008558, Japan

³Center for Diversity and Inclusion, Okayama University Hospital, Okayama, 7008558, Japan

***Corresponding author**

Yoshito, NISHIMURA MD, PhD, MPH

Department of General Medicine, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Kita-ku, Okayama, Japan.

Tel.: +81-86-235-7342; Fax: +81-86-235-7345

E-mail: nishimura-yoshito@okayama-u.ac.jp

Abstract

The COVID-19 pandemic has negatively affected medical education. However, few data are available about medical students' distress during the pandemic. This study aimed to provide details on how medical students had been affected by the pandemic. In this cross-sectional study, 717 medical students participated in the web-based survey. The questions included how their mental status had changed before and after the Japanese nationwide state of emergency (SOE). 65.9% (473/717) participated in the study. 29.8% (141/473) reported concerns about the shift toward online education, mostly because they thought online education could have been ineffective compared with in-person learning. Participants' subjective mental health status significantly worsened after the SOE was lifted ($p < .001$). Those who had concerns about a shift toward online education had higher odds of having generalized anxiety and being depressed (OR 1.97, 95% CI 1.19 – 3.28), as did those who requested food aid and mental health care resources (OR 1.99, 95% CI 1.16 – 3.44; OR 3.56, 95% CI 2.07 – 6.15, respectively). Given our findings, the sudden shift to online education might have overwhelmed medical students. Thus, we recommend educators to inform learners that online learning is non-inferior to in-person learning, which could attenuate potential depression and anxiety.

Keywords: coronavirus disease 2019 (COVID-19); online education; depression; pandemic

Introduction

The coronavirus disease 2019 (COVID-19) global pandemic has drastically changed our lives, with more than eight million cases and 400,000 deaths reported globally as of June 18, 2020, according to the World Health Organization statistics(1). In Japan, 17,668 cases have been confirmed as of June 18, 2020, with an explosion in the number of cases in early April(2). While Japan seems to be bringing the outbreak under control through active cluster tracing, restriction of mass gatherings, and advocating universal masking and hand hygiene, there were no simple solutions to this issue(3). After the surge in COVID-19 cases, on April 16, 2020, the Government of Japan (GoJ) declared a state of emergency (SOE) in all 47 prefectures, which lasted until May 25, 2020(4). In Okayama, a prefecture in the western part of mainland Japan(5) with approximately 1.9 million people (ranked 20/47 among the 47 prefectures), only 25 confirmed cases had been reported by the end of May 2020. Behind the scenes of successful COVID-19 mitigation, however, those in education-related jobs and medical students struggled with the rapid change in the socio-educational system.

Although the SOE was lifted on May 14 in Okayama based on the low incidence of COVID-19, even before the SOE, from March 25, Okayama University School of Medicine (OUSM), one of the largest national universities in Japan, requested its students to stay home to prevent the

possible community spread of COVID-19 within the medical school and hospital. Until the stay-home request was lifted on May 22, medical students were required to cope with this sudden change in their lifestyle, and shifted to online education amid the fear and considerable uncertainty surrounding COVID-19. In the early 2000s, the severe acute respiratory syndrome coronavirus (SARS-CoV) outbreak had a devastating impact on academic education, including sudden curriculum changes and rapid integration of information technology(6-8). Similarly, the current COVID-19 pandemic has provoked significant turmoil in society. In particular, mental health problems due to the pandemic have drawn attention worldwide as studies have suggested the need for mental health care interventions during the outbreak(9-14). Medical students, who essentially need clinical exposure, may have been impacted even further by the pandemic.

In Japan, medical schools have different admission practices and curricula than in the United States. Students typically enter medical school immediately after high school graduation, often at 18 years old, and they go through six years of medical education before graduation. Despite the differences in education system, the fundamental philosophy of academic medicine is the same; to provide quality educational experience. To date, few published studies have investigated the status of medical students' living environment, socio-educational, and mental health status during the COVID-19 pandemic. To address this, we conducted a total population survey of OUSM

medical students to comprehensively clarify what they require and how they have been mentally affected by COVID-19.

Methods

Study Design, setting, and participants

We performed a cross-sectional study that employed an anonymous, self-administered voluntary web-based survey. Participants comprised medical students in all years of study at OUSM. We used purposive sampling to conduct a total population sample for the 717 medical students who belonged to the OUSM as of April 1, 2020 (the first day of the academic year in Japan). The participants' consent was implied by their completion of the survey.

The survey was developed through consultation with a medical education expert panel at OUSM and piloting, and administered with Qualtrics (Qualtrics International Inc., Provo, Utah), a web-based survey platform. We provided survey instructions and instruments in Japanese. We distributed survey links to the students using OUSM official mailing lists. All participants were invited to complete the survey within one week (June 8–14, 2020, in Japan Standard Time). No financial incentives were provided for their participation in the survey. In the survey, we included entries on demographics (age, gender, education before entering medical school, employment

status on date of response, changes in employment status due to the COVID-19 pandemic, marital status, living environment, household size, and comorbidities) as well as COVID-19 related items (e.g., “Chance of contracting COVID-19 during the current pandemic”), self-learning associated activities (e.g., Average time of self-learning/day), validated depression and anxiety scale instruments, and financial situations. To protect participants’ anonymity as much as possible, respondents were not prompted to enter their year of study.

Measurements

COVID-19-related questions

To evaluate the extent of participants’ concern and preparedness for COVID-19, they were asked: “What do you think the chances are that you will contract COVID-19 during the current pandemic?”; “What is your degree of concern about the health status of your family?”; “Do you think you have enough information about the symptoms of COVID-19?”; “Do you think you have enough information about prevention and treatment of COVID-19?”; and “Do you feel worried about COVID-19?” To further assess students’ concerns, the participants were prompted to answer “I am concerned because my future career formation may be negatively affected due to the COVID-19 pandemic”; “I am concerned because the COVID-19 pandemic may attenuate our relationship to teachers”; “I am concerned because of the disruption to ongoing research or

extracurricular activities”; and “I am concerned about the shift toward online education.” All questions were evaluated on a 5-point Likert scale except for the item on the health status of the family (3-point Likert scale). Those who answered “Very concerned” or “Concerned” in the entry on concern about the shift toward online education were prompted to provide reasons. To describe participants’ needs, they were asked to note the types of support they wish to receive from the university if there is a resurgence of COVID-19. These responses were mandatory.

Self-learning and related activity

For subjective mental health status and the average time per day that participants stayed at home, read books, played video games, and learned by themselves, respondents were prompted to answer “How many hours a day did you... / stay at home / read books / play video games / self-learn?” before the SOE order (April 16, 2020) or during the last two weeks (with the base date of when participants completed the survey).

Depression and anxiety disorders

We assessed the presence of depression using the 9-item Patient Health Questionnaire (PHQ-9), a common screening tool for mood disorders. We used the validated Japanese translation of the scale(15). The total scores of PHQ-9 range from 0 to 27 and we defined scores of 10 or more as

having “depression.” We screened for anxiety disorders using the Japanese version of the 7-item Generalized Anxiety Disorder Scale (GAD-7), which was validated in 2010(16). The score ranges from 0 to 21, and we defined scores of 10 or more as having “anxiety.” Both instruments ask respondents about their mental health status during the last two weeks.

Financial situation

In Japan, university students typically live on a monthly allowance from parents. According to the latest statistics by the National Federation of University Co-operative Associations, students receive 72,810 JPY (approximately 680 USD) a month on average(17). Participants were prompted to give their monthly allowance from the following options: “None,” “0,” “< 30,000 JPY (280 USD),” “30,000 – 49,999 JPY (280 – 467 USD),” “50,000 – 69,999 JPY (467 – 654 USD),” “70,000 – 99,999 JPY (654 – 935 USD),” and “≥ 100,000 JPY (≥ 935 USD).” Respondents were also asked to answer if they were on scholarship or student loan.

Statistical analysis

We analyzed the data using JMP version 13.1.0 (SAS Institute Inc., Cary, North Carolina). We used the Wilcoxon signed-rank test to examine differences in the time participants spent on self-learning related activities based on non-normal distribution. For associations between

categorical variables containing small sample sizes, we employed Fisher’s exact test. To examine the predictive factors of categorical dependent variables, we used univariate logistic regression analyses. The threshold for significance was defined as $p < 0.05$.

Ethical approval

This study protocol was approved by the institutional review board of Okayama University Hospital (reference number 2006-029, approved on June 5, 2020).

Data Availability Statement

The datasets generated and analysed during the current study are available from the corresponding author on reasonable request.

Results

Response rates to the survey were 473 out of 717 (65.9%) OUSM students in all six years combined. Participants’ demographic characteristics are summarized in **Table 1**. Of note, 250 (52.9%) respondents reported that they were engaged in part-time work, while 44 (9.3%) reported having resigned or been fired due to the COVID-19 pandemic. Eight (1.7%) and 6 (1.3%) noted that they had a past medical history of anxiety disorders and depression,

respectively.

COVID-19 related survey items

Table 2 and **Supplementary Figure 1** show respondents' answers to the COVID-19 related survey items. Eighty-one (17.2%) responded that they were either "likely" or "very likely" to contract COVID-19 during the ongoing pandemic; 275 (58.1%) agreed or strongly agreed to a prompt "I feel worried about COVID-19." Regarding the breakdown of students' concerns about COVID-19, 182 (38.5%), 121 (25.6%), and 235 (49.7%) acknowledged that they were concerned about the negative impacts of COVID-19 on their future career formation, relationship with teachers, and ongoing research or extracurricular activities, respectively, while 141 (29.8%) also reported concern about a shift toward online education. The reasons for concerns included ineffective educational effect compared with on-site education ($n = 92$, 65.2%), possible resurgence of the COVID-19 outbreak leading to a sudden change in the curriculum ($n = 74$, 52.5%), and decreased clinical exposure ($n = 92$, 65.2%).

Of the participants, 298 (63.0%) answered that they would request financial aid if a stay-home order recurred due to a COVID-19 resurgence, followed by request for food aid ($n = 100$, 21.1%), technical support for online education ($n = 100$, 21.1%), and mental health care resources ($n = 85$,

18.0%).

Change in self-learning related time before and after the SOE

We compared the change in average time respondents spent at home, reading books, playing video games, and self-learning per day before the SOE and within two weeks prior to the survey completion (after the SOE). As shown in **Table 3** and **Supplementary Figure 2**, the participants spent significantly longer on all the activities mentioned above after the SOE than before the SOE ($p < .001$). There were also significant differences in their subjective mental health status before and after the SOE based on Wilcoxon signed-rank test ($p < .001$).

Regression analyses of factors associated with depression and anxiety

Of the participants, 75 (15.8%) had PHQ-9 scores of 10 or more, and 34 (7.2%) had GAD-7 scores of 10 or more. **Table 4** presents all the results of univariate regression analyses. The odds of being depressed were significantly higher in those who had concerns about a shift toward online education (OR 1.97, 95% CI 1.19 – 3.28), and in those who requested food aid (OR 1.99, 95% CI 1.16 – 3.44) and mental health care resources (OR 3.56, 95% CI 2.07 – 6.15) from the university in the event of resurgence of COVID. Regarding generalized anxiety, the odds were higher in respondents who requested food aid (OR 2.50, 95% CI 1.21 – 5.20) and mental health

care resources (OR 3.16, 95% CI 1.51 – 6.59).

Prevalence of financial hardship

Table 5 and **Supplementary Figure 3** show the self-reported amounts of monthly allowance from parents and situations regarding financial aid. A total of 332 (70.2%) respondents had a monthly allowance that was lower than the national average (72,810 JPY, approximately 680 USD)(17), while 141 (27.7%) received some sort of financial aid for their living and education.

Discussion

To the best of our knowledge, this study is the first survey of Japanese medical students regarding their life circumstances and challenges due to COVID-19. Although Japan has had fewer COVID-19 cases than the US, and Okayama has had comparatively fewer cases than other Japanese prefectures, the study results underscore that the pandemic caused a profound negative socio-educational impact on our society and medical students. We found that approximately 10% of the students on part-time jobs had been fired due to the pandemic. Furthermore, a considerable number of medical students had concerns around their basic life security, demanding support from the university in the form of financial aid, food aid, technical support, and mental health care resources due to the SOE. Regarding students' subjective psychological distress, those who

expressed concerns about the rapid shift toward online education and fear around basic life security were more likely to be depressed and anxious after the SOE.

Concerns around future career disruption, attenuated relationship to medical teachers, and disruption of ongoing extracurricular activities were prevalent among the participants, which underlines the considerable uncertainty amid the COVID-19 pandemic (Table 2). In particular, 63% of respondents reported the need for financial aid in the event of a second wave of the pandemic. These data correspond to the fact that more than 70% of participants received no more than the national average monthly allowance from their parents (Table 5). Contrary to the general notion in Japan that “medical students are financially well-off,” many may have experienced financial hardship. Recently, the theory of willpower has gathered public attention. Financial stability is considered a prerequisite of willpower, which aids appropriate decision-making(18). Combined with previous research findings that financial instability could lead to worse mental health outcome(19, 20), educators are expected to develop a strategy to offer financial support to those in need.

Regarding lifestyle changes and psychological distress, after the SOE was lifted, the participants experienced significantly worse mental health status, and spent significantly longer at home,

reading books, playing video games, and learning by themselves (Table 3) than before the announcement of SOE. Our results are consistent with prior longitudinal and cross-sectional studies that the COVID-19 pandemic led to a sedentary life and psychological distress(9, 12, 13, 21). As suggested by a previous study, the increase in gaming could be a coping behavior against psychological stress(22).

As for regression analyses of factors associated with depression and anxiety, in our study population, those who requested food aid and mental health care resources from the university upon the future resurgence of the COVID-19 outbreak had significantly higher odds of depression and anxiety. Furthermore, concerns around the shift toward online education were identified as a factor associated with depression (Table 4). Surprisingly, 65.2% of those who had concerns about the shift toward online education thought online education was less effective than in-person education. While previous studies have reported the utility and noninferiority of online learning compared to offline(23, 24), in-person learning, the results of the current study revealed a potential gap in perception regarding the effectiveness of online education between medical students and educators. Educators should not assume that students know the potential benefits of online learning, and it is essential to inform learners that online learning is non-inferior to in-person learning, which could attenuate potential depression and anxiety. While in-person

communication has become difficult nowadays due to the fear for COVID-19, medical schools may need to reach out to students to find out those suffering from underlying life insecurity and to provide multilateral support, including early mental health care interventions.

Several limitations to this study should be noted. First, due to the single-center cross-sectional survey design, we may not conclude causal relationships. Second, the cross-sectional research contains a limitation in terms of addressing changes over time. While we illustrated the changes in students' average time spent on self-learning related activities before and after the SOE in Table 3, a longitudinal study design would be more appropriate to examine the differences in the study cohort. Third, we asked participants to provide their mental health status and time spent on self-learning related activities before the SOE (approximately six weeks before the survey implementation), both of which are subject to recall bias. Also, PHQ-9 and GAD-7 scores were obtained only after the SOE. Thus, it is possible that they might have not changed during the period. Lastly, due to the nature of the survey topic, those who were interested in the public health emergency or mental health may have been more likely to respond, which would lead to self-selection bias. Despite these limitations, a total population sampling strategy coupled with higher-than-usual response rates(25) contributed to high internal validity.

271 In conclusion, through the study, we have provided graphical data and evidence regarding the
 272 socio-educational impacts of the COVID-19 pandemic on medical students. In circumstances of
 273 considerable uncertainty, both educators and medical students need to be flexible, patient, and
 274 resilient. The uncertainty and drastic change triggered substantial psychological distress in
 275 students, which was greater than we assumed. As discussed, Japan has had a relatively small
 276 number of COVID-19 cases compared to the US and other European countries. Although
 277 Okayama prefecture survived the pandemic with fewer confirmed COVID-19 cases than other
 278 prefectures, medical students experienced significant impacts due to the public health emergency.
 279 The educational institution should recognize the prevalence of basic needs insecurity, such as
 280 financial difficulties and a shortage of staples, including food, etc. As medical educators, we
 281 need to be accountable for the advantages of online education in the field of medicine to alleviate
 282 students' psychological distress, in addition to providing multilateral support to those in need,
 283 including early mental health care interventions. While we targeted medical students in a single
 284 Japanese national university, the survey results warrant further research and analysis if the
 285 distress is amplified by existing anxiety, depression, burnout, etc., or is a wholly
 286 COVID-19-related phenomenon. We call for increased research in populations with more
 287 COVID-19 cases than Japan, to figure out the challenges medical students have been facing amid
 288 the pandemic in different cultures and backgrounds.

289

290 **References**

291 1. World Health Organization. Coronavirus disease (COVID-19) Situation Report - 150

292 2020 [Available from:

293 <https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200618-covid-19-sitre>

294 [p-150.pdf?sfvrsn=aa9fe9cf_2](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200618-covid-19-sitre-p-150.pdf?sfvrsn=aa9fe9cf_2).

295 2. Ministry of Health LaWJ. About Coronavirus Disease 2019 (COVID-19) 2020

296 [Available from: https://www.mhlw.go.jp/stf/seisakunitsuite/bunya/newpage_00032.html.

297 3. Looi MK. Covid-19: Japan ends state of emergency but warns of "new normal". BMJ.

298 2020;369:m2100.

299 4. Government of Japan. Basic Policies for Novel Coronavirus Disease Control by the

300 Government of Japan (Summary) 2020 [Available from:

301 <https://www.mhlw.go.jp/content/10900000/000634753.pdf>.

302 5. Statistics Bureau of Japan. Current Population Estimates as of October 1, 2019 2019

303 [Available from: <https://www.stat.go.jp/english/data/jinsui/2019np/index.html>.

304 6. Clark J. Fear of SARS thwarts medical education in Toronto. BMJ.

305 2003;326(7393):784.

306 7. Patil NG, Chan Y, Yan H. SARS and its effect on medical education in Hong Kong. Med

307 Educ. 2003;37(12):1127-8.

308 8. Sherbino J, Atzema C. "SARS-Ed": severe acute respiratory syndrome and the impact
309 on medical education. *Ann Emerg Med.* 2004;44(3):229-31.

310 9. Huckins JF, DaSilva AW, Wang W, Hedlund E, Rogers C, Nepal SK, et al. Mental
311 Health and Behavior During the Early Phases of the COVID-19 Pandemic: A Longitudinal
312 Mobile Smartphone and Ecological Momentary Assessment Study in College Students. *J Med
313 Internet Res.* 2020.

314 10. Liu S, Yang L, Zhang C, Xiang YT, Liu Z, Hu S, et al. Online mental health services in
315 China during the COVID-19 outbreak. *Lancet Psychiatry.* 2020;7(4):e17-e8.

316 11. Shigemura J, Kurosawa M. Mental health impact of the COVID-19 pandemic in Japan.
317 *Psychol Trauma.* 2020.

318 12. Talevi D, Socci V, Carai M, Carnaghi G, Faleri S, Trebbi E, et al. Mental health
319 outcomes of the CoViD-19 pandemic. *Riv Psichiatri.* 2020;55(3):137-44.

320 13. Wang C, Pan R, Wan X, Tan Y, Xu L, Ho CS, et al. Immediate Psychological Responses
321 and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19)
322 Epidemic among the General Population in China. *Int J Environ Res Public Health.* 2020;17(5).

323 14. Zhang Y, Zhang H, Ma X, Di Q. Mental Health Problems during the COVID-19
324 Pandemics and the Mitigation Effects of Exercise: A Longitudinal Study of College Students in

China. Int J Environ Res Public Health. 2020;17(10).

15. Muramatsu K, Miyaoka H, Kamijima K, Muramatsu Y, Tanaka Y, Hosaka M, et al. Performance of the Japanese version of the Patient Health Questionnaire-9 (J-PHQ-9) for depression in primary care. Gen Hosp Psychiatry. 2018;52:64-9.

16. Muramatsu K, Miyaoka H, Ueshima K, Muramatsu Y, Fuse K, Yoshimine H, et al. Validation and utility of a Japanese version of the GAD-7. Jpn J Psychosom Med. 2010;50(6):592.

17. National Federation of University Co-operative Associations. Summary of Campus Life Data 2019 [Article in Japanese] 2019 [Available from: <https://www.univcoop.or.jp/press/life/report.html>].

18. Baumeister RF, Tice DM, Vohs KD. The Strength Model of Self-Regulation: Conclusions From the Second Decade of Willpower Research. Perspect Psychol Sci. 2018;13(2):141-5.

19. Richardson T, Elliott P, Roberts R, Jansen M. A Longitudinal Study of Financial Difficulties and Mental Health in a National Sample of British Undergraduate Students. Community Ment Health J. 2017;53(3):344-52.

20. McCloud T, Bann D. Financial stress and mental health among higher education students in the UK up to 2018: rapid review of evidence. J Epidemiol Community Health.

343 2019;73(10):977-84.

344 21. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The
345 psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*.
346 2020;395(10227):912-20.

347 22. Balhara YPS, Kattula D, Singh S, Chukkali S, Bhargava R. Impact of lockdown
348 following COVID-19 on the gaming behavior of college students. *Indian J Public Health*.
349 2020;64(Supplement):S172-S6.

350 23. Pei L, Wu H. Does online learning work better than offline learning in undergraduate
351 medical education? A systematic review and meta-analysis. *Med Educ Online*.
352 2019;24(1):1666538.

353 24. Seymour-Walsh AE, Weber A, Bell A. Practical approaches to pedagogically rich online
354 tutorials in health professions education. *Rural Remote Health*. 2020;20(2):6045.

355 25. Cunningham CT, Quan H, Hemmelgarn B, Noseworthy T, Beck CA, Dixon E, et al.
356 Exploring physician specialist response rates to web-based surveys. *BMC Med Res Methodol*.
357 2015;15:32.

358

359

360

361

362

363

364

365

366 **Acknowledgments**

367 We would like to thank the Academic Affairs Division of Okayama University Medical School

368 for its cooperation in conducting the survey.

369

370 **Author contributions**

371 YN wrote the manuscript, designed the study, and analysed the data. KO, KT, MO, HH, and HK

372 analyzed the data and revised the manuscript. FO designed and supervised the research.

373

374 **Conflicts of interest**

375 The authors declare no conflicts of interest in association with the present study.

376

377 **Funding**

378 None.

379

380

381

382

383

384 **Figure Legend**

385 **Supplementary Figure 1**

386 (A) The breakdown of students' concerns about COVID-19. 182 (38.5%), 121 (25.6%), and 235
387 (49.7%) acknowledged that they were concerned about the negative impacts of COVID-19 on
388 their future career formation, relationship with teachers, and ongoing research or extracurricular
389 activities, respectively, while 141 (29.8%) also reported concern about a shift toward online
390 education. (B) Of the participants, 298 (63.0%) answered that they would request financial aid if
391 a stay-home order recurred due to a COVID-19 resurgence, followed by request for food aid (n =
392 100, 21.1%), technical support for online education (n = 100, 21.1%), and mental health care
393 resources (n = 85, 18.0%).

394

395 **Supplementary Figure 2**

396 The bar charts show the change in average time respondents spent at home, reading books,

397 playing video games, and self-learning per day before the SOE and within two weeks prior to the
398 survey completion (after the SOE). Error bars show the 95% confidence intervals.

399

400 **Supplementary Figure 3**

401 The self-reported amounts of monthly allowance from parents and situations regarding financial
402 aid are shown.

Table 1. Demographic Characteristics of the Study Participants Comprising 473

Japanese Medical Students

Characteristic	Value	SD	95% CI
Age in years^a			
Mean	22.0	3.3	21.7-22.3
Gender, no. (%)			
Female	161 (34.0)		
Male	311 (65.8)		
Non-confirming	1 (0.2)		
Education before medical school, no. (%)			
High school	434 (91.8)		
Career college	1 (0.2)		
4-year university	31 (6.6)		
Master's degree	7 (1.5)		
Employment status, no. (%)			
Part-time job	250 (52.9)		
Schoolwork only: no extra work	221 (46.7)		
Self-owned business	2 (0.4)		

Resigned or fired due to the pandemic	44 (9.3)		
Marital status, no. (%)			
Single	467 (98.7)		
Married	5 (1.1)		
Missing	1 (0.2)		
Living environment (%)			
Living alone	308 (65.1)		
Living with family	127 (26.9)		
Living with a partner	20 (4.2)		
Living alone: family or relatives nearby	18 (3.8)		
Household size			
Mean	1.7	1.3	1.6-1.8
Comorbidity, no. (%)			
None	401 (84.8)		
Asthma	37 (7.8)		
Anxiety disorder	8 (1.7)		
Depression	6 (1.3)		
Hypertension	5 (1.1)		

Heart diseases	3 (0.6)
Dyslipidemia	2 (0.4)
Other	24 (5.1)
<hr/>	
Total number of participants	473
<hr/>	

Abbreviations: CI, confidence interval; SD, standard deviation.

^a Of the 473 respondents, 2 did not specify their age.

Table 2. Results of the COVID-19-related Survey Items

Survey Entry	No. (%)
“Chance of contracting COVID-19 during the current pandemic”	
Very likely	15 (3.2)
Likely	66 (14.0)
Neutral	238 (50.3)
Unlikely	121 (25.6)
Very unlikely	33 (6.9)
“Degree of concern about the health status of your family”	
Very concerned	198 (41.9)
Somewhat concerned	237 (50.1)
Not concerned	38 (8.0)
“I have enough knowledge about symptoms of COVID-19”	
Strongly agree	8 (1.7)
Agree	158 (33.4)
Neutral	169 (35.7)
Disagree	116 (24.5)
Strongly disagree	22 (4.7)

“I have enough knowledge about the treatment of COVID-19”

Strongly agree	6 (1.3)
Agree	65 (13.7)
Neutral	138 (29.2)
Disagree	196 (41.4)
Strongly disagree	68 (14.4)

“I feel worried about COVID-19”

Strongly agree	46 (9.7)
Agree	229 (48.4)
Neutral	106 (22.4)
Disagree	77 (16.3)
Strongly disagree	15 (3.2)

Students’ concern	Future Career	Attenuated	Disruption of	Shift towards
	Formation due	Relationship to	Ongoing	Online
	to COVID-19	Teachers	Research or	Education
			Extracurricular	
			Activities	

	No. (%)	No. (%)	No. (%)	No. (%)
Very concerned	46 (9.7)	20 (4.2)	66 (14.0)	18 (3.8)
Concerned	136 (28.8)	101 (21.4)	169 (35.7)	123 (26.0)
Neutral	130 (27.5)	136 (28.7)	91 (19.2)	115 (24.3)
Not concerned	85 (18.0)	129 (27.3)	67 (14.2)	121 (25.6)
Not at all concerned	76 (16.0)	87 (18.4)	80 (16.9)	96 (20.3)

“The Reasons for Concern about Shift toward Online Education”^a

Online education may not be as effective as on-site education	92 (65.2)
Fear of possible resurgence of COVID-19 outbreak, which would cause a sudden change in the curriculum	74 (52.5)
Less clinical exposure	92 (65.2)
Other	16 (11.3)

“What supports do you want from the university if a stay-home order takes

place again due to the resurgence of COVID-19?”^b

Financial aid (e.g., tuition waiver, no-interest student loan)	298 (63.0)
Food aid	100 (21.1)
Resources for mental health care	85 (18.0)
Technical support (e.g., free rental of PC, pocket WiFi)	100 (21.1)

Basic livelihood support (e.g., resources for health maintenance or preventing DV)	57 (12.1)
Parenting support	6 (1.3)
Other	55 (11.6)

Abbreviations: COVID-19, coronavirus disease 2019; PC, personal computer.

^a The survey entry was explicitly answered by participants who answered “Very concerned” or “Concerned” in the previous entry (“Concerned about Shift toward Online Education,” $n = 141$). Multiple answers permitted.

^b Multiple answers permitted.

Table 3. Comparison of Age-weighted Self-learning and Related Activity Time of

Medical Students Before and After the National State of Emergency in Japan

	Before SOE		Last 2 Weeks ^a		<i>P</i> value ^b
	Time (h)	95% CI	Time (h)	95% CI	
Measure					
Average Time at Home / Day	16.1	15.7-16.6	18.8	18.5-19.2	< .001
Average Time Reading Books / Day	0.70	0.58-0.82	1.1	0.96-1.2	< .001
Average Time Playing Video Games / Day	1.5	1.3-1.7	2.1	1.9-2.3	< .001
Average Time Self-learning / Day	2.6	2.4-2.8	4.0	3.7-4.2	< .001
	Before SOE		Last 2 Weeks		
	No. (%)		No. (%)		
Subjective Mental Health Status ^c					
Very good	85 (18.0)		39 (8.2)		
Good	171 (36.1)		58 (12.2)		
Fair	193 (40.8)		286 (60.5)		
Bad	18 (3.8)		76 (16.1)		
Very bad	6 (1.3)		14 (3.0)		

Abbreviations: CI, confidence interval; h, hour; SOE, state of emergency.

^a The base date of the “last 2 weeks” is the day when the participants answered the

survey. The survey period was from June 8 to 14, 2020.

^b P value is calculated with Wilcoxon signed-rank test.

^c $P < .001$ with Wilcoxon signed-rank test. There is a significant difference in the subjective mental health status of participants before the Japanese national state of emergency and the last two weeks prior to the date of answering the survey.

Table 4. Results of Univariate Regression Analyses of Factors Associated with Depression and Generalized Anxiety in Japanese Medical Students

Dependent Variable	Independent Variable	OR (95% CI) ^a	P value
Depression (<i>n</i> = 75) (PHQ-9 score ≥ 10)	Female (vs. Male)	1.08 (0.58 – 2.01)	.81
	Age	1.06 (0.96 – 1.16)	.45
	Concerns around Future Career Formation ^b	1.40 (0.85 – 2.30)	.18
	Concerns around Attenuated Relationship to Teachers ^b	1.46 (0.85 – 2.50)	.17
	Concerns around Disruption of Ongoing Research or Extracurricular Activities ^b	0.81 (0.50 – 1.33)	.41
	Concerns around Shift toward Online Education ^b	1.97 (1.19 – 3.28)	.009
	Request for Financial Aid	0.92 (0.55 – 1.53)	.74
	Request for Food Aid	1.99 (1.16 – 3.44)	.013
	Request for Mental Health Care Resources	3.56 (2.07 – 6.15)	<.0001
Generalized Anxiety (<i>n</i> = 34) (GAD-7 score ≥ 10)	Female (vs. Male)	0.81 (0.37 – 1.74)	.58
	Age	0.96 (0.85 – 1.09)	.57
	Concerns around Future Career Formation ^b	0.75 (0.36 – 1.58)	.45
	Concerns around Attenuated Relationship to Teachers ^b	1.43 (0.68 – 3.03)	.35

Concerns around Disruption of Ongoing Research or Extracurricular Activities ^b	0.53 (0.26 – 1.09)	.09
Concerns around Shift toward Online Education ^b	1.50 (0.73 – 3.10)	.27
Request for Financial Aid	1.25 (0.59 – 2.62)	.56
Request for Food Aid	2.50 (1.21 – 5.20)	.014
Request for Mental Health Care Resources	3.16 (1.51 – 6.59)	.002

Abbreviations: CI, confidence interval; GAD-7, Generalized Anxiety Disorder-7; OR, odds ratio; PHQ-9, Patient Health Questionnaire-9.

^a Univariate regression analysis was performed for each dependent variable.

^b Participants who answered, “Very concerned” or “Concerned” in the respective survey entries listed in Table 2 were considered the “Positive concerns group.” On the other hand, those who answered “Neutral,” “Not concerned,” or “Not at all concerned” were considered the “No concerns group” in this analysis.

Table 5. Description of the Financial Situation of Japanese Medical Students

Characteristic	No. (%)
Monthly allowance from parents (JPY) ^a	
No allowance	127 (26.9)

< 30,000 JPY	43 (9.1)
30,000 – 49,999 JPY	81 (17.1)
50,000 – 69,999 JPY	81 (17.1)
70,000 – 99,999 JPY	78 (16.5)
≥ 100,000 JPY	63 (13.3)
<hr/>	
Financial Aid	
None	342 (72.3)
Student loan only	74 (15.6)
Scholarship only	44 (9.3)
Both student loan and scholarship	13 (2.8)
<hr/>	

Abbreviations: JPY, Japanese Yen.

^a According to the Bank of Japan, the exchange rates of 1 US dollar to JPY was 107.39-41 on the morning of June 17, 2020.