

**eTable 5. Association between levels of the biological inflammatory markers NLR, LCRP, IL-6 and lactates and the endpoint of death in the subsamples of patients with COVID-19 who had at least two measures of each marker during the visit.**

	First measure		Last measure	
	Number of events / Number of patients	Crude Cox regression analysis <sup>α</sup>	Number of events / Number of patients	Crude Cox regression analysis <sup>α</sup>
	Events / N (%)	HR (95% CI; p-value)	Events / N (%)	HR (95% CI; p-value)
<b>Biological inflammatory markers</b>				
LCRP	426 / 1,879 (22.7%)	0.83 (0.76 - 0.90; <0.001*)	423 / 1,876 (22.5%)	0.47 (0.43 - 0.60; <0.001*)
NLR	492 / 2,158 (22.8%)	1.26 (1.11 - 1.43; <0.001*)	493 / 2,159 (22.8%)	2.71 (2.47 - 2.96; <0.001*)
IL-6	53 / 315 (16.8%)	1.29 (1.06 - 1.57; 0.010*)	53 / 3,15 (16.8%)	1.53 (1.33 - 1.76; <0.001*)
Lactates	319 / 1,159 (27.5%)	1.27 (0.79 – 2.03; 0.326)	823 / 1,163 (27.8%)	2.75 (2.43 – 3.13; <0.001*)

<sup>α</sup> Values were log-transformed to address non-normal distribution of the biological variables.