

**eTable 6. Course of the levels of the biological inflammatory markers NLR, LCRP, IL-6 and lactates during the visit according to hydroxyzine use.**

	<b>Full population</b>	<b>Exposed to hydroxyzine</b>	<b>Not exposed to hydroxyzine</b>	<b>Two-way ANOVA: Interaction term</b>
	N / Mean (SD)	N / Mean (SD)	N / Mean (SD)	F (p) <sup>α</sup>
<b>Biological inflammatory markers</b>				
LCRP				16.33 (<0.001*)
<i>First measurement</i>	1,879 / 695.1 (2,732.0)	63 / 308.4 (666.3)	1,816 / 708.5 (2,775.3)	
<i>Last measurement</i>	1,879 / 1,396.2 (3,831.7)	63 / 1,640.5 (2,673.9)	1,816 / 1,387.7 (3,865.9)	
NLR				14.80 (<0.001*)
<i>First measurement</i>	2,161 / 7.1 (6.8)	72 / 8.3 (6.8)	2,089 / 7.1 (6.8)	
<i>Last measurement</i>	2,161 / 6.8 (9.1)	72 / 4.9 (5.8)	2,089 / 6.8 (9.2)	
IL-6				5.87 (0.016*)
<i>First measurement</i>	315 / 182.4 (405.7)	10/ 216.2 (230.1)	305 / 181.2 (410.4)	
<i>Last measurement</i>	315 / 238.1 (800.2)	10 / 51.9 (68.6)	305 / 244.2 (812.4)	
Lactates				0.94 (0.333)
<i>First measurement</i>	1,142 / 1.5 (0.8)	43 / 1.4 (0.7)	1,099 / 1.5 (0.8)	
<i>Last measurement</i>	1,142 / 1.5 (1.3)	43 / 1.2 (0.5)	1,099 / 1.6 (1.3)	

<sup>α</sup> Values were log-transformed to address non-normal distribution of the biological variables.

\* p-value is significant (p<0.05).