

eTable 7. Course of the levels of the biological inflammatory markers NLR, LCRP, and IL-6 during the visit according to hydroxyzine dose.

	Full population	Low dosage	High dosage	Two-way ANOVA: Interaction term
	N / Mean (SD)	N / Mean (SD)	N / Mean (SD)	F (p) ^α
Biological inflammatory markers				
LCRP				4.57 (0.037*)
<i>First measurement</i>	55 / 329.2 (708.6)	25 / 273.5 (399.4)	30 / 375.6 (893.4)	
<i>Last measurement</i>	55 / 1,423.8 (2,016.8)	25 / 957.2 (1,728.3)	30 / 1,812.7 (2,181.2)	
NLR				5.62 (0.021*)
<i>First measurement</i>	60 / 8.6 (7.2)	26 / 6.3 (4.1)	34 / 10.4 (8.4)	
<i>Last measurement</i>	60 / 4.6 (5.0)	26 / 5.7 (7.0)	34 / 3.8 (2.6)	
IL-6				0.62 (0.454)
<i>First measurement</i>	10 / 216.2 (230.1)	5 / 95.5 (148.4)	337.0 (246.3)	
<i>Last measurement</i>	10 / 51.9 (68.6)	5 / 35.5 (52.3)	68.4 (84.7)	

^α Values were log-transformed to address non-normal distribution of the biological variables.

* p-value is significant (p<0.05).