

**Rapid Support for Older Adults during the initial stages of the COVID-19
Pandemic: Results from a Geriatric Psychiatry helpline**

Anna-Sophia Wahl, MD PhD^{1*†}, Gloria S. Benson, PhD^{1*}, Lucrezia Hausner, MD¹, Sandra Schmitt, MSc¹, Annika Knoll, MSc¹, Adriana Feretti Bondy, MSc¹, Dimitri Hefter, MD¹, Lutz Frölich, MD¹

Supplementary Material

Supplementary Table 1. Skewness and Kurtosis for mental health variables of depression and anxiety. SE Standard Error. Both variables were sufficiently normally distributed.

Supplementary Table 2. Table depicting the different features of help provided and subsequent appointments scheduled by us for psychiatric or psychotherapeutic counseling or support by our social worker. Total N represents the number of callers which participated in the survey. N represents the number of callers which affirmed the different features asked. The percentage was calculated as $(N/\text{total } N) \times 100$.

	N	Skewness		Kurtosis	
		Value	SE	Value	SE
Anxiety	35	-0.058	0.398	-1.271	0.778
Depression	44	0.293	0.357	-0.908	0.702

Supplementary Table 1 SD= Standard deviation. SE= Standard error

	Total N	N or M (SD)	(%)
Result of the hotline call			
Providing information of Covid-19	51	30	58.8
Practical help/information for daily living	51	1	1.9
Providing social contacts	51	1	1.9
Providing help for somatic diseases/issues	51	7	13.7
Recommendation for psychiatric counseling	51	15	29.4
Providing relieving psychological support	51	29	56.9
Providing contact to welfare work	51	1	1.9
Subsequent treatments			
No subsequent treatments	51	38	74.5
Subsequent treatments required	51	10	19.6
Subsequent psychiatric treatment provided	51	3	5.9
Subsequent psychotherapy provided	51	7	13.7
Subsequent welfare work provided	51	1	1.9

Supplementary Table 2