Using COVID-19 deaths as a surrogate to measure the progression of the pandemics.

Carlos Hernandez-Suarez^{a,*}, Efren Murillo-Zamora^b

^a*Facultad de Ciencias, Universidad de Colima, Bernal Diaz del Castillo 340, Colima, Colima, 28040, MEXICO*

^b*Departamento de Epidemiología, Unidad de Medicina Familiar No. 19, IMSS, Av. Javier Mina 301, 28000, Colima, Colima. MEXICO*

Abstract

The IFR (Infection Fatality Risk) is one of the most important parameters of an infectious disease. If properly estimated, the observed number of deaths divided by the IFR can be used to estimate the current number of infections and, if immunity is permanent, we can estimate the fraction of susceptible which can be used to plan reopening of activities and vaccine deployment, when these become available. Here we suggest how to use the observed deaths by COVID-19 in an arbitrary population as a surrogate for the progression of the epidemic with the purpose of decision making. We compare several estimates of IFR for SARS-CoV-2 with our estimate that uses the number of additional deaths in households in a database population of 159,150 laboratory-confirmed (RT-qPCR) COVID-19 by SARS-COV-2 in Mexico. The main result is that if the number of deaths in a region is close to 2 per thousand individuals, the fraction of remaining susceptible may be too small for the vaccine to make a difference in the total number of infected.

Keywords: COVID-19, SARS-CoV-2, IFR, Asymptomatic, Immunity, Total Infections, Lethality, Vaccines

*Corresponding author

Email addresses: carlosmh@mac.com (Carlos Hernandez-Suarez),
efren.murilloza@imss.gob.mx (Efren Murillo-Zamora)

Preprint submitted to medRxiv

October 29, 2020

1. Introduction

COVID-19 pandemic has shown to have a low lethality, nevertheless, the burden of the disease so far is huge. Economical activities are suspended or reduced and there is pressure for reopening of schools which requires more information, not only available to policymakers but also to the general public, to ameliorate civic unrest. The competition for developing a vaccine is keen, with about 321 vaccine candidates with 32 in clinical trials in progress (Le et al., 2020), raising safety concerns (Harrison and Wu, 2020; Peeples, 2020), and the additional economic cost for the purchase and deployment of vaccines has not been yet added to the burden of the disease.

At the moment, without vaccines or effective pharmaceutical treatments available, the decision on whether reopening activities or not depends mostly on the number of additional infections/deaths that will be caused because of some policy, say, opening schools, allowing public gatherings, opening touristic places, increasing the current density allowed in some places as cinemas, restaurants, buses, and planes, etc. For a highly infectious virus as SARS-CoV-2, the decision must be rooted in the amount of remaining susceptible in the region affected for some change in policy.

Using the number of individuals that access private or public hospitals as a surrogate of the current number of infections, or even the number of confirmed cases, is not accurate, because those quantities strongly depend on the availability of medical services that are not always accessible to the bulk of a population in many countries, or simply because of lack of resources for test deployment or because of policies that disregard testing. The number of deaths is less dubious. if we manage to calculate the average number of infections that will result in the death of the average individual, then, we can use this number to estimate the number of infections that were required to observe the current number of deaths, and from here, the share of susceptible available in a well-defined population. Even although deaths were not reported as COVID-19 related, records may exist that shed light on the likelihood that death was or not caused by

SARS-CoV-2 infection, which is part of the policy adopted by some countries like Belgium, which is, without doubt, one of the reasons why this country keeps one of the highest number of deaths per capita in Europe.

Before proceeding, we need to deploy two facts. The first one attempts to
 35 establish that, with few exceptions, it is almost impossible to stop the pandemic by mitigation and control measures, and the most we can do is reducing the infection rate (at a huge economic and social cost) which is known as *flattening the curve*. The second establishes that we are in the condition now to have reliable lower bound estimates for IFR, the Infection Fatality Risk of SARS-CoV-2. It
 40 is the confluence of these two facts that allow us to establish a surrogate for the progression of the epidemic in a region and thus, for the calculation of the share of susceptible available, which is the basis to decide on reopening activities and vaccine deployment.

Fact 1: SARS-CoV-2 is highly contagious and will infect most of the 45 population

Several studies have shown that the basic reproductive number R_0 is high (Sanche et al., 2020; D'Arienzo and Coniglio, 2020; Najafimehr et al., 2020; Liu et al., 2020; Alimohamadi et al., 2020), and it has been calculated as high as 5.7. It is important to remember that R_0 is the potential of disease transmission
 50 in the absence of any control or mitigation measures, and it is the potential that comes into play as soon as control or mitigation measures are suspended. We calculated the R_0 for 40 countries using only the first two weeks of data, analyzing the progression in the number of deaths. These 40 countries were the first countries affected by pandemics except for China. The R_0 calculated
 55 is shown in Figure 1. The estimation of the true potential of the disease is more evident for those countries that were affected first (e.g. Italy and Spain), where our estimate of R_0 is about the same as the reported by Sanche et al. (2020). The learning curve is evident in the reduction of R_0 when the pandemic advances. Estimates of R_0 in countries where the pandemic arrived at later

60 stages will not reflect the true potential of the virus alone, but the preparedness of a population.

Figure 1: Age of the pandemic vs. R_0 . For each country, the baseline was taken as the date when the accumulated number of deaths was at least 5. Day zero is 02/24/2020. See Table 1 for expanded data.

Table (1) in Appendix contains the main statistics derived from the analysis of λ and R_0 for the countries analyzed. From the estimated λ 's, we can see that the first countries to be affected had an R_0 larger than 5 whereas this was decreasing to achieve values just above 2.5, this implies that at the beginning λ was as high as one effective contact every three days to one effective contact every 5 days, on average.

There are some facts we need to consider when a virus spreads with that strength: first, forecasting the size of the epidemics (total number of infected) is simpler, since social contact structures are bypassed and become irrelevant, and thus the contact network resembles more a random mixing pattern. For this random mixing, the estimated fraction of infected f , can be approximated

(Kermack and McKendrick, 1927) with:

$$f = 1 - e^{-R_0 f} \quad (1)$$

(for a probabilistic derivation, see Hernandez-Suarez and Mendoza-Cano, 2009).

75 For instance, if $R_0 = 5.7$ the fraction of infected is $f = 0.99$. However, if we manage to reduce R_0 by half, $f = 0.93$, which is not a big difference in the size of the epidemic. Nevertheless, reducing R_0 by half is an incredible task that involves mainly lockdowns and face-masks, and the former has a huge economic cost that can not hold for long except for rich countries or countries whose
80 political or cultural organization allows its implementation. Nevertheless, even if a country manages to implement actions to reduce R_0 to a value smaller to one, and can support citizens economically to maintain the lockdown for long periods, a handful of infected individuals that enter the country from abroad can restart the infection process, if lockdowns are lifted. From here, the observed waves
85 (see Figure 2 for examples in Japan, Cuba, S. Korea and New Zealand). This is particularly true for a disease where the number of asymptomatic individuals is by far, larger than that of symptomatic and thus, difficult to control. As a consequence, equation (1) can be seen as a quota that must be fulfilled, as long as the fraction of remaining susceptible is larger than f .

90 **Fact 2: the overall IFR of SARS-CoV-2 is somewhere between 0.2 and 0.3 %**

All calculations in this section are done with the data on deaths reported to 10/29/2020 (Dong et al., 2020). Also, in this section we use IFR to indicate the true overall Infection Fatality Risk of SARS-CoV-2, that is, it is the fraction of
95 individuals that die if infected with SARS-CoV-2, whereas IFR^* indicates some estimate of IFR for a region of interest. Many attempts have been made to estimate this important parameter that would be a remarkable surrogate to measure the progress of the epidemics in an arbitrarily defined region or population, but the main inconvenient is that while the number of deaths from SARS-CoV-2

Figure 2: **The appearance of a second wave.** Cumulated deaths per day. The use of deaths instead of confirmed cases delays the detection of a wave but is more efficient in the sense that does not depend on testing policies, availability or panic. The current IFR* (per million inhabitants) is Japan: 14; Cuba: 11; S. Korea: 9; New Zealand: 5. Data updated to 10/29/2020.

100 can be approximated, the denominator, the total number of infected from the disease is elusive, especially considering that a large fraction of individuals that are infected are asymptomatic. Besides, the efficacy of immunity tests to detect who has been infected and recovered has been challenged with the findings that the ability to detect antibodies is reduced in a few days, especially in those with
105 mild or no symptoms (Long et al., 2020; Ibarrondo et al.). Recently, Eyre et al. (2020) reported that a high fraction of individuals with none or mild symptoms may be undiagnosed mainly due to the calibration strategies of some standardized tests and concluded that samples from individuals with mild/asymptomatic infection should be included in SARS-CoV-2 immunoassay evaluations.

110 Some studies to estimate the IFR have been recently released for Iceland (Gudbjartsson et al.), India (Mukhopadhyay and Chakraborty, 2020), Germany (Streeck et al., 2020) and Denmark (Erikstrup et al., 2020) among others. In a review of 23 studies where the IFR was estimated Ioannidis (2020) a median of

0.25% is reported, which is about 2.5 deaths for each 1000 individuals. The study
 115 in Iceland, (Gudbjartsson et al.), the most comprehensive to date, proposes an
 IFR of 0.3% (95% CI, 0.2 to 0.6). Special care must be taken with this later
 estimate, since Iceland has a population of 321,641, with only 29 deaths so far.
 In other studies that used transmission models fed with serological data, the IFR
 estimates where as high as 2.5-5 (Thorne, 2020) and 3-8 (Roques et al., 2020)
 120 per thousand infections. Wilson (2020) estimated the IFR using serological
 data from New York City and arrived to an IFR estimate of 8.6. We believe
 this particular estimate is too high, and will support our claim analyzing the
 behavior of the epidemic in some regions. The reason of such a high estimate
 may be the sampling scheme, which was not random at all, and would explain
 125 the high discrepancy between the observed differences in the number of infected
 between males and females (15.9 and 12 percent respectively) which the authors
 acknowledge they can't explain.

Estimation of the IFR using an alternative method

Hernandez-Suarez et al. (2020) proposed an alternative method to estimate
 130 the IFR based on the assumption that all those living with an infected individual
 will be infected. We use the same data set that has grown from 3,193 households
 to 131,240 households. The essence of the method is as follows: let n_j be the
 number of individuals living in a house where an infected was confirmed infected
 with SARS-CoV-2, whose symptoms started on day d_j . Let x_j the number of
 135 deaths among the remaining $n_j - 1$ individuals whose symptoms started after
 day d_j , then, an estimate of the IFR is:

$$\text{IFR} = \frac{\sum_j x_j}{\sum_j (n_j - 1)} \quad (2)$$

In this example we build an approximation to (2) using a database from
 Mexico's IMSS (Instituto Mexicano de Seguro Social), the Mexican Institute
 for Social Insurance. The database has 212,935 (laboratory-confirmed (quantitative
 140 reverse transcription polymerase chain reaction, RT-qPCR) COVID- 19
 by SARS-COV-2 cases from April 1 to September 17, 2020. In an attempt to

Table 1: Observed frequencies of deaths per household. The index case is excluded.

Deaths	Households
0	130,248
1	921
2	41
3	22
4	4
5	2
6	1
7	1

consider only households with final outcomes we excluded cases with symptoms onset in the last 6 weeks before the final case recorded, that is, we considered only cases with symptom onset from April 1 to August 6, 2020. The final data set has 135,855 cases. The mean age of this final set was 47.9 years with a standard deviation of 16.5 years and median 47 years. From these, there were 54 % males and 46 % females. Tables 2 and 3 contain summary statistics from the final data set.

Every individual has an associated NSS (Social Security Number) that is shared among all members of a family. As an approximation, we assumed all individuals sharing the same NSS live in the same household, and then grouped the cases in households. If there were more than one case in a household, we only considered the household if all cases were already solved a deaths or recoveries. In every household with more than one case we consider the index case as the individual with the earliest symptom onset and counted the number of deaths among the remaining members of the house. From the final set of 131,240 households, there were 130,248 with no deaths among the remaining members of the household and 992 houses with at least one additional death (see Table 1).

Table 2: Observed frequencies of deaths by gender

Age	Males	Females
0-9	0	0
10-19	546	578
20-29	7,956	8,324
30-39	13,413	12,818
40-49	13,770	12,399
50-59	12,728	9,688
60+	18,716	13,329
Total	67,129	57,136

Table 3: Summary statistics of data set

	N	%
Total	135,855	
Gender		
Male	73,256	53.9
Female	62,599	46.1
Condition		
Ambulatory	75,725	55.7
Hospitalized	60,130	44.3
Outcome		
Recovered	103,432	76.1
Death	32,423	23.9

160 From Table 1 we have $\sum_j x_j = 1,108$. We do not have $\sum_j n_j$ and we build an estimate with the number of households times the average household size, that is, $131,240 \times \mu$ where $\mu = 3.7$ (Encuesta Intercensal, 2015). Using (2) we have our estimate of IFR is

$$\text{IFR}^* = \frac{1,108}{131,240(\mu - 1)} = 0.0031 \quad (3)$$

that is, 3.1 cases per thousand infected, which is close to 2.5, the median of 23 studies reported in Ioannidis (2020) and very similar to the 3.0 recently derived from the study in Iceland (Gudbjartsson et al.) Special care must be taken with our estimate (3) since we are building a rough approximation to the total number of additional individuals living in each household.

170 If the disease will affect most of the population, as suggested previously, then, the most effective way to estimate the overall IFR is by measuring the fraction of deaths in a population in which the epidemic has evolved for a long time and it is over or near the end. It is evident that we need to focus in regions with: *a)* high observed IFR^* and *b)* lack of new waves. Such is the case of regions as New Jersey and New York states (see figure 3). The current IFR^* calculated for these states is 1.854 and 1.729 respectively.

The lack of new waves in these two states indicates that the virus has infected and killed a fraction of individuals close to the IFR, and thus the epidemic rampaged those states and it stopped due to what is commonly called *herd immunity*, that actually conveys no immunity at all, but a lack of a significant mass of susceptible for the R_0 to be effective. Thus, the progression of the epidemic in a region can be estimated with IFR^*/IFR .

2. Consequences from the previous facts for vaccines and the return to normality

185 As a consequence of the two previous facts, we can conclude that it is possible to monitor the development of the epidemic in a region by following the

Figure 3: **Absence of second waves in the states of New Jersey and New York.** Cumulated deaths per day. The current IFR* (per million inhabitants) is New Jersey: 1.854; New York: 1.729. Data updated to 10/29/2020.

number of deaths. One can estimate the fraction of total infected at time t , asymptomatic or not, with:

$$\theta_t = \frac{x_t}{N \times IFR}$$

where x_t is the number of observed deaths at time t and N is the population size. The estimated fraction of susceptible as a function of the IFR in New Jersey and New York is shown in figure 4. We can see that if $IFR=2/1000$, the fraction of remaining susceptible in New Jersey and New York is 7.3% and 13.5% respectively, nevertheless, if it is as high as 3, the fraction of susceptible

raises to 38% and 42% for those states.

Figure 4: The fraction of remaining susceptible in the states of New Jersey and New York as a function of the lethality of SARS-CoV-2.

3. Discussion

Because of the aforementioned facts, it is very important to monitor the COVID-19 related deaths and every effort has to be made to improve current estimates of the IFR. Hernandez-Suarez et al. (2020) suggested that in light of the high contagiousness of SARS-CoV-2, the secondary cases resulting in fatalities in a household with at least one confirmed infected, may be useful to estimate the IFR, which would provide a large amount of data and minimal testing requirements. Antibody testing in New York shows to date (Data updated to 10/29/2020) that out of 2,608,997 tests, 642,977 were positive for antibodies (NYC-Health) which implies 24% of the population may be immune, contradicting our findings that for an IFR as high as 2.5 per thousand the percentage of infected should be close to 70% already (see Figure 4). Nevertheless, the fraction of people with antibodies is not a good surrogate of the fraction of infected with SARS-CoV-2 because it underestimates the number of infected

for two reasons: first, antibody testing is voluntary and it is natural to expect that individuals with no symptoms are less compelled to be tested than those symptomatic, so, a large fraction of infected is not tested, and second, we already mentioned some failures in testing reported, apparently due to calibration procedures that tend to fail in individuals with none or mild symptoms (Eyre et al., 2020).

The idea that the share of susceptible in New Jersey is relatively small, is supported on the fact that the number of active cases reported to date (10/29/2020) is close to 42,000 and unless they are fully quarantined the infectious pressure from these individuals and those unreported must be huge, nevertheless, no new significant peak is observed. The same happens in New York, although the share of susceptible is higher.

Everything leads to the following question: what will be the purpose of a vaccine in a region where the IFR* is already large enough to suggest there is only a small fraction of susceptible with limited chance of infection? This question is especially valid from the comprehensive study in Iceland that strongly suggests the existence of immunity due to infection (Gudbjartsson et al.). Unless it is proven that immunity wanes beyond a protective level after some time, everything seems to indicate that those regions where the share of estimated susceptible is already low, should have less priority in the distribution of vaccines. If facts 1 and 2 are not taken into account, a vaccination campaign in a region where the proportion of deaths among the population is close to the IFR, will just give the impression of being effective but will not play a role, regardless of the efficacy of the vaccine.

Acknowledgements

This work is part of the DFID-UNHCR-World Bank program “Building the Evidence on Protracted Forced Displacement: A Multi-Stakeholder Partnership”. The program is funded by UK aid from the United Kingdom’s Department for International Development (DFID), it is managed by the World Bank

Group (WBG) and was established in partnership with the United Nations High Commissioner for Refugees (UNHCR). The scope of the program is to expand the global knowledge on forced displacement by funding quality research and disseminating results for the use of practitioners and policy makers. This work does not necessarily reflect the views of DFID, the WBG or UNHCR.

4. Bibliography

- Alimohamadi, Y., Taghdir, M., Sepandi, M., 2020. The estimate of the basic reproduction number for novel coronavirus disease (COVID-19): a systematic review and meta-analysis. *Journal of Preventive Medicine and Public Health* .
- D'Arienzo, M., Coniglio, A., 2020. Assessment of the sars-cov-2 basic reproduction number, r_0 , based on the early phase of covid-19 outbreak in italy. *Biosafety and Health* 2, 57 – 59. doi:10.1016/j.bsheal.2020.03.004.
- Dong, E., Du, H., Gardner, L., 2020. An interactive web-based dashboard to track covid-19 in real time. *The Lancet infectious diseases* 20, 533–534.
- Encuesta Intercensal, 2015. INEGI. Recovered from: [http://www. beta. inegi. org. mx/proyectos/enchogares/especiales/intercensal](http://www.beta.inegi.org.mx/proyectos/enchogares/especiales/intercensal) .
- Erikstrup, C., Hother, C.E., Pedersen, O.B.V., Mølbak, K., Skov, R.L., Holm, D.K., Sækmose, S., Nilsson, A.C., Brooks, P.T., Boldsen, J.K., Mikkelsen, C., Gybel-Brask, M., Sørensen, E., Dinh, K.M., Mikkelsen, S., Møller, B.K., Haunstrup, T., Harritshøj, L., Jensen, B.A., Hjalgrim, H., Lillevang, S.T., Ullum, H., 2020. Estimation of SARS-CoV-2 infection fatality rate by real-time antibody screening of blood donors. *medRxiv* doi:10.1101/2020.04.24.20075291.
- Eyre, D.W., Lumley, S.F., O'Donnell, D., Stoesser, N.E., Matthews, P.C., Howarth, A., Hatch, S.B., Marsden, B.D., Cox, S., James, T., et al., 2020. Stringent thresholds for sars-cov-2 igg assays result in under-detection of cases reporting loss of taste/smell. *medRxiv* .

- 265 Guðbjartsson, D.F., Norddahl, G.L., Melsted, P., Gunnarsdottir, K., Holm, H., Eythorsson, E., Arnthorsson, A.O., Helgason, D., Bjarnadottir, K., Ingvarsson, R.F., Thorsteinsdottir, B., Kristjansdottir, S., Birgisdottir, K., Kristinsdottir, A.M., Sigurdsson, M.I., Arnadottir, G.A., Ivarsdottir, E.V., Andresdottir, M., Jonsson, F., Agustsdottir, A.B., Berglund, J., Eiriksdottir, B., Fridriksdottir, R., Gardarsdottir, E.E., Gottfredsson, M., Gretarsdottir, O.S., Gudmundsdottir, S., Gudmundsson, K.R., Gunnarsdottir, T.R., Gylfason, A., Helgason, A., Jensson, B.O., Jonasdottir, A., Jonsson, H., Kristjansson, T., Kristinsson, K.G., Magnusdottir, D.N., Magnusson, O.T., Olafsdottir, L.B., Rognvaldsson, S., le Roux, L., Sigmundsdottir, G., Sigurdsson, A., Sveinbjornsson, G., Sveinsdottir, K.E., Sveinsdottir, M., Thorarensen, E.A., Thorbjornsson, B., Thordardottir, M., Saemundsdottir, J., Kristjansson, S.H., Josefsdottir, K.S., Masson, G., Georgsson, G., Kristjansson, M., Moller, A., Palsson, R., Gudnason, T., Thorsteinsdottir, U., Jonsdottir, I., Sulem, P., Stefansson, K., . Humoral immune response to sars-cov-2 in ice-land. New England Journal of Medicine doi:10.1056/NEJMoa2026116.
- 270
- Harrison, E.A., Wu, J.W., 2020. Vaccine confidence in the time of covid-19. European Journal of Epidemiology 35, 325–330. URL: <https://doi.org/10.1007/s10654-020-00634-3>, doi:10.1007/s10654-020-00634-3.
- Hernandez-Suarez, C., Mendoza-Cano, O., 2009. Applications of occupancy urn models to epidemiology. Mathematical Biosciences and Engineering 6, 509–520.
- 285
- Hernandez-Suarez, C., Verme, P., Murillo-Zamora, E., 2020. Estimation of the infection fatality rate and the total number of sars-cov-2 infections. medRxiv doi:10.1101/2020.04.23.20077446.
- 290
- Ibarrondo, F.J., Fulcher, J.A., Goodman-Meza, D., Elliott, J., Hofmann, C., Hausner, M.A., Ferbas, K.G., Tobin, N.H., Aldrovandi, G.M., Yang, O.O., . Rapid decay of anti-SARS-CoV-2 antibodies in persons with mild COVID-19. New England Journal of Medicine doi:10.1056/NEJMc2025179.

- Ioannidis, J., 2020. The infection fatality rate of COVID-19 inferred from sero-
295 prevalence data. medRxiv doi:10.1101/2020.05.13.20101253.
- Kermack, W.O., McKendrick, A.G., 1927. A contribution to the mathematical
theory of epidemics. Proceedings of the Royal Society of London. Series A
115, 700–721.
- Le, T.T., Cramer, J., Chen, R., Mayhew, S., 2020. Evolution of the covid-19
300 vaccine development landscape. Nat Rev Drug Discov .
- Liu, Y., Gayle, A.A., Wilder-Smith, A., Rocklöv, J., 2020. The reproductive
number of COVID-19 is higher compared to SARS coronavirus. Journal of
Travel Medicine 27. doi:10.1093/jtm/taaa021. taaa021.
- Long, Q.X., Tang, X.J., Shi, Q.L., Li, Q., Deng, H.J., Yuan, J., Hu, J.L., Xu,
305 W., Zhang, Y., Lv, F.J., Su, K., Zhang, F., Gong, J., Wu, B., Liu, X.M.,
Li, J.J., Qiu, J.F., Chen, J., Huang, A.L., 2020. Clinical and immunolog-
ical assessment of asymptomatic SARS-CoV-2 infections. Nature Medicine
doi:10.1038/s41591-020-0965-6.
- Mukhopadhyay, S., Chakraborty, D., 2020. Estimation of undetected covid-19
310 infections in india. medRxiv doi:10.1101/2020.04.20.20072892.
- Najafimehr, H., Mohamed Ali, K., Safari, S., Youseffard, M., Hosseini, M.,
2020. Estimation of basic reproduction number for covid-19 and the reasons
for its differences. International Journal of Clinical Practice 74. doi:10.1111/
ijcp.13518.
- 315 NYC-Health, . COVID-19: Data. [https://www1.nyc.gov/site/doh/covid/
covid-19-data.page](https://www1.nyc.gov/site/doh/covid/covid-19-data.page). Accessed: 2020-09-03.
- Peeples, L., 2020. News feature: Avoiding pitfalls in the pur-
suit of a covid-19 vaccine. Proceedings of the National
Academy of Sciences 117, 8218–8221. URL: [https://www.
320 pnas.org/content/117/15/8218](https://www.pnas.org/content/117/15/8218), doi:10.1073/pnas.2005456117,
arXiv:<https://www.pnas.org/content/117/15/8218.full.pdf>.

- Roques, L., Klein, E.K., Papaix, J., Sar, A., Soubeyrand, S., 2020. Using early data to estimate the actual infection fatality ratio from covid-19 in france. *Biology* 9, 97.
- 325 Sanche, S., Lin, Y.T., Xu, C., Romero-Severson, E., Hengartner, N., Ke, R., 2020. High contagiousness and rapid spread of severe acute respiratory syndrome coronavirus 2. *Emerging Infectious Diseases* 26, 1470.
- Streeck, H., Schulte, B., Kuemmerer, B., Richter, E., Hoeller, T., Fuhrmann, C., Bartok, E., Dolscheid, R., Berger, M., Wessendorf, L., Eschbach-Bludau, 330 M., Kellings, A., Schwaiger, A., Coenen, M., Hoffmann, P., Noethen, M., Eishuebinger, A.M., Exner, M., Schmithausen, R., Schmid, M., Kuemmerer, B., 2020. Infection fatality rate of SARS-CoV-2 infection in a german community with a super-spreading event. medRxiv doi:10.1101/2020.05.04.20090076.
- Thorne, R.S., 2020. Inferring the infection fatality rate (ifr) of covid-19 from 335 the behaviour of lombardy, madrid and london relative to the remainder of italy, spain and england. arXiv preprint arXiv:2005.00495 .
- Wilson, L., 2020. SARS-CoV-2, COVID-19, infection fatality rate (IFR) implied by the serology, antibody, testing in New York City. COVID-19, Infection Fatality Rate (IFR) Implied by the Serology, Antibody, Testing in New York 340 City (May 1, 2020) .

5. Appendix

Table A1: Contact rate λ and R_0 for the first 40 countries with diagnosed SARS-CoV-2. R_0 was calculated at the beginning of the epidemic in each country.

Country	λ	R_0	R^2	Country	λ	R_0	R^2
Italy	0.366	5.79	0.997	Philippines	0.237	3.748	0.988
Spain	0.352	5.569	0.995	Colombia	0.235	3.708	0.996
Netherlands	0.329	5.196	0.998	Denmark	0.234	3.696	0.994
Germany	0.322	5.083	0.999	Poland	0.234	3.694	0.992
United Kingdom	0.319	5.036	0.995	Norway	0.233	3.681	0.997
Belgium	0.306	4.844	0.996	Romania	0.229	3.627	0.987
France	0.294	4.645	0.979	Chile	0.226	3.569	0.996
US	0.29	4.579	0.993	Greece	0.224	3.545	0.994
Peru	0.289	4.571	0.996	Russia	0.223	3.528	0.996
Mexico	0.284	4.494	0.998	Pakistan	0.218	3.45	0.993
Turkey	0.283	4.48	0.998	Israel	0.218	3.444	0.995
Brazil	0.282	4.461	0.997	Morocco	0.216	3.418	0.993
Switzerland	0.273	4.323	0.995	Ukraine	0.215	3.393	0.995
Sweden	0.273	4.31	0.982	Panama	0.211	3.333	0.991
Ecuador	0.26	4.116	0.997	Hungary	0.206	3.252	0.994
Austria	0.254	4.009	0.992	Slovenia	0.204	3.222	0.998
India	0.248	3.925	0.99	Finland	0.201	3.18	0.994
Portugal	0.248	3.922	0.992	UA Emirates	0.185	2.926	0.993
Canada	0.247	3.903	0.996	Nigeria	0.173	2.732	0.995
Ireland	0.246	3.888	0.992	Sudan	0.173	2.73	0.995