Transmission routes of SARS-CoV-2 among healthcare workers of a French university hospital in Paris, France: a case-control study

Adrien Contejean^{1,2*}, Jérémie Leporrier^{3*}, Etienne Canoui², Jacques Fourgeaud^{4,5}, Alice-Andrée Mariaggi⁶, Fanny Alby-Laurent^{1,3}, Emmanuel Lafont^{1,3}, Lauren Beaudeau², Claire Rouzard³, Fabienne Lecieux⁷, Agnès Greffet⁸, Anne-Sophie L'Honneur⁶, Jean-Marc Tréluyer^{9,10,11}, Fanny Lanternier^{1,3}, Anne Casetta¹², Pierre Frange^{5,13}, Marianne Leruez-Ville^{4,5}, Flore Rozenberg^{6,14}, Olivier Lortholary^{1,3,15}, Solen Kernéis^{2,16,17}

*Contributed equally

¹ Université de Paris, Faculté de Médecine, F-75006 Paris, France

² Equipe Mobile d'Infectiologie, AP-HP, Hôpital Cochin, F-75014 Paris, France

³ Service de maladies infectieuses et tropicales, AP-HP, IHU Imagine, Hôpital Necker-Enfants malades, F-75015 Paris, France

⁴ Laboratoire de virologie, AP-HP, Hôpital Necker-Enfants malades, F-75015 Paris, France

⁵ EHU 7328 PACT, Institut Imagine, Université de Paris, Faculté de Médecine, F-75006 Paris, France

⁶ Département de virologie, AP-HP, Hôpital Cochin, F-75014 Paris, France

⁷ Service de santé au travail, AP-HP, Hôpital Cochin, F-75014 Paris, France

⁸ Service de santé au travail, AP-HP, Hôpital Necker-Enfants malades, F-75015 Paris, France

⁹ Département de soins intensifs pédiatriques, AP-HP, Hôpital Necker-Enfants malades, F-75015 Paris, France

¹⁰ Pharmacology and Drug Evaluation in Children and Pregnant Women EA7323, Université de Paris, Faculté de Médecine, F-75006 Paris, France

¹¹ Département de pharmacologie clinique AP-HP, Hôpital Cochin, F-75014 Paris, France

¹² Equipe opérationnelle d'hygiène hospitalière, AP-HP, Hôpital Cochin, F-75014 Paris, France

¹³ Laboratoire de microbiologie clinique, AP-HP, Hôpital Necker-Enfants malades, F-75015 Paris, France

¹⁴ Université de Paris, Institut Cochin, INSERM U1016, CNRS UMR8104, F-75006 Paris, France

¹⁵ Institut Pasteur, Molecular Mycology Unit, National Reference Centre for Invasive Mycoses and Antifungals, CNRS UMR 2000, Paris, France

¹⁶ Université de Paris, INSERM, IAME, F-75006 Paris, France

¹⁷ Institut Pasteur, Epidemiology and Modelling of Antibiotic Evasion (EMAE), F-75015 Paris, France

Corresponding author:

Solen Kernéis

Equipe Mobile d'Infectiologie, Hôpital Cochin, 27, rue du Faubourg Saint Jacques, 75014, Paris, France

Phone: +33158414111

Mail: solen.kerneis@aphp.fr

Key words: COVID-19, SARS-CoV-2, Healthcare Workers, Determinants, Case-Control study

Abstract

In this case-control study on 564 healthcare workers of a university hospital in Paris (France), contacts without protection with COVID-19 patients or with colleagues were associated with infection with SARS-CoV-2, while working in a COVID-dedicated unit, using public transportation and having children kept in childcare facilities were not.

Introduction

Effective protection of healthcare workers (HCW) against severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) requires precise assessment of the respective role of in-hospital and out-of-hospital exposures on transmission routes in this at high-risk population.

We previously published an observational multicenter cohort study on HCW infected by SARS-CoV-2 during the first French coronavirus disease-19 (COVID-19) breakthrough in 2020 spring[1]. Only 20% of the HCW screened positive for SARS-CoV-2 reported close contact with a suspected or confirmed COVID19 patient without Personal Protective Equipment (PPE), and 78% were not regularly posted in a COVID-19-dedicated unit. On the other hand, 54% declared frequent close contacts during occupational activities with colleagues without protection. We were however unable to compare positive HCW to a robust control group gathering personnel not infected with SARS-CoV-2. Diagnosis sensitivity of reverse transcriptase polymerase chain reaction (rtPCR) on nasopharyngeal swab for COVID-19 is imperfect[2] and serological assessment was not available at this time. IgG serological test has since been proven to be reliably associated with a COVID-19 past-infection[3].

We present here a case-control study which aimed to compare COVID-19 positive and negative HCW regarding their occupational activity, symptoms and in-hospital and out-of-hospital exposures to SARS-CoV-2, to determine the respective roles of different SARS-CoV-2 transmission routes.

Material and methods

Patients and design

This case-control study was led among HCW of a 2,100-bed tertiary-care university hospital (AP-HP, Centre, Université de Paris) located in central Paris, France, employing 13,278 personnel. From 24th February, to 10th April 2020, symptomatic staff were referred to dedicated on-site screening centers where trained medical staff collected a nasopharyngeal swab for SARS-CoV-2 rtPCR. HCW who tested positive on rtPCR were included as cases (HCW+). For each confirmed case, we included a control symptomatic HCW tested for SARS-CoV-2 on the same day, who had a negative rtPCR and a negative serological assessment performed at least 1 month after symptoms onset (HCW-). Both cases and controls were questioned in detail on their professional activity, symptoms, occupational and non-

occupational exposures to SARS-CoV-2 immediately after screening[1]. During this period, social distancing measures were implemented in France. On March 12th all school and childcare facilities closed, except for children of hospital staff, and a nationwide lockdown started on March 17th. Lift of containment measures occurred on May 11th.

Virology methods

SARS-CoV-2 rtPCR technique has been described elsewhere[1]. SARS-CoV-2 serology was determined by the Abbott® SARS-CoV-2 IgG assay, a chemiluminescent microparticle immunoassay intended for the qualitative detection of IgG antibodies to SARS-CoV-2.

Statistical analysis

Continuous variables are presented as median (interquartile range) and categorical variables as number (percentage). Fisher exact tests were used for comparisons of qualitative variables and Mann-Whitney tests for quantitative variables. All tests were 2-sided with a .05 value for significance. Factors associated with SARS-CoV-2 infection were assessed using multivariate logistic regression models. To account for the impact of the lockdown on community exposures, we considered two periods before and after March 22nd, 2020 (March 17th [date of national lockdown] + mean incubation period of 5 days). For each period, we first entered all exposures with a p-value < 0.40 in a multivariate model then used a backward stepwise selection procedure (removal criteria: $p \geq 0.05$) to build the final model. Statistical analyses were performed using R-software (3.3.2, R Foundation for Statistical Computing, Vienna, Austria).

Results

Between February 24th and April 10th, 2020, 1344 symptomatic HCW were screened for SARS-CoV-2 by rtPCR on a nasopharyngeal swab. Among them, 373 had positive rtPCR results (28%), 336 (90%) completed the questionnaire, and were included as cases (HCW+). Among 338 matched HCW with negative rtPCR, 247 (73%) had a serological assessment, and 228 (92%) tested negative. This group of 228 HCW with both negative rtPCR and serology constituted the control group (HCW-).

Exposures reported by cases and controls are displayed in the Table. Cases and controls were comparable in terms of age, sex and professional category. Cases presented more frequently with

anosmia or ageusia and with asthenia, fever, muscle pain, dyspnea, headaches and diarrhea than controls. Frequency of cough or rhinorrhea did not differ between groups.

In the univariate analysis (table), occupational activities with direct patient facing or assignment to a COVID-19-dedicated unit were similar in cases and controls. However, cases reported more close contacts with suspected or confirmed COVID-19 patients without PPE. Controls declared better compliance to mask wearing during occupational activities or in the presence of colleagues. Out-of-hospital exposures did not differ between cases and controls except for mask wearing outside home. In particular, using public transportation, having a child at home or a child kept out of the household were not associated with a significant increased risk of COVID-19.

Multivariate analysis showed that in the pre-lockdown period, close contacts with colleagues without protection (OR 2.58 [1.49-4.60]) was independently associated with an increased risk of COVID-19 in HCW, whereas systematic mask wearing outside of home was found independently protective against COVID-19 in this population (OR 0.43 [0.21-0.85]). Conversely, in the post-lockdown period, only close contacts with suspected or confirmed COVID-19 patients without PPE (OR 3.87 [1.73- 9.89]) was independently associated with an increased risk of COVID-19 in HCW.

Discussion

This case-control study among HCW highly exposed to SARS-CoV-2 suggests that PPE and especially medical mask is efficient in preventing COVID-19 transmission in HCW, including its systematic use outside hospital and home. To our knowledge, it is the first study to report both occupational and non-occupational exposures associated with SARS-CoV-2 infection in HCW since previous reports mainly focused on professional exposures and did not use IgG serology to contribute ruling-out COVID-19 in controls[4,5].

Unsurprisingly, symptoms were significantly different between cases and controls. Anosmia and ageusia appeared to be strongly associated to SARS-CoV-2 infection as reported elsewhere[6,7], whereas cough or rhinorrhea were not. Clinical scores including number of symptoms statistically associated with COVID-19 have been developed and might be a useful tool to quickly sort symptomatic HCW before results of SARS-CoV-2 testing[7].

Occupational risk factors for COVID-19 in HCW are dominated by exposure to suspected or confirmed COVID-19 patients without PPE or repeated contacts with numerous colleagues without medical mask. Indeed, SARS-CoV-2 infectiousness starts up to two days before symptoms onset[8], thus strict compliance to universal masking and social distancing measures are critical to prevent SARS-CoV-2 transmissions from asymptomatic patients or colleagues. Moreover, adherence to PPE during patient care is crucial to prevent COVID-19 infection in HCW, as we previously suggested[1]. Conversely, in our cohort, direct patient care was not associated with an increased risk of COVID-19 (including in COVID-19-dedicated wards), as long as HCW wear PPE. Of note, PPE supplies were immediately and fully available in our center, which was not the case in all French healthcare settings. Compliance to protective measures may also have been better among highly trained HCW of COVID-19-dedicated units. Recently, a large cohort study on 99,795 HCW suggested that frontline HCW were at increased risk of COVID-19 compared to community individuals, especially in case of exposition to patients with inadequate PPE[4].

Analysis of non-occupational exposures suggest that wearing a mask outside home may provide protection against COVID-19. The specific effect of mask wearing in the community is controversial. Indeed, HCW who reported wearing a mask outside home in our cohort (17%) were also probably more cautious regarding social activities and other suspected sources of SARS-CoV-2, which were not assessed in the questionnaire. Community use of facemasks have failed to demonstrate efficiency in prevention of influenzae transmission in a meta-analysis[9]. In Hong-Kong, a study suggested that the number of COVID-19 clusters were reduced when universal masking was recommended[10], and a study conducted in the USA concluded that mask wearing mandatory permitted to reduce daily COVID-19 growth rates[11]. However, to our knowledge, no study with high level of evidence has been published yet on that question.

As suggested in our first report[1] and here confirmed by the case-control study, HCW who reported to have children kept outside the family home did not have a higher risk of COVID-19 infection. However, childcare facilities that remained open for HCW's children during the national lockdown gathered a limited number of children simultaneously (<10). This question of SARS-CoV-2 transmission from children is highly debated since the beginning of the pandemic, but accumulating

data suggest that children are not significant drivers for COVID-19 pandemic[12]. Therefore, our results suggest that keeping schools and childcare facilities open for HCW is acceptable especially in case of second containment period, when HCW should be fully available at hospital.

We acknowledge several limitations, in particular recall bias, but cases and controls were interrogated prospectively and shortly after PCR assay. Additionally, our questionnaire included declarative data maybe not fully compressive regarding SARS-CoV-2 exposures.

In conclusion, profession category, direct patient care or assignment to a COVID-19-dedicated unit were not associated with COVID-19 infection in our cohort of HCW. Conversely, close contacts with suspected or confirmed COVID-19 patients without PPE or multiplication of contacts with colleagues without mask may be risk factors for COVID-19 infection in HCW. Systematic mask wearing outside home was associated with a reduction of this risk during the pre-lockdown period.

Authors' Contributions

AC, JL, OL and SK designed the study and drafted the paper. AC, JL, ML, FR, OL and SK contributed to data analysis and interpretation. All authors critically revised the manuscript for important intellectual content and gave final approval for the version to be published. SK had full access to all the data in the study and had final responsibility for the decision to submit for publication.

Declaration of interests

All authors declare no conflict of interest in relation with the submitted work

Role of the funding source

This study had no funding source or sponsor implicated in the study design, in the collection, analysis, and interpretation of data, in the writing of the report, and in the decision to submit the article for publication.

Ethical Considerations

This study was approved by the Ethical Review Committee for publications of the Cochin University Hospital (number AAA-2020–08012). According to French policy, a non-opposition statement was obtained for all participants, meaning that all had received written detailed information on the objectives of the study and were free to request withdrawal of their data at any time.

Acknowledgments

The authors warmly thank medical students involved in data collection: Laurence Clastres, Mathilde Lehmann, Aline Pellegrini, Marine Sisouvan, Ilana Slotine, Diem Soubou and Abigaëlle Vergnet, the following physicians and nurses who actively contributed to the screening: Claire Aguilar, Caroline Charlier, Gérard Chéron, Claudine Duvivier, Muriel Fortier, Remy Gauzit, Béatrice Grandordy, Marc Lecuit, Léonie Meyer, Gabrielle Paluszczek, Perrine Parize, Guillemette Thin and Christine Vinter.

References

1. Contejean A, Leporrier J, Canoui E, et al. Comparing dynamics and determinants of SARS-CoV-2 transmissions among health care workers of adult and pediatric settings in central Paris. *Clin Infect Dis* **2020**;
2. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA* **2020**;
3. Bryan A, Pepper G, Wener MH, et al. Performance Characteristics of the Abbott Architect SARS-CoV-2 IgG Assay and Seroprevalence in Boise, Idaho. *J Clin Microbiol* **2020**; 58.
4. Nguyen LH, Drew DA, Graham MS, et al. Risk of COVID-19 among front-line health-care workers and the general community: a prospective cohort study. *Lancet Public Health* **2020**; 5:e475–e483.
5. Çelebi G, Pişkin N, Çelik Bekleviç A, et al. Specific risk factors for SARS-CoV-2 transmission among health care workers in a university hospital. *Am J Infect Control* **2020**; 48:1225–1230.
6. Yan CH, Faraji F, Prajapati DP, Boone CE, DeConde AS. Association of chemosensory dysfunction and COVID-19 in patients presenting with influenza-like symptoms. *International Forum of Allergy & Rhinology* **2020**; 10:806–813.
7. Lan F-Y, Filler R, Mathew S, et al. COVID-19 symptoms predictive of healthcare workers' SARS-CoV-2 PCR results. *PLOS ONE* **2020**; 15:e0235460.

8. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med* **2020**;
9. Xiao J, Shiu EYC, Gao H, et al. Nonpharmaceutical Measures for Pandemic Influenza in Nonhealthcare Settings-Personal Protective and Environmental Measures. *Emerging Infect Dis* **2020**; 26:967–975.
10. Cheng VC-C, Wong S-C, Chuang VW-M, et al. The role of community-wide wearing of face mask for control of coronavirus disease 2019 (COVID-19) epidemic due to SARS-CoV-2. *Journal of Infection* **2020**; 81:107–114.
11. Lyu W, Wehby GL. Community Use Of Face Masks And COVID-19: Evidence From A Natural Experiment Of State Mandates In The US. *Health Affairs* **2020**; 39:1419–1425.
12. Lee B, Raszka WV. COVID-19 Transmission and Children: The Child Is Not to Blame. *Pediatrics* **2020**; 146.

Table: Cases and controls comparisons regarding demography, profession, symptoms and occupational or out-of-hospital exposures.

	Overall n=564	Controls (HCW-) n=228	Cases (HCW+) n=336	Univariate analysis Full period OR [95% CI]	Multivariate analysis Pre-lockdown OR [95% CI]	Multivariate analysis Post-lockdown OR [95% CI]
Median age (IQR), years	40 (31-53)	41 (33-52)	40 (30-53)	0.99 [0.98- 1.00]		
Sex (female)	448 (79)	183 (80)	265 (79)	0.91 [0.59-1.42]		
Professional category						
Physicians	155 (28)	64 (28)	91 (27)	ref		
Paramedic staff (nurses, care assistants...)	331 (59)	126 (55)	205 (61)	1.14 [0.77-1.69]		
Administrative staff (secretaries)	56 (10)	28 (12)	28 (8)	0.70 [0.38-1.30]		
Other employees (housekeepers...)	22 (4)	10 (4)	12 (4)	0.84 [0.34-2.11]		
Symptoms						
Headaches	420 (75)	158 (69)	262 (78)	1.57 [1.05-2.34]		
Asthenia	405 (72)	133 (58)	272 (81)	3.03 [2.04-4.52]		
Fever (measured or reported)	369 (65)	123 (54)	246 (73)	2.33 [1.61-3.38]		
Cough	363 (64)	136 (60)	227 (68)	1.41 [0.98-2.03]		
Muscle pain	323 (57)	103 (45)	220 (66)	2.30 [1.61-3.30]		
Rhinorrhoea	284 (50)	113 (49)	171 (51)	1.05 [0.74-1.50]		
Anosmia	244 (43)	15 (7)	229 (68)	30.2 [16.7-57.7]		
Agueusia	233 (41)	18 (8)	215 (64)	20.6 [12.0-37.3]		
Diarrhoea	197 (35)	68 (30)	129 (38)	1.47 [1.01-2.14]		
Dyspnoea	187 (33)	61 (27)	126 (38)	1.64 [1.12-2.42]		
Exposures						
In-hospital exposures						
Exposure to patients						
Occupational activities with direct patient facing	388 (69)	154 (68)	234 (70)	1.10 [0.75-1.60]		
Regularly posted in a unit dedicated to COVID19-patients	123 (22)	48 (21)	75 (22)	1.08 [0.70-1.66]		
Had on average ≥ 1 /day close contact with suspected or confirmed COVID-19 patients without PPE*	100 (18)	28 (12)	72 (21)	1.94 [1.19-3.25]	-	3.87 [1.73- 9.89]
Had on average ≥ 1 /day close contact with suspected or confirmed COVID-19 patients with PPE	204 (38)	93 (45)	111 (34)	0.61 [0.41-0.88]		
Exposure to colleagues						
Wears a medical mask always/most of the time at hospital	372 (67)	163 (73)	209 (63)	0.63 [0.42-0.92]		
Spends on average >1 hour/day with colleagues without mask	153 (31)	42 (24)	111 (34)	1.68 [1.09-2.62]		

Had on average >10 close contacts/day with colleagues without mask	101 (18)	26 (12)	75 (23)	2.24 [1.36-3.80]	2.58 [1.49-4.60]	-
Out-of-hospital exposure						
Uses public transports	328 (58)	127 (56)	201 (60)	1.18 [0.83-1.69]	0.43 [0.21-0.85]	-
Systematically wears a mask outside home	96 (17)	50 (22)	46 (14)	0.57 [0.36-0.90]		
Systematically washes hands when back home	509 (91)	210 (92)	299 (90)	0.80 [0.41-1.52]		
Leaves home on average ≥ 1 times a week	495 (88)	205 (90)	290 (86)	0.71 [0.40-1.23]		
Lives with ≥ 2 additional household members	281 (50)	109 (48)	172 (51)	1.14 [0.80-1.61]		
Lives with ≥ 1 child aged 0-4 years	82 (15)	35 (15)	47 (14)	0.90 [0.54-1.49]		
Lives with ≥ 1 child aged 5-15 years	138 (25)	60 (26)	78 (23)	0.85 [0.56-1.27]		
Lives with ≥ 1 adult aged 70+ years	12 (3)	6 (3)	6 (2)	0.67 [0.18-2.55]		
Has ≥ 1 child kept outside the household (school, nursery)	105 (19)	45 (20)	60 (18)	0.88 [0.56-1.39]		
Nursery	31/105 (30)	13/45 (29)	18/60 (30)	1.05 [0.42-2.71]		
School	70/105 (67)	28/45 (62)	42/60 (70)	1.41 [0.58-3.47]		
Children kept in a facility with >5 other children	89/100 (89)	35/41 (85)	54/59 (92)	1.84 [0.43-8.25]		
<hr/>						
*PPE: Personal Protective Equipment						