

SUPPLEMENTAL MATERIAL

SARS-CoV-2 Infection Hospitalization Rate and Infection Fatality Rate among the Non-Congregant Population in Connecticut

Table of contents:

eMethods. COVID-19 case classification criteria

eTable 1. ICD-10 codes to identify COVID-19 visits by Connecticut Hospital Association.

eTable 2. Estimated number of individuals with SARS-CoV-2-specific antibodies among the non-congregate population in Connecticut, between March 1 and June 1, 2020, by sociodemographic characteristics.

eTable 3. Estimates for infection hospitalization rate and infection fatality rate at 95% margin of error among the non-congregate population in Connecticut between March 1 and June 1, 2020, by sociodemographic characteristics

eTable 4. Confirmed, probable, and total infection fatality rate among the non-congregate population in Connecticut between March 1 and June 1, 2020, by sociodemographic characteristics.

eMethods. COVID-19 case classification criteria

Case classifications –

- Confirmed:
 - Meets confirmatory laboratory evidence.
- Probable:
 - Meets clinical criteria AND epidemiologic evidence with no confirmatory laboratory testing performed for COVID-19.
 - Meets presumptive laboratory evidence AND either clinical criteria OR epidemiologic evidence.
 - Meets vital records criteria with no confirmatory laboratory testing performed for COVID19.

Source: Standardized surveillance case definition and national notification for 2019 novel coronavirus disease (COVID-19). Council of State and Territorial Epidemiologists.

https://cdn.ymaws.com/www.cste.org/resource/resmgr/2020ps/interim-20-id-01_covid-19.pdf.

Published 2020. Accessed September 8, 2020.

eTable 1. ICD-10 codes to identify COVID-19 visits by Connecticut Hospital Association.

ICD-10 Codes	Description
J1289 and (B9729 or U071)	COVID-related Pneumonia
(J208 or J40) and (B9729 or U071)	COVID-related Bronchitis
(J988 or J22) and (B9729 or U071)	COVID-related respiratory infection
J80 and (B9729 or U071)	COVID-related ARDS
U071	COVID-19

Note: the above diagnosis codes can be in any position, in other words, they do not have to be the principle diagnosis code.

eTable 2. Estimated number of individuals with SARS-CoV-2-specific antibodies among the non-congregate population in Connecticut, between March 1 and June 1, 2020, by sociodemographic characteristics.

Characteristics	Total Population*, N	Seroprevalence[†] of SARS-CoV-2 antibodies, % (±MOE at 90% CI)	Estimated number of individuals with antibodies, N (90% CI)	Seroprevalence[†] of SARS-CoV-2 antibodies, % (±MOE at 95% CI)	Estimated number of individuals with antibodies, N (95% CI)
State total	2,837,877	4.0% (±2.0)	113,515 (56,758 – 170,273)	4.0% (±2.3)	113,515 (48,244 – 178,786)
Sex					
Men	1,365,019	2.5% (±2.4%)	34,125 (1,365 – 66,886)	2.5% (±2.9%)	34,125 (NE – 73,711)
Women	1,472,858	5.3% (±2.9%)	78,061 (35,349 – 120,774)	5.3% (±3.5%)	78,061 (26,511– 129,612)
Age group					
18-29	564,738	6.4% (±7.7%)	36,143 (NE – 79,628)	6.4% (±9.2%)	36,143 (NE – 88,099)
30-44	649,874	4.9% (±4.6%)	31,844 (1,950 – 61,738)	4.9% (±5.5%)	31,844 (NE – 67,587)
45-54	496,628	6.6% (±4.7%)	32,777 (9,436 – 56,119)	6.6% (±5.6%)	32,777 (NE – 60,589)
55-64	513,656	2.6% (±2.4%)	13,355 (1,027 – 25,683)	2.6% (±2.9%)	13,355 (NE – 28,251)
≥65	612,981	0.8% (±1.2%)	4,904 (NE – 12,260)	0.8% (±1.4%)	4,904 (NE – 13,486)
Race/Ethnicity					
Non-Hispanic Black	278,112	6.4% (±5.5%)	17,799 (2,503 – 33,095)	6.4% (±6.5%)	17,799 (NE – 35,876)
Non-Hispanic White	1,969,487	2.7% (±1.7%)	53,176 (19,695 – 86,657)	2.7% (±2.0%)	53,176 (13,786 – 92,566)
Hispanic	408,654	19.9% (±6.7%)	81,322 (53,942 – 108,702)	19.9% (±8.0%)	81,322 (48,630 – 114,015)
County					
New Haven	683,928	3.4% (±3.2%)	23,254 (1,368 – 45,139)	3.4% (±3.8%)	23,254 (NE – 49,243)
New London	215,679	1.7% (±3.3%)	3,667 (NE – 10,784)	1.7% (±4.0%)	3,667 (NE – 12,294)
Middlesex	133,380	-	-	-	-
Fairfield	732,172	5.7% (±5.4%)	41,734 (2,197 – 81,271)	5.7% (±6.4%)	41,734 (NE – 88,593)
Hartford	706,631	4.0% (±3.6)	28,265 (2,827 – 53,704)	4.0% (±4.3%)	28,265 (NE – 58,650)
Litchfield	147,570	1.6% (±3.2%)	2,361 (NE – 7,083)	1.6% (±3.8%)	2,361 (NE – 7,969)
* Source: 2018 American Community Survey					
† Source: Post-Infection Prevalence Study					
Abbreviations: CI, confidence interval; MOE, margin of error					

eTable 3. Estimates for infection hospitalization rate and infection fatality rate at 95% margin of error among the non-congregate population in Connecticut between March 1 and June 1, 2020, by sociodemographic characteristics

	COVID-19-related Hospitalizations, N	Infection Hospitalization Rate, Weighted % (95% CI)	Total* COVID-19-related Deaths, N	Total Infection Fatality Rate, Weighted % (95% CI)
State Total	7792	6.86 (4.36–16.15)	1079	0.95 (0.60–2.24)
Sex				
Men	4166	12.21 (5.67–NE)	815	2.39 (1.11–NE)
Women	3625	4.64 (2.80–14.21)	264	0.34 (0.20–1.00)
Age-group				
18-29	288	0.80 (0.33–NE)	5	0.01 (0.01–NE)
30-44	855	2.68 (1.27–NE)	34	0.11 (0.05–NE)
45-54	1013	3.09 (1.67–NE)	59	0.18 (0.10–NE)
55-64	1660	12.43 (5.88–NE)	174	1.30 (0.62–NE)
≥65	3918	79.89 (29.05–NE)	807	16.46 (5.98–NE)
Race/Ethnicity				
Non-Hispanic Black	1762	9.90 (1.90–NE)	259	1.46 (0.72–NE)
Non-Hispanic White	3948	7.42 (11.00–28.64)	593	1.12 (0.64–4.30)
Hispanic	845	1.04 (0.74–1.74)	190	0.23 (0.17–0.39)
County				
New Haven	2101	9.04 (4.27–NE)	270	1.16 (0.55–NE)
New London	148	4.04 (1.20–NE)	21	0.57 (0.17–NE)
Middlesex	231	NE	32	NE
Fairfield	2665	6.39 (3.01–NE)	399	0.96 (0.45–NE)
Hartford	1755	6.21 (2.99–NE)	294	1.04 (0.50–NE)
Litchfield	195	8.26 (2.45–NE)	39	1.65 (0.49–NE)
Abbreviations: CI, confidence interval; NE, non-estimable				
*Total COVID-19-related deaths includes both confirmed and probable COVID-19 deaths.				

eTable 4. Confirmed, probable, and total infection fatality rate among the non-congregate population in Connecticut between March 1 and June 1, 2020, by sociodemographic characteristics.

	Confirmed deaths, N	Confirmed IFR, weighted % (90% CI)	Confirmed IFR, weighted % (95% CI)	Probable deaths, N	Probable IFR, weighted % (90% CI)	Probable IFR, weighted % (95% CI)
State Total	860	0.76 (0.51–1.52)	0.76 (0.48–1.78)	219	0.19 (0.13–0.39)	0.19 (0.12–0.45)
Sex						
Men	598	1.75 (0.89–43.81)	1.75 (0.81–NE)	217	0.64 (0.32–15.90)	0.64 (0.29–NE)
Women	262	0.34 (0.22–0.74)	0.34 (0.20–0.99)	2	0.00 (0.00–0.01)	0.00 (0.00–0.01)
Age group						
18-29	5	0.01 (0.01–NE)	0.01 (0.01–NE)	0	0.00 (0.00–0.00)	0.00 (0.00–0.00)
30-44	30	0.09 (0.05–1.54)	0.09 (0.04–NE)	4	0.01 (0.01–0.21)	0.01 (0.01–NE)
45-54	47	0.14 (0.08–0.50)	0.14 (0.08–NE)	12	0.04 (0.02–0.13)	0.04 (0.02–NE)
55-64	148	1.11 (0.58–14.41)	1.11 (0.52–NE)	26	0.19 (0.10–2.53)	0.19 (0.09–NE)
≥65	630	12.85 (5.14–NE)	12.85 (4.67–NE)	177	3.61 (1.44–NE)	3.61 (1.31–NE)
Race/ethnicity						
Non-Hispanic Black	220	1.24 (0.66–8.79)	1.24 (0.61–NE)	39	0.22 (0.12–1.56)	0.22 (0.11–NE)
Non-Hispanic White	450	0.85 (0.52–2.28)	0.85 (0.49–3.26)	143	0.27 (0.17–0.73)	0.27 (0.15–1.04)
Hispanic	160	0.20 (0.15–0.30)	0.20 (0.14–0.33)	30	0.04 (0.03–0.06)	0.04 (0.03–0.06)
County						
New Haven	236	1.01 (0.52–17.25)	1.01 (0.48–NE)	34	0.15 (0.08–2.49)	0.15 (0.07–NE)
New London	15	0.41 (0.14–NE)	0.41 (0.12–NE)	6	0.16 (0.06–NE)	0.16 (0.05–NE)
Middlesex	26	NE	NE	6	NE	NE
Fairfield	315	0.75 (0.39–14.34)	0.75 (0.36–NE)	84	0.20 (0.10–3.82)	0.20 (0.09–NE)
Hartford	224	0.79 (0.42–7.92)	0.79 (0.38–NE)	70	0.25 (0.13–2.48)	0.25 (0.12–NE)
Litchfield	28	1.19 (0.40–NE)	1.19 (0.35–NE)	11	0.47 (0.16–NE)	0.47 (0.14–NE)
Abbreviations: CI, confidence interval; IFR, infection fatality rate; NE, non-estimable						