SUPPLEMENTAL MATERIAL

Racial and Ethnic Disparities in Health of Adults in the United States: A Twenty-Year National Health Interview Survey Analysis, 1999-2018

Table of contents:

eFigure 1. Flowchart of study population.


eFigure 2. Distribution of hours of sleep per day by race/ethnicity.


eTable 1. Sociodemographic and clinical characteristics of the study population during the first

two years (1999 and 2000) and the last two years (2017 and 2018) of the study period by

race/ethnicity.

eFigure 1. Flowchart showing selection of the study population.





eFigure 2. Distribution of hours of sleep per day by race/ethnicity.

	Non-Hispanic White		Hispanic		Non-Hispanic Black		Non-Hispanic Asian	
	1999-2000	2017-2018	1999-2000	2017-2018	1999-2000	2017-2018	1999-2000	2017-2018
	n (%, 95%		n (%, 95%	n (%, 95%	n (%, 95%	n (%, 95%	n (%, 95%	n (%, 95%
Characteristics	CI)	n (%, 95% CI)	CI)	CI)	CI)	CI)	CI)	CI)
Total	41,882	36,410	10,305	6,423	8,758	5,884	1,597	2,640
Age (years)								
18–39	15,169 (39.5%	10,050 (33.3%	3,610 (37.1%	2,886 (50.1%	3,862 (49.8%	1,994 (43.7%	844 (52.4%	1,085 (41.9%
	[38.9-40.2])	[32.6–34.1])	[35.0-39.2])	[48.4–51.8])	[48.3–51.3])	[41.8-45.6])	[49.2–55.6])	[39.3-44.6])
40–64	17,246 (42.2% [41.5–42.8])	14,919 (42.3% [41.6–43.0])	6,617 (62.1% [60.0–64.2])	2,492 (39.1% [37.6–40.6])	3,539 (38.5% [37.1–39.8])	2,434 (41.1% [39.3–42.9])	581 (38.3% [35.1–41.4])	1,005 (42.5% [40.2–44.9])
≥65	9,467 (18.3% [17.8–18.8])	11,441 (24.4% [23.7–25.1])	78 (0.8% [0.6–1.1])	1,045 (10.8% [9.9–11.8])	1,357 (11.8% [11.0–12.6])	1,456 (15.3% [14.2–16.3])	172 (9.4% [7.8–11.4])	550 (15.5% [13.9–17.4])
Sex	[17.0–10.0])	[23.7-23.1])	[0.0-1.1])	[9.9–11.8])		[14.2-10.3])	[7.0-11.4])	[13.7-17.4])
Men	18,512 (48.1% [47.6–48.7])	16,798 (48.6% [48.0–49.3])	4,399 (49.4% [48.2–50.6])	2,788 (49.7% [48.1–51.4])	3,255 (44.4% [43.0–45.9])	2,435 (45.5% [43.7–47.2])	726 (48.7% [45.4–51.9])	1,270 (46.7% [44.4–49.0])
Women	23,370 (51.9% [51.3–52.4])	19,612 (51.4% [50.7–52.0])	5,906 (50.6% [49.4–51.8])	3,635 (50.3% [48.7–51.9])	5,503 (55.6% [54.1–57.0])	3,449 (54.5% [52.8–56.3])	871 (51.4% [48.1–54.6])	1,370 (53.3% [51.1–55.6])
Citizenship status								
US citizen	41,177 (98.3% [98.2–98.5])	35,943 (98.4% [98.2–98.6])	6,617 (62.1% [60.0–64.2])	4,694 (71.5% [69.8–73.3])	8,385 (95.2% [94.5–95.8])	5,655 (94.8% [93.6–95.8])	921 (57.1% [53.5–60.6])	1,844 (69.9% [67.2–72.4])
Non-US citizen	672 (1.6% [1.5–1.8])	446 (1.5% [1.3– 1.7])	3,610 (37.1% [35.0–39.2])	1,664 (27.4% [25.7–29.1])	360 (4.6% [4.0–5.3])	220 (5.1% [4.1–6.2])	658 (41.6% [38.0–45.2])	775 (29.2% [26.7–31.8])
Unknown	33 (0.0% [0.0– 0.0])	21 (0.0% [0.0– 0.1])	78 (0.8% [0.6–1.1])	65 (1.1% [0.8–1.6])	$ \begin{array}{c} 13 (0.2\% \\ [0.0-0.4]) \end{array} $	9 (0.1% [0.0– 0.3])	18 (1.3% [0.7–2.4])	21 (0.9% [0.5–1.7])
Education level	~··~ 1/	1/	<i>/</i>				[])	[]
Less than high school	5,770 (13.3% [12.8–13.8])	2,708 (7.4% [6.9–7.9])	4,810 (44.0% [42.4–45.7])	1,831 (27.5% [25.7–29.4])	2,228 (24.1% [22.8–25.5])	948 (14.1% [12.8–15.6])	179 (11.2% [9.3–13.5])	221 (7.9% [6.6–9.4])

eTable 1. Sociodemographic and clinical characteristics of the study population during the first two years (1999 and 2000) and the last two years (2017 and 2018) of the study period by race/ethnicity.

	1		T	T	1			
High school diploma /GED	12,918 (31.5% [30.8–32.1])	8,710 (23.6% [22.8–24.3])	2,394 (24.0% [22.9–25.0])	1,615 (26.4% [24.8–28.0])	2,569 (31.1% [29.8–32.4])	1,646 (28.7% [27.1–30.3])	276 (17.7% [15.8–19.8])	383 (15.2% [13.4–17.3])
Some college	12,147 (29.3% [28.8–29.8])	11,509 (31.3% [30.5–32.0])	2,086 (21.4% [20.2–22.6])	1,737 (27.8% [26.3–29.4])	2,558 (29.3% [28.0–30.6])	1,905 (32.7% [31.0–34.5])	367 (23.8% [21.3–26.5])	546 (20.6% [18.7–22.8])
≥Bachelor's	10,685 (25.2%	13,386 (37.5%	888 (9.2%	1,193 (17.3%	1,300 (14.3%	1,347 (23.9%	750 (45.6%	1,478 (55.6%
degree	[24.5–25.9])	[36.4–38.6])	[8.4–10.1])	[15.9–18.8])	[13.2–15.5])	[22.0–25.9])	[42.6–48.6])	[52.5–58.6])
Unknown	362 (0.8% [0.7–0.9])	97 (0.3% [0.2– 0.4])	127 (1.4% [1.1–1.8])	47 (1.0% [0.7–1.5])	103 (1.3% [1.0–1.6])	38 (0.6% [0.4–0.9])	25 (1.8% [1.2–2.7])	12 (0.7% [0.3–1.5])
Family income	[0.7-0.7])	0.4])	[1.1-1.0])		[1.0-1.0])	[0.4-0.7])		[0.3-1.3])
•	20 642 (77 00/	27 211 (79 90/	4 412 (40 20/	2165(5250)	4 272 (54 50)	2 0 6 1 (55 20/	1 002 (71 00/	1.006 (72.00/
High/middle	30,643 (77.0%	27,211 (78.8%	4,412 (49.3%	3,165 (53.5%	4,372 (54.5%	2,961 (55.3%	1,082 (71.8%	1,886 (73.8%
income	[76.4–77.6])	[77.9–79.6])	[47.6–50.9])	[51.4–55.5])	[52.5–56.5])	[52.9–57.6])	[68.3–75.1])	[71.0–76.5])
Low income	11,239 (23.0% [22.4–23.6])	9,199 (21.2% [20.4–22.1])	5,893 (50.7% [49.1–52.4])	3,258 (46.5% [44.5–48.6])	4,386 (45.5% [43.6–47.5])	2,923 (44.7% [42.4–47.1])	515 (28.2% [24.9–31.7])	754 (26.2% [23.5–29.0])
Insurance	[22.4-23.0])	[20.4-22.1])	[+7.1-52.4])	[++]-+0.0])	[43.0-47.3])	[+2.+-+7.1])	[24.7-31.7])	[23.3-27.0])
status								
	37,216 (88.7%	34,050 (93.2%	6,669 (63.4%	4,937 (75.4%	7,036 (78.9%	5,216 (87.0%	1,329 (82.3%	2,469 (93.2%
Insured	[88.3–89.1])	[92.8–93.6])	[61.6–65.2])	[73.4–77.2])	[77.8–80.0])	[85.6-88.3])	[80.0-84.5])	[91.8–94.3])
Uninsured	4,522 (11.0%	2,270 (6.5%	3,591 (36.1%	1,454 (23.8%	1,663 (20.1%	632 (11.9%	260 (17.3%	163 (6.3%
Uninsuled	[10.6–11.4])	[6.2–6.9])	[34.3–37.9])	[22.0–25.8])	[19.1–21.2])	[10.7–13.4])	[15.1–19.7])	[5.2–7.6])
Unknown	144 (0.3%	90 (0.3% [0.2–	45 (0.5%	32 (0.8%	59 (1.0%	36 (1.1%	8 (0.4%	8 (0.6%
UIIKIIOWII	[0.3–0.4])	0.4])	[0.4–0.7])	[0.5–1.3])	[0.7–1.3])	[0.7-1.7])	[0.2–0.9])	[0.3–1.2])
Region								
Northeast	8,168 (20.2%)	6,431 (19.2%	1,639 (15.7%	789 (13.5%	1,582 (18.0%	733 (16.0%	333 (21.2%	493 (20.8%
normeast	[19.5–21.0])	[17.5–21.1])	[14.5–17.1])	[11.2–16.2])	[16.7–19.4])	[13.5–18.8])	[18.7–24.0])	[16.9–25.4])
Midwest	12,012 (29.4%	10,273 (27.4%	637 (7.9%	640 (9.4%	1,698 (18.8%	914 (15.2%	214 (14.7%	325 (11.8%
Midwest	[28.5–30.3])	[25.7–29.3])	[6.9–9.1])	[7.6–11.6])	[17.2–20.5])	[13.0–17.8])	[11.9–18.0])	[9.5–14.6])
South	13,935 (33.9%	11,884 (33.0%	3,729 (35.4%	2,614 (37.1%	4,645 (55.7%	3,754 (60.5%	298 (18.9%	683 (25.5%
South	[33.0–34.8])	[30.9–35.2])	[33.2–37.8])	[32.5-42.0])	[53.4–57.9])	[56.7–64.3])	[16.4-21.8])	[21.6-30.0])
West	7,767 (16.5%	7,822 (20.4%	4,300 (40.9%	2,380 (40.0%	833 (7.6%	483 (8.3%	752 (45.2%	1,139 (41.8%
	[15.8–17.2])	[18.3–22.7])	[38.6–43.3])	[35.3-44.8])	[6.7–8.5])	[6.9–9.9])	[41.2–49.2])	[36.7–47.1])
Marital status								

Unmarried	20,259 (38.3%	19,152 (43.6%	4,953 (41.3%	3,569 (50.6%	6,232 (62.2%	4,364 (67.2%	664 (35.5%	1,123 (34.9%
	[37.5–39.0])	[42.8–44.3])	[40.2–42.5])	[48.9–52.2])	[60.8–63.7])	[65.5–68.9])	[32.5–38.6])	[32.5–37.4])
Married/Living	21,431 (61.4%	17,179 (56.3%	5,302 (58.3%	2,840 (49.1%	2,459 (37.1%	1,502 (32.6%	923 (64.0%	1,509 (64.9%
with partner	[60.6–62.1])	[55.5–57.0])	[57.1–59.4])	[47.5–50.7])	[35.7–38.6])	[30.9–34.3])	[61.0–67.0])	[62.4–67.4])
Unknown	192 (0.3%	79 (0.2% [0.1–	50 (0.4%	14 (0.4%	67 (0.6%	18 (0.3 [0.2–	10 (0.5%	8 (0.2%
UIIKIIOWII	[0.3–0.4])	0.2])	[0.3–0.5])	[0.2–0.6])	[0.5–0.9])	0.4])	[0.3–0.9])	[0.0–0.5])
Employment								
Status								
With a	26,251 (65.6%	20,990 (62.1%	6,430 (66.1%	4,075 (66.5%	5,410 (63.9%	3,248 (61.0%	1,029 (65.9%	1,667 (67.1%
job/Working	[65.0–66.3])	[61.3–62.9])	[64.9–67.4])	[64.7–68.3])	[62.4–65.4])	[59.0–63.1])	[63.2–68.5])	[64.9–69.3])
Not in labor	15,037 (32.9%	14,581 (35.4%	3,579 (31.1%	2,138 (29.8%	3,020 (31.9%	2,308 (32.4%	530 (31.7%	907 (30.0%
force	[32.2–33.6])	[34.6–36.2])	[29.9–32.3])	[28.0-31.8])	[30.5–33.4])	[30.5–34.3])	[29.1–34.4])	[27.8–32.3])
Unomployed	562 (1.4%	824 (2.5% [2.3–	276 (2.5%	208 (3.6%	308 (3.9%	324 (6.5%	35 (2.2%	64 (2.8%
Unemployed	[1.2–1.5])	2.7])	[2.2–2.9])	[3.0–4.3])	[3.4–4.6])	[5.7–7.4])	[1.5–3.1])	[2.1–3.7])
Unknown	32 (0.1% [0.1–	15 (0.0% [0.0–	20 (0.3%	2 (0.02%	20 (0.3%	4 (0.1% [0.0–	3 (0.2%	2 (0.1%
	0.1])	0.1])	[0.2–0.4])	[0.0-0.1])	[0.2–0.4])	0.3])	[0.0-0.8])	[0.0-0.4])
Comorbidities								
Unantonoion	10,466 (23.0%	13,280 (32.7%	1,762 (15.3%	1,673 (22.3%	2,857 (28.8%	2,687 (37.6	237 (14.9%	670 (24.2%
Hypertension	[22.5–23.4])	[31.9–33.4])	[14.4–16.2])	[21.0-23.8])	[27.6–30.1])	[35.9–39.3])	[12.9–17.2])	[22.2–26.3])
Diabetes	2,303 (5.2%	3,698 (9.1%	720 (6.3%	766 (10.5%	803 (8.2%	888 (11.7%	57 (3.9%	247 (8.9%
Diabetes	[5.0–5.4])	[8.7–9.4])	[5.8–6.9])	[9.6–11.4])	[7.5-8.9])	[10.8–12.7])	[3.0–5.1])	[7.7–10.3])
Prior stroke/MI	2,404 (5.1%	2,762 (6.4%	307 (2.5%	266 (3.3%	456 (4.5%	484 (6.0%	35 (1.7%	97 (3.2%
PHOI SUOKE/IVII	[4.9–5.3])	[6.1–6.8])	[2.2–2.9])	[2.8–3.9])	[4.0–5.1])	[5.4–6.7])	[1.2–2.2])	[2.5–4.0])
Cancer	3,628 (7.8%	5,164 (12.3	235 (2.2%	289 (3.4%	301 (2.9%	379 (4.8%	31 (1.7%	126 (4.3%
Cancer	[7.5–8.1])	[11.9–12.8])	[1.9–2.5])	[2.9–3.8])	[2.5–3.4])	[4.3–5.4])	[1.1–2.7])	[3.4–5.3])
Emphysema /	2,785 (6.2%	2,163 (5.3%	330 (2.9%	231 (2.8%	444 (4.3%	332 (4.4%	35 (1.6%	58 (2.0%
Chronic	[5.9–6.5])	[5.0–5.6])	[2.5-3.4])	[2.4-3.3])	[3.8–4.7])	[3.7-5.1])	[1.1-2.4])	[1.4–2.7])
bronchitis	[3.9-0.3])	[5.0-5.0])	[2.3-3.4])	[2.4-3.3])	[3.0-4.7])	[3.7-3.1])	[1.1-2.4])	[1.4-2.7])
Current	10,085 (24.2%	5,712 (15.1%	1,915 (18.3%	690 (9.8%	2,124 (23.8%	973 (14.7%	247 (14.9%	201 (7.2%
smoker	[23.7–24.8])	[14.6–15.7])	[17.4–19.3])	[9.0–10.8])	[22.6–25.1])	[13.5–16.1])	[12.9–17.2])	[6.0-8.5])
Flu vaccine in	13,642 (30.7%	17,708 (47.1%	1,982 (18.6%	2,379 (34.9%	1,879 (20.6%	2,198 (35.2%	426 (26.1%	1,270 (48.2%
past 12m	[30.0–31.3])	[46.4–47.9])	[17.5–19.6])	[33.3–36.5])	[19.7–21.5])	[33.4–37.1])	[23.3–29.2])	[45.6–50.7])

Obese (BMI	8,006 (20.2%	10,627 (30.3%	2,339 (23.0%	2,093 (34.0%	2,542 (29.1%	2,312 (39.4%	84 (6.1%	289 (12.0%	
\geq 30 kg/m ²)	[19.7–20.6])	[29.6–31.0])	[21.9–24.1])	[32.5–35.6])	[28.0–30.2])	[37.8–41.1])	[4.9–7.7])	[10.5–13.6])	
Abbreviations: BMI, body mass index; CI, confidence interval; MI, myocardial infarction; US, United States									