Title

The smartphone: an evolution or revolution in virtual patient healthcare during and beyond the COVID-19 pandemic ? An evaluation and comparison of the smartphone against other currently available wearable technologies in a secondary care setting during the COVID-19 pandemic.

Corresponding Author:

Professor Asif Raza,

Dept of Urology: London Northwest University Healthcare NHS Trust (LNWUH NHS Trust)

London, UK, HA1 3UJ

Plymouth University Peninsula Medical and Dental School

Imperial Medical College, London UK

Email: asifraza@nhs.net

Tel +44 2089675152, Fax: +44 207 112 8283

Other authors:

Mr Subhabrata Mukherjee

Dept of Urology: London Northwest University Healthcare NHS Trust (LNWUH NHS Trust)

London, UK, HA1 3UJ

Dr Vikash Patel

Dept of Urology: London Northwest University Healthcare NHS Trust (LNWUH NHS Trust)

London, UK, HA1 3UJ

Dr Naila Kamal

Dept of Obstetrics and Gynaecology: London Northwest University Healthcare NHS Trust (LNWUH NHS Trust)

London, UK, HA1 3UJ

Dr Ewa – Lichtarowicz-Krynska

Dept of Paediatrics : London Northwest University Healthcare NHS Trust (LNUWH NHS Trust), River Island Academic Centre for Paediatrics and Child Health, Imperial College, London, UK, HA1 3UJ

Key Words: Telemedicine, MHealth, COVID-19, Smartphone, Cross infection

Word Count: 1497

Summary Box: 294

Abstract

Smartphones are now commonly used, for virtual outpatient consultations, to help reduce disease transmission during the COVID-19 pandemic. Nosocomial spread of COVID 19 and hospital acquired infections are usually from staff or students to patients. Reducing non-essential staff numbers on ward rounds may reduce the risk.

We describe the novel use of smartphones, with Microsoft Teams, to live stream inpatient interactions, radiological images, pathology results, charts and patient review between an office-based and ward team (virtual ward round) and for teaching medical students in secondary care. After Research and Ethics, Digital services and Information Governance approval we compared a smartphone and head-worn device (Realwear HMT-1). Data collection was by participant questionnaire. Statistical analysis was performed using the Mann – Whitney test.

There was no statistically significant difference in audio and video feed quality between the smart phone (p value = 0.3) and Realwear device (p value = 0.41). However the smartphone was preferred during ward rounds and was 85% cheaper than the Realwear device. Urology medical staff numbers on the ward were reduced by 50%. Ward round efficiency improved as administrative tasks could be performed by the office team during the virtual ward round.

Virtual ward rounds using smartphones can facilitate remote communication between staff, students and patients. Staff in isolation or shielding can also assist front line colleagues from home. Smarter use of the smart phone may help reduce staff numbers on wards and reduce the number COVID-19 and nosocomial infections, potentially reducing morbidity and mortality locally and globally.

Title

The smartphone: an evolution or revolution in virtual patient healthcare during and beyond the COVID-19 pandemic ? An evaluation and comparison of the smartphone against other currently available wearable technologies in a secondary care setting during the COVID-19 pandemic.

Introduction

There has been an exponential increase in telemedicine, [1,2] mobile health platforms (mhealth) [3-5] and virtual clinics during the COVID-19 pandemic.[6] Up to 87% of doctors use their smartphone for work related tasks.[7]

The necessity for certain patients or staff to shield or isolate during the first wave of COVID-19 e.g. higher risk Black, Asian and Minority Ethnic (BAME) staff, highlighted limitations in communication platforms between staff, and between doctors and patients. This could have contributed to higher patient morbidity and mortality particularly in care homes.[8] Medical student teaching suffered as hospital access was restricted during the height of the pandemic. Newer but often costly technologies are playing an increasing role in medical education.[9]

However concerns remain about smart devices and data privacy and security, network and Wi-Fi capabilities, costs, user acceptance and bacterial contamination. The Information Commissioner's Office (UK) enforce General Data Protection Regulations (GDPR) [10] and NHS Digital has approved Microsoft teams (MS Teams), a secure platform, with 256-bit Advanced Encryption Standard (AES) protection of data for routine use by clinicians.

Staff smartphones may be colonised with MRSA and VRE.[11-13] However smartphone and tablet touchscreens are associated with lower infection levels than standard touchpad keyboards in hospitals.[14] Decontamination methods include antibacterial screens (silver

coated), alcohol-based wipes (70% isopropyl alcohol), Clinell wipes and Ultraviolet light(UVC). [15-17]

Performing virtual ward rounds may help reduce nosocomial infection risk including COVID 19 transmission. Up to 1.4 million patients are affected by non-COVID-19 hospital acquired nosocomial infections annually increasing morbidity and mortality particularly in the elderly and frail. [18] Up to 20%-70% patient files and charts harbour infection due to frequent handling by staff [19-21] COVID-19 spreads by droplet transmission, poor hand hygiene and contact between staff and patients [22] and up to 80% of staff may be asymptomatic carriers of COVID-19 [23] risking nosocomial spread from staff to patients. [24 – 27]

Virtual ward rounds have been reported previously, with less staff on the ward, using a double telepresence robot (iPad on a trolley),[28] Microsoft Hololens [29] and laptops carried onto the ward.[30] The adoption of an office based team (OBT) and remote ward team (RWT) in our practice reduced staff numbers by 50% on the ward round.

We wished to determine the most practical, user friendly and cost-effective wearable device that would allow staff to safely communicate remotely during virtual ward rounds, assess device practicality, use in theatres by surgeons, and for remote teaching of medical students during and after COVID 19.

Method

A google search revealed several commercially available wearable devices that live stream video feed. Devices required MS Teams functionality (without using third party apps for connection to MS teams) to enable live video streaming and two-way audio communication. Selected devices were tested in mock clinical scenarios (Paediatric Resuscitation and Sim-Man simulation training) and subsequently tested with patients. Fig 1

Our local Research and Ethics committee approved the study as a service evaluation and proof of concept study. The local Information Governance Officer (IGO) and local Digital services team approved live streaming via MS Teams. No recording or storing of video or images during or after the mock or real-life patient interaction was allowed for the study unless separate written consent was obtained for video or images to be kept for teaching purposes or publication. Informed consent was taken from patients, students and staff prior to streaming.

Inclusion criteria included all genders, over 18 years old, all socio-economic groups and medical or surgical condition not requiring emergency treatment. Exclusion criteria included patients unable to give valid informed consent due to a language barrier or incapacity, patients with life threatening emergencies, or those who were septic or confused on admission.

An online questionnaire was completed by the wearer of the device (RWT) and receivers of the live video feed stream (OBT, students) after streaming the interaction. (Table 1). Likert scoring was used to assess wearer and receiver satisfaction with the devices and streaming quality. Patients were asked to complete a post streaming satisfaction questionnaire.

The (OBT) and students were blinded to the device being used by the (RWT). The device worn was randomly allocated to the (RWT) by using an online randomisation programme (Graph Pads: Quickcalc) The same mock clinical scenario was used for teaching medical students (Fig 1.) for each device to ensure devices could be compared in the same location ensuring Wi-Fi signal strength and network characteristics were similar for both devices.

A sim free hospital approved smartphone was used at the request of our Digital Services department in preference to a personal smartphone. The selected devices were connected to

the staff only hospital Wi-Fi, registered to a dedicated network via MAC address and Wi-Fi password was WPA2 encrypted. Device access was password protected.

To ensure privacy of ward patients not in the study bedside curtains were drawn around the patient, in the ward, before live streaming started and undrawn when streaming stopped. The wearer of the device would place the device in standby mode, or switch it off before curtains were withdrawn, and moving to the next patient.

The statistical analyses compared responses for the smartphone to the Realwear device. Due to the ordinal nature of the recommendation measure, the analyses were performed using the Mann-Whitney test. Separate analyses were performed for wearers and receivers.

Results

Table 1 summary of results : type of device used, questionnaire results, different and use of devices with PPE.

The smartphone was most practical in a ward setting. Most respondents preferred ease of use, the audio and video feed, larger screen, being able to easily see radiological images and results on the smartphone compared to the Realwear device. Most of the negative comments when using a smartphone were due to the harness or mount rather than smartphone.

The Realwear device was preferred for theatre use but not suitable for wards or teaching as the noise cancellation on the device did not allow patients or other staff voices to be picked up by its microphone and therefore heard by the OBT or students. The resolution of the images being viewed remotely were not high enough definition to teach surgeons comprehensively however enough anatomical detail was visible to provide advice to the trainee during a basic surgical procedure.

The results of the online questionnaire suggested that there was no statistically significant difference between the Phone and Realwear for either group of study participants. None of the patients objected to being live streamed or the concept of the virtual ward round.

Discussion

Limitations in our study included the small number of study participants, the use of older model hospital iphones without a sim, with lower screen resolution, smaller screen size and shorter battery life than current models. Evaluation of the Realwear device with proposed future software updates, with better image resolution and a fix for noise cancellation, would have allowed a better comparison with the smartphone.

The local information governance and Digital services team did not approve video recording or taking pictures on the device. This limited the full evaluation of devices as teaching aids. Our local digital health services team had concerns about the amount of data that would be generated from multiple device use, saving video clips and where and how long the data would be kept i.e. cloud or local servers.

Network coverage was patchy in certain parts of the hospital leading to poor streaming quality and occasional dropped calls. The local trust was conducting a network survey during the study with plans to install multiple wireless access points in areas of poor network coverage.

Our infection control team raised concerns about the disinfection of chest harnesses however these can be worn under sterile gowns however would still need to be cleaned as per local trust guidelines. We used lanyards as an alternative as these are easier to disinfect after use and less bulky to wear.

Conclusion

The involvement of the local digital services and Information governance teams is essential before implementing virtual streaming of patients and their clinical information. A local agreement at the outset to record as well as live stream would be beneficial for student teaching and medical records.

Further research is needed on how remote viewing of live streams by staff or students off site can be kept secure ensuring no breaches in patient privacy, whether virtual ward rounds will improve ward round efficiency, allow better experiential learning for junior staff performing consultant free ward rounds and procedures, whether the adoption of office and ward-based teams using a smartphone for secure streaming of information in real time can help reduce the risk of nosocomial infection transmission. Further development of more infection friendly harnesses for use with smart phones needs to be developed.

Smarter use of the smartphone during and after COVID -19 pandemic could save not only overall costs for smart technologies but also reduce mortality and morbidity.

However for the benefits to be realised clinician engagement of a new model of remote based management of patients and teaching of students is essential.

References:

1. Monaghesh E, Hajizadeh A. The role of telehealth during COVID-19 outbreak: a systematic review based on current evidence. *BMC Public Health*. 2020;20:1193. Published 2020 Aug 1. doi:10.1186/s12889-020-09301-4.
2. Wilson K. Mobile cell phone technology puts the future of health care in our hands. *CMAJ*. 2018;190:E378-E379. doi:10.1503/cmaj.180269.
3. Ventola CL. Mobile devices and apps for health care professionals: uses and benefits. *P T*. 2014;39:356-64.
4. Wu M, Luo J. Wearable technology applications in healthcare: A literature review. *Online Journal of Nursing Informatics*. 2019. Available from: <https://www.himss.org/resources/wearable-technology-applications-healthcare-literature-review> [Accessed 27 September 2020].
5. Yang Q, Van Stee SK. The Comparative Effectiveness of Mobile Phone Interventions in Improving Health Outcomes: Meta-Analytic Review. *JMIR Mhealth Uhealth*. 2019;7:e11244. Published 2019 Apr 3. doi:10.2196/11244.
6. Gilbert, Anthony William et al. "Rapid implementation of virtual clinics due to COVID-19: report and early evaluation of a quality improvement initiative." *BMJ open quality* vol. 9,2 (2020): e000985. doi:10.1136/bmjopen-2020-000985
7. Chase J. Healthcare Professionals: The iPad and other drugs. *Medical Marketing & Media*. Available from: <https://www.mmm-online.com/home/channel/features/healthcare-professionals-the-ipad-and-other-drugs/> [Accessed 27 September 2020].
8. Burki T. England and Wales see 20 000 excess deaths in care homes. *Lancet*. 2020;395:1602. doi:10.1016/S0140-6736(20)31199-5.

9. Remtulla R. The Present and Future Applications of Technology in Adapting Medical Education Amidst the COVID-19 Pandemic. *JMIR Med Educ.* 2020;6:e20190. Published 2020 Jul 17. doi:10.2196/20190.
10. McCall B. What does the GDPR mean for the medical community? *Lancet.* 2018;391:1249-50. doi:10.1016/S0140-6736(18)30739-6.
11. Simmonds R, Lee D, Hayhurst E. Mobile phones as fomites for potential pathogens in hospitals: microbiome analysis reveals hidden contaminants. *J Hosp Infect.* 2020;104:207-13. doi:10.1016/j.jhin.2019.09.010.
12. Bhardwaj N, Khatri M, Bhardwaj SK, et al. A review on mobile phones as bacterial reservoirs in healthcare environments and potential device decontamination approaches. *Environ Res.* 2020;186:109569. doi:10.1016/j.envres.2020.109569.
13. Malhotra S, Wlodarczyk J, Kuo C, et al. Shining a light on the pathogenicity of health care providers' mobile phones: Use of a novel ultraviolet-C wave disinfection device [published online ahead of print, 2020 Jun 5]. *Am J Infect Control.* 2020;S0196-6553(20)30358-8. doi:10.1016/j.ajic.2020.05.040.
14. Pal P, Roy A, Moore G, et al. Keypad mobile phones are associated with a significant increased risk of microbial contamination compared to touch screen phones. *Journal of Infection Prevention.* 2013;14:65-68. <https://doi.org/10.1177/1757177413475903>.
15. Inagaki H, Saito A, Sugiyama H, et al. Rapid inactivation of SARS-CoV-2 with deep-UV LED irradiation. *Emerg Microbes Infect.* 2020;9:1744-7. doi:10.1080/22221751.2020.1796529.

16. Panigrahi SK, Pathak VK, Kumar MM, et al. Covid-19 and mobile phone hygiene in healthcare settings. *BMJ Glob Health*. 2020;5:e002505. Published 2020 Apr 22.
doi:10.1136/bmjgh-2020-002505.
17. Mark D, Leonard C, Breen H, et al. Mobile phones in clinical practice: reducing the risk of bacterial contamination. *Int J Clin Pract*. 2014;68:1060-4. doi:10.1111/ijcp.12448
18. Luangasanatip N, Hongsuwan M, Limmathurotsakul D, et al. Comparative efficacy of interventions to promote hand hygiene in hospital: systematic review and network meta-analysis. *BMJ*. 2015;351:h3728. Published 2015 Jul 28. doi:10.1136/bmj.h3728.
19. Cheng VC, Chau PH, Lee WM, et al. Hand-touch contact assessment of high-touch and mutual-touch surfaces among healthcare workers, patients, and visitors. *J Hosp Infect*. 2015;90:220-5. doi:10.1016/j.jhin.2014.12.024.
20. Panhotra BR, Saxena AK, Al-Mulhim AS. Contamination of patients' files in intensive care units: an indication of strict handwashing after entering case notes. *Am J Infect Control*. 2005;33:398-401. doi:10.1016/j.ajic.2004.12.009.
21. Teng SO, Lee WS, Ou TY, et al. Bacterial contamination of patients' medical charts in a surgical ward and the intensive care unit: impact on nosocomial infections. *J Microbiol Immunol Infect*. 2009;42:86-91.
22. Wong SCY, Kwong RT, Wu TC, et al. Risk of nosocomial transmission of coronavirus disease 2019: an experience in a general ward setting in Hong Kong. *J Hosp Infect*. 2020;105:119-27. doi:10.1016/j.jhin.2020.03.036.
23. Evans S, Agnew E, Vynnycky E, et al. The impact of testing and infection prevention and control strategies on within-hospital transmission dynamics of COVID-19 in English hospitals. *medRxiv*. 2020 May 20. doi:10.1101/2020.05.12.20095562.
<https://www.medrxiv.org/content/10.1101/2020.05.12.20095562v2>.

24. Meredith LW, Hamilton WL, Warne B, et al. Rapid implementation of SARS-CoV-2 sequencing to investigate cases of health-care associated COVID-19: a prospective genomic surveillance study [published online ahead of print, 2020 Jul 14]. *Lancet Infect Dis.* 2020;S1473-3099(20)30562-4. doi:10.1016/S1473-3099(20)30562-4.
25. Arons MM, Hatfield KM, Reddy SC, et al. Presymptomatic SARS-CoV-2 Infections and Transmission in a Skilled Nursing Facility. *N Engl J Med.* 2020;382:2081-2090. doi:10.1056/NEJMoa2008457.
26. Rivett L, Sridhar S, Sparkes D, et al. Screening of healthcare workers for SARS-CoV-2 highlights the role of asymptomatic carriage in COVID-19 transmission. *Elife.* 2020;9:e58728. Published 2020 May 11. doi:10.7554/eLife.58728.
27. Treibel TA, Manisty C, Burton M, et al. COVID-19: PCR screening of asymptomatic health-care workers at London hospital. *Lancet.* 2020;395:1608-1610. doi:10.1016/S0140-6736(20)31100-4.
28. Croghan SM, Carroll P, Reade S, et al. Robot Assisted Surgical Ward Rounds: Virtually Always There. *J Innov Health Inform.* 2018;25:982. Published 2018 May 2. doi:10.14236/jhi.v25i1.982.
29. Cafferkey JJ, Hampson DOP, Ross C, et al. Using HoloLens™ to reduce staff exposure to aerosol generating procedures during a global pandemic. *medRxiv.* 2020 May 26. doi: 10.1101/2020.05.24.20107193.
<https://www.medrxiv.org/content/10.1101/2020.05.24.20107193v1>.
30. Schwamm LH, Estrada J, Erskine A, et al. Virtual care: new models of caring for our patients and workforce. *Lancet Digit Health.* 2020;2:e282-e285. doi:10.1016/S2589-7500(20)30104-7.

Acknowledgements

I would like to thank Mr Conrad Jaggwe Technical Lead and Facilities officer, Postgraduate Medical Education Dept, London Northwest Healthcare University NHS Trust for his technical assistance and advice on how to set up these devices, the Digital services team for providing the smartphones for evaluation, the Education fellows, Urology trainees and Medical students who assisted in testing and assessing devices during mock and real-life patient interactions. I would also like to thank the Realwear company for providing a device for us to evaluate during the study.

Fig 1

Office Based Team (OBT) place a call through MS Teams to Ward based team (WBT) to begin the virtual clinical encounter

Office based team placing a call to remote based team

Office based team select the window they wish to share with the ward based team

View from office based team PC

WBT can see images or results on their smartphone shared from OBT immediately via MS Teams

CT scan image on smartphone of ward based team - images can be scrolled through in real time

Pathology results shared with the ward based team

Smartphone + Realwear HMT-1 (Ward based team stream to Office based team through MS Teams by toggling camera on)

Smartphone on chest harness

Realware HMT-1 Ward use

Stream from Smartphone - different patient

Stream from Realwear HMT-1

Realwear HMT-1 Theatre use

Trainee in operating theatre performing surgery

Consultant watching live surgery stream via MS Teams remotely.

Able to have 2 way audio and video communication with trainee during the procedure and provide advice if needed

Smartphone teaching simulation (sim man)

Smartphone on a chest harness during a sim man teaching simulation

Realwear HMT-1 teaching simulation

Medical student watching simulation via live stream through MS Teams

Table 1– Device survey, summary of results of smart devices evaluated for virtual patient interaction and teaching simulations, PPE and devices

Devices evaluated during the study											
Device	Wi-Fi enabled	2-way audio	MS Teams	Battery life	Resolution	Hands free	Head worn	Body worn	Screen size	Ease of cleaning	Cost of device
Smart Phone	Yes	Yes	Yes	8-10 hrs	720p, 1080p and 4K	N/A	Yes	Yes	4 in x 10.2 in	Easiest to clean	£300 - £1000
Realware HMT-1 Device	Yes	Yes	Yes	8 hrs	Video up to 1080p	Partly	Yes	No	0.68 in	yes	£2000 (\$2520)
Alternative devices not evaluated as did not meet inclusion criteria											
Go Pro Hero 8	Yes	No	No (unless connected directly to computer)	2-3 hrs	1080p	Yes	Yes	Yes	1.95in (one-way video)		£350
Vuzix blade smart glass	Yes	Yes	No	8 hrs	720p 30fps 1080p 24fps 8 MP	Partly	Yes	N/A	N/A		£900
Microsoft HoloLens 2	Yes	Yes	Yes	3 hrs	1080p	Yes	Yes	N/A	N/A	yes	£3349
Google Glass 2 Enterprise Edition	Yes	Yes	No unless use third part app from Zapier	8 hrs	8MP	Partly	Yes	N/A	N/A		£780 (\$999)
Survey Results for device wearers											
Device Wearer (user)	Number of evaluations (device users)	Most liked feature	Least liked feature	Quality of audio stream between office team and ward team	Video stream quality did not hinder the quality of patient interaction	Likert 1= not recommend	Likert 2= recommend with major changes	Likert 3= recommend with minor changes	Likert 4= recommend without any changes		
Smartphone	10	Small size and compact	Not completely hands free	Excellent	8 agree 2 disagree	n=1 (10%)	n=3 (30%) 2 – Head worn. 1 – Poor connection and small screen.	N=1 (10%)	n =5 (50%)		

Realwear HMT-1	10	Ease of wearing on head	Screen too small to view information clearly	Excellent	N/A	n=0	n = 4 (40%) Audio issues, screen too small, stream resolution	n = 6 (60%) short battery life when streaming continuously device heating up	n=0

Smartphones and body mount survey results of wearers of device and receivers of audio and video live stream

Smartphone Mount used	Number of evaluations of wearer of device	Most liked feature	Least liked feature	Quality of audio stream from ward / theatre team to office team		1= not recommend	2= recommend with major changes	3 = recommend with minor changes	4 = recommend without any changes
Smartphone and head mount	2	-	Discomfort when on head	Good		2	-	-	-
Smartphone and chest mount	6	-	Movement of phone	Good		0	1	1	4
Smartphone and lanyard	2	-	Not completely hands free	Good		-	0	1	1

Office based team / medical student evaluation of quality of audio and video feed from remote ward team / staff in simulation

	Quality of audio feed good / excellent	Quality of image on video stream good / excellent	Device video stream quality allowed safe communication	Device mount whilst being used Stable image from	Recommend device for routine use	P- value
Smart Phone	9 yes, 1 no, 1 undecided, 4 no response	9 yes, 3 no, 3 no response	9 yes, 2 no, 4 no response	9 yes, 2 no, 4 no response	10 yes, 1 no, 4 un decided	0.30
Realwear HMT-1 device	7 yes, 3 no, 4 no response	7 yes, 5 no, 2 no response	6 yes, 4 undecided, 4 no response	10 yes, 1 no, 3 no response	7 yes, 6 undecided, 1 no	0.41

Patient questionnaire responses

Numbers of patients answering questionnaire	Patient prefer consultant to be physically present rather than virtual	Did you feel your privacy was maintained during streaming	Would you be filmed again in virtual ward round
4	Yes = 0, 4 = no	Yes = 4, no = 0	4 = yes, 0 = no

Devices use with PPE

Device	Facemask ± goggles	Facemask and visor	Advantages	Disadvantages
yes	yes	Mainly hands free	Difficult to wear with visor and goggles

yes	yes	Mainly hands free	Difficult to wear with visor
Head worn mounts and devices with PPE, not used during study due to lack of availability or not meeting inclusion criteria				
GoPro Hero 8	yes	yes	Easy to clean after use	-
Microsoft HoloLens 2	yes	yes	Can be worn with mask. Tailor-made visor developed for use with device. Remain sterile	Device has to be disassembled for cleaning after each use, time consuming
Google Glass 2 Enterprise	-	-	Can be worn easily under visor and with face mask	Not assessed
Chest worn Lanyard with mobile device included in the study				
Device	iPhone on lanyard	Long-sleeve apron	Advantages	Disadvantages
Smartphone			Can be cleaned easily after use	-
Chest worn harness of devices included in the study with PPE				
Device	Surgical gown (Not for use in theatre)	Harness under long-sleeve apron	Advantages	Disadvantages
Smartphone			Lightweight can be cleaned as per infection control	Chest harness has to be cleaned / disinfected after each use
Alternative to phone / realware HMT-1				
Computer on wheels			-	Difficult to keep clean as multiple users