Race/Ethnic Disparities in Mild Cognitive Impairment and Dementia: The Northern Manhattan Study

(title character count=92)

Clinton B. Wright, MD, MS¹, Janet T. DeRosa, MPH², Michelle P. Moon, MS², Kevin Strobino, MPH², Charles DeCarli, MD³, Ying Kuen Cheung, PhD⁴, Stephanie Assuras, PhD², Bonnie Levin, PhD^{5,6}, Yaakov Stern PhD², Xiaoyan Sun, MD, PhD⁶, Tatjana Rundek, MD, PhD^{6,7}, Mitchell S.V. Elkind, MD, MS², Ralph L. Sacco, MD, MS^{5,6}

¹ National Institute of Neurological Disorders and Stroke, national Institutes of Health, Bethesda, MD

² Department of Neurology, Vagelos College of Physicians & Surgeons, Columbia University, New York, NY

³ Department of Neurology, University of California at Davis, Davis, CA

⁴ Department of Biostatistics, Mailman School of Public Health, Columbia University, New York, NY

⁵ Evelyn F McKnight Brain Institute, University of Miami Miller School of Medicine, Miami, FL

⁶ Department of Neurology, University of Miami Miller School of Medicine, Miami, FL

Corresponding author:

Clinton Wright, MD, 6001 Executive Boulevard, Bethesda, MD 20852

Tel: 301-402-1660

ABSTRACT

OBJECTIVE: Estimate the prevalence of mild cognitive impairment (MCI) and probable dementia in the racially and ethnically diverse community-based Northern Manhattan Study cohort and examine sociodemographic, vascular risk factor, and brain imaging correlates.

METHODS: Cases of MCI and probable dementia were adjudicated by a team of neuropsychologists and neurologists and prevalence was estimated across race/ethnic groups. Ordinal proportional odds models were used to estimate race/ethnic differences in prevalence rates for MCI or probable dementia adjusting for sociodemographic variables (model 1), model 1 plus potentially modifiable vascular risk factors (model 2), and model 1 plus structural imaging markers of brain integrity (model 3).

RESULTS: There were 989 participants with cognitive outcome determinations (mean age 69 ± 9 years; 68% Hispanic, 16% Black, 14% White; 62% women; mean ($\pm$ SD) follow-up five (± 0.6) years). Prevalence rates for MCI (20%) and probable dementia (5%) were significantly different by race/ethnicity even after accounting for age and education difference across race-ethnic groups; Hispanic and Black participants had greater prevalence rates than Whites. Adjusting for sociodemographic and brain imaging factors explained the most variance in the race/ethnicity associations. White matter hyperintensity burden explained much of the disparity between Black and White, but not between Hispanic and White, participants.

CONCLUSIONS: In this diverse community-based cohort, white matter hyperintensity burden partially explained disparities in MCI and dementia prevalence in Black but not Hispanic participants compared to Whites. Longer follow-up and incidence data are needed to further clarify these relationships.

KEY WORDS: dementia, mild cognitive impairment, cohort studies, Hispanic American, African American

INTRODUCTION

Variability in dementia rates across racial and ethnic groups has been estimated at 60%, underscoring the importance of work to understand ethnoracial disparities.¹ Improving our understanding and reducing rates of cognitive impairment and dementia disparities is a major goal of the National Alzheimer's Plan (NAPA). Understanding the factors underlying these disparities, including the importance of modifiable risk factors and susceptibility to pathological processes and resistance to them, is a priority to appropriately target interventional strategies.

Research in diverse cohorts plays an important role in clarifying race/ethnic differences, and disparities in the risk of mild cognitive impairment (MCI) and dementia have been reported for Black people in multiple studies over the past decades,¹⁻⁴ but data on Hispanics is more limited. Diverse cohorts that include multiple ethnic and racial groups with deep phenotyping of key behavioral and modifiable risk factor exposures, intermediate markers of brain integrity, and cognitive outcomes are needed to improve our understanding of disparities, especially in vascular cognitive impairment and dementia (VCID) risk. White matter hyperintensities, subclinical brain infarcts, and brain atrophy measures detected by MRI can be particularly helpful in this regard. We examined ethnoracial disparities in prevalence rates of mild cognitive impairment (MCI) and dementia in a diverse population-based cohort of Hispanic, Black, and White people living in the Northern Manhattan community of New York City.

MATERIAL AND METHODS

Study Population

The NOMAS sample enrolled adults aged 40 and older (range 40-94) who had resided in Northern Manhattan for more than three months at the time of recruitment and had never been diagnosed with a stroke¹⁸. Participants were recruited between 1993-2001 using random-digit dialing to participate in a baseline interview and assessment (enrollment response rate was 75%), with ongoing annual telephone and in-person follow-up (N=3,298; loss to follow up <5%). In 2003 the MRI sub-study began recruitment

during annual telephone follow-up and included NOMAS participants who were clinically stroke-free, age 50 and older, and had no contraindications to MRI. An additional 199 household members were recruited to yield a final sample size of 1,290 by 2008. The study is approved by the IRBs of Columbia University Medical Center and the University of Miami School of Medicine and all subjects provided written informed consent.

Baseline Evaluation

Participants had a thorough evaluation of vascular risk factors and medical history at the time of enrollment, including a physical/neurological examination by study physicians. Race/ethnicity was defined by self-identification using a series of questions modeled after the US census and conforming to standard definitions outlined by Directive 15. Years of educational attainment, including degree achieved, were self-reported. Standardized questions were adapted from the CDC Behavioral Risk Factor Surveillance System regarding history of hypertension, diabetes mellitus, smoking, and cardiovascular conditions, including congestive heart failure, angina, coronary artery disease, atrial fibrillation, valvular heart disease, and peripheral vascular disease^{20,21}. Data on medication use was collected. Smoking was categorized as current (within the past year), former, or never smoker of cigarettes, cigars, or pipes. Leisure-time physical activity was assessed by self-report using a questionnaire adapted from the National Health Interview Survey and moderate to heavy physical activity was defined as participation in at least one of several rigorous physical activities in a typical 14-day period.²² Moderate alcohol use was defined as current drinking of one drink per month up to two drinks per day. Fasting blood specimens were analyzed at the Core Laboratory of the Irving Center for Clinical Research to determine blood glucose and lipid levels, including total cholesterol and HDL. Waist and hip circumferences were measured in inches with a flexible tape measure while participants were standing and wearing no heavy outer garments. Waist circumference (WC) was measured at the level of the umbilicus, and hip circumference was measured at the level of the bilateral greater trochanters, as previously described.⁵

Systolic and diastolic blood pressures were calculated by averaging two measurements (before and after the physical examination) from the right brachial artery after a 10-minute rest in a seated position (Dinamap Pro100, Critikon Inc).

Brain MRI

Imaging was performed on a 1.5-Tesla MRI system (Philips Medical Systems, Best, The Netherlands) at the Columbia University Hatch Research Center. To segment white matter hyperintensities, semiautomated measurements of pixel distributions using mathematical modeling of pixel-intensity histograms for cerebrospinal fluid and brain white and gray matter were used to identify the optimal pixel-intensity threshold to distinguish cerebrospinal fluid from brain matter, using a custom-designed image analysis package (QUANTA 6.2 using a Sun Microsystems Ultra 5 workstation).⁶ For subclinical brain infarct readings, methods to identify and classify MRI-defined subclinical infarcts (SBI) have been published.⁷ Two independent raters used a superimposed image of the subtraction, fluid attenuated inversion recovery (FLAIR), proton density, and T2-weighted images at 3× magnified view for interpretation of lesion characteristics. Agreement among raters has been good (published kappa values, 0.73–0.90).⁸ Processing of MRI scans to calculate total intracranial volumes (ICV), total cerebral volumes, and white matter hyperintensity volumes (WMHV), has been previously described.⁹ To correct for head size, WMHV was calculated as percent total ICV and log-transformed to a normal distribution (log-WMHV). Proportion of total cerebral volume to total ICV was examined as brain parenchymal fraction (BPF). Subclinical infarcts (SBI) were rated as present or absent.

Cognitive and Functional Assessment

Participants in the MRI sub-study were recruited between 2003 to 2008 and assessed in-person (defined as visit 1) and were interviewed by a trained research assistant, who administered structured questionnaires and a neuropsychological (NP) battery. All tests were administered in English or Spanish

based on participant preference. The Mini Mental State Examination and Cognitive Failures Questionnaire were administered at the time of the NP battery in a designated quiet room. Most testing was on the day of MRI.^{10, 11} Episodic memory was measured using three sub-scores derived from a 12-word five trial list-learning task: list learning total score, delayed recall score, and delayed recognition score. Executive function was assessed with two sub-scores: the difference in time to complete the Color Trails test Form 1 and Form 2, and the sum of the Odd-Man-Out subtests 2 and 4.¹² Processing speed was assessed with the Grooved Pegboard task non-dominant hand time and the Color Trails test Form 1.^{13, 14} Working memory was assessed with the Digit Ordering and Letter Number Sequencing tests.^{15, 16} Semantic memory was measured using three tests: picture naming (modified Boston Naming), category fluency (Animal Naming) and phonemic fluency (C, F, L in English speakers and P, S, V in Spanish speakers).^{17, 18} The Visual-Motor Integration test and the Symbol Digit Modalities tests were not part of the domains but were also available for review.^{13, 19} At the initial visit we estimated premorbid intelligence and literacy with the Peabody Picture Vocabulary Test (Test de Vocabulario en Imagenes Peabody for Spanish speakers), the Wide Range Achievement Test (English speakers) and the Word Accentuation Test (Spanish speakers).²⁰⁻²² Depressive symptoms were quantified with the Center for Epidemiological Studies Depression scale (CES-D), a 20-item scale assessing depressive affect, somatic complaints, positive affect, and interpersonal relations.²³ Research assistants interviewed participants to assess cognitive functional status.

A mean of 5.0±0.6 years after the initial neuropsychological assessment, a second in-person assessment was conducted between 2008 and 2015 (visit 2). At the second visit, the neuropsychological battery was repeated with the addition of the Letter Number Sequencing and Symbol Digit Modalities tests, and the Informant Questionnaire of Cognitive Decline in the Elderly (IQCODE) was administered to a family member or friend of the participant as close in time to the neuropsychological assessment as possible.²⁵ The relationship of the informant to the participant was documented and aided adjudication.

Ascertainment of mild cognitive impairment and probable dementia

In 2015 NOMAS entered its fifth consecutive grant cycle and for the first time was funded to ascertain MCI and dementia status. A team of neuropsychologists and neurologists with dementia expertise used established criteria for case ascertainment.²⁶⁻²⁸ Each pair of adjudicators reviewed data from both visit 1 and visit 2 and completed a visit 2 grading form documenting cognitive status as normal, MCI (with notation of amnestic and non-amnestic sub-types), probable dementia, other psychiatric disorder, or unable to classify.

To assess cognitive performance, we used individual neuropsychological test scores and cognitive domain-specific scores. Cognitive domains (episodic memory, semantic memory, processing speed and executive function) were created based on an exploratory factor analysis and prior findings. Because NOMAS participants have low educational attainment, literacy, and socio-economic status, and there are limited established norms that consider these factors, we constructed NOMAS-specific norms using neuropsychological test scores from visit 1. The normative values for each NP test were calculated from age (50-60, 61-70, 71-80 and >80 years old) and education (0-6, 7-12 and >12 years of education) group-specific means and standard deviations. Each NP test score was standardized against its normative values, and each cognitive domain Z score was obtained by taking the average of the standardized NP test scores for the available component tests in a given domain.

Adjudicators accessed a Redcap web portal that displayed demographic information, visit eyesight and hearing status, neuropsychological domain and literacy Z-scores, dementia rating, MMSE and CES-D scores, most recent medication list and prior history of antidepressant and other psychiatric medications, psychiatric history, cognitive failures questionnaire, IQCODE score (cutoff for probable dementia >3.6), and stroke history between visits (if applicable).¹¹ Downloadable case-specific test forms were available for review to aid in assessments. Adjudicators were randomly assigned cases blinded to

each other's ratings based on an algorithm that identified participants with possible cognitive impairment. We derived an algorithm to segregate cases who had neuropsychological testing at Visit 2 into one of two pools: 1) those requiring adjudication, and 2) those rated as cognitively normal based on an algorithm. The criteria for consensus adjudication were:

- Missing at least one of the 11 NP battery tests; or
- Two or more age and education normalized NP test Z-scores < -1.5 ; or
- One or more domain Z-scores < -1.5 .

All available information pertaining to medical history, informant reports, self-rated questionnaires, educational attainment, literacy, notes from the participant interview at the visit, and other data were used for case ascertainment, but rule-based criteria helped guide cognitive status determination as follows:

- Normal: no cognitive domain Z-scores below -1.5 , no evidence of dementia on the participant interview or IQCODE (score < 3.6).
- MCI: any domain Z-score below -1.5 and no evidence of dementia on the IQCODE (score < 3.6).
- Probable dementia: any domain Z-score below -1.5 and an IQCODE score 3.6 or greater.

Discordant ratings were resolved by consensus of the neuropsychologist/neurologist adjudication pair, and disagreements were resolved through interdisciplinary Dementia Consensus Committee (DCC) reviews.

Statistical Analysis

Participant characteristics were compared across race/ethnicities. Characteristics between participants who only attended visit 1 and those who returned for visit 2 were also compared. We used ordinal proportional odds models to examine race/ethnic differences in prevalence using a three-level ordinal dependent variable of normal, MCI, and probable dementia, and difference of association across the

three ethnic groups collectively was tested using a likelihood ratio test with 2 degrees of freedom (LRT 2 df). Less than 2% of the sample self-identified as other race (Asian or Pacific Islander, American Indian or Alaskan Native) and were excluded from this analysis. Hispanic and non-Hispanic Black (referred to as Black) were entered into the model (reference=White). Model 1 was adjusted for age, sex, years of educational attainment, medical insurance status (proxy for income), and literacy; model 2 was adjusted for variables in model 1 plus systolic and diastolic blood pressure, antihypertensive medication use, diabetes mellitus, smoking status, moderate alcohol use, physical activity, low and high density lipoprotein levels, and waist-to-hip ratio; model 3 was adjusted for variables in model 1 plus brain parenchymal fraction, log-WMHV, and SBI status. We tested for potential effect modifiers of race and ethnicity by entering multiplicative terms for significant covariates into the model. We performed a sensitivity analysis among participants with normal cognition at visit 1 (MMSE >26) to limit confounding by pre-morbid cognitive problems. Analyses were conducted using SAS 9.4 (SAS Institute, Cary, NC).

RESULTS

Characteristics of the study sample are shown in Table 1 (mean age=69, IQR 63-74; 68% Hispanic, 16% Black, 14% White, 2% other race). Compared to participants who only attended visit 1, those who returned for visit 2 (flow across visits shown in the Figure) were younger (OR=0.96, 95%CI=0.93 to 0.99), more likely to be women (OR=1.6, 95% CI 1.0 to 2.3), had lower systolic (OR=0.7, 95% CI=0.6 to 0.9) and higher diastolic (OR=1.3, 95% CI=1.1 to 1.6) blood pressures (BP), and had better global cognitive domain scores at visit 1 (OR=1.8 per SD, 95% CI=1.4 to 2.5; Supplementary Table 1).

We ascertained the cognitive status of 989 NOMAS participants who attended visit 2. One hundred ninety-nine participants were adjudicated as having MCI (20%), 45 as having probable dementia (5%), 8 as Other or Unable to Classify (<1%), and 737 participants were adjudicated or identified by algorithm as having no cognitive impairment (74%). The breakdown of cognitive status by race/ethnicity is shown in

Table 2. The ordinal odds of MCI or probable dementia was significantly greater for Black and Hispanic participants as a group compared to Whites adjusting for sociodemographic variables (LRT 2df, $p < 0.0001$; Table 3). Table 3 shows that each race/ethnicity had significantly greater odds of MCI or probable dementia than Whites (two-fold for Blacks and four-fold for Hispanics). Adjusting further for vascular risk factors in model 2, the association across race/ethnicity remained significant (LRT 2df, $p < 0.0001$), but the effect was slightly attenuated by about 6% for Black participants and was unchanged for Hispanics (Table 3). No individual risk factor was significantly associated with the odds of MCI or probable dementia in this model (data not shown). When we adjusted for sociodemographic variables and brain imaging markers in model 3, the odds of MCI or probable dementia remained significantly greater for Black and Hispanic participants than Whites (LRT 2df, $p < 0.0001$), but the association was attenuated by almost 25% for Blacks compared to model 1 (OR=1.7, 95% CI=0.9 to 3.32) and again did not change for Hispanics (OR=4.2, 95% CI=2.0 to 8.8). In this model, each unit greater log-WMHV burden was associated with 1.3 times greater odds of MCI or probable dementia (OR=1.3, 95% CI=1.1 to 1.6), but neither smaller BPF nor SBI status was significantly associated with odds of MCI or probable dementia in model 3. There was no significant interaction between Black race and log-WMHV (p for interaction 0.473). Restricting the primary analyses to NOMAS participants with normal MMSE scores (27 or greater) at visit 1, we found similar results (Supplemental Table 2).

DISCUSSION

This analysis of MCI and dementia prevalence in NOMAS represents the first estimate in the sample since expert adjudication of cognitive status was initiated in 2016. We found prevalence rates of MCI and dementia similar to a number of other cohorts that have examined Hispanic, Black, and White people.²⁹⁻³¹ For dementia the relatively low prevalence of 5% is similar to other cohorts that included people below the age of 65, and studies have generally reported that the proportion of people with dementia rises dramatically with age.⁴ We found marked race/ethnic differences in this study. Black

participants were twice as likely as White participants, and Hispanic participants more than four times as likely than White participants, and twice as likely as Black participants, to have MCI or dementia adjusting for sociodemographic factors.

Racial and ethnic disparities in dementia risk have been known for many years and are well supported by studies in different groups, including in a sample of Medicare participants in Northern Manhattan.³² However, more recent data reported temporal trends showing sharp reductions in dementia rates across race/ethnic groups with greater declines for non-Hispanic Blacks and Whites.³³ The biracial Atherosclerosis Risk in Communities (ARIC) Neurocognitive Study also found a greater prevalence of dementia, but not MCI, in Blacks compared to Whites.²⁹ Few studies have included Hispanics to be able to compare MCI and dementia rates across racial and ethnic groups.⁴ Continued follow-up in NOMAS will provide incidence data in a sample that included people as young as 50 at the time of the first evaluation of MCI and dementia status (visit 1).

Given the importance of midlife as opposed to late life vascular risk factors in the development of VCID, opportunities to study these relationships in diverse cohorts are needed. Especially those where participants live in the same community, allowing comparison across groups without confounding introduced by different environments and heterogeneity in other local factors. The greater burden of some vascular risk factors that has been found in studies that have included Blacks or Hispanics might be expected to explain some of these disparities. Interestingly, in this cross-sectional analysis, adjusting for potentially modifiable vascular risk factors did not explain much of the variance in the association of race/ethnicity with MCI and dementia status, although the association for Blacks was attenuated slightly. Not all studies have confirmed link with vascular risk factor exposures. In the ARIC study, Blacks were at elevated risk of dementia compared to Whites but midlife risk factor burden did not play a key role.³ Similarly, in the Health, Aging, and Body Composition study, Blacks were at elevated risk of

dementia compared to Whites and adjusting for vascular and other comorbidities did not attenuate this effect.³⁴ Among Black and Caribbean Hispanic Medicare recipients in Northern Manhattan, diabetes contributed significantly to MCI and dementia risk.^{35, 36} Diabetes was also a risk factor for MCI in the Hispanic Communities Health Study, which includes Caribbean Hispanic participants such as in NOMAS and other groups.³⁷ In NOMAS, we did not find that diabetes was a significant contributor to the odds of MCI or dementia prevalence, but continued follow-up is needed. Midlife hypertension has also been associated with MCI and dementia risk in diverse cohorts, but the risk for dementia was greater for White than Black participants in a biracial study.^{38, 39} There are a number of reasons why midlife risk factors would not explain later cognitive outcomes, including variability in duration of exposure and competing risk of death.

Brain imaging studies can help clarify disparities in the downstream effects of exposure to behavioral and vascular risk factors as well as capture evidence of neurodegeneration. White matter lesion burden has been associated with conversion from normal to MCI and rate of decline from MCI to dementia.^{40, 41} This is of particular interest because WMH are related to cerebral small vessel disease, providing an opportunity for prevention through risk factor control especially hypertension.⁴² Both Black and Hispanic participants have been found to have a greater burden of WMH in several studies, and WMH burden was related to ideal cardiovascular health in NOMAS.^{43, 44} In a study among Mexican Americans both hippocampal volume and WMH burden were independent predictors of dementia.⁴⁵ In keeping with this, the current analysis found that WMH burden explained the disparity in odds of MCI and dementia between Black and non-Hispanic Whites. However, sociodemographic variables, vascular risk factors, and brain imaging markers did not explain the disparity between Hispanic and White participants. The reason for this is not certain but degree of risk factor control and length of exposure are not captured by risk factor adjustment and could be of explanatory value.

Strengths of this study include the well-phenotyped diverse cohort that includes behavioral and risk factor data as well as imaging markers of brain integrity. Survival of participants to participate in the MRI sub-cohort yielded a younger, healthier, sample with better cognitive performance than the original population-based NOMAS sample. Given the likelihood of differential dropout due to mortality leading up to the waves of cognitive assessments it is likely that some bias was introduced. We did not conduct analyses of the competing risk of mortality for this study. Prospective data collection is underway and will allow the current prevalence estimates to establish a baseline for incidence studies where competing risk models will be of value. As with all cross-sectional observational studies, unmeasured confounding of risk factors for cognitive and brain health are another potential source of bias.

In conclusion, data from this racially and ethnically diverse cohort study show that Black and Hispanic people were more likely than Whites to have MCI or dementia at the second visit even if they had no more than mild cognitive problems at the first. These disparities in MCI and dementia prevalence were independent of sociodemographic and vascular risk factors, but WMH burden to a large extent explained the disparity between Black and White participants. Prospective data from NOMAS will help clarify these findings in the future.

ACKNOWLEDGEMENTS

This study was supported by the National Institute of Neurological Disorders and Stroke (grant NS 29993)

DISCLOSURES

Clinton Wright reports royalties from two chapters on vascular dementia in UpToDate.com.

Janet DeRosa is supported by NINDS grant NS29993 and reports no other relevant disclosures.

Stephanie Assuras reports no relevant disclosures.

Michelle Moon is supported by NINDS grant NS29993 and reports no other relevant disclosures.

Kevin Strobino is supported by NINDS grant NS29993 and reports no other relevant disclosures.

Charles DeCarli is a consultant to Novartis for a safety study of heart failure treatment.

Ying Kuen Cheung is supported by NINDS grant NS29993 and reports no other relevant disclosures.

Xiaoyan Sun is supported by NINDS grant NS29993 and reports no relevant disclosures.

Bonnie Levin is supported by NINDS grant NS29993 and reports no relevant disclosures.

Yaakov Stern is supported by NINDS grant NS29993 and consults for Eisai, Lilly, and Arcadia.

Tatjana Rundek is supported by NINDS grant NS29993, NIA, and NCATS.

Mitchell Elkind is supported by NINDS grant NS29993 and reports royalties from chapters on stroke in

UpToDate.com.

Ralph Sacco is supported by NINDS grant NS29993 and by NCATS for a Clinical Translational Science Award.

REFERENCES

1. Mayeda ER, Glymour MM, Quesenberry CP, Whitmer RA. Inequalities in Dementia Incidence between Six Racial and Ethnic Groups over 14 Years. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2016;12:216-224
2. Gurland BJ, Wilder DE, Lantigua R, Stern Y, Chen J, Killeffer EHP, et al. Rates of Dementia in Three Ethnoracial Groups. *International Journal of Geriatric Psychiatry*. 1999;14:481-493
3. Gottesman RF, Albert MS, Alonso A, Coker LH, Coresh J, Davis SM, et al. Associations between Midlife Vascular Risk Factors and 25-Year Incident Dementia in the Atherosclerosis Risk in Communities (Aric) Cohort. *JAMA neurology*. 2017;74:1246-1254
4. Mehta KM, Yeo GW. Systematic Review of Dementia Prevalence and Incidence in United States Race/Ethnic Populations. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2017;13:72-83
5. Caunca MR, Gardener H, Simonetto M, Cheung YK, Alperin N, Yoshita M, et al. Measures of Obesity Are Associated with Mri Markers of Brain Aging: The Northern Manhattan Study. *Neurology*. 2019;93:e791-e803
6. DeCarli C, Murphy DG, Teichberg D, Campbell G, Sobering GS. Local Histogram Correction of Mri Spatially Dependent Image Pixel Intensity Nonuniformity. *Journal of magnetic resonance imaging : JMRI*. 1996;6:519-528
7. Prabhakaran S, Wright CB, Yoshita M, Delapaz R, Brown T, DeCarli C, et al. Prevalence and Determinants of Subclinical Brain Infarction: The Northern Manhattan Study. *Neurology*. 2008;70:425-430
8. DeCarli C, Fletcher E, Ramey V, Harvey D, Jagust WJ. Anatomical Mapping of White Matter Hyperintensities (Wmh): Exploring the Relationships between Periventricular Wmh, Deep Wmh, and Total Wmh Burden. *Stroke*. 2005;36:50-55

9. Dong C, Nabizadeh N, Caunca M, Cheung YK, Rundek T, Elkind MS, et al. Cognitive Correlates of White Matter Lesion Load and Brain Atrophy: The Northern Manhattan Study. *Neurology*. 2015;85:441-449
10. Folstein MF, Folstein SE, McHugh PR. "Mini-Mental State". A Practical Method for Grading the Cognitive State of Patients for the Clinician. *Journal of psychiatric research*. 1975;12:189-198
11. Broadbent DE, Cooper PF, FitzGerald P, Parkes KR. The Cognitive Failures Questionnaire (Cfq) and Its Correlates. *Br J Clin Psychol*. 1982;21:1-16
12. D'Elia L, Satz P, Uchiyama C, T W. *Color Trails Test Professional Manual*. Odessa: Psychological Assessment Resources; 1994.
13. Beery K. *The Vmi, Developmental Test of Visual Motor Integration*. Modern Curriculum Press, Cleveland, OH; 1989.
14. Ruff RM, Parker SB. Gender- and Age-Specific Changes in Motor Speed and Eye-Hand Coordination in Adults: Normative Values for the Finger Tapping and Grooved Pegboard Tests. *Percept Mot Skills*. 1993;76:1219-1230
15. Weschler D. *Wechsler Adult Intelligence Scale - Third Edition*. Psychological Corporation; 1997.
16. Hoppe CD, Muller UD, Werheid KD, Thone AD, von Cramon YD. Digit Ordering Test: Clinical, Psychometric, and Experimental Evaluation of a Verbal Working Memory Test. *Clin Neuropsychol*. 2000;14:38-55
17. Goodglass H, Kaplan E. *The Assessment of Aphasia and Related Disorders*. Philadelphia: Lea & Febiger; 1983.
18. Benton A HK, Sivan A. . *Manual for the Multilingual Aphasia Examination*. 3rd Ed., Iowa City: AJA Associates, Inc.; 1994.
19. Smith A. *Symbol Digit Modalities Test: Manual*.: Western Psychological Services; 1982.
20. Dunn L DD. *Peabody Picture Vocabulary Test*. Circle Pines: American Guidance Service; 1997.

21. Stone M, Jastak S, Wilkinson G. . *Wide Range Achievement Test*. Rasch Measurement Transactions. 1995;8:403.; 1995.
22. Del Ser T, Gonzalez-Montalvo JI, Martinez-Espinosa S, Delgado-Villapalos C, Bermejo F. Estimation of Premorbid Intelligence in Spanish People with the Word Accentuation Test and Its Application to the Diagnosis of Dementia. *Brain Cogn*. 1997;33:343-356
23. Radloff LS. The Ces-D Scale: A Self-Report Depression Scale for Research in the General Population. *Applied Psychological Measurement*. 1977;1:385-401
24. Morris JC. The Clinical Dementia Rating (Cdr): Current Version and Scoring Rules. *Neurology*. 1993;43:2412-a-
25. Jorm AF. A Short Form of the Informant Questionnaire on Cognitive Decline in the Elderly (Iqcode): Development and Cross-Validation. *Psychol Med*. 1994;24:145-153
26. Albert MS, DeKosky ST, Dickson D, Dubois B, Feldman HH, Fox NC, et al. The Diagnosis of Mild Cognitive Impairment Due to Alzheimer's Disease: Recommendations from the National Institute on Aging-Alzheimer's Association Workgroups on Diagnostic Guidelines for Alzheimer's Disease. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. 2011;7:270-279
27. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Jr., Kawas CH, et al. The Diagnosis of Dementia Due to Alzheimer's Disease: Recommendations from the National Institute on Aging-Alzheimer's Association Workgroups on Diagnostic Guidelines for Alzheimer's Disease. *Alzheimers Dement*. 2011;7:263-269
28. American Psychiatric Association. *Dsm-5 Task Force. Diagnostic and Statistical Manual of Mental Disorders: Dsm-5*. . Washington, DC; 2013.
29. Knopman DS, Gottesman RF, Sharrett AR, Wruck LM, Windham BG, Coker L, et al. Mild Cognitive Impairment and Dementia Prevalence: The Atherosclerosis Risk in Communities Neurocognitive Study (Aric-Ncs). *Alzheimers Dement (Amst)*. 2016;2:1-11

30. Petersen RC, Roberts RO, Knopman DS, Geda YE, Cha RH, Pankratz VS, et al. Prevalence of Mild Cognitive Impairment Is Higher in Men. The Mayo Clinic Study of Aging. *Neurology*. 2010;75:889-897
31. Lopez OL, Jagust WJ, DeKosky ST, Becker JT, Fitzpatrick A, Dulberg C, et al. Prevalence and Classification of Mild Cognitive Impairment in the Cardiovascular Health Study Cognition Study: Part 1. *Archives of neurology*. 2003;60:1385-1389
32. Gurland BJ, Wilder DE, Lantigua R, Stern Y, Chen J, Killeffer EH, et al. Rates of Dementia in Three Ethnoracial Groups. *Int J Geriatr Psychiatry*. 1999;14:481-493
33. Noble JM, Schupf N, Manly JJ, Andrews H, Tang MX, Mayeux R. Secular Trends in the Incidence of Dementia in a Multi-Ethnic Community. *Journal of Alzheimer's disease : JAD*. 2017;60:1065-1075
34. Yaffe K, Falvey C, Harris TB, Newman A, Satterfield S, Koster A, et al. Effect of Socioeconomic Disparities on Incidence of Dementia among Biracial Older Adults: Prospective Study. *BMJ : British Medical Journal*. 2013;347:f7051
35. Luchsinger JA, Reitz C, Patel B, Tang MX, Manly JJ, Mayeux R. Relation of Diabetes to Mild Cognitive Impairment. *Arch Neurol*. 2007;64:570-575
36. Cheng D, Noble J, Tang MX, Schupf N, Mayeux R, Luchsinger JA. Type 2 Diabetes and Late-Onset Alzheimer's Disease. *Dementia and geriatric cognitive disorders*. 2011;31:424-430
37. Gonzalez HM, Tarraf W, Gonzalez KA, Fornage M, Zeng D, Gallo LC, et al. Diabetes, Cognitive Decline, and Mild Cognitive Impairment among Diverse Hispanics/Latinos: Study of Latinos- Investigation of Neurocognitive Aging Results (Hchs/Sol). *Diabetes Care*. 2020;43:1111-1117
38. Walker KA, Sharrett AR, Wu A, Schneider ALC, Albert M, Lutsey PL, et al. Association of Midlife to Late-Life Blood Pressure Patterns with Incident Dementia. *Jama*. 2019;322:535-545

39. Reitz C, Tang MX, Manly J, Mayeux R, Luchsinger JA. Hypertension and the Risk of Mild Cognitive Impairment. *Arch Neurol*. 2007;64:1734-1740
40. Bangen KJ, Preis SR, Delano-Wood L, Wolf PA, Libon DJ, Bondi MW, et al. Baseline White Matter Hyperintensities and Hippocampal Volume Are Associated with Conversion from Normal Cognition to Mild Cognitive Impairment in the Framingham Offspring Study. *Alzheimer disease and associated disorders*. 2018;32:50-56
41. Tosto G, Zimmerman ME, Carmichael OT, Brickman AM, Alzheimer's Disease Neuroimaging I. Predicting Aggressive Decline in Mild Cognitive Impairment: The Importance of White Matter Hyperintensities. *JAMA neurology*. 2014;71:872-877
42. Group SMIftSR, Nasrallah IM, Pajewski NM, Auchus AP, Chelune G, Cheung AK, et al. Association of Intensive Vs Standard Blood Pressure Control with Cerebral White Matter Lesions. *Jama*. 2019;322:524-534
43. Brickman AM, Schupf N, Manly JJ, Luchsinger JA, Andrews H, Tang MX, et al. Brain Morphology in Older African Americans, Caribbean Hispanics, and Whites from Northern Manhattan. *Arch Neurol*. 2008;65:1053-1061
44. Gardener H, Caunca M, Dong C, Cheung Ying K, Alperin N, Rundek T, et al. Ideal Cardiovascular Health and Biomarkers of Subclinical Brain Aging: The Northern Manhattan Study. *Journal of the American Heart Association*. 2018;7:e009544
45. Wu CC, Mungas D, Petkov CI, Eberling JL, Zrelak PA, Buonocore MH, et al. Brain Structure and Cognition in a Community Sample of Elderly Latinos. *Neurology*. 2002;59:383-391

Table 1: Participant characteristics

	Total (N=989)	Hispanic (N=672)	Black (N=162)	White (N=134)	p-value*
Age, years	69 ± 9	68 ± 8	72 ± 9	72 ± 9	<.0001
Years of Education	10 ± 5	8 ± 5	13 ± 3	15 ± 3	<.0001
Health Insurance Status					
Medicaid or no insurance (%)	484 (49)	434 (65)	35 (22)	12 (9)	<.0001
Medicare or private insurance	505 (51)	238 (35)	127 (78)	122 (91)	
Women	610 (62)	425 (63)	111 (69)	67 (50)	0.002
Current Smoker	155 (16)	97 (14)	39 (24)	18 (13)	0.003
Moderate or Heavy Physical Activity	97 (10)	45 (7)	19 (12)	28 (21)	<.0001
Diabetes mellitus	267 (27)	207 (31)	42 (26)	15 (11)	<.0001
Systolic BP, mm Hg	136 ± 17	136 ± 17	139 ± 18	132 ± 17	0.0003
Diastolic BP, mm Hg	79 ± 9	79 ± 9	80 ± 10	76 ± 10	0.0032
Anti-hypertensive use	590 (60)	414 (62)	109 (67)	60 (45)	<.0001
HDL, mg/dL	53 ± 17	50 ± 14	61 ± 20	59 ± 20	<0.0001
LDL, mg/dL	116 ± 36	118 ± 36	111 ± 35	114 ± 35	0.314

Waist Hip Ratio	0.9 (0.1)	0.9 (0.1)	0.9 (0.1)	0.9 (0.1)	0.016
Literacy/vocabulary					
WRAT and WAT, Z-score	0.01 ± 1.0	-0.01 ± 1.0	-0.25 ± 1.0	0.410 ± 0.8	<.0001
Peabody, Z-score	0.05 ± 1.0	0.02 ± 1.0	-0.19 ± 1.0	0.478 ± 0.8	<.0001
Percent WMHV**	0.5 ± 0.7	0.5 ± 0.7	0.9 ± 1.0	0.6 ± 0.8	0.0007
TCB ***	0.73 ± 0.04	0.74 ± 0.04	0.71 ± 0.04	0.71 ± 0.05	<.0001

* p-values for age adjusted global comparison among 3 race-ethnicity groups (2 degrees of freedom)

** Percent Total Cranial Volume adjusted WMHV

*** Total Cranial Volume adjusted (proportion)

Table 2: Mild cognitive impairment and probable dementia status by race/ethnicity at visit 2

Cognitive status	Total (N=989)	Hispanic (N=672)	Black (N=162)	White (N=134)	P-value
Normal, N (%) [*]	737 (74)	488 (74)	116 (72)	114 (86)	<.0001 ^{**}
MCI	199 (20)	141 (20)	40 (25)	16 (12)	
Probable dementia	45 (5)	36 (5)	6 (3)	3 (2)	

^{*} Rounded to whole numbers

^{**}Chi-Squared with 4 degrees of freedom

Table 3: Multivariable model of race/ethnicity and MCI/probable dementia outcomes

Parameter	OR	95% CI of OR	p-value*	p-value†
Model 1				
Black vs. White	1.9	1 3.7	0.048	<0.0001
Hispanic vs. White	4.5	2.3 8.9	<0.0001	
Hispanic vs. Black	2.3	1.3 4	0.003	
Model 2				
Black vs. White	1.8	0.9 3.6	0.213	<0.0001
Hispanic vs. White	4.5	2.2 9.2	0.0002	
Hispanic vs. Black	2.6	1.4 4.9	0.003	
Model 3				
Black vs. White	1.7	0.9 3.3	0.113	<0.0001
Hispanic vs. White	4.3	2.2 8.5	<0.0001	
Hispanic vs. Black	2.5	1.4 4.4	0.001	

* p-value for pairwise comparison

† p-value for Likelihood Ratio Test with 2 degrees of freedom

Model 1 adjusted for age, education, medical insurance status, literacy.

Model 2 adjusted for variables in model 1 and smoking, leisure time physical activity, diabetes mellitus, systolic and diastolic blood pressure, antihypertensive use, low- and high-density lipoprotein levels, waist hip ratio, and reported ethanol use.

Model 3 adjusted for variables in model 1 and brain parenchymal fraction, subclinical infarct status, and log white matter hyperintensity volume.

FIGURE LEGEND

Flow chart showing NOMAS MRI sample participation in first and second in-person assessments.

* Loss to follow-up may be temporary as attempts to recontact continue.

FIGURE

Supplemental Material

Supplemental Table 1: Comparison of participants who did or did not return for visit 2

Variable	Returned for Visit 2		p-value
	Yes (n=989)	No (n=301)	
Global cognitive domain Z-score at visit 1 (SD)	0.1 (0.7)	-0.3 (0.8)	<0.0001
Age, visit 1	69 (9)	75 (9)	<0.0001
Education, years	10 (5)	10 (5)	0.905
Race/ethnicity*			
White	134 (14)	57 (19)	0.007
Black	162 (16)	61 (20)	
Hispanic	672 (68)	175 (58)	
Sex			
Women	610 (62)	170 (57)	0.106
Men	379 (38)	131 (44)	
Medical insurance status			
Medicare or private insurance	505 (51)	172 (57)	0.064
Medicaid or no insurance	484 (49)	129 (43)	
Literacy Z-scores			
WRAT/WAT	0.01 (0.9)	-0.05 (1.04)	0.387
Vocabulary	0.05 (1.0)	-0.16 (1.0)	0.003
Current smoker	155 (16)	48 (16)	0.909
Moderate or Heavy Physical Activity	97 (10)	34 (11)	0.464
Diabetes mellitus	267 (27)	86 (29)	0.592
Systolic BP, mm Hg	136 (17)	139 (18)	0.004

Diastolic BP, mm Hg	79 (9)	76 (10)	0.0005
HDL, mg/dL	53 (17)	54.9 (17)	0.079
LDL, mg/dL	116 (36)	114.2 (34)	0.442

* Column percentages sum to 98 because data for Other races are not included.

Supplemental Table 2: Multivariable model of race/ethnicity and MCI/probable dementia outcomes among participants that were cognitively normal at visit 1

Parameter		OR	95% CI of OR		p-value*	p-value**
Model 1						
Black	vs. White	1.8	0.8	4	0.175	<.0001
Hispanic	vs. White	6	2.7	13.6	<.0001	
Hispanic	vs. Black	3.4	1.7	7.1	0.0008	
Model 2						
Black	vs. White	1.8	0.7	4.1	0.364	<.0001
Hispanic	vs. White	6.9	2.9	16.8	0.0001	
Hispanic	vs. Black	3.9	1.7	9.0	0.0011	
Model 3						
Black	vs. White	1.7	0.7	3.8	0.215	<.0001
Hispanic	vs. White	5.9	2.6	13.4	<.0001	
Hispanic	vs. Black	3.5	1.7	7.2	0.0009	

Cognitively normal defined as MMSE score of 27 or greater.

* p-value for testing pairwise comparison

** p-value for Likelihood Ratio Test with 2 degrees of freedom

Model 1 adjusted for age, education, medical insurance status, literacy.

Model 2 adjusted for variables in model 1 and smoking, leisure time physical activity, diabetes mellitus, systolic and diastolic blood pressure, antihypertensive

use, low and high density lipoprotein levels, waist hip ratio, and reported ethanol

use.

Model 3 adjusted for variables in model 1 and brain parenchymal fraction,

subclinical infarct status, and log white matter hyperintensity volume.