Symptoms suggestive of COVID-19 in households with and without children: a descriptive survey

Short running title: COVID-19 symptoms in households

Grace Grove^{1,2}, Nida Ziauddeen^{1*}, Nisreen A Alwan^{1,2,3*}

¹School of Primary Care, Population Sciences and Medical Education, Faculty of Medicine,
University of Southampton, Southampton SO16 6YD, UK

²NIHR Applied Research Collaboration Wessex, Southampton, UK

³NIHR Southampton Biomedical Research Centre, University of Southampton and University
Hospital Southampton NHS Foundation Trust, Southampton SO16 6YD, UK

*These authors contributed equally to this work.

Corresponding author: Grace Grove

School of Primary Care, Population Sciences and Medical Education, Faculty of Medicine,
University of Southampton, Southampton SO16 6YD, UK

Email: grace.grove@nhs.net

Synopsis

- Study question.

What is the nature and duration of symptoms suggestive of COVID-19 in UK households with and without children during March and May 2020? Do the symptoms vary between adults and children?

- What's already known.

There has been uncertainty about the extent to which children get and transmit SARS-CoV-2 within households. Symptoms associated with SARS-CoV-2 infection are well described in adults but symptoms and their duration are less well-characterised in children.

- What this study adds

On average, children had shorter and milder illness than adults, but still symptoms lasted more than 2 weeks in a significant proportion of children. In over a third of both adult and child symptomatic households, the child was the first to become ill.

Abstract

Background: Exploring transmission and symptoms of COVID-19 in children is vital, given that schools have recently fully reopened.

Objectives: This study aimed to characterise the nature and duration of symptoms suggestive of COVID-19 in UK households, and examine whether the symptoms varied between households with and without children and between adults and children from March to May 2020 in the UK.

Methods: An online questionnaire posted on social media (Mumsnet, Twitter, Facebook) was used to gather demographic and symptom information within UK households.

Results: Results from 508 households (1057 adults and 398 children) were available for analysis. 64.1% of respondent households with children and 59.1% of households without children had adults with symptoms suggestive of COVID-19. The proportion of adults that reported being symptomatic was 46.1% in households with children (and 36.7% in households without children). In 37.8% of households with at least one adult and one child with symptoms, the child's onset of symptoms started before the adult. Of all children, 35.7% experienced symptoms, with almost a quarter experiencing fluctuating symptoms for more than 2 weeks compared to almost half of symptomatic adults. In general, children had a shorter (median 5 days) and milder illness course than adults (median 10 days). Fatigue was the most common symptom in adults (79.7%) and cough was the most common symptom in children (53.5%). Chest tightness, shortness of breath, fatigue, muscle ache and diarrhoea were more common in adults than children, while cough and fever were equally common.

Conclusion: Children had shorter and milder illness than adults, but in almost a quarter of children symptoms lasted more than 2 weeks. In over a third of both adult-child

symptomatic households, the child was the first to become ill. Child to adult transmission and clinical presentation in children need to be further characterised.

Keywords: COVID-19; Adult; Child; United Kingdom

Introduction

Eight months on from the start of the pandemic, there is still ongoing debate around how COVID-19 manifests in children and whether they transmit the infection more or less efficiently than adults. This is particularly important now that schools across the world are have reopened at full capacity. Some suggest that school closures during spring 2020 may have had a modest effect on preventing deaths from COVID-19, much less than other social distancing measures.¹ However, evidence that children do transmit the virus to each other and to adults is emerging.²

A recent COVID-19 outbreak of 260 people in Georgia, USA included 180 children aged 11-17 years and 51 aged 6-10 years who tested positive, with attack rates of 44% and 51% respectively.³ A large study of almost 60,000 contacts and 5,700 index cases in South Korea in the period between January and March 2020 found that 17% of contacts have COVID-19 when the index case in the household was aged 10-19 years, compared to an average of 12% for cases of all ages.⁴ A study of 192 children presenting to hospital or urgent care in Massachusetts found nasopharyngeal viral loads in the first days of displaying symptoms to be higher than adults with severe disease.⁵

There is now a wide range of symptoms associated with SARS-CoV-2 infection.⁶ COVID-19 symptoms in children are less well-characterised than in adults. In general, children appear to be more likely to have a mild course than adults.^{7,8} In a recent review, up to 35% of children were asymptomatic. Amongst those who did develop symptoms, the most common was cough, followed by fever and pharyngitis. Other symptoms reported in children include; diarrhoea and vomiting, headache, upper respiratory tract symptoms, wheeze, fatigue, skin signs and less frequently, limb and chest pains.^{9,10} In the Massachusetts study, children with positive and negative SARS-CoV-2 tests had similar symptoms with fever, cough,

rhinorrhoea and headache being the most common.⁵ Children can present with severe illness, *Paediatric inflammatory multisystem syndrome associated with SARS-CoV-2* (PIMS-TS) and sadly there have been reported deaths in children.¹⁰

There is increasing evidence that COVID-19 symptoms in cases who did not require hospitalisation can be protracted and fluctuating. A US study that tracked 292 outpatients who tested positive for SARS-CoV-2 found that 35% of symptomatic respondents reported not returning to their usual state of health by 14-21 days after test date, including 26% among those aged 18-34 years. Continuing symptoms were commonly reported (43% for cough, 35% for fatigue and 29% for shortness of breath).¹¹ The COVID Symptom Study which uses a smartphone app to collect COVID-19 symptoms from millions of people found that 1 in 10 of those reporting symptoms is sick for more than three weeks.¹² A survey in the Netherlands of more than 1,600 mostly non-hospitalised people with COVID-19 symptoms, found that symptoms such as fatigue (88%), shortness of breath (75%), chest pressure (45%), headache (40%), muscle pain (36%) and palpitations (32%) last for months after initial infection.¹³

This study aimed to characterise the nature and duration of symptoms suggestive of COVID-19 in households experiencing mild symptoms, if the symptoms varied between households with and without children, and between adults and children from March to May 2020 in the UK.

Methods

This is a cross-sectional study using an online questionnaire. Adults aged 18 years and over resident in the UK were eligible to take part in this study. The questionnaire was developed

by the research team and published using iSurvey, the University of Southampton's online survey generation and research tool for distributing online questionnaires. iSurvey stores all information on the University server, which uses secure encryption and ensures that any data cannot be accessed by third parties. Survey responses were anonymous but participants were asked to give consent to be re-contacted in the future regarding COVID-19 testing status and result. Participants who consented were asked to provide an email address and contact telephone number. Ethical approval was granted by the University of Southampton, Faculty of Medicine Ethics Committee (Reference 56462).

The questionnaire was posted on Mumsnet, a parenting website run by parents, and on social media websites (Twitter and Facebook). These websites were chosen to post the questionnaire because it reaches a large audience. Permission was not required to post the questionnaire; however, it could only be posted in one place on Mumsnet (not-for-profit NFP surveys) and could not be shared elsewhere on the website by the researchers. The post contained a link to the questionnaire and brief information about the study. On opening the link, potential participants were taken to an in-depth information sheet. Participants gave their consent by ticking a box and confirmed as part of the consent that they were aged 18 years and over and resident in the UK. They then proceeded on to the questionnaire. Participants were able to share the questionnaire via other social networking sites and email with other people. The questionnaire was posted for a period of two weeks, from May 13th to 25th 2020.

The questionnaire included questions about the adult responding to the questionnaire as well as other adults and children in their household, all of whom are referred to as 'respondents' throughout. Questions included demographic information, activities that

involved contact with other people, contact with COVID-19 cases and symptoms suggestive of COVID-19 experienced for each household member and their duration.

Data was downloaded from iSurvey once the survey was taken offline. Raw data were checked and blank data rows (indicating participants who had viewed but not completed the questionnaire) were deleted. Participants were asked to report if they were ill on or after 16th March but a large number reported being ill in early March so all participants that reported experiencing symptoms indicative of COVID-19 from the 1st of March were included in the analysis. Households with children were categorised into early years, primary school aged and secondary school aged based on the age of the oldest child in the household.

Statistical analysis

Statistical analysis was undertaken using Stata 15.0. Descriptive percentages and summary statistics were generated. Univariable comparisons were carried out using an independent t-test for parametric continuous variables, Mann-Whitney test for non-parametric continuous variables and chi square test for categorical variables.

Results

Responses from a total of 508 households were available for analysis. Data were available on a total of 1057 adults and 398 children (aged 0-18 years). There were 232 households without children, 234 households with children and a further 42 households that were stated as households with children but did not provide information on the children. These 43 households were excluded from analyses comparing households with and without

children. Adults in households with children were younger than those in households without children. 36.6% of households without children had at least one member leaving the house for work during lockdown compared to 42.7% of households with children.

The mean number of children in households who reported individual data on children was 1.73 (SD 0.71). There were more respondents in the south than the north of the UK. There were similar proportions of respondents from the north and the south in households with and without children (Table 1). 64.1% of households with children and 59.1% of households without children reported having at least one adult with symptoms indicative of COVID-19. Among households with children, 153 of 234 households reported at least one individual experiencing symptoms, 90 of them had at least one adult and one child experiencing symptoms. Within these 90 households, an adult was first to experience symptoms in 51.1%, a child in 37.8% and an adult and child on the same day in 11.1% (Table S1). In the instances where a child was the first to exhibit symptoms, the proportion was similar in households with primary and secondary children.

When looking at individuals, the mean age of adults was 43.9 years (SD 13.4), and the mean age of children was 9.2 years (SD 4.7). 54.0% of the adult respondents were women and 54.8% of the child respondents were boys. Of the total 1057 adults and 398 children, 41.4% (438) of adults and 35.7% (142) children reported being unwell (Table 2). A higher proportion of children (24.1%) than adults (10.5%) described not leaving the home at all during lockdown. The most common reason for leaving the home was exercise for both adults and children. The vast majority of respondents described themselves as being from a white ethnic background (90.7% of adults and 86.9% of children). Out of all adults, 22.9% were leaving home for work.

The proportion of adults in households with children that reported being symptomatic (229 of 496, 46.1%) was higher than adults in households without children (179 of 488, 36.7%) (Table 3). Of those adults who experienced symptoms, more were women (60.9% in households without children and 59.4% in households with children). Shortness of breath and sneezing were more common in adults in households without children compared to those with children, with the rest of symptoms being similar (Table 3). A similar pattern was observed between the two household types after excluding households in which at least one individual was leaving home for work during lockdown (Table S2).

Overall, 41.4% of adults and 35.7% of children were reported to have symptoms indicative of COVID-19. Of those symptomatic adults, 22.6% were reported to be leaving for work during lockdown while asymptomatic. Fatigue was the most common symptom in adults (79.7% of symptomatic adults) and cough was the most common symptom in children (53.5% of symptomatic children) (Table 4, Figure 1). Presence of cough and fever was similar between adults (55.9% and 48.6% respectively) and children (53.5% and 51.4% respectively). Abdominal pain and vomiting were also similar between adults (20.8%, 5.0%) and children (18.3%, 6.3%). Chest pain, chest tightness and shortness of breath were all more common in adults (26.5%, 45.7%, 44.1% respectively) than in children (4.2%, 7.0%, 11.3% respectively). Adults were also more likely to report muscle ache, loss of appetite, diarrhoea, and loss of sense of taste or smell than children. After excluding households where at least one adult was leaving for work (adults n=258 and children n=69), similar patterns are observed (Table S3).

The median duration of symptoms was longer in adults in households without children (14 days, IQR 5.0, 21.0) than in adults in households with children (10 days, IQR 6.0, 20.0) (Table 3). 42.5% of the adults in household without children and 47.6% in those with children

reported symptoms fluctuating for more than two weeks. Overall, adults (regardless of household status) were unwell for longer (median 10 days, IQR 6.0, 21.0) than children (median 5 days, IQR 2.5, 8.5) (Table 4). Adults were also more likely to report a fluctuating illness for more than two weeks than children (45.0% in adults, 22.5% in children). Of the adults that reported being tested for SARS-CoV-2, 17 tested positive and 53 tested negative (Table 5). One adult who tested positive reported not experiencing symptoms. Only 7 children were tested, and all received a negative result. Nearly a quarter of those who tested positive were from an ethnic minority background. The majority of individuals who tested positive reported fluctuation in symptoms (31.3% for <2 weeks and 50.0% for >2 weeks, compared to 14.7% and 52.9% respectively in those who tested negative). However, the duration from symptom onset to test was longer in those that tested negative (median 11.5 days, IQR 4, 30) than in those that tested positive (median 4.5 days, IQR 2.5, 6.5). The most common symptoms in those testing positive were cough, fever, fatigue, muscle ache, loss of sense of smell and loss of sense of taste. A high proportion of those who tested negative also reported cough, fever and fatigue as symptoms but muscle ache, loss of sense of smell and loss of sense of taste were less common.

Discussion

In this survey, 64.1% of households with children and 59.1% of households without children had adults with symptoms suggestive of COVID-19 in March-April 2020. The proportion of adults that reported being symptomatic was higher in households with children (46.1%) than without children (36.7%). In 37.8% of households with at least one adult and one child with symptoms, the child's onset of symptoms was before the adult. Of all children, 35.7%

experienced symptoms, with almost a quarter of those with symptoms experiencing fluctuating symptoms for more than 2 weeks. In general, children had a shorter and milder illness course than adults. Chest tightness, shortness of breath, fatigue, muscle ache and diarrhoea were more common in adults than children, while cough and fever were equally common.

The main driver for the spread of COVID-19 is person-to-person transmission¹⁴ but until recently there has been scientific uncertainty around the extent to which children transmit the virus. The evidence that children transmit SARS-CoV-2 is increasing. There was no evidence of difference in rate of infection by age in China¹⁵ and England.¹⁶ Household contacts of index cases aged 10-19 years were most likely to test positive (18.6%) followed by household contacts of older adults (17.0% in 60-69 and 18.0% in 70-79 years) and 11.8% overall.⁴ A meta-analysis of nine contact tracing studies concluded that children are half as likely as adults to catch COVID-19. However, the included studies were all considered low to medium quality, included few children, high variation among studies and the adult category included older adults who are known to be at increased risk.¹⁷

True infection and transmission rate cannot be accurately identified without proactive case finding, mass testing and contact tracing. Children have been tested less frequently as they exhibit milder or no clinical symptoms.^{9,18,19} Two percent of 255,549 identified cases in Italy²⁰ and 9.3% of 4,766,825 identified cases in USA²¹ were in children aged 0-18 years. However, the number of child cases in USA showed a 90% increase over 4 weeks from 200,184 on 9th July to 380,174 on 6th August 2020.²¹

Our findings could be supportive of possible viral transmission from children to adults as the symptoms experienced in households were during the weeks around the schools shut down date in England (20th March 2020), when it is likely that community transmission of COVID-

19 was reasonably high. In over a third of the surveyed households where at least one adult and one child had symptoms, the child was the first member of the household to display symptoms. Leaving for work during lockdown was more common in households with children, so the index case could have been the person leaving for work in households where there was more than one person unwell, with the child being a secondary case. However, we repeated this analysis excluding the households where adults left the house for work during lockdown as this could be the possible source of infection and the findings were similar. The majority of the clinical research to date has focused on adults, and thus our understanding of the clinical picture of COVID-19 in children is more limited.^{7,22} Additionally, the research that is available on children tends to focus on more severe cases, as these are the group most likely to have been admitted to hospital and tested for COVID-19 early in the pandemic.²³

Our finding support what has been reported elsewhere that children have shorter and less severe illness than adults.^{7,8,22,24,25} The pathophysiology underlying these differences is unclear, and may include differing immune responses, reduced expression of ACE2 in children (which may be important for viral entry into cells) and increased levels of other viral illnesses in children, leading to competition with SARS-CoV-2²⁶. Additionally, children are less likely to have concomitant comorbidities that increase risk of hospitalisation and death from COVID-19²⁷, and differences in reporting may impact on symptoms recorded for paediatric cases.

It seems reasonable to assume that young children may find it difficult to clearly vocalise their symptoms, particularly those that are unusual or may be subtle, for example, changes in sense of smell or taste. Interestingly, prevalence of cough and fever was similar between children and adults. Cough and fever are both classic of COVID-19 and clearly detectable

symptoms, even for young children. Other studies have also found cough and fever to be common symptoms in children.^{22,23,25} A striking finding in our study has been that almost a quarter of symptomatic children had fluctuating symptoms for longer than two weeks. This survey provided detailed information on symptom type and duration for UK-based households during the time of the COVID-19 pandemic lockdown (which started on 23 March 2020). The household approach allowed exploration of whether adults or children were the first to experience symptoms in their households. However, due to the nature of the online and social media platform this survey may not be representative of the general population, in terms of exposure to SARS-CoV-2 and factors associated with severity of infection such as ethnicity and deprivation.²⁸ It is likely those who experienced possible COVID-19 symptoms may have felt more compelled to undertake the survey than those who had not.

One limitation of the household approach was that asking one member of the household to complete the survey for all household members may have resulted in some inaccuracies in responses, if the survey completer relied on their memory rather than asking other household members directly. Additionally, the survey asked for symptom data for those unwell on 16th March or later. However, whilst some participants only gave data for the requested timeframe, many participants provided data for early March (70% in households without children and 75% in households with children reported symptoms on or after 16th March). The decision was made to include these responses in our analysis, but this means that some respondents who were unwell in early March and did not provide data on their symptoms will have been included in the asymptomatic group for the analysis of the data. It was not possible to confirm COVID-19 infection status in most respondents, due to very limited community testing at the time that the survey was undertaken. However, a large

number of respondents report both classic COVID-19 symptoms^{9,25,29}, and a prolonged, fluctuating course, which is unusual in other infectious illnesses, especially in children. A fluctuating course of respiratory illness in children could be due to co-infection or successive infections. However, this is unlikely to explain the fluctuating course of illness observed in this sample as households were asked to isolate for 14 days and this was followed by lockdown thus restricting exposure to another infection. This prolonged and fluctuating illness, coupled with high community transmission at the time, suggest that it is likely that many of the respondents may truly have had COVID-19. In those who were tested, it is important to consider the timing of test. Those who tested negative were tested later, on average, than those who tested positive. The median time of test for those testing negative was 11.5 days after symptom onset, well beyond the UK government's recommended 5 days³⁰. This may have resulted in some individuals who truly did contract COVID-19 receiving a negative test result as a result of reducing viral load over time.

Conclusion

Over a third of the children included in this survey experienced symptoms suggestive of COVID-19, and were the first to become unwell in over a third of households where at least one adult and one child were symptomatic. Cough and fever were common symptoms in both adults and children, with almost half of the adults and almost a quarter of the children who became unwell experiencing fluctuating symptoms for more than 2 weeks.

Acknowledgements

We thank all survey participants for responding to the survey.

Funding

The authors have no funding to report.

References:

1. Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, et al. School closure and management practices during coronavirus outbreaks including COVID-19: a rapid systematic review. *Lancet Child Adolesc Health*. 2020 May;4(5):397–404.
2. European Centre for Disease Prevention and Control. COVID-19 in children and the role of school settings in COVID-19 transmission [Internet]. 2020 Aug [cited 2020 Sep 1]. Available from: <https://www.ecdc.europa.eu/sites/default/files/documents/COVID-19-schools-transmission-August%202020.pdf>
3. Szablewski CM, Chang KT, Brown MM, Chu VT, Yousaf AR, Anyalechi N, et al. SARS-CoV-2 Transmission and Infection Among Attendees of an Overnight Camp — Georgia, June 2020. *MMWR Morb Mortal Wkly Rep*. 2020 Aug 7;69(31):1023–5.
4. Park YJ, Choe YJ, Park O, Park SY, Kim Y-M, Kim J, et al. Contact Tracing during Coronavirus Disease Outbreak, South Korea, 2020. *Emerg Infect Dis* [Internet]. 2020 Oct [cited 2020 Sep 1];26(10). Available from: http://wwwnc.cdc.gov/eid/article/26/10/20-1315_article.htm
5. Yonker LM, Neilan AM, Bartsch Y, Patel AB, Regan J, Arya P, et al. Pediatric SARS-CoV-2: Clinical Presentation, Infectivity, and Immune Responses. *J Pediatr*. 2020 Aug;S0022347620310234.

6. Centers for Disease Control and Prevention. Coronavirus Disease 2019 (COVID-19) – Symptoms [Internet]. 2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>
7. Chang T-H, Wu J-L, Chang L-Y. Clinical characteristics and diagnostic challenges of pediatric COVID-19: A systematic review and meta-analysis. J Formos Med Assoc. 2020 May;119(5):982–9.
8. Swann OV, Holden KA, Turtle L, Pollock L, Fairfield CJ, Drake TM, et al. Clinical characteristics of children and young people admitted to hospital with covid-19 in United Kingdom: prospective multicentre observational cohort study. BMJ. 2020 Aug 27;m3249.
9. Zimmermann P, Curtis N. Coronavirus Infections in Children Including COVID-19: An Overview of the Epidemiology, Clinical Features, Diagnosis, Treatment and Prevention Options in Children. Pediatr Infect Dis J. 2020;39(5):355–68.
10. Royal College of Paediatrics and Child Health. COVID-19 - research evidence summaries [Internet]. Available from: <https://www.rcpch.ac.uk/resources/covid-19-research-evidence-summaries>
11. Tenforde MW, Kim SS, Lindsell CJ, Billig Rose E, Shapiro NI, Files DC, et al. Symptom Duration and Risk Factors for Delayed Return to Usual Health Among Outpatients with COVID-19 in a Multistate Health Care Systems Network — United States, March–June 2020. MMWR Morb Mortal Wkly Rep. 2020 Jul 31;69(30):993–8.
12. COVID Symptom Study. How long does COVID last? [Internet]. 2020 Aug. Available from: <https://covid.joinzoe.com/post/covid-long-term>
13. Peiling schetst schokkend beeld thuiszitters | Coronalongplein [Internet]. Available from: <https://coronalongplein.nl/informatie/peiling-schetst-schokkend-beeld-thuiszitters>

14. Chan JF-W, Yuan S, Kok K-H, To KK-W, Chu H, Yang J, et al. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster. *Lancet Lond Engl*. 2020 15;395(10223):514–23.
15. Bi Q, Wu Y, Mei S, Ye C, Zou X, Zhang Z, et al. Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *Lancet Infect Dis*. 2020 Apr;S1473309920302875.
16. Bovill H. Coronavirus (COVID-19) Infection Survey pilot: England, 12 June 2020 [Internet]. Office for National Statistics; 2020 Jun. Available from: <https://www.ons.gov.uk/peoplepopulationandcommunity/healthandsocialcare/conditionsanddiseases/bulletins/coronaviruscovid19infectionsurveypilot/12june2020#characteristics-of-people-testing-positive-for-covid-19>
17. Viner RM, Mytton OT, Bonell C, Melendez-Torres GJ, Ward JL, Hudson L, et al. Susceptibility to and transmission of COVID-19 amongst children and adolescents compared with adults: a systematic review and meta-analysis [Internet]. *Public and Global Health*; 2020 May [cited 2020 Jun 1]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2020.05.20.20108126>
18. Chen Z-M, Fu J-F, Shu Q, Chen Y-H, Hua C-Z, Li F-B, et al. Diagnosis and treatment recommendations for pediatric respiratory infection caused by the 2019 novel coronavirus. *World J Pediatr WJP*. 2020 Feb 5;
19. Han MS, Choi EH, Chang SH, Jin B-L, Lee EJ, Kim BN, et al. Clinical Characteristics and Viral RNA Detection in Children With Coronavirus Disease 2019 in the Republic of Korea. *JAMA Pediatr* [Internet]. 2020 Aug 28 [cited 2020 Sep 1]; Available from: <https://jamanetwork.com/journals/jamapediatrics/fullarticle/2770150>

20. The COVID-19 Task force of the Department of Infectious Diseases and the IT Service. Integrated surveillance of COVID-19 in Italy [Internet]. Istituto Superiore di Sanità; 2020 May. Available from:
https://www.epicentro.iss.it/en/coronavirus/bollettino/Infografica_18maggio%20ENG.pdf
21. American Academy of Pediatrics. Children and COVID-19: State Data Report. A joint report from the American Academy of Pediatrics and the Children's Hospital Association [Internet]. 2020 Jun. Available from:
<https://downloads.aap.org/AAP/PDF/AAP%20and%20CHA%20-%20Children%20and%20COVID-19%20State%20Data%20Report%208.6.20%20FINAL.pdf>
22. Götzinger F, Santiago-García B, Noguera-Julián A, Lanaspá M, Lancella L, Calò Carducci FI, et al. COVID-19 in children and adolescents in Europe: a multinational, multicentre cohort study. *Lancet Child Adolesc Health*. 2020 Sep;4(9):653–61.
23. Qiu H, Wu J, Hong L, Luo Y, Song Q, Chen D. Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *Lancet Infect Dis*. 2020 Mar;S1473309920301985.
24. Ludvigsson JF. Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults. *Acta Paediatr*. 2020 Jun;109(6):1088–95.
25. Du W, Yu J, Wang H, Zhang X, Zhang S, Li Q, et al. Clinical characteristics of COVID-19 in children compared with adults in Shandong Province, China. *Infection*. 2020 Jun;48(3):445–52.
26. Yuki K, Fujiogi M, Koutsogiannaki S. COVID-19 pathophysiology: A review. *Clin Immunol Orlando Fla*. 2020;215:108427.

27. Atkins JL, Masoli JAH, Delgado J, Pilling LC, Kuo C-L, Kuchel GA, et al. Preexisting Comorbidities Predicting COVID-19 and Mortality in the UK Biobank Community Cohort. Newman AB, editor. J Gerontol Ser A. 2020 Jul 20;glaa183.
28. Public Health England. Disparities in the risk and outcomes from COVID-19 [Internet]. 2020 Aug. Available from: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/908434/Disparities_in_the_risk_and_outcomes_of_COVID_August_2020_update.pdf
29. Menni C, Valdes AM, Freidin MB, Sudre CH, Nguyen LH, Drew DA, et al. Real-time tracking of self-reported symptoms to predict potential COVID-19. Nat Med. 2020 Jul;26(7):1037–40.
30. Get a free NHS test today to check if you have coronavirus [Internet]. Available from: <https://www.gov.uk/get-coronavirus-test>

Figure legend

Figure 1: Prevalence of symptoms amongst symptomatic adults (n=438) and children (n=142)

Table 1: Household characteristics of the study sample

	Households without children		Households with children	
	n	%	n	%
n of households	232	-	234	-
Total n of adults	488	-	496	-
n of adults in household				
1	63	27.2	19	8.1
2	108	46.6	178	76.1
3	36	15.5	28	12.0
4	24	10.3	7	3.0
5	1	0.4	2	0.9
Total n of children	-	-	398	-
n of children in household				
1	-	-	94	40.2
2	-	-	119	50.9
3	-	-	18	7.7
4	-	-	3	1.3
Households categorised by school group				
Early years	-	-	25	10.7
Primary	-	-	106	45.3
Secondary	-	-	103	44.0

Ethnicity of respondent				
White	215	92.7	217	92.7
BAME	13	5.6	15	6.4
Unspecified	4	1.7	2	0.9
Household ethnicity				
White	205	88.4	200	85.5
BAME	6	2.6	6	2.6
White and BAME	19	8.2	26	11.1
Unspecified	2	0.9	2	0.9
Leaving for work				
No	147	63.4	134	57.3
Yes	85	36.6	100	42.7
1 person	57	24.6	70	29.9
2 people	27	11.6	30	12.8
3 people	1	0.4	-	-
Region				
South East	53	22.8	47	20.1
South West	28	12.1	16	6.8
London	31	13.4	52	22.2
East of England	16	6.9	22	9.4
North East	9	3.9	2	0.9
North West	26	11.2	14	6.0
East Midlands	3	1.3	16	6.8

West Midlands	22	9.5	15	6.4
Yorkshire & Humber	17	7.3	13	5.6
Scotland	13	5.6	24	10.3
Northern Ireland	3	1.3	5	2.1
Wales	9	3.9	5	2.1
Unspecified	2	0.9	3	1.3
n of households with at least one symptomatic adult	137	59.1	150	64.1

Table 2: Individual characteristics of the study sample for adults and children (aged <18 years)

Variables	Adults		Children	
	n	%	n	%
n	1057		398	
n with symptoms	438	41.5	142	35.7
Age, years (mean $\pm$ SD)	43.9 $\pm$ 13.4		9.2 $\pm$ 4.7	
Gender				
Male	477	45.1	218	54.8
Female	571	54.0	178	44.7
Other	3	0.3	-	-
Unspecified	6	0.6	2	0.5
Work status during lockdown				
Not leaving for work	815	77.1	395	99.2

Leaving for work	242	22.9	1	0.3
Unspecified	-	-	2	0.5
Leaving home during lockdown				
Not leaving the house	111	10.5	96	24.1
Shopping for essentials	706	66.8	36	9.0
Exercise	749	70.9	299	75.1
Caring responsibilities	67	6.3	-	-
Attending school/ nursery	-	-	29	7.3
Ethnicity				
White	959	90.7	346	86.9
BAME	63	6.0	42	10.6
Other/Unspecified	35	3.3	10	2.5
Smoking status				
Non smoker	821	77.7	-	-
Ex-smoker	120	11.4	-	-
Current smoker	115	10.9	-	-
Unspecified	1	0.1	-	-

Table 3: Individual characteristics of symptomatic adults stratified by households with and without children*

	Without children		With children		p-value**
	n	%	n	%	

n	179/488	36.7	229/496	46.1	
Age, years (mean $\pm$ SD)	45.4 $\pm$ 14.5		42.7 $\pm$ 8.4		0.02
Gender					
Male	67	37.4	92	40.2	0.25
Female	109	60.9	136	59.4	
Other	2	1.1	-	-	
Unspecified	1	0.6	1	0.4	
Ethnicity					
White	156	87.2	209	91.3	0.60
BAME	19	10.6	18	7.9	
Unspecified	4	2.2	2	0.9	
Smoking status					
Non-smoker	141	78.8	182	79.5	0.98
Ex-smoker	22	12.3	27	11.8	
Current smoker	16	8.9	20	8.7	
Leaving home during lockdown					
None	14	7.8	11	4.8	0.21
Shopping	126	70.4	171	74.7	0.34
Exercise	125	69.8	177	77.3	0.09
Work	38	21.2	56	24.5	0.45

Caring responsibilities	15	8.4	14	6.1	0.38
Duration of symptoms, days (median, IQR)	14.0 (5.0 to 21.0)		10.0 (6.0 to 20.0)		0.85
Nature of symptoms					
Constant <1 week	27	15.1	46	20.1	0.17
Constant >1 week	24	13.4	26	11.4	
Fluctuating <2 weeks	50	27.9	46	20.1	
Fluctuating >2 weeks	76	42.5	109	47.6	
Unspecified	2	1.1	2	0.9	
Sought medical advice for symptomatic adults					
No	94	52.5	145	63.3	0.03
Yes	85	47.5	84	36.7	
Type of symptoms					
Cough	107	59.8	119	52.2	0.13
Fever	90	50.3	107	46.7	0.48
Chest pain	50	27.9	59	25.9	0.64
Chest tightness	83	46.4	103	45.2	0.81

Shortness of breath	89	49.7	91	39.9	0.05
Fatigue	143	79.9	185	81.1	0.75
Muscle ache	102	57.0	127	55.7	0.80
Loss of appetite	67	37.4	85	37.3	0.98
Diarrhoea	55	30.7	55	24.1	0.14
Abdominal Pain	42	23.5	40	17.5	0.14
Vomiting	11	6.2	9	4.0	0.31
Hoarse Voice	34	19.0	28	12.3	0.06
Sneezing	47	26.3	30	13.2	0.001
Loss of sense of smell	48	26.8	58	25.4	0.75
Loss of sense of taste	45	25.1	67	29.4	0.34
Confusion	17	9.5	14	6.1	0.21
Sore throat	76	42.5	118	51.8	0.06
Other: Headache	11	6.2	18	7.9	0.49
Other: Skin	11	6.2	12	5.3	0.71

*Only includes adults who have reported symptoms within the study period excluding 43 households who have reported having children in the household but the respondent provided no data on children)

**Chi square test for categorical variables, Independent t-test or Mann-Whitney test for continuous variables

Table 4: Characteristics of symptomatic adults and children (aged <18 years)

Variables	Adults		Children		p- value*
	n	%	n	%	
n	438/1057	41.4	142/398	35.7	
Age, years (mean $\pm$ SD)	43.6 $\pm$ 11.5		8.7 $\pm$ 4.5		<0.001
Gender					
Male	166	37.9	76	53.5	0.004
Female	268	61.2	66	46.5	
Other	2	0.5	-	-	
Unspecified	2	0.5	-	-	
Work status during lockdown					
Not leaving for work	339	77.4	-	-	
Leaving for work	99	22.6	-	-	
Leaving home					
None	28	6.4	25	17.6	<0.001
Shopping	316	72.1	18	12.7	<0.001
Exercise	325	74.2	110	77.5	0.44
Caring responsibilities	30	6.8	-	-	-
Attending school/ nursery	-	-	18	12.7	-
Duration of symptoms, days	10.0		5.0		<0.001

(median, IQR)	(6.0-21.0)		(2.5-8.5)		
Nature of Symptoms					
Constant <1 week	75	17.1	47	33.1	
Constant >1 week	53	12.1	38	26.8	
Fluctuating <2 weeks	108	24.7	23	16.2	<0.001
Fluctuating >2 weeks	197	45.0	32	22.5	
Unspecified	5	1.1	2	2	
Sought medical advice					
No	262	59.8	108	76.1	<0.001
Yes	176	40.2	30	21.1	
Symptoms					
Cough	245	55.9	76	53.5	0.60
Fever	213	48.6	73	51.4	0.57
Chest pain	116	26.5	6	4.2	<0.001
Chest tightness	200	45.7	10	7.0	<0.001
Shortness of breath	193	44.1	16	11.3	<0.001
Fatigue	349	79.7	68	47.9	<0.001
Muscle ache	243	55.5	28	19.7	<0.001
Loss of appetite	160	36.5	27	19.0	<0.001
Diarrhoea	121	27.6	21	14.8	0.002
Abdominal Pain	91	20.8	26	18.3	0.52
Vomiting	22	5.0	9	6.3	0.55
Hoarse Voice	67	15.3	7	4.9	0.001

Sneezing	87	19.9	16	11.3	0.02
Loss of sense of smell	115	26.3	11	7.8	<0.001
Loss of sense of taste	123	28.1	14	9.9	<0.001
Confusion	35	8.0	2	1.4	0.005
Sore throat	208	47.5	49	34.5	0.006
Other: Headache	32	7.3	10	7.0	0.97
Other: Skin	24	5.5	11	7.7	0.25

*Chi square test for categorical variables, Independent t-test or Mann-Whitney test for continuous variables

Table 5: Characteristics of the adult study sample who were tested for SARS-CoV-2

	Tested for SARS-CoV-2			
	Tested positive		Tested negative	
	n	%	n	%
n	17		53	
Age, years (mean $\pm$ SD)	44 $\pm$ 11.0		46.2 $\pm$ 12.1	
Gender				
Male	4	23.5	20	37.7
Female	13	76.5	33	62.3
Other	-	-	-	-
Unspecified	-	-	-	-
Ethnicity				
White	13	76.5	53	100.0

BAME	4	23.5	-	-
Unspecified	-	-	-	-
Symptomatic	16	94.1	34	64.2
Duration from symptom onset to test, median (IQR)	4.5 (2.5 to 6.5)		11.5 (4.0 to 30.0)	
Duration of symptoms, median (IQR)	16.5 (9.5 to 27.5)		14 (5.0 to 30.0)	
Nature of symptoms				
Constant <1 week	1	6.3	3	8.8
Constant >1 week	1	6.3	8	23.5
Fluctuating <2 weeks	5	31.3	5	14.7
Fluctuating >2 weeks	8	50.0	18	52.9
Unspecified	1	6.3	-	-
Sought medical advice				
No	12	75.0	8	23.5
Yes	4	25.0	26	76.5
Symptoms				
Cough	11	68.8	21	61.8
Fever	11	68.8	23	67.6
Chest pain	5	31.3	13	38.2
Chest tightness	6	37.5	19	55.9
Shortness of breath	10	62.5	19	55.9
Fatigue	16	100.0	28	82.4

Muscle ache	13	81.3	16	47.1
Loss of appetite	8	50.0	10	29.4
Diarrhoea	8	50.0	9	26.5
Abdominal Pain	3	18.8	6	17.6
Vomiting	-	-	2	5.9
Hoarse Voice	4	25.0	9	26.5
Sneezing	5	31.3	5	14.7
Loss of sense of smell	12	75.0	4	11.8
Loss of sense of taste	11	68.8	5	14.7
Confusion	2	12.5	3	8.8
Sore throat	9	56.3	16	47.1
Other: Headache	3	18.8	3	8.8
Other: Skin	1	6.3	4	11.8