**Maternal perception of child weight and concern about child overweight mediates the
relationship between child weight and non-responsive feeding practices**

Jian Wang¹, Daqiao Zhu^{1,*}, Xuwen Cheng², Yicong Liu³, Bingqian Zhu¹, Scott Montgomery^{4,5,6},
Yang Cao⁴

¹ School of Nursing, Shanghai Jiao Tong University, 200025 Shanghai, China;
wj-brittany@hotmail.com (J.W.); zhubq@shsmu.edu.cn (B.Z.)

² Department of Children's Disease Prevention, Jinyang Community Health Service Center, 200136
Shanghai, China; jyebcheng@126.com

³ Department of Nursing, Shanghai Children's Medical Center, 200127 Shanghai, China;
liuzhouyicong@hotmail.com

⁴ Clinical Epidemiology and Biostatistics, School of Medical Sciences, Örebro University, 70182
Örebro, Sweden; scott.montgomery@oru.se (S.M.); yang.cao@oru.se (Y.C.)

⁵ Clinical Epidemiology Division, Department of Medicine, Karolinska Institutet, 17177 Stockholm,
Sweden

⁶ Department of Epidemiology and Public Health, University College London, WC1E 6BT London,
United Kingdom

* Correspondence: zhudaqiao@aliyun.com

Abstract

We aimed to examine the mediating effects of maternal perception of child weight (weight perception) and concern about overweight (weight concern) on the paths between child weight and non-responsive feeding practices. We recruited a convenience sample of 1164 mothers who were primary caregivers of preschool children. Child body mass index (BMI) Z-score was calculated to assess child weight. The Chinese version of the Child Feeding Questionnaire (C-CFQ) was used to measure four common non-responsive feeding practices, weight perception and weight concern. Structural equation modeling (SEM) was used to examine the associations between child BMI Z-scores, maternal feeding practices, and other covariates. Sixty percent of the mothers perceived their overweight/obese children as normal weight or even underweight. The disagreement between actual child weight and maternal weight perception was statistically significant ($Kappa = 0.212, P < 0.001$). SEM indicated that weight perception fully mediated the relationship between child BMI Z-scores and pressure to eat. Weight concern fully mediated the relationships between child BMI Z-scores and the other three feeding practices. The serial mediating effects of weight perception and concern were statistically significant for the paths between child BMI Z-score and monitoring ($\beta = 0.035, P < 0.001$), restriction ($\beta = 0.022, P < 0.001$), and food as a reward ($\beta = -0.017, P < 0.05$).

Conclusion: Child weight may influence maternal feeding practices through weight perception and concern. Thus, interventions are needed to increase the accuracy of weight perception, which may influence several maternal feeding practices and thereby contribute to child health.

Keywords: preschool children; maternal concern; weight perception; feeding practices

38 **What is Known:**

- 39 • Non-responsive feeding practices may contribute to childhood obesity or eating disorders.
- 40 • Relationships between maternal weight perception and concern, child weight, and feeding
- 41 practices have been mixed.

42 **What is New:**

- 43 • Child weight may influence maternal non-responsive feeding practices through maternal weight
- 44 perception and concern.
- 45 • Interventions are needed to increase the accuracy of caregivers' perception of child weight which
- 46 may influence maternal feeding practices and thereby contribute to child health.

47 **Abbreviations**

- 48 AIC: Akaike information criterion
- 49 BMI: Body mass index
- 50 BMI Z-score: body mass index Z-score
- 51 C-CFQ: Chinese version of the Child Feeding Questionnaire
- 52 CFI: comparative fit index
- 53 CNY: Chinese Yuan
- 54 GFI: goodness of fit index
- 55 RMR: root mean square residual
- 56 RMSEA: root mean square error of approximation
- 57 SD: standard deviation
- 58 SE: standard error
- 59 SEM: structural equation modeling
- 60 Weight concern: maternal concern about child overweight
- 61 Weight perception: maternal perception of child weight

1. Introduction

Feeding practices refer to specific practices or strategies that parents employ to manage what, when and how much their children eat and shape their children's eating patterns [1,2]. There are two types of parental feeding practices: non-responsive and responsive feeding [3-5]. Non-responsive feeding practices (also known as coercive control), such as applying pressure to eat, restricting food, monitoring consumption, and using food as a reward, is very common, especially in Chinese families [6-8]. These non-responsive feeding practices may contribute to the development of childhood obesity or long-term eating disorders [9-12].

Many studies have identified factors related to non-responsive feeding practices [13-17]. Among these factors, child weight has been an important factor [13,15-17]. Longitudinal studies indicated that mothers of children with higher body mass index (BMI) Z-scores were more likely to apply restrictive feeding practices [18-20] and monitor their children's diet [18] and less likely to apply pressure to eat [18,20] than mothers of children with lower BMI Z-scores. However, it has been reported that parental perception of and concern about child weight (weight perception and weight concern) rather than actual child weight are associated with parental feeding practices [21-24]. For instance, Payne et al. [23] found that parents who were highly concerned about their child's weight were more likely to use restrictive feeding than parents with lower levels of concern, while there was no significant association between child weight and feeding practices. Yilmaz et al. [24] found that parents were less likely to encourage their children to eat if they perceived them as overweight than if they perceived them as normal weight or underweight. In comparison, Freitas et al. [25] recently reported that maternal restrictive feeding practices were independently associated with mothers' weight perception and concern as well as their child's weight status. Overall, current evidence of the relationships between

maternal weight perception and concern, actual child weight, and maternal non-responsive feeding practices has been inconsistent.

The inconsistent findings might be due to differences in factors included in the analyses (especially weight-related variables). Some studies did not examine maternal weight perception and/or weight concern and accordingly discovered only correlations between maternal feeding practices and child weight [15,26]. Other studies performed univariate analyses to investigate associations between factors of interest and parental feeding practices [15,24,27,28]. Collectively, there is a need to examine the intercorrelations between these variables.

Evidence-based theories of information processing and behaviorism learning, such as Gagne's Information Processing Model [29], have suggested that cognitive changes (e.g., perception or concern) occur only when an objective stimulus (e.g., child weight) exists. The changes may then lead to specific behaviors (e.g., feeding practices). Similarly, Marengo proposed a middle-range explanatory theory in which the following elements occurred in order: parental beliefs and values about child body weight, parental perception of child weight, parental concern, and family lifestyle changes [30]. Therefore, caregivers' weight perception and weight concern might mediate the association between child weight and feeding practices. A study of 7- to 9-year-old children in the U.K. reported that maternal concern about overweight fully mediated the relationship between child adiposity and the use of restrictive feeding practices [31]. In addition, previous research has indicated that child weight plays an important role in caregivers' feeding practices [13,15-17,25].

Based on the above theories and empirical evidence, we proposed a hypothesized model (Figure 1). In this model, (1) child weight had a direct effect on four common non-responsive feeding practices, and (2) the associations between child weight status and these non-responsive feeding practices were

mediated by maternal weight perception and weight concern. To explore the relationships proposed in the hypothesized model, we used structural equation modeling (SEM) while controlling for potential covariates, such as child age [14,16] and sex [14,16]. We also controlled for caregivers' age [16,32] and weight classification based on BMI [16,17], education level [14,17], and family income [32,33].

Figure 1. Hypothesized model for relationships between variables of interest

Note. Weight perception: perception of child weight; Weight concern: concern about child overweight.

2. Materials and Methods

2.1. Study design and participants

A correlational and cross-sectional design was used. This study was conducted between December 2017 and May 2020 in Shanghai, China. A convenience sample was selected from three public kindergartens located in three areas representing different economic levels in Pudong District, Shanghai. Only mothers were recruited because they are the primary caregivers of children typically in China. The required minimum sample size was 346, calculated using the formula below [34]:

$$N_{\varepsilon} = \frac{\delta_{1-\beta}}{\varepsilon^2 df} + 1,$$

where $\alpha = 0.05$; $\beta = 0.90$; $\varepsilon = 0.05$; $\delta_{1-\beta} \approx 33.6$; and $df = 39$.

A total of 1164 mothers were recruited. We excluded unqualified questionnaires, mothers who were not the primary caregivers, and children with diseases related to nutrition or extreme age-standardized BMI Z-scores. Finally, 1106 (95.02%) responses were included in this report.

Ethical approval was obtained from the Research Ethics Committee of Shanghai Jiao Tong University. Written informed consent was obtained from all participants.

2.2. Demographic and socioeconomic data

Demographic and socioeconomic data, including children's age, sex and mothers' age, weight, height, education level, and annual household income, were measured using a self-reported questionnaire. Children's weight and height were measured and collected by trained health teachers in the kindergartens. Maternal weight status was classified as underweight ($\text{BMI} < 18.5 \text{ kg/m}^2$), normal weight ($18.5 \text{ kg/m}^2 \leq \text{BMI} < 24.0 \text{ kg/m}^2$), or overweight or obese ($\text{BMI} \geq 24.0 \text{ kg/m}^2$) [35]. According to the World Health Organization (WHO) guidelines, child age-standardized BMI Z-scores were calculated using the software WHO Anthro (for 2- to 5-year-old children) and Young Growth Curve (for 5- to 6-year-old children). BMI Z-scores were categorized into three groups: underweight (Z-score < -2), normal weight ($-2 \leq \text{Z-score} \leq 1$), and overweight or obese (Z-score > 1) [36].

2.3. Maternal feeding practices

Maternal feeding practices were evaluated using the Chinese version of the Child Feeding Questionnaire (C-CFQ). The C-CFQ assesses four types of non-responsive feeding practices: four items of monitoring (the extent to oversee their child's eating [37]), six items of restriction (strict limitations on the child's access to foods or opportunities to consume unhealthy foods [38]), four items of pressure to eat (insists, demands, or physically struggles with the child in order to get the child to eat more food [37-39]), and two items of food as a reward (use of desired foods as a way to regulate the

child's eating or behaviors [39]). Each item was rated on a 5-point Likert scale. The response options for each item were "always," "usually," "sometimes", "almost never", and "never". Each subscale was calculated by averaging the scores of all the items in that subscale. The C-CFQ demonstrated good internal consistency reliability in the current study (Cronbach's $\alpha = 0.748-0.890$). The questionnaire also showed good construct validity [40,41].

2.4. Maternal perception of child weight (weight perception) and concern about overweight (weight concern)

Weight perception was assessed by asking "How would you describe your child's weight?". The responses included "very underweight," "slightly underweight," "normal weight," "slightly overweight," and "very overweight." Weight concern was assessed by asking "How concerned are you about your child becoming or staying overweight in the future". The responses included "unconcerned," "slightly concerned," "concerned," "fairly concerned," and "very concerned." Both items have been used and validated in previous studies [22,31].

2.5. Statistical analysis

SPSS Statistics 24.0 and SPSS Amos 24.0 for Windows (IBM Corp, Armonk, NY, USA) were used for data coding, cleaning, and analysis. Descriptive statistics were used to describe the participants' characteristics. The agreement between maternal weight perception and reported child weight was assessed using the Kappa statistic. Pearson's correlation analysis was used to explore the relationships between continuous variables. All the studied variables were approximately normally distributed with skewness and kurtosis below 2.0. Parameters for the hypothetical path model were estimated using structural equation modeling (SEM) via maximum likelihood estimation. The path analysis was conducted, and goodness of fit of the models was evaluated using the following indices and cut-offs: χ^2

between 1.0 and 2.0, goodness of fit index (GFI) and comparative fit index (CFI) > 0.90, root mean square error of approximation (RMSEA) < 0.06, root mean square residual (RMR) < 0.08, and a small Akaike information criterion (AIC) [40,41]. In the final model, we controlled for demographics that were significantly associated with maternal feeding practices ($P < 0.05$). The following conditions suggest a significant mediating effect: (1) the initial predictor (child BMI Z-score) was associated with the outcome (feeding practices) and the proposed mediators (perception or concern); (2) the mediators were associated with the outcome; and (3) the effect of the predictor on the outcome decreased substantially once the mediator was added as a second predictor in the model. To control for type I errors, the bootstrap method was used when examining the mediating effect of weight perception and weight concern [42]. Statistical significance was set at $P < 0.05$ (two-sided).

3. Results

3.1. Demographic and socioeconomic characteristics

The demographic characteristics of the participants are shown in Table 1. Two hundred nineteen (19.8%) preschool children were reported to be overweight or obese. A few demographic variables were statistically significantly related to mothers' feeding practices (not shown). Maternal age was negatively associated with the use of food as a reward ($r = -0.099$, $P = 0.001$). Maternal educational level was negatively associated with applying pressure to eat ($r = -0.121$, $P = 0.002$) and the use of food as a reward ($r = -0.087$, $P = 0.025$). Child age was negatively associated with maternal monitoring of the child's diet ($r = -0.111$, $P < 0.001$), applying pressure to eat ($r = -0.067$, $P = 0.025$) and restriction of food ($r = -0.060$, $P = 0.045$). Annual household income showed a positive relationship with maternal monitoring of the child's diet ($r = 0.084$, $P = 0.005$). These variables were added to the

model as covariates.

Table 1. Demographic characteristics of the participants ($n = 1106$)

	n (%) or (mean $\pm$ SD)
Child characteristics	
Age	4.56 $\pm$ 1.35
Sex	
Boys	588 (53.3)
Girls	516 (46.7)
Weight status	
Underweight: BMI Z-score < -2	44 (4.0)
Normal weight: $-2 \leq$ BMI Z-score ≤ 1	843 (76.2)
Overweight or Obesity: BMI Z-score > 1	219 (19.8)
Mother characteristics	
Age	34.66 $\pm$ 3.96
Weight status	
Underweight: BMI < 18.5 kg/m ²	111 (10.2)
Normal weight: $18.5 \text{ kg/m}^2 \leq$ BMI < 24.0 kg/m ²	784 (71.8)
Overweight or Obesity: BMI $\geq 24.0 \text{ kg/m}^2$	197 (18.0)
Education level	
College or higher	872 (78.8)
Senior high school or below	234 (21.2)
Household income/year	
Above average (> CNY 300,000)	638 (57.7)
Below average ($\leq$ CNY 300,000)	468 (42.3)

Notes. SD: standard deviation; BMI: body mass index; BMI Z-score: body mass index Z-score; CNY: Chinese Yuan

3.2. Agreement between maternal perception of child weight (weight perception) and actual child

weight status

Since few mothers perceived their children's weight as very overweight or very underweight, these two statuses were combined with overweight and underweight, respectively. Three categories (perceived underweight, normal weight, and overweight) were used in the final analysis. 14.1% of mothers felt that their children weighed too much, 59.0% reported that their children weighted a normal

weight, and 26.9% felt that their children were underweight. The agreement between maternal weight perception and actual child weight status was weak ($Kappa = 0.212$, $P < 0.001$) (Table 2). A total of 660 (59.67%) mothers properly perceived their children's weight status, and 378 (34.18%) mothers underestimated their children's weight status. Most mothers of overweight/obese children felt their children were normal weight ($n = 106$, 48.4%) or even underweight ($n = 9$, 4.1%) (Table 2).

Table 2. Agreement of actual child weight status with maternal weight perception

Actual child weight status (BMI Z-score)	Maternal weight perception ($n = 1106$)			
	Underweight	Normal weight	Overweight/Obesity	Total
Underweight	26 (59.1%)	16 (36.4%)	2 (4.5%)	44 (3.98%)
Normal weight	263 (31.2%)	530 (62.9%)	50 (5.9%)	843 (76.22%)
Overweight/Obesity	9 (4.1%)	106 (48.4%)	104 (47.5%)	219 (19.80%)
Total	298 (26.9%)	652 (59.0%)	156 (14.1%)	1106

Notes. BMI Z-score: body mass index Z-score; Weight perception: maternal perception of child weight; $Kappa = 0.212$, $P < 0.001$.

3.3. Description of variables included in the path analysis

Of the mothers, 74.1% were not concerned about their child becoming overweight, 18.4% reported some concern, and 7.5% reported great concern. The descriptions of weight perception, concern, and feeding practices are shown in Table 3. The mean scores of weight perception (2.84) and concern (1.85) were both lower than the median level of 3. Pressure to eat and food as a reward were approximately at the median level. Slightly higher mean scores were reported for restriction and monitoring.

Table 3. Description of weight perception, concern and feeding practices ($n = 1106$)

Variables	Range	Item numbers	Mean	SD
Weight perception	1-5	1	2.84	0.70

Weight concern	1-5	1	1.85	1.03
Pressure to eat	1-5	4	2.97	0.79
Food as a reward	1-5	2	3.38	0.76
Restriction	1-5	6	3.88	0.61
Monitoring	1-5	4	3.97	0.82

Notes. Weight perception: maternal perception of child weight; Weight concern: maternal concern about child overweight.

3.4. Relationships between variables in the path analysis

The bivariate associations between the four feeding practices (monitoring, pressure to eat, use food as a reward, and restriction), BMI Z-score, and weight perception and concern are shown in Table

4. Most of the associations were statistically significant.

Table 4. Correlations of the key variables ($n = 1106$)

	Monitoring	Pressure to eat	Food as a reward	Restriction	BMI Z-score	Weight perception
Pressure to eat	0.142**					
Food as a reward	-0.097**	0.144**				
Restriction	0.445**	0.254**	0.026			
BMI Z-score	0.042	-0.084**	-0.025	0.052		
Weight perception	0.032	-0.098**	-0.038	0.065*	0.514**	
Weight concern	0.124**	-0.058	-0.065*	0.108**	0.390**	0.528**

Notes: BMI Z-score: body mass index Z-score; Weight perception: maternal perception of child weight; Weight concern: maternal concern about child overweight. * $P < 0.05$, ** $P < 0.01$.

3.5. Path analysis

Based on the hypothesized model (Figure 1) and the above bivariate correlation analyses, the following variables were included in the initial model: BMI Z-score, weight perception and concern, monitoring, restriction, pressure to eat, food as a reward, child age, maternal age, maternal education level, and annual family income (Figure 2). The structural model revealed a good fit of the data:

$\chi^2_{(32)}=86.297$ ($P < 0.001$), GFI = 0.987, CFI = 0.961, RMSEA = 0.046, RMR = 0.035, and AIC =

176.297. In the initial model, three direct paths were not significant. We removed the three non-significant paths respectively to obtain the most parsimonious model. After simplifying the model, all the remaining statistically significant paths were entered into a final model to assess the relative importance of the direct and indirect effects of BMI Z-score on maternal feeding practices.

Figure 2. Initial model of the relationships between the study variables

Notes: Paths were adjusted for child age, maternal age, maternal education level, and annual household income. Arrows represent directions. Significant paths are shown as solid lines, and non-significant paths are shown as dashed lines.

BMI Z-score: body mass index Z-score; Weight perception: maternal perception of child weight; Weight concern: maternal concern about child overweight. *** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$.

The final model is illustrated in Figure 3. It showed a better fit than the initial model, with the following parameters: $\chi^2_{(34)}=89.109$ ($P < 0.001$), GFI = 0.986, CFI = 0.960, RMSEA = 0.038, and RMR = 0.046. The AIC value was lower in the current model (153.109) than in the previous model. The model accounted for 1.5%, 4.1%, 3.4%, and 1.5% of the variance in food as a reward, monitoring, restriction, and pressure to eat, respectively. All the direct paths in the final model were statistically

significant. Specifically, weight concern was associated with monitoring (standardized $\beta = 0.13$, $P < 0.001$), restriction (standardized $\beta = 0.12$, $P < 0.001$), and food as a reward (standardized $\beta = -0.07$, $P < 0.05$). The standardized β of the direct path from weight perception to pressure to eat was -0.10 ($P < 0.001$).

Figure 3. Final model of the relationships between the study variables

Notes: BMI Z-score: body mass index Z-score; Weight perception: maternal perception of child weight; Weight concern: maternal concern about child overweight. *** $P < 0.001$; ** $P < 0.01$; * $P < 0.05$.

3.6. Indirect effects of child weight and maternal perception of child weight (weight perception) on maternal feeding practices

The indirect effects of child weight and weight perception on maternal feeding practices are presented in Table 5. When all the variables were included in the model, both paths through a single mediation (indirect effects 1-4 in Table 5) and a serial mediation (indirect effects 5-7 in Table 5) were statistically significant. The total indirect impact on maternal feeding practices was also statistically

significant ($P < 0.05$). The mediation of weight concern (i.e., M2 in Table 5) between weight perception (i.e., M1 in Table 5) and food as a reward, restriction, and monitoring (i.e., Y1, Y2, and Y3 in Table 5) was also statistically significant.

278

Table 5. Indirect effects of child weight and weight perception on maternal feeding practices

Effect	β	Boot SE	Bootstrapping 95% Confidence Interval		Z
			Lower	Upper	
Indirect 1: $X \rightarrow M1 \rightarrow Y4$	-0.034	0.010	-0.055	-0.014	3.4**
Indirect 2: $M1 \rightarrow M2 \rightarrow Y1$	-0.033	0.016	-0.066	-0.002	2.06*
Indirect 3: $M1 \rightarrow M2 \rightarrow Y2$	0.044	0.011	0.022	0.067	4***
Indirect 4: $M1 \rightarrow M2 \rightarrow Y3$	0.069	0.015	0.040	0.100	4.6***
Indirect 5: $X \rightarrow M1 \rightarrow M2 \rightarrow Y1$	-0.017	0.008	-0.034	-0.001	2.125*
Indirect 6: $X \rightarrow M1 \rightarrow M2 \rightarrow Y2$	0.022	0.006	0.011	0.034	3.67***
Indirect 7: $X \rightarrow M1 \rightarrow M2 \rightarrow Y3$	0.035	0.008	0.020	0.052	4.375***

Notes: N = 1106. Number of bootstrap samples for bias-corrected bootstrap confidence intervals: 10000. X = BMI Z-score, M1 = Weight perception, M2 = Weight concern, Y1 = Reward, Y2 = Restriction, Y3 = Monitoring, Y4 = Pressure to eat; SE, standard error; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

4. Discussion

In this study, we examined the mediating roles of maternal perception of child weight and concern about overweight on the association between child weight and non-responsive feeding practices. The results from the SEM largely supported our hypotheses.

In this study, although the child BMI Z-score was significantly related to pressure to eat in the bivariate analysis, it was not a direct predictor in the SEM. Based on the bootstrapping results, child BMI Z-score influenced mothers' feeding practices via three paths: two full single mediating effects of weight perception and concern and a serial multiple mediating effects of weight perception and concern. Specifically, the relationship between child BMI Z-score and pressure to eat was fully mediated by

weight perception. This finding suggested that mothers of children with higher BMI Z-scores were more likely than those of children with lower BMI Z-scores to perceive their children as having a high weight, which in turn decreased the likelihood of them forcing their children to eat. Additionally, weight concern fully mediated the associations between child BMI Z-score and maternal restriction, monitoring, and use of food as a reward. In this case, mothers whose children were overweight or obese were more likely than those whose children were normal or underweight to report concerns and monitor or restrict their children's intake rather than reward them for eating. In addition, mothers of children with higher BMI Z-scores were more likely to perceive their children as having a high weight status and express concern about their children being overweight or obese, which ultimately affected their feeding practices, such as restricting the consumption of unhealthy food.

Previous studies have assessed the correlations between caregivers' feeding practices and weight-related variables [13,21,24,27]; however, few have explored their interrelationships. A population-based cohort study showed that mothers' concerns about their children's weight mediated the relationship between child BMI Z-score at 4 years and restrictive feeding at 10 years [43]. Our findings are consistent with previous evidence [21,22,31] and in accordance with Costanzo and Woody's suggestion [44] that parents should use controlled feeding practices when they are concerned about their child's weight. Therefore, although the results of this study supported our hypotheses, the mediating effect of weight perception and concern warrants further investigation.

In the current study, maternal perception of child weight was a mediator of child BMI and maternal feeding practices. This finding emphasizes the importance of accurate perception of child weight. Only when mothers recognize their preschool children's weight problems do they implement effective feeding practices to control their children's weight. In this study, we found a high percentage

of misperception of child weight, with mothers tending to underestimate their children's weight. The results are consistent with recent findings [45,46]. The inability to accurately identify child overweight/obesity is problematic given that substantial evidence suggests that weight perception is critical in motivating health-related behavioral changes [47-49]. Thus, mothers who consider their overweight/obese children as normal weight or even underweight might not be concerned about their children being overweight or obese and therefore adopt inappropriate feeding practices, such as forcing children to eat more. This practice may further influence a child's development and growth, especially their weight.

Although our model showed a good fit of the data, the variance in feeding practices accounted for by the predictors was relatively low. The low variance was in line with other studies [21,31]. For instance, Gregory et al. [21] found that maternal concern about child overweight was a significant predictor, explaining an additional 2% of the variance in restriction, while food responsiveness explained 7%. It is likely that weight perception and concern were not major contributing factors of maternal feeding practices. Another possible explanation is that important factors related to maternal feeding practices might have been missed [50]. Compared to mothers recruited in other studies [17,25,31], mothers in our study showed a relatively low level of weight concern. Additionally, the correlations between weight concern and maternal feeding practices were weak. These findings indicate that mothers in our study might have used specific feeding practices to address their concerns about their children's undereating habits and underweight [13,21,51]. For instance, it has been indicated that maternal use of pressure to eat is independently associated with maternal concern about their child eating too little [13,51]. Similarly, Gregory et al. [21] found that maternal concern about child underweight was the strongest independent predictor of pressure to eat, contributing to 15% of

the variance. In contrast, mothers tend to consider a relatively high child weight as an indicator of good health, especially in China [52-54]. They thus may be less concerned about child overweight than mothers in other countries. Moreover, monitoring or restricting unhealthy food intake may be associated with maternal concern about pediatric health problems, resulting in a decreased/lack of appetite or chronic diseases, such as obesity [49,55,56].

To the best of our knowledge, this study was among the first to examine the structural relationships between child weight, mothers' weight perception and weight concern, and non-responsive feeding practices. The significant paths and well-fitted model provided further empirical support for the hypothetical model. However, this study has limited generalizability because of the sampling method used (e.g., a convenience sampling of only mothers). Although the prevalence of overweight or obese children in our study was similar to that in a large-scale study conducted in Shanghai, China [57], random samples from multiple sites are needed in the future. Another limitation was the correlational design, which precluded us from making causal inferences. Additionally, self-reported information may be subject to recall bias. Finally, we focused on only non-responsive feeding practices, whereas other positive approaches of feeding (e.g., rules, limits, and modeling) were not assessed. Thus, longitudinal studies using comprehensive questionnaires are needed to confirm findings from this study.

5. Conclusion

In conclusion, maternal perception of child weight and concern about overweight rather than actual child weight may be related to mothers' feeding practices. Mothers' perception of their children's weight not only influenced their non-responsive feeding practices directly but also had an indirect effect via concern about child overweight. Thus, non-responsive feeding practices associated

360 with childhood obesity should first address maternal perception of child weight. Managing concern
361 about child overweight may be a potential intervention. Future studies should focus on correcting
362 caregivers' misperception about child weight. Interventions designed to improve caregivers' accurate
363 assessments of child weight are needed so that they can adopt appropriate feeding practices, thereby
364 optimizing child health.

Acknowledgements

We would like to thank all of the child healthcare workers at Jinyang Community Health Service Center for their help with data collection.

Author contributions

Data collection: X.C., Y.C., L.Z., and D.Z.; Formal analysis: J.W.; Investigation: X.C., Y.C., L.Z., and D.Z.; Methodology: D.Z., and Y.C.; Project administration: D.Z.; Resources: D.Z.; Software: J.W.; Supervision: D.Z., S.M. and Y.C.; Visualization: J.W.; Writing – original draft: D. Z, J.W., and Y.C.; and Writing – review and editing: J.W., X.C., B.Z., D.Z., S.M., and Y.C.

Funding

D.Z. and J.W.’s work was supported by a grant from the National Social Science Foundation of China (No.: 19BSH070). The funding body was not involved in the design of the study, data collection or analysis, interpretation of the results, or writing of the manuscript.

Compliance with ethical standards

Ethical approval

Ethical approval was obtained from the Research Ethics Committee of Shanghai Jiao Tong University. Written informed consent was obtained from the mothers.

Conflicts of interest

The authors declare no conflicts of interest.

Consent for publication

All caregivers understood and agreed that their information could be published in any journal. The authors declare that there is no conflict of interest regarding the publication of this article.

Reference

1. Boots, S. B., Tiggemann, M., & Corsini, N. (2018). "That's enough now!": A prospective study of the effects of maternal control on children's snack intake. *Appetite*, 126, 1-7. doi: 10.1016/j.appet.2018.03.008
2. Ventura, A. K., & Birch, L. L. (2008). Does parenting affect children's eating and weight status? *International Journal of Behavioral Nutrition and Physical Activity*, 5, 15. doi: 10.1186/1479-5868-5-15
3. Jansen, E., Williams, K. E., Mallan, K. M., Nicholson, J. M., & Daniels, L. A. (2018). Bidirectional associations between mothers' feeding practices and child eating behaviours. *International Journal of Behavioral Nutrition and Physical Activity*, 15(1), 3. doi: 10.1186/s12966-018-0644-x
4. Savage, J. S., Rollins, B. Y., Kugler, K. C., Birch, L. L., & Marini, M. E. (2017). Development of a theory-based questionnaire to assess structure and control in parent feeding (SCPF). *International journal of behavioral nutrition and physical activity*, 14(1), 9. doi: 10.1186/s12966-017-0466-2
5. Shi, C., Li, N., Dong, J., Wang, L., Li, X., Ji, C., Wang, X., Chi, X., Guo, X., Tong, M., & Zhang, M. (2017). Association between maternal nonresponsive feeding practice and child's eating behavior and weight status: children aged 1 to 6 years. *European journal of pediatrics*, 176(12), 1603-1612. doi: 10.1007/s00431-017-3007-8
6. Heller, R. L., & Mobley, A. R. (2019). Instruments assessing parental responsive feeding in children ages birth to 5 years: A systematic review. *Appetite*, 138, 23-51. doi: 10.1016/j.appet.2019.03.006
7. Zhang, X., & Benton, L. (2019). The Association of Acculturation and Complementary Infant and Young Child Feeding Practices Among New Chinese Immigrant Mothers in England: A Mixed Methods Study. *International journal of environmental research and public health*, 16(18), 3282. doi: 10.3390/ijerph16183282

- 409 8. Zhou, N., Cheah, C. S. L., Van Hook, J., Thompson, D. A., & Jones, S. S. (2015). A cultural
410 understanding of Chinese immigrant mothers' feeding practices. *Appetite*, 87,
411 160-167. doi: 10.1016/j.appet.2014.12.215
- 412 9. Harris, H., Mallan, K. M., Nambiar, S., & Daniels, L. A. (2014). The relationship between
413 controlling feeding practices and boys' and girls' eating in the absence of hunger. *Eating behaviors*,
414 15(4),519-522. doi: 10.1016/j.eatbeh.2014.07.003
- 415 10. Rollins, B. Y., Loken, E., Savage, J. S., & Birch, L. L. (2014). Maternal controlling feeding
416 practices and girls' inhibitory control interact to predict changes in BMI and eating in the absence of
417 hunger from 5 to 7 y. *The American journal of clinical nutrition*, 99(2), 249-257. doi:
418 10.3945/ajcn.113.063545
- 419 11. Shloim, N., Edelson, L. R., Martin, N., & Hetherington, M. M. (2015). Parenting Styles, Feeding
420 Styles, Feeding Practices, and Weight Status in 4-12 Year-Old Children: A Systematic Review of the
421 Literature. *Frontiers in psychology*, 6, 1849. doi: 10.3389/fpsyg.2015.01849
- 422 12. Stang, J., & Loth, K. A. (2011). Parenting style and child feeding practices: potential mitigating
423 factors in the etiology of childhood obesity. *Journal of the American Dietetic Association*, 111(9),
424 1301-1305. doi: 10.1016/j.jada.2011.06.010
- 425 13. de Souza Rezende, P., Bellotto de Moraes, D. E., Mais, L. A., Warkentin, S., & Augusto de Aguiar
426 Carrazedo Taddei, J. (2019). Maternal pressure to eat: Associations with maternal and child
427 characteristics among 2-to 8-year-olds in Brazil. *Appetite*, 133, 40-46. doi:
428 10.1016/j.appet.2018.10.014
- 429 14. Lo, K., Cheung, C., Lee, A., Keung, V., & Tam, W. (2016). Associated Demographic Factors of
430 Instrumental and Emotional Feeding in Parents of Hong Kong Children. *Journal of the Academy*

of Nutrition and Dietetics, 116(12), 1925-1931. doi: 10.1016/j.jand.2016.06.006

15. Powers, S. W., Chamberlin, L. A., van Schaick, K. B., Sherman, S. N., & Whitaker, R. C. (2006). Maternal feeding strategies, child eating behaviors, and child BMI in low-income African-American preschoolers. *Obesity (Silver Spring)*, 14(11), 2026-2033. doi: 10.1038/oby.2006.237

16. Russell, C. G., Haszard, J. J., Taylor, R. W., Heath, A.-L. M., Taylor, B., & Campbell, K. J. (2018). Parental feeding practices associated with children's eating and weight: What are parents of toddlers and preschool children doing? *Appetite*, 128, 120-128. doi: 10.1016/j.appet.2018.05.145

17. Warkentin, S., Mais, L. A., Latorre, M. d. R. D. d. O., Carnell, S., & de Aguiar CarrazedoTaddei, J. A. (2018). Relationships between parent feeding behaviors and parent and child characteristics in Brazilian preschoolers: a cross-sectional study. *BMC Public Health*, 18(1), 704-704. doi: 10.1186/s12889-018-5593-4

18. Eichler, J., Schmidt, R., Poulain, T., Hiemisch, A., Kiess, W., & Hilbert, A. (2019). Stability, Continuity, and Bi-Directional Associations of Parental Feeding Practices and Standardized Child Body Mass Index in Children from 2 to 12 Years of Age. *Nutrients*, 11(8), 1751. doi: 10.3390/nu11081751

19. Jansen, P. W., Tharner, A., van der Ende, J., Wake, M., Raat, H., Hofman, A., Verhulst F. C., van Ijzendoorn, M. H., Jaddoe, V. W V., & Tiemeier, H. (2014). Feeding practices and child weight: is the association bidirectional in preschool children? *The American journal of clinical nutrition*, 100(5), 1329-1336. doi: 10.3945/ajcn.114.088922

20. Quah, P. L., Ng, J. C., Fries, L. R., Chan, M. J., Aris, I. M., Lee, Y. S., Yap, F., Godfrey, K. M., Chong, Y-S., Shek, L. P., Tan, K. H., Forde, C. G., & Chong, M. F. F. (2019). Longitudinal Analysis Between Maternal Feeding Practices and Body Mass Index (BMI): A Study in Asian Singaporean Preschoolers. *Frontiers in nutrition*, 6, 32-32. doi: 10.3389/fnut.2019.00032

- 453 21. Gregory, J. E., Paxton, S. J., & Brozovic, A. M. (2010). Pressure to eat and restriction are
454 associated with child eating behaviours and maternal concern about child weight, but not child body
455 mass index, in 2- to 4-year-old children. *Appetite*, 54(3), 550-556. doi:
456 <https://doi.org/10.1016/j.appet.2010.02.013>
- 457 22. May, A. L., Donohue, M., Scanlon, K. S., Sherry, B., Dalenius, K., Faulkner, P., & Birch, L. L.
458 (2007). Child-feeding strategies are associated with maternal concern about children becoming
459 overweight, but not children's weight status. *Journal of the American Dietetic Association*, 107(7),
460 1167-1175. doi: 10.1016/j.jada.2007.04.009
- 461 23. Payne, L. O., Galloway, A. T., & Webb, R. M. (2011). Parental use of differential restrictive feeding
462 practices with siblings. *Int J Pediatr Obes*, 6(2-2), e540-e546. doi: 10.3109/17477166.2011.575144
- 463 24. Yilmaz, R., Erkorkmaz, U., Ozcetin, M., & Karaaslan, E. (2013). How does parents' visual
464 perception of their child's weight status affect their feeding style? *Nutrición hospitalaria*, 28(3),
465 741-746. doi: 10.3305/nh.2013.28.3.6358
- 466 25. Freitas, F. R., Moraes, D. E. B., Warkentin, S., Mais, L. A., Ivers, J. F., & Taddei, J. A. A. C. (2019).
467 Maternal restrictive feeding practices for child weight control and associated characteristics. *Jornal de*
468 *pediatria*, 95(2), 201-208. doi: 10.1016/j.jpmed.2017.12.009
- 469 26. Brann, L. S., & Skinner, J. D. (2005). More controlling child-feeding practices are found among
470 parents of boys with an average body mass index compared with parents of boys with a high body mass
471 index. *Journal of the American Dietetic Association*, 105(9), 1411-1416. doi:
472 [10.1016/j.jada.2005.06.005](https://doi.org/10.1016/j.jada.2005.06.005)
- 473 27. Lydecker, J. A., & Grilo, C. M. (2016). The apple of their eye: Attitudinal and behavioral correlates
474 of parents' perceptions of child obesity. *Obesity (Silver Spring)*, 24(5), 1124-1131. doi:

475 10.1002/oby.21439

476 28. Payas, N., Budd, G. M., & Polansky, M. (2010). Exploring relationships among maternal BMI,
477 family factors, and concern for child's weight. *J Child Adolesc Psychiatr Nurs*, 23(4), 223-230. doi:
478 10.1111/j.1744-6171.2010.00248.x

479 29. Gagne, R., & Driscoll, M. (1988). *Essentials of learning for instruction* (2nd ed.). Englewood Cliffs,
480 NJ: Prentice-Hall

481 30. Mareno, N. (2014). Parental perception of child weight: a concept analysis. *Journal of advanced*
482 *nursing*, 70(1), 34-45. doi: 10.1111/jan.12143

483 31. Webber, L., Hill, C., Cooke, L., Carnell, S., & Wardle, J. (2010). Associations between child weight
484 and maternal feeding styles are mediated by maternal perceptions and concerns. *European journal of*
485 *clinical nutrition*, 64(3), 259-265. doi: 10.1038/ejcn.2009.146

486 32. Moreira, I., Severo, M., Oliveira, A., Durão, C., Moreira, P., Barros, H., & Lopes, C. (2016). Social
487 and health behavioural determinants of maternal child-feeding patterns in preschool-aged children.
488 *Maternal & Child Nutrition*, 12(2), 314-325. doi: 10.1111/mcn.12132

489 33. Yang, W. Y., Burrows, T., MacDonald-Wicks, L., Williams, L. T., Collins, C. E., & Chee, W. S. S.
490 (2018). Parent-child feeding practices in a developing country: Findings from the Family Diet Study.
491 *Appetite*, 125, 90-97. doi: 10.1016/j.appet.2018.01.037

492 34. Kim, & H., K. (2005). The Relation Among Fit Indexes, Power, and Sample Size in Structural
493 Equation Modeling. *Structural Equation Modeling A Multidisciplinary Journal*, 12(3), doi:
494 368-390.10.1207/s15328007sem1203_2

495 35. Zhou, B.-F., & Cooperative Meta-Analysis Group of the Working Group on Obesity in, C. (2002).
496 Predictive values of body mass index and waist circumference for risk factors of certain related

497 diseases in Chinese adults--study on optimal cut-off points of body mass index and waist circumference
498 in Chinese adults. Biomedical and environmental sciences : BES, 15(1), 83-96. PMID: 12046553

499 36. de Onis, M., Garza, C., Onyango, A. W., & Rolland-Cachera, M. F. (2009). WHO growth standards
500 for infants and young children. Archives de pédiatrie, 16(1), 47-53. doi: 10.1016/j.arcped.2008.10.010

501 37. Birch, L. L., Fisher, J. O., Grimm-Thomas, K., Markey, C. N., Sawyer, R., & Johnson, S. L. (2001).
502 Confirmatory factor analysis of the Child Feeding Questionnaire: a measure of parental attitudes,
503 beliefs and practices about child feeding and obesity proneness. Appetite, 36(3), 201-210. doi:
504 10.1006/appe.2001.0398

505 38. Vaughn, A. E., Ward, D. S., Fisher, J. O., Faith, M. S., Hughes, S. O., Kremers, S. P.,
506 Musher-Eizenman, D. R., O'Connor, T. M., Patrick, H., & Power, T. G. (2016). Fundamental constructs
507 in food parenting practices: a content map to guide future research. Nutrition reviews, 74(2), 98-117.
508 doi: 10.1093/nutrit/nuv061

509 39. Arlinghaus, K. R., Hernandez, D. C., Eagleton, S. G., Chen, T. A., Power, T. G., & Hughes, S. O.
510 (2019). Exploratory factor analysis of The Comprehensive Feeding Practices Questionnaire (CFPQ) in
511 a low-income hispanic sample of preschool aged children. Appetite, 140, 82-90. doi:
512 10.1016/j.appet.2019.04.029

513 40. Hu, L. t., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis:
514 Conventional criteria versus new alternatives. Structural Equation Modeling: A Multidisciplinary
515 Journal, 6(1), 1-55. doi: 10.1080/10705519909540118

516 41. Byrne, B. (2016). Structural equation modeling with AMOS: Basic concepts, applications, and
517 programming (third edition). New York, NY: Taylor and Francis Group

518 42. Hayes, A. (2013). Introduction to mediation, moderation, and conditional process analysis. Journal

519 of Educational Measurement, 51(3), 335-337. doi: 10.1111/jedm.12050

520 43. Derks, I. P., Tiemeier, H., Sijbrands, E. J., Nicholson, J. M., Voortman, T., Verhulst, F. C., Jaddoe, V.

521 W., & Jansen, P. W. (2017). Testing the direction of effects between child body composition and

522 restrictive feeding practices: results from a population-based cohort. *The American journal of clinical*

523 *nutrition*, 106(3), 783-790. doi: 10.3945/ajcn.117.156448

524 44. Costanzo, P. R., & Woody, E. Z. (1985). Domain-Specific Parenting Styles and Their Impact on the

525 Child's Development of Particular Deviance: The Example of Obesity Proneness. *Journal of Social &*

526 *Clinical Psychology*, 3(4), 425-445. doi: 10.1521/jscp.1985.3.4.425

527 45. AlHasan, D. M., Breneman, C. B., Lynes, C. L., & Callahan-Myrick, K. (2018). Factors that

528 Influence Parental Misperception of Their Child's Actual Weight Status in South Carolina. *Maternal*

529 *and child health journal*, 22(7), 1077-1084. doi: 10.1007/s10995-018-2491-4

530 46. Berggren, S., Roswall, J., Alm, B., Bergman, S., Dahlgren, J., & Almquist-Tangen, G. (2018).

531 Parents with overweight children two and five years of age did not perceive them as weighing too

532 much. *Acta paediatrica*, 107(6), 1060-1064. doi: 10.1111/apa.14174

533 47. Almoosawi, S., Jones, A. R., Parkinson, K. N., Pearce, M. S., Collins, H., & Adamson, A. J. (2016).

534 Parental Perception of Weight Status: Influence on Children's Diet in the Gateshead Millennium Study.

535 *PLoS One*, 11(2), e0144931. doi: 10.1371/journal.pone.0144931

536 48. Daddario, D. K. (2007). A review of the use of the health belief model for weight management.

537 *Medsurg Nursing*, 16(6), 363-366. PMID: 18390255

538 49. Rhee, K. E., De Lago, C. W., Arscott-Mills, T., Mehta, S. D., & Davis, R. K. (2005). Factors

539 associated with parental readiness to make changes for overweight children. *Pediatrics*, 116(1),

540 e94-e101. doi: 10.1542/peds.2004-2479

541 50. Miller, A. L., Miller, S. E., & Clark, K. M. (2018). Child, Caregiver, Family, and Social-Contextual
542 Factors to Consider when Implementing Parent-Focused Child Feeding Interventions. *Current nutrition*
543 *reports*, 7(4), 303-309. doi: 10.1007/s13668-018-0255-9

544 51. Haines, J., Downing, K. L., Tang, L., Campbell, K. J., & Hesketh, K. D. (2018). Associations
545 between maternal concern about child's weight and related behaviours and maternal weight-related
546 parenting practices: a cross-sectional study. *International Journal of Behavioral Nutrition and Physical*
547 *Activity*, 15(1), 104-104. doi: 10.1186/s12966-018-0738-5

548 52. Gupta, N., Goel, K., Shah, P., & Misra, A. (2012). Childhood obesity in developing countries:
549 epidemiology, determinants, and prevention. *Endocrine Reviews*, 33(1), 48-70. doi:
550 10.1210/er.2010-0028

551 53. Hossain, M. S., Siddiquee, M. H., Ferdous, S., Faruki, M., Jahan, R., Shahik, S. M., Raheem, E., &
552 Okely, A. D. (2019). Is Childhood Overweight/Obesity Perceived as a Health Problem by Mothers of
553 Preschool Aged Children in Bangladesh? A Community Level Cross-Sectional Study. *International*
554 *journal of environmental research and public health*, 16(2), 202. doi: 10.3390/ijerph16020202

555 54. Jiang, J., Rosenqvist, U., Wang, H., Greiner, T., Lian, G., & Sarkadi, A.(2007).Influence of
556 grandparents on eating behaviors of young children in Chinese three-generation families. *Appetite*,
557 48(3), 377-383. doi: 10.1016/j.appet.2006.10.004

558 55. Gibson, E. L., Wardle, J., & Watts, C. J. (1998). Fruit and vegetable consumption, nutritional
559 knowledge and beliefs in mothers and children. *Appetite*, 31(2), 205-228. doi: 10.1006/appe.1998.0180

560 56. Lee, W. S., Tee, C. W., Tan, A. G., Wong, S. Y., Chew, K. S., Cheang, H. K., Lim, T., Lum, L. C.,
561 Aw, M. M. (2019). Parental concern of feeding difficulty predicts poor growth status in their child.
562 *Pediatrics and neonatology*, 60(6), 676-683. doi: 10.1016/j.pedneo.2019.04.004

563 57. Chen, C., Jin, Z., Yang, Y., Jiang, F., Huang, H., & Liu, S. (2019). Association of low birth weight
564 with thinness and severe obesity in children aged 3-12 years: a large-scale population-based
565 cross-sectional study in Shanghai, China. *BMJ open*, 9(5), e028738. doi:
566 10.1136/bmjopen-2018-028738