Comparative efficacy and safety of current drugs against COVID-19: A systematic review and network meta-analysis

Yi Li^a, Wei He^{b,a,*}

^a*School of Pharmacy, China Pharmaceutical University, Nanjing 210009, China*

^b*Shanghai Skin Disease Hospital, Tongji University School of Medicine, Shanghai 200443, China*

*Corresponding author.

E-mail addresses: Wei HE (weihe@cpu.edu.cn).

Abstract

The rapid spread of coronavirus disease (COVID-19) has greatly disrupted the livelihood of many people around the world. To date, more than 35.16 million COVID-19 cases with 1.037million total deaths have been reported worldwide. Compared with China, where the disease was first reported, cases of COVID-19, the number of confirmed cases for the disease in the rest of the world have been incredibly high. Even though several drugs have been suggested to be used against the disease, the said interventions should be backed by empirical clinical evidence. Therefore, this paper provides a systematic review and a meta-analysis of efficacy and safety of different COVID-19 drugs.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

Research in context

Evidence before this study

Currently, Covid-19 is one of the most urgent and significant health challenge, globally. However, so far there is no specific and effective treatment strategy against the disease. Nonetheless, there are numerous debates over the effectiveness and potential adverse effects of different COVID-19 antivirals. In general, there is invaluable need to continually report on new advances and successes against COVID-19, apparently to aid in managing the pandemic.

Added value of this study

This study provides a comprehensive, evidence-based guide on the management of multiple COVID-19 symptoms. In particular, we provide a review of 14 drugs, placebos and standard treatments against COVID 19. Meanwhile, we also performed a meta-analysis based on four clinical outcome indicators, to measure and compare the efficacy and safety of current interventions.

Implications of all the available evidence

Findings of this research will guide clinical decision in COVID-19 patients. It will also provide a basis for predicting clinical outcomes such as efficacy, mortality and safety of interventions against the disease.

Introduction

Despite stringent containment and isolation measures, the global incidence of COVID-19 continues to rise. Globally, there have been more than 35,165,808 confirmed cases and more than 1,037,153 deaths from COVID-19 pandemic, as of October 2, 2020¹. Currently, even though there are numerous researches on effective drug against COVID-19, no specific drugs have so far been recommended. Most of the used drugs are based on the pathogenesis of the disease, however, doubts continues to mount over their safety and effectiveness². One concern is that short-term benefits are on average minimal, and more worse, the balance between long-term benefits and

harms is still poorly studied. Although many drugs are available for COVID 19, their use should be guided by integrated evidence to inform the best choice for each patient³.

Network meta-analysis provides a comparative efficacy and a broader evidence-based picture of several interventions, the reason it was selected for exploring relative merits of multiple COVID-19 control modalities⁴. This is critical for COVID-19 because the view on best drugs is constantly changing, and in particular, in the race against managing this pandemic. Because it can provide a complete, extensive and updated viewpoint of evidence, network meta-analysis is an ideal tool to guide practical development and application of drugs against COVID 19. Our study provides a thorough comparison on effectiveness and safety of currently available interventions against COVID 19, likely not to have been experienced in some clinical settings. This network meta-analysis was performed using Bayesian Markov chain Monte Carlo (MCMC) random effect model that is capable of fitting models of virtually unlimited complexity, and as such have revolutionized the practice of Bayesian data analysis⁵. Specifically, we evaluated the safety and effectiveness of 14 drugs, along with placebos and standard care, according to four clinical outcomes: average recovery time, response rate, mortality and the rate of adverse events. Findings of this research aimed at guiding the rationale of clinical decisions among COVID-19 patients.

Methods

Search strategy and selection criteria

The systematic review and network meta-analysis was performed on numerous reports in the Cochrane Central Register of Controlled Trials, PubMed, Embase and Web of Science. We included relevant studies published since inception the above databases till August 11, 2020, regardless of the language. The search terms were “severe acute respiratory syndrome coronavirus 2*” (MeSH) OR “2019nCoV*” OR “SARSCoV2*” OR “2019 novel coronavirus*” OR “COVID19 virus disease” OR “COVID19 virus infection”, combined with a list of all included drugs.

Randomized controlled trials (RCTs) were included if they evaluated comparable outcome of drugs with standard care, placebos or safety and effectiveness of a monotherapy in adults (≥ 18 years old and of both sexes) with COVID-19. The interventional modules must have been in line with NIH COVID-19 Treatment Guidelines⁶. Notable though, the included reports were not limited by dosage and mode of administration. Additionally, included studies must have reported on at least one of the aforementioned outcomes. Finally, quasi-randomised controlled clinical trials (CCTs) and incomplete trials, non-drug treatment such as ventilation or trials on less than 10 participants were all excluded from the meta-analysis. For duplicated publications, we only included the latest and most complete edition. Reviews, case reports, letters and conference abstract were also excluded from our analyses.

Relevant studies were selected independently by two reviewers. In particular, they reviewed the main reports and supplementary materials, extracted the relevant information from the included trials and assessed the risk of bias according to modified Jadad scale⁷. The quality of included studies was rated based on four items: (1). generation of random sequence, (2). blindness of the trials, (3). concealment and (4). withdrawals and dropouts. Any discrepancies were resolved by a third arbitrator. Duplicates were first excluded using EndNote (version 9.3.3). Studies with irrelevant title and abstract were then removed. The remaining papers were read in whole to assess their suitability. Finally, relevant data were extracted independently according to pre-determined criteria.

Outcomes

The efficacy and safety of interventional drugs were assessed based on four outcomes: the average recovery time (average duration of hospital stay or days to reach clinical rehabilitation standard) and response rate (based on the proportion of patients discharged or reaching clinical rehabilitation standard during the test) for efficacy outcomes. On the other hand, safety outcomes included mortality (the proportion of patients who died during the test) and the rate of adverse events (the proportion of patients exhibiting adverse effect(s) associated with the drug).

Data analysis

Network meta-analysis was performed using a Bayesian Markov chain Monte Carlo (MCMC) random effect model. Key parameters for dichotomous outcomes included odds ratios (RR) and standardized mean difference (SMD) both at 95% CI for continuous outcomes⁸. Node splitting model tested the consistency in closed-loop interventions with both direct and indirect evidence with $P < 0.05$ considered statistically significant. If direct and indirect evidence were statistical indifferent, then the consistency model was used in the network meta-analysis; otherwise the source and significance of inconsistency was assessed⁵. Additionally, potential scale reduction factors (PSRFs) were calculated to evaluate the convergence of the model. A PSRF value close to 1 is indicative of good convergence, thus the results are high reliability⁹. $PSRF \leq 1.02$ was acceptable. Finally, comparative effectivity and safety of drugs was analyzed according to ranking probability under the Bayesian network model.

Bayesian network model analysis was performed using ADDIS software¹⁰, whereas the diagrams were plotted using Stata V. 14.2⁴.

Results

Data collection and risk of bias

In general, we identified 1137 articles, and after careful review, we extracted 128 of them for in depth review (Figure 1). We constituted the final analyses via including 28 trial studies, 14 drugs together with placebos or standard care and 17078 patients. Of the trial studies, 21 (75%) were standard care- or placebo- controlled trials, whereas the remaining 7 (25%) performed drug-drug comparisons. In addition, 4 (14%), 16 (57%) and 8 (28%) of the trials comprised of patients from North and South America, Asia and Europe, respectively. Characteristics of each report are listed in Table 2. Meanwhile, the risks of publication bias were assessed based on the modified Jadad scale. Therefore, studies with a score of 0-3 were considered high degree of bias to publication, converse to those with a score of more than 4 (Figure 2).

Bayesian evidence network and convergence evaluation

The network of relevant comparisons for the four outcomes is shown in Figure 3. The relationship between all interventional methods was plotted based on the direct comparison data. Briefly, each cycle represents an interventional drug. The width of the connection lines is proportional to the number of trials comparing every pair of treatments, whereas the size of each circle is proportional to the number of randomly assigned participants (i.e, sample size). The direct or indirect evidence among different interventional methods provided the basic conditions for the network meta-analysis. Specifically, except for Arbidol, Chloroquine, Febuxostat and Favipiravir, the rest of the drugs were evaluated against at least one standard care- or placebo.

Convergence PSRF of the four outcomes was acceptable, implying minor fluctuations in Markov chain and stable iteration trajectory. As such, the convergence of the model was within the expected distribution in the iterative process and, as a result, can be used in analyzing the data¹¹. Specifically, our model contained 4 chains, featuring 20,000 tuning iterations, 50,000 simulation iterations, 10 000 thinning interval inference samples, with a variance of 2.5. The number of iterations and annealing was sufficient; therefore, no additional extension was needed.

Consistency test

Node splitting analysis was performed using ADDIS software. There was no significant difference ($P > 0.05$) in average recovery time (20 trials, comprising 14169 patients) and mortality (16 trials, comprising 16117 patients) across studies. This points to coherence between direct and indirect comparison. However, there was a significant difference in response rate (24 trials, comprising 15120 patients) between two pairs of intervention drugs ($P < 0.05$). In particular local inconsistency was found between chloroquine and hydroxychloroquine as well as between lopinavir–ritonavir and chloroquine. Meanwhile, inconsistencies were also observed between interferon β and lopinavir–ritonavir, and between lopinavir–ritonavir and standard care with regard to rate of adverse events (21 trials, comprising 4198 patients). These significant differences suggest of potential uncertainty

of predicted effect regarding the above drugs, and as such, their clinical applications against COVID 19 need further evaluation¹².

Effects and rank probability of the interventions

Average recovery days

Average duration of hospital stays or days up clinical rehabilitation standard was reported in 20 trials incorporating 14169 participants. Treatment nodes included in the network meta-analysis were arbidol, azvudine, chloroquine, colchicine, favipiravir, hydroxychloroquine, interferon β , ruxolitinib, lopinavir-ritonavir, methylprednisolone and remdesivir, evaluated alongside placebos and standard care. There was no significant difference in average recovery days between interventions (Table 2). However, rank probability test showed favipiravir (rank probability: 0.46) is most likely the best intervention against COVID 19, followed by arbidol (rank probability: 0.18) and azvudine (rank probability: 0.15) (Figure 4 and Table 3).

Response rate

Of the 28 included reports, 24 trials encompassing 15120 participants reported on the number of patients discharged or who reached clinical rehabilitation standard during the test. The treatment nodes involved in the network meta-analysis were azvudine, chloroquine, dexamethasone, colchicine, favipiravir, febuxostat, hydroxychloroquine, interferon β , ruxolitinib, lopinavir-ritonavir, remdesivir, placebo and standard care. As shown in Table 2, there was no convincing evidence that any of the interventions significantly enhanced the response rate of patients with COVID 19. Nonetheless, rank probability test showed that the drugs, colchicine (rank probability: 0.42), chloroquine (rank probability: 0.28) and favipiravir (rank probability: 0.11), are most likely to be as the best interventions against the disease. Meanwhile, the treatment effects of febuxostat (0.04) dexamethasone (0.03) and ruxolitinib (0.03) were comparable to that of placebos (0.03) (Figure 4 and Table 3).

Mortality

Sixteen trials featuring 16117 participants reported on the number of patients who died of COVID 19 during the trials. Treatment nodes used in the analysis are as follows: colchicine, hydroxychloroquine, interferon β , ruxolitinib, ribavirin, lopinavir-ritonavir, remdesivir, placebo and standard care. Ruxolitinib was closely associated with the reduced mortality-rates [(2.24E+13, 20.03-7.78E+45 for placebo and 2.00E+13, 17.9-6.01E+45 for standard care (Table 2)]. Moreover, rank probability test showed ruxolitinib (rank probability: 0.99) is most likely the safest intervention against COVID 19 (Figure 4 and Table 3). Other interventions did not inhibited significant difference in mortality.

Rate of adverse events

Meanwhile 21 trials contained 4198 participants demonstrated the adverse effect is related with COVID 19 treatment with drugs such as arbidol, azvudine, chloroquine, colchicine, favipiravir, hydroxychloroquine, interferon β , ruxolitinib, lopinavir-ritonavir, methylprednisolone, remdesivir, placebo and standard care. As shown in Table 2, moderate certainty evidence showed that compared with standard care, hydroxychloroquine is significantly linked to the adverse events (OR=6.87, 1.52-41.44). Nonetheless, other drugs did not significantly affect the mortality ($P > 0.05$). Rank probability test showed that azvudine (rank probability: 0.97) may be an intervention associated with least adverse effect. Even so, randomized-controlled trials in patients with COVID-19 has so far provided little definitive evidence about adverse effects for most interventions (Figure 4 and Table 3)

Discussion

COVID-19 pandemic is currently the most catastrophic public health emergency in the world, with the disease agent displaying one of the fastest transmission rates ever recorded. Throughout human history, severe acute respiratory syndrome strain 2 (SARS Cov 2) virus has been the most difficult infectious agent to control or manage, shown also to cause the most diverse range of complications. Finding effective and safe treatment is not only the key but also the major challenge in controlling the pandemic. There have been numerous investment by research community across the world, exploring the pathogenic mechanism of COVID-19. Continuous reporting on the disease has elucidated the molecular mechanism and potential infection pathway of COVID-19, offering the chance for the rationalized use of clinical treatment. However, there is still concern over the effectiveness and safety of various interventions. Meanwhile, short-term treatment benefits are averagely minimal, and the balance between long-term benefits and harms is still poorly understood. Therefore, although clinicians have a wide pool of drugs to select from, however, the best choice for each patient should be backed by empirical evidence.

Accordingly, this network meta-analysis was performed to understand the relative balance between merits and potential shortcomings of multiple COVID-19 interventions. In general, 17078 individuals reported in 28 papers were randomly assigned to one of 14 individual drugs against COVID-19, placebo or standard care. Response of patients to various COVID 19 infections across the world was obtained thorough search of published reports in any language. The key parameters of interest were average recovery days, response and mortality rates and rate of occurrence of adverse events. This provided a solid base to comprehensively evaluate the effectiveness and reliability of various drugs against COVID-19. Our analysis found the efficacy of commonly proposed drugs to be insignificantly indifferent when compared with placebos or standard care. Moreover, the therapeutic effects of the drugs is always modest. For instance, compared with other drugs, patients on remdesivir, favipiravir and azvudine exhibited higher recovery rate and lower occurrence of adverse events, though statistically insignificant. By contrast, compared with other

drugs, chloroquine and hydroxychloroquine are not effective against COVID 19, and thus, may not be a first choice in the clinic. On the other hand, this meta-analysis demonstrated that compared with other interventions, ruxolitinib is associated with lower mortality rate, but the curative effect and other benefits of this drug in patients with COVID 19 remains unclear. In addition, according to a randomized-controlled trial, hydroxychloroquine did not reduce COVID 19-associated mortalities, and moreover, promoted the occurrence of adverse events. In general, so far there is no sufficient evidence demonstrated that the proposed COVID 19 drugs have improved efficacy and safety compared with placebo or standard care.

Our findings notwithstanding, this report has several limitations. First, the study focused on hospitalized patients, thus it may have suffered a possible selection bias. Therefore, it is not clear how asymptomatic individuals with a positive nucleic acid test result or mild COVID 19 symptoms would respond to the proposed drugs. Second, given we included trials from the whole world, the majority of studies involved in this meta-analysis were basically on Chinese population, thus further universal studies should be reviewed. Meanwhile, the heterogeneity between studies included in this report is critical. Sources of heterogeneity included broad inclusion criteria, differences in clinical trial design, patient cooperation and measurement of outcome. Additionally, differences in individual samples with regard to severity of the disease, intervention time, age and gender composition may have contributed to heterogeneity. Thus given the significant heterogeneity between studies related to COVID-19, our findings should be interpreted with caution, random effect model notwithstanding¹³.

Despite these limitations, this network meta-analysis provides preliminary understanding on current COVID-19 treatment. In this report, we first conducted a comprehensive search in five databases including Cochrane, EmBase, PubMed and Web of Science, incorporating a relative a large sample of COVID 19 patients across the world. Although the reports on several drugs such as colchicine are limited, findings of this study may still provide a preliminary reflection of their safety and efficacy. In addition, selection, inclusion and data extraction were performed by two independent

researchers, thus minimized the selection bias and strengthened the reliability of the results. Additionally, all included studies were tested for publication bias through sensitivity analysis. The elaborate comparison of advantages of interventions against COVID 19 fully recognizes the inherent limitations of the method used in this study, the complexity of the patient population and the uncertainty caused by the choice of dosage and treatment.

So far, there is no safe and effective drug against COVID-19¹⁴. Meanwhile vaccination is seen as the most effective approach to control the spread of the viral. However, vaccine development is an entirely challenging process. Therefore, before successful development and popularization of effective vaccines against COVID 19, drugs remain the most effective method against the global pandemic¹⁵. Current therapeutics include antiviral drugs (remdesivir, ribavirin, favipiravir, lopinavir-ritonavir, arbidol etc.), antimalarial drugs (hydroxychloroquine and chloroquine), glucocorticoid (dexamethasone) and immune modulation drugs (interferon- β). Our results demonstrated favipiravir, an RNA-dependent RNA polymerase, may be a potential effective drug against COVID-19. Pathologically, coronavirus infection can induce an imbalance of immune regulation network via cytokine storm syndrome (CSS) and diffuse damage to target cells such as alveolar epithelial cells¹⁶. This results in acute respiratory distress syndrome (ARDS), septic shock, multiple organ dysfunction and even death. Notably, compared with placebos and standard care, immune modulation drugs such as interferon- β increases the mortality of patients with COVID 19 infection. One possible reason is that immune agents may aggravate the degree of immune imbalance, thus causing death. Our findings are thus instructive for selecting the most effective intervention of managing COVID-19 patients. When implemented, our findings will rationalize clinical decisions among COVID-19 patients.

Contributors

All authors collectively and equally participated in all aspects of the research, from preliminary literature assesment, data extraction, statistical analysis as well as writing and editing of the manuscript.

Declarations of Competing Interest

There are no conflicts on interest.

Acknowledgment

This study was supported by the National Natural Science Foundation of China (Nos. 81872823 and 82073782), the Double First-Class (CPU2018PZQ13, China) of the China Pharmaceutical University, the Shanghai Science and Technology Committee (No. 19430741500), the Key Laboratory of Modern Chinese Medicine Preparation of Ministry of Education of Jiangxi University of Traditional Chinese Medicine (TCM-201905, China), and the Start-up Grant from City University of Hong Kong (No. 9610472).

References

1. University JH. COVID-19 Map - Johns Hopkins Coronavirus Resource Center. <https://coronavirus.jhu.edu/map.html> (accessed October 5 2020).
2. Kim AHJ, Sparks JA, Liew JW, et al. A Rush to Judgment? Rapid Reporting and Dissemination of Results and Its Consequences Regarding the Use of Hydroxychloroquine for COVID-19. *Annals of Internal Medicine* 2020; **172**(12): 819-21.
3. Djulbegovic B, Guyatt G. Evidence-based medicine in times of crisis. *Journal of Clinical Epidemiology* 2020; **126**: 164-6.
4. Chaimani A, Higgins JPT, Mavridis D, Spyridonos P, Salanti G. Graphical tools for network meta-analysis in STATA. *PLoS One* 2013; **8**(10): e76654-e.
5. Carlin BP, Chib S. Bayesian Model Choice Via Markov Chain Monte Carlo Methods. *Journal of the Royal Statistical Society: Series B (Methodological)* 1995; **57**(3): 473-84.
6. Health NIo. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. <https://www.covid19treatmentguidelines.nih.gov>.
7. Olivo SA, Macedo LG, Gadotti IC, Fuentes J, Stanton T, Magee DJ. Scales to Assess the Quality of Randomized Controlled Trials: A Systematic Review. *Physical Therapy* 2008; **88**(2): 156-75.
8. Egger M, Smith GD, Phillips AN. Meta-analysis: Principles and procedures. *BMJ* 1997; **315**(7121): 1533.
9. Friedrich JO, Adhikari NK, Beyene J. Ratio of means for analyzing continuous outcomes in meta-analysis performed as well as mean difference methods. *J Clin Epidemiol* 2011; **64**(5): 556-64.
10. van Valkenhoef G, Tervonen T, Zwinkels T, de Brock B, Hillege H. ADDIS: A decision support system for evidence-based medicine. *Decision Support Systems* 2013; **55**(2): 459-75.
11. Garbolino P, Taroni F. Evaluation of scientific evidence using Bayesian networks. *Forensic Science International* 2002; **125**(2): 149-55.
12. Higgins JPT, Jackson D, Barrett JK, Lu G, Ades AE, White IR. Consistency and inconsistency in network meta-analysis: concepts and models for multi-arm studies. *Research Synthesis Methods* 2012; **3**(2): 98-110.
13. Turner RM, Jackson D, Wei Y, Thompson SG, Higgins JP. Predictive distributions for between-study heterogeneity and simple methods for their application in Bayesian meta-analysis. *Stat Med* 2015; **34**(6): 984-98.
14. Rome BN, Avorn J. Drug Evaluation during the Covid-19 Pandemic. *New England Journal of Medicine* 2020; **382**(24): 2282-4.
15. Dong L, Hu S, Gao J. Discovering drugs to treat coronavirus disease 2019 (COVID-19). *Drug Discoveries & Therapeutics* 2020; **14**(1): 58-60.
16. Channappanavar R, Perlman S. Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Seminars in Immunopathology* 2017; **39**(5): 529-39.
17. Beigel JH, Tomashek KM, Dodd LE, et al. Remdesivir for the Treatment of Covid-19 - Preliminary Report. *N Engl J Med* 2020.
18. Cao B, Wang Y, Wen D, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *N Engl J Med* 2020; **382**(19): 1787-99.
19. Cao Y, Wei J, Zou L, et al. Ruxolitinib in treatment of severe coronavirus disease 2019 (COVID-19): A multicenter, single-blind, randomized controlled trial. *J Allergy Clin Immunol* 2020; **146**(1): 137-46.e3.
20. Chen C, Zhang Y, Huang J, et al. Favipiravir versus Arbidol for COVID-19: A Randomized Clinical Trial. *medRxiv* 2020: 2020.03.17.20037432.
21. Chen C-P, Lin Y-C, Chen T-C, et al. A Multicenter, randomized, open-label, controlled trial to evaluate the efficacy and tolerability of hydroxychloroquine and a retrospective study in adult patients with mild to moderate Coronavirus disease 2019 (COVID-19). *medRxiv* 2020: 2020.07.08.20148841.

22. Chen L, Zhang Z-y, Fu J-g, et al. Efficacy and safety of chloroquine or hydroxychloroquine in moderate type of COVID-19: a prospective open-label randomized controlled study. *medRxiv* 2020: 2020.06.19.20136093.
23. Corral L, Bahamonde A, Arnaiz delas Revillas F, et al. GLUCOCOVID: A controlled trial of methylprednisolone in adults hospitalized with COVID-19 pneumonia. *medRxiv* 2020: 2020.06.17.20133579.
24. Davoodi L, Abedi SM, Salehifar E, et al. Febuxostat therapy in outpatients with suspected COVID-19: A clinical trial. *Int J Clin Pract* 2020: e13600.
25. Davoudi-Monfared E, Rahmani H, Khalili H, et al. Efficacy and safety of interferon beta-1a in treatment of severe COVID-19: A randomized clinical trial. *medRxiv* 2020: 2020.05.28.20116467.
26. Devereux SG, Giannopoulos G, Vrachatis DA, et al. Effect of Colchicine vs Standard Care on Cardiac and Inflammatory Biomarkers and Clinical Outcomes in Patients Hospitalized With Coronavirus Disease 2019: The GRECCO-19 Randomized Clinical Trial. *JAMA Netw Open* 2020; 3(6): e2013136.
27. Horby P, Lim WS, Emberson JR, et al. Dexamethasone in Hospitalized Patients with Covid-19 - Preliminary Report. *N Engl J Med* 2020.
28. Horby P, Mafham M, Linsell L, et al. Effect of Hydroxychloroquine in Hospitalized Patients with COVID-19: Preliminary results from a multi-centre, randomized, controlled trial. *medRxiv* 2020: 2020.07.15.20151852.
29. Huang M, Tang T, Pang P, et al. Treating COVID-19 with Chloroquine. *J Mol Cell Biol* 2020; 12(4): 322-5.
30. Hung IF-N, Lung K-C, Tso EY-K, et al. Triple combination of interferon beta-1b, lopinavir-ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomised, phase 2 trial. *The Lancet* 2020; 395(10238): 1695-704.
31. Li Y, Xie Z, Lin W, et al. An exploratory randomized controlled study on the efficacy and safety of lopinavir/ritonavir or arbidol treating adult patients hospitalized with mild/moderate COVID-19 (ELACOI). *medRxiv* 2020: 2020.03.19.20038984.
32. Mitjà O, Corbacho-Monné M, Ubals M, et al. Hydroxychloroquine for Early Treatment of Adults with Mild Covid-19: A Randomized-Controlled Trial. *Clin Infect Dis* 2020.
33. Ren Z, Luo H, Yu Z, et al. A Randomized, Open-Label, Controlled Clinical Trial of Azvudine Tablets in the Treatment of Mild and Common COVID-19, a Pilot Study. *Advanced Science* 2020; n/a(n/a): 2001435.
34. Skipper CP, Pastick KA, Engen NW, et al. Hydroxychloroquine in Nonhospitalized Adults With Early COVID-19: A Randomized Trial. *Ann Intern Med* 2020.
35. Tang W, Cao Z, Han M, et al. Hydroxychloroquine in patients with mainly mild to moderate coronavirus disease 2019: open label, randomised controlled trial. *BMJ* 2020; 369: m1849.
36. Wang Y, Zhang D, Du G, et al. Remdesivir in adults with severe COVID-19: a randomised, double-blind, placebo-controlled, multicentre trial. *Lancet* 2020; 395(10236): 1569-78.
37. Geleris J, Sun Y, Platt J, et al. Observational study of hydroxychloroquine in hospitalized patients with COVID-19. *New England Journal of Medicine* 2020; 382(25): 2411-8.
38. Spinner CD, Gottlieb RL, Criner GJ, et al. Effect of Remdesivir vs Standard Care on Clinical Status at 11 Days in Patients With Moderate COVID-19: A Randomized Clinical Trial. *JAMA* 2020.
39. Kim JW, Kim EJ, Kwon HH, et al. Lopinavir-ritonavir versus hydroxychloroquine for viral clearance and clinical improvement in patients with mild to moderate coronavirus disease 2019. *The Korean journal of internal medicine* 2020.
40. Cavalcanti AB, Zampieri FG, Rosa RG, et al. Hydroxychloroquine with or without Azithromycin in Mild-to-Moderate Covid-19. *The New England journal of medicine* 2020.
41. Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *International Journal of Antimicrobial Agents* 2020; 56(1).

42. Paccoud O, Tubach F, Baptiste A, et al. Compassionate use of hydroxychloroquine in clinical practice for patients with mild to severe Covid-19 in a French university hospital. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 2020.
43. Tong S, Su Y, Yu Y, et al. Ribavirin therapy for severe COVID-19: a retrospective cohort study. *International Journal of Antimicrobial Agents* 2020.
44. Xu P, Huang J, Fan Z, et al. Arbidol/IFN- α 2b therapy for patients with corona virus disease 2019: a retrospective multicenter cohort study. *Microbes and Infection* 2020; **22**(4-5): 200-5.

Study	Country	Male (%)	Average year		Number of subjects	Sample size		Type of research	Treatment		Outcome
			Experiment group	Controlled group		Experiment group	Controlled group		Experiment group	Controlled group	
Beigel ¹⁷	USA	64.3	58.6	59.2	1059	538	521	RCT	Remdesivir	Placebo	1234
Cao ¹⁸	China	60.3	58.0		199	94	100	RCT	Lopinavir–ritonavir	Standard care	1234
Cao ¹⁹	China	58.5	64.0	63.0	41	20	21	RCT	Ruxolitinib	Standard care	1234
Chen ²⁰	China	49.1	NR		236	120	116	RCT	Arbidol	Favipiravir	24
Chen ²¹	China	57.6	32.9		30	19	11	RCT	Hydroxychloroquine	Standard care	124
Chen ²²	China	45.5	42.2	44.5	48	18	18	RCT	Chloroquine	Hydroxychloroquine	124
Corral-Gudino ²³	Spain	58.0	69.0		85	56	29	RCT	Methylprednisolone	Standard care	24
Davoodi ²⁴	Iran	59.3	58.0	57.3	54	29	25	RCT	Febuxostat	Hydroxychloroquine	12
Davoudi-Monfared ²⁵	Iran	53.8	56.1	59.5	71	42	39	RCT	Interferonβ-1a	Standard care	1234
Deftereos ²⁶	Greece	60.0	65.0	63.0	105	50	55	RCT	Colchicine	Standard care	1234
Horby ²⁷	UK	64.0	66.1		6425	2104	4321	RCT	Dexamethasone	Standard care	123
Horby ²⁸	UK	62.0	65.2	65.4	4716	1561	3155	RCT	Hydroxychloroquine	Standard care	123
Huang ²⁹	China	59.1	41.5	53.0	22	10	12	RCT	Chloroquine	Lopinavir/Ritonavir	124
Hung ³⁰	China	52.0	51.0	52.0	127	86	41	RCT	Interferonβ-1b	Lopinavir/Ritonavir	134
Li ³¹	China	50.0	50.7	50.5	69	34	5	RCT	Lopinavir-ritonavir	Arbidol	124
Mitjà ³²	Spain	65.6	41.7	41.6	293	136	157	RCT	Hydroxychloroquine	Standard care	234
Ren ³³	China	NR	52.0	50.5	20	10	10	RCT	Azvudine	Standard care	124
Skipper ³⁴	US and Canada	42.0	41.0	39.0	423	212	211	RCT	Hydroxychloroquine	Placebo	34
Tang ³⁵	China	55.0	48.0	44.1	150	79	80	RCT	Hydroxychloroquine	Standard care	124
Wang ³⁶	China	56.0	66.0	64.0	237	158	79	RCT	Remdesivir	Placebo	1234
Geleris ³⁷	US	56	66.0	64.0	1376	811	565	Observational Study	Hydroxychloroquine	Standard care	3

Spinner ³⁸	Germany	61.0	58.0	57.0	393	193	200	RCT	Remdesivir	Standard care	234
Kim ³⁹	Korea	25.0	64.3		65	31	34	RCT	Lopinavir-ritonavir	Standard care	124
Cavalcanti ⁴⁰	Brazil	45.8	49.8	53.5	448	221	227	RCT	Hydroxychloroquine	Standard care	1234
Gautret ⁴¹	France	41.7	51.2	37.3	46	20	26	Open-label	Hydroxychloroquine	Standard care	2
Paccoud ⁴²	France	NR	66.0		84	38	46	Observational Study	Hydroxychloroquine	Standard care	234
Tong ⁴³	China	41.7	51.2	37.3	115	44	71	RCT	Ribavirin	Standard care	13
Xu ⁴⁴	China	57.7	53.2	50.9	141	70	71	retrospective	Interferon β -1b	Arbidol	12

Table 1. Characteristics of research objects. Outcomes: 1. average recovery days; 2. response rate; 3. mortality; 4. rate of adverse events.

Figure 2. Risk level evaluated results according the modified Jadad scale

Figure 3. The network of eligible comparisons for the four outcomes. (a). average recovery days; (b). response rate; (c). Mortality; (D). rate of adverse events. Each cycle represented an intervention drug, the width of the connection lines is proportional to the number of trials comparing every pair of treatments, and the size of every circle is proportional to the number of randomly assigned participants.

Average recovery days

Comparsion

Response rate

Arbidol	1.54 (-7.96, 10.55)	4.44 (-3.74, 12.48)	3.48 (-5.80, 12.60)	3.43 (-6.32, 13.11)	-0.99 (-7.80, 5.54)	-	5.80 (-1.87, 12.93)	2.94 (-2.60, 8.19)	4.51 (-5.53, 13.96)	3.21 (-6.21, 11.99)	5.21 (-2.03, 12.03)	-	-	4.47 (-2.86, 11.43)
2.26 (0.04, 150.20)	Azvdine	2.89 (-4.63, 10.47)	1.96 (-5.98, 9.92)	1.87 (-6.66, 10.44)	-2.46 (-13.73, 8.94)	-	4.31 (-2.06, 10.67)	1.40 (-5.95, 8.95)	2.98 (-5.67, 11.78)	1.61 (-6.60, 9.70)	3.71 (-3.19, 10.57)	-	-	2.94 (-2.89, 8.74)
0.26 (0.00, 4.89)	0.11 (0.00, 5.02)	Chloroquine	-0.95 (-8.32, 6.77)	-1.08 (-8.85, 7.16)	-5.41 (-15.77, 5.31)	-	1.38 (-3.08, 5.99)	-1.52 (-7.54, 4.78)	0.07 (-8.33, 8.33)	-1.27 (-8.75, 6.18)	0.77 (-3.88, 5.52)	-	-	0.01 (-4.88, 5.04)
2.01 (0.04, 86.21)	0.90 (0.01, 62.10)	7.74 (0.30, 602.40)	Dexamethasone	-0.07 (-8.38, 8.38)	-4.48 (-15.43, 6.98)	-	2.34 (-3.91, 8.44)	-0.59 (-7.70, 6.84)	1.06 (-7.64, 9.56)	-0.31 (-8.33, 7.47)	1.68 (-4.95, 8.35)	-	-	0.99 (-4.65, 6.45)
0.20 (0.00, 13.47)	0.09 (0.00, 7.11)	0.81 (0.01, 85.12)	0.11 (0.00, 7.12)	Colchicine	-4.42 (-16.37, 7.62)	-	2.43 (-4.37, 9.05)	-0.54 (-8.37, 7.33)	1.13 (-8.18, 10.26)	-0.25 (-8.92, 7.99)	1.77 (-5.66, 9.00)	-	-	1.06 (-5.18, 7.10)
0.73 (0.05, 11.11)	0.32 (0.00, 45.26)	2.73 (0.06, 397.35)	0.36 (0.00, 38.86)	3.71 (0.03, 855.76)	Favipiravir	-	6.80 (-3.50, 16.52)	3.88 (-4.75, 12.41)	5.47 (-6.54, 17.35)	4.13 (-7.51, 15.25)	6.16 (-3.66, 15.80)	-	-	5.53 (-4.59, 15.09)
1.35 (0.02, 55.97)	0.60 (0.00, 42.72)	5.32 (0.15, 343.95)	0.69 (0.01, 34.95)	6.79 (0.07, 954.07)	1.88 (0.01, 175.26)	Febuxostat	-	-	-	-	-	-	-	-
1.31 (0.08, 15.75)	0.57 (0.02, 13.82)	5.07 (0.54, 98.76)	0.67 (0.03, 11.51)	6.16 (0.20, 454.39)	1.81 (0.04, 68.56)	0.97 (0.06, 18.01)	Hydroxychloroquine	-2.96 (-7.84, 2.19)	-1.27 (-8.59, 5.78)	-2.66 (-8.96, 3.39)	-0.63 (-4.39, 3.30)	-	-	-1.37 (-3.98, 1.24)
1.03 (0.10, 10.89)	0.46 (0.01, 22.21)	3.95 (0.24, 211.99)	0.51 (0.01, 19.50)	5.05 (0.10, 636.54)	1.42 (0.04, 49.80)	0.75 (0.02, 43.05)	0.79 (0.07, 11.27)	Interferon β	1.59 (-6.61, 9.62)	0.25 (-7.35, 7.34)	2.34 (-2.39, 6.79)	-	-	1.57 (-3.20, 6.09)
1.60 (0.03, 76.14)	0.70 (0.01, 55.36)	6.34 (0.16, 646.66)	0.79 (0.01, 46.91)	7.87 (0.10, 1343.49)	2.17 (0.02, 267.10)	1.16 (0.02, 111.39)	1.23 (0.05, 33.55)	1.53 (0.03, 68.80)	Ruxolitinib	-1.37 (-10.15, 7.24)	0.70 (-6.85, 8.38)	-	-	-0.08 (-6.67, 6.65)
-	-	-	-	-	-	-	-	-	-	Ribavirin	2.02 (-4.56, 8.95)	-	-	1.30 (-4.27, 7.00)
1.18 (0.12, 12.56)	0.53 (0.01, 20.85)	4.64 (0.48, 138.43)	0.60 (0.02, 16.47)	5.72 (0.15, 500.30)	1.61 (0.05, 60.31)	0.89 (0.04, 33.62)	0.92 (0.15, 7.17)	1.16 (0.09, 14.79)	0.75 (0.02, 28.21)	-	Lopinavir–ritonavir	-	-	-0.72 (-4.51, 2.92)
1.12 (0.03, 45.92)	0.49 (0.01, 33.99)	4.38 (0.14, 376.99)	0.55 (0.01, 24.92)	5.46 (0.08, 702.17)	1.55 (0.01, 149.62)	0.83 (0.02, 58.70)	0.85 (0.05, 18.93)	1.10 (0.03, 40.23)	0.69 (0.01, 42.40)	-	0.93 (0.03, 26.93)	Remdesivir	-	-
1.62 (0.02, 104.99)	0.71 (0.01, 69.27)	6.29 (0.13, 828.46)	0.80 (0.01, 59.95)	7.97 (0.08, 1588.30)	2.23 (0.01, 316.37)	1.20 (0.01, 126.59)	1.23 (0.04, 46.97)	1.61 (0.02, 85.38)	1.00 (0.01, 86.33)	-	1.39 (0.03, 56.99)	1.45 (0.22, 10.14)	Placebo	-
2.31 (0.18, 30.25)	1.02 (0.04, 25.25)	9.15 (0.97, 259.14)	1.16 (0.08, 17.30)	11.23 (0.44, 721.36)	3.22 (0.07, 140.88)	1.75 (0.09, 44.38)	1.79 (0.64, 6.24)	2.27 (0.21, 22.90)	1.44 (0.08, 29.36)	-	1.97 (0.30, 11.87)	2.09 (0.14, 34.21)	1.45 (0.05, 42.46)	Standard care

Mortality
 Comparison
 Rate of adverse events

Arbidol	-	-	-	-	-	-	-	-	-	-	-	-	-
8.96E+18 (19.11,1.17E+61)	Azvudine	-	-	-	-	-	-	-	-	-	-	-	-
0.23 (0.00, 39.34)	0.00 (0.00, 0.01)	Chloroquine	-	-	-	-	-	-	-	-	-	-	-
0.35 (0.00, 242.99)	0.00 (0.00, 0.02)	1.53 (0.01, 342.65)	Colchicine	-	0.18 (0.00, 6.08)	0.05 (0.00, 3.57)	0.00 (0.00, 0.01)	0.08 (0.00, 4.87)	0.11 (0.00, 6.36)	-	0.16 (0.00, 4.66)	0.16 (0.00, 6.00)	0.14 (0.01, 3.07)
0.66 (0.01, 28.69)	0.00 (0.00, 0.05)	2.81 (0.01, 1424.20)	1.84 (0.00, 1723.16)	Favipiravir	-	-	-	-	-	-	-	-	-
0.28 (0.00, 35.04)	0.00 (0.00, 0.01)	1.20 (0.06, 27.68)	0.78 (0.01, 40.61)	0.42 (0.00, 195.39)	Hydroxychloroquine	0.31 (0.01, 8.75)	0.00 (0.00, 0.05)	0.46 (0.02, 13.17)	0.60 (0.03, 15.60)	-	0.90 (0.10, 7.09)	0.90 (0.09, 7.11)	0.81 (0.13, 5.34)
0.59 (0.00, 289.59)	0.00 (0.00, 0.03)	2.67 (0.03, 406.82)	1.68 (0.01, 388.53)	0.89 (0.00, 1438.31)	2.20 (0.05, 151.79)	Interferon β	0.00 (0.00, 0.16)	1.53 (0.03, 63.67)	1.96 (0.04, 85.41)	-	3.06 (0.09, 60.93)	2.98 (0.08, 74.32)	2.62 (0.16, 41.21)
1.11 (0.00, 750.62)	0.00 (0.00, 0.06)	4.85 (0.03, 918.30)	3.10 (0.01, 667.40)	1.66 (0.00, 3304.08)	4.05 (0.06, 303.23)	1.85 (0.01, 304.47)	Ruxolitinib	1.21E+13 (7.62,4.94E+45)	1.42E+13 (11.57,6.51E+45)	-	2.23E+13 (21.6,48E+45)	2.24E+13 (20.03,7.78E+45)	2.00E+13 (17.9,6.01E+45)
-	-	-	-	-	-	-	-	Ribavirin	1.27 (0.03, 54.58)	-	1.98 (0.06, 41.30)	1.97 (0.05, 46.03)	1.73 (0.11, 25.65)
0.30 (0.01, 14.92)	0.00 (0.00, 0.01)	1.29 (0.06, 26.99)	0.84 (0.01, 57.63)	0.45 (0.00, 106.21)	1.07 (0.07, 10.52)	0.48 (0.00, 17.97)	0.26 (0.00, 21.98)	-	Lopinavir–ritonavir	-	1.48 (0.06, 29.55)	1.50 (0.05, 34.79)	1.32 (0.10, 18.72)
11.52 (0.04, 5264.32)	0.00 (0.00, 0.59)	46.85 (0.38, 9098.12)	29.85 (0.12, 6118.34)	17.04 (0.02, 25155.95)	40.76 (0.70, 2803.29)	18.97 (0.08, 3116.64)	10.02 (0.04, 2333.94)	-	38.00 (0.51, 4740.98)	Methylprednisolone	-	-	-
1.42 (0.01, 448.48)	0.00 (0.00, 0.06)	6.02 (0.11, 497.49)	4.03 (0.02, 444.81)	2.13 (0.00, 2494.74)	5.20 (0.23, 126.38)	2.39 (0.02, 204.01)	1.25 (0.01, 180.37)	-	4.81 (0.13, 276.20)	0.13 (0.00, 15.74)	Remdesivir	1.01 (0.18, 5.11)	0.8 (0.19, 6.13)
1.08 (0.01, 328.49)	0.00 (0.00, 0.05)	4.61 (0.08, 348.63)	2.99 (0.02, 334.22)	1.64 (0.00, 1415.90)	3.76 (0.20, 74.04)	1.77 (0.01, 157.84)	0.95 (0.01, 128.64)	-	3.62 (0.10, 201.66)	0.10 (0.00, 11.12)	0.74 (0.07, 7.72)	Placebo	0.88 (0.13, 8.89)
1.89 (0.02, 283.63)	0.00 (0.00, 0.07)	8.27 (0.37, 274.00)	5.39 (0.11, 222.41)	2.90 (0.01, 1554.09)	6.87 (1.52, 41.44)	3.23 (0.07, 123.15)	1.68 (0.03, 90.21)	-	6.45 (0.66, 114.39)	0.17 (0.00, 8.24)	1.33 (0.07, 29.59)	1.78 (0.10, 42.79)	Standard care

Table 2. Directly compare the results of 14 drugs, placebo and standard therapies. Compared with the row definition process, the data is the SMD or RR (95% CI) of the column definition process. In the average recovery day, if the SMD is greater than 1, the column definition processing takes precedence. For the remission rate, mortality and the incidence of adverse events, RR less than 0 is beneficial to define the treatment. In order to get the reverse comparison of RR, you need to take the inverse number. Important results are shown in bold.

Figure 4. Rank probability stacked graph of four outcomes in a percentage scale. (a). average recovery days; (b). response rate; (c). Mortality; (D). rate of adverse events.

	Average recovery days		Response rate		Mortality		Rate of adverse events	
	Prob.	Rank	Prob.	Rank	Prob.	Rank	Prob.	Rank
Arbidol	0.18	2	0.01	10	-	-	0	3
Azvudine	0.15	3	0.02	8	-	-	0.97	1
Chloroquine	0.01	9	0.28	2	-	-	0	3
Colchicine	0.06	4	0.42	1	0	2	0	3
Dexamethasone	0.04	7	0.01	10	-	-	-	-
Favipiravir	0.46	1	0.11	3	-	-	0	3
Febuxostat	-	-	0.04	4	-	-	-	-
Hydroxychloroquine	0	10	0	11	0	2	0	3
Interferon β	0.01	8	0.02	8	0	2	0	3
Lopinavir–ritonavir	0	10	0	11	0	2	0	3
Methylprednisolone	-	-	-	-	-	-	0.02	2
Remdesivir	-	-	0.03	5	0	2	0	3
Ribavirin	0.05	5	-	-	0	2	-	-
Ruxolitinib	0.04	6	0.03	5	0.99	1	0	3
Placebo	-	-	0.03	5	0	2	0	3
Standard care	0	10	0	11	0	2	0	3

Table 3. Rank probability table of four outcomes