Projections and fractional dynamics of COVID-19 with optimal control analysis

Khondoker Nazmoon Nabi^{a,*}, Pushpendra Kumar^b, Vedat Suat Erturk^c

^aDepartment of Mathematics, Bangladesh University of Engineering and Technology (BUET), Dhaka, Bangladesh

^bDepartment of Mathematics and Statistics, School of Basic and Applied Sciences, Central University of Punjab, Bathinda, Punjab-151001, India

^cDepartment of Mathematics, Ondokuz Mayıs University, Atakum-55200, Samsun, Turkey

Abstract

When the entire world is eagerly waiting for a safe, effective and widely available COVID-19 vaccine, unprecedented spikes of new cases are evident in numerous countries. To gain a deeper understanding about the future dynamics of COVID-19, a compartmental mathematical model has been proposed in this paper incorporating all possible non-pharmaceutical intervention policies. Model parameters have been calibrated using sophisticated trust-region-reflective algorithm and short-term projection results have been illustrated for Argentina, Bangladesh, Brazil, Colombia and India. Control reproduction numbers ($\mathcal{R}_c$) have been calculated in order to get insights about the current epidemic scenario in the above-mentioned countries. Forecasting results depict that the aforesaid countries are having downward trends in daily COVID-19 cases. However, it is highly recommended to use efficacious face coverings and maintain strict physical distancing, as the pandemic is not over in any country. Global sensitivity analysis enlightens the fact that efficacy of face coverings is the most significant parameter, which could significantly control the transmission dynamics of the novel coronavirus compared to other non-pharmaceutical measures. In addition, reduction in effective contact rate with isolated patients is also essential in bringing down the epidemic threshold ($\mathcal{R}_c$) below unity. All necessary graphical simulations have been performed with the help of Caputo-Fabrizio fractional derivatives. In addition, optimal control problem for fractional system has been designed and the existence of unique solution has also been showed by using Picard-Lindelof technique. Finally, the unconditionally stability of the given fractional numerical technique has been proved.

Keywords: COVID-19, mathematical model, calibration, optimal control, Caputo-Fabrizio fractional derivative

2010 MSC: 26A33, 37N25, 92C60, 92D30

*Corresponding author: khnabi@math.buet.ac.bd (Khondoker Nazmoon Nabi)

1. Introduction

To date, as there is no world-wide accepted vaccine that can provide full immunity to the human body against COVID-19, non-pharmaceutical intervention strategies are the realistic and effective solutions to control second wave of the pandemic. However, around 40 different coronavirus vaccines are experiencing critical clinical trials and nine already in the final stage of testing on thousands of people. Importantly, a leading vaccine candidate developed by the University of Oxford is already in an advanced stage of testing and it has been found in trials that it can trigger an immune response [1]. Generally, an effective vaccine would take years, if not decades, to develop. As research in this field is happening at a breakneck speed, scientists believe that an effective vaccine is likely to become widely available by mid-2021. Lack of transparency could be a vital issue in upcoming days and a false sense of security could evolve among general people if the vaccine doesn't work effectively.

Mathematical models can always provide considerable insights of the transmission dynamics and complexities of any infectious diseases, which eventually help government officials design overall epidemic planning. Importantly, mathematical analysis always plays a notable role in making vital public health decisions, resource allocation and implementation of social distancing measures and other non-pharmaceutical interventions. From the beginning of the COVID-19 outbreak, mathematicians and researchers are working relentlessly and have already done tremendous contributions in limiting the spread of the coronavirus in different parts of the world [2, 3, 4, 5, 6]. In an early contribution, Ferguson *et al.* [2] showed the impact of different non-pharmaceutical intervention strategies on COVID-19 mortality by developing an agent-based model. In another study, Ngonghala *et al.* showed that effective and comprehensive usage of face coverings can significantly limit the spread of the virus and reduce the COVID-induced mortality in different states of the USA in general in the absence of community lockdown measures and stringent social distancing practice. On the other hand, Nabi [5] projected the future dynamics of COVID-19 for various COVID-19 hotspots by proposing a compartmental mathematical model and concluded that early relaxation of lockdown measures and social distancing could bring a second wave in no time. As a matter of reality, inhabitants in several countries compelled to violate containment measures due to prolonged lockdown measures and severe economic recession [4]. Netherlands having one of the best health care systems in the world, is grappling with continuous spikes in daily cases due to aversion to masks.

In this study, in the absence of a safe, effective and world-wide approved vaccine, a new compartmental mathematical COVID-19 model has been designed incorporating all possible non-pharmaceutical intervention strategies such as wearing face coverings, social distancing, home or self-quarantine and self or institutional isolation. In addition, the impacts of different interventions scenarios have been analysed rigorously. The aim of this work is to project the future dynamics of COVID-19 outbreak in five countries namely Argentina, Bangladesh, Brazil, Colombia and India which are worst-hit countries in the world. Estimation of parameters

has been performed by using real-time data, followed by a projection of the evolution of the disease. Global sensitivity analysis is applied to determine the influential mechanisms in the model that drive the transmission dynamics of the disease. For fractional simulations, we used the well known non-integer order derivative called Caputo-Fabrizio (CF) fractional derivative. Since last few decades, there are so many epidemic models have been solved by non-integer order derivatives. Recently some applications of non-integer order derivatives in mathematical epidemiology can be seen from [7, 8, 9, 10, 11]. There are so many research papers have been come to study the outbreaks of coronavirus, in which some are [12, 13, 14, 15, 16, 17, 18, 19, 20, 21]. We performed the optimal control problem in CF derivative sense and provided the existence of unique solution by well-known technique named as Picard-Lindelof technique. We also proved the unconditionally stability of the given fractional numerical technique. We used the numerical data of all five given countries and perform the all necessary graphical simulations.

The entire chapter is organized as follows. Materials and methods are presented in Section 2. Section 3 is solely devoted to asymptotic stability of the proposed model. In section 4, estimation of model parameters and projection results have been discussed using daily COVID-19 data of Argentina, Bangladesh, Brazil, Colombia and India. In section 5, to quantify the impact of different model mechanisms, LHS-PRCC global sensitivity scheme has been performed. In section 6, numerical and graphical simulations have been illustrated using Caputo-Fabrizio fractional derivatives. Later, optimal control problem has been designed in fractional sense in section 7. The chapter ends with some insightful findings and strategies, which could significantly control the transmission dynamics of COVID-19.

2. Materials and methods

2.1. Model formulation

A compartmental mathematical has been designed to describe the transmission dynamics of the COVID-19 incorporating all possible real-life interactions. Considering different infection status, the entire human population (denoted by $N(t)$ at time t) has been stratified into nine mutually-exclusive compartments of susceptible individuals ($S(t)$), early-exposed individuals ($E_1(t)$), pre-symptomatic individuals ($E_2(t)$), symptomatically-infectious ($I(t)$), asymptotically-infectious or infectious individuals with mild-symptoms ($A(t)$), quarantined infectious ($Q(t)$), hospitalised or isolated individuals ($L(t)$), recovered individuals ($R(t)$), disease-induced death cases ($D(t)$). Hence,

$$N(t) = S(t) + E_1(t) + E_2(t) + I(t) + A(t) + Q(t) + L(t) + R(t) + D(t)$$

The following assumptions have been considered to formulate the above-mentioned model.

- Vital dynamic (birth and natural deaths) has been ignored as the main objective of this study is to observe the short-term dynamics of COVID-19 pandemic.

Figure 1: Flow diagram of the COVID-19 transmission dynamics.

- Recovered individuals are immune to the disease which means that they cannot get reinfected. However, in a recent study of Tillett *et al.* [22], genomic evidence for reinfection with SARS-CoV-2 has been found. Nevertheless, it is important to note this singular finding does not provide generalisability of this phenomenon and hence it is not considered in our study.

The flow diagram of the proposed model is illustrated in Figure 1, where susceptible individuals can become infected by an effective contact with individuals in the pre-symptomatic ($E_2(t)$), symptomatically-infectious ($I(t)$), asymptomatically-infectious ($A(t)$), quarantined-infectious ($Q(t)$) and isolated-infectious ($L(t)$). Effective contact rates are λ_{E_2} , λ_I , λ_A , λ_Q , and λ_L respectively and the expressions are defined in (2). Importantly, the compartment $E_1(t)$ consists of early-infected individuals who are still not infectious, whereas the individuals in pre-symptomatic cohort $E_2(t)$ have the capability of transmitting coronavirus before the end of the disease incubation period. A proportion of individuals in newly-exposed compartment ($E_1(t)$) progress to pre-symptomatic class ($E_2(t)$) at a rate κ_1 . After the completion of disease mean incubation period, at a rate $\rho\kappa_2$, a fraction of individuals who have clear clinical symptoms of COVID-19 progress to $I(t)$ compartment. Individuals in $E_2(t)$ class who do not have any clear symptoms progress to $A(t)$ class at a rate $(1 - \rho)\kappa_2$. Pre-symptomatic individuals are assumed to be self-quarantined at a rate q . With the help of diagnostic or surveillance testing approaches, symptomatically-infectious individuals and asymptomatically-infectious individuals are brought under institutional or home isolation at rates τ_A and τ_I respectively. Moreover, the parameter $\gamma_I(\gamma_A)(\gamma_Q)(\gamma_L)$ represents the recovery rate for individuals in the $I(A)(Q)(L)$ class. Finally, the disease-induced mortality rate for individuals in the $I(Q)(L)$ compartment is defined by the parameter $\delta_I(\delta_Q)(\delta_L)$. Considering all the above-mentioned interactions, the transmission dynamics of COVID-19 can

be described by the following system of nonlinear ordinary differential equations.

$$\left\{ \begin{array}{l} \frac{dS}{dt} = -(\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2}) S, \\ \frac{dE_1}{dt} = (\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2}) S - \kappa_1 E_1, \\ \frac{dE_2}{dt} = \kappa_1 E_1 - (\kappa_2 + q) E_2, \\ \frac{dI}{dt} = \rho \kappa_2 E_2 - (\tau_I + \gamma_I + \delta_I) I, \\ \frac{dA}{dt} = (1 - \rho) \kappa_2 E_2 - (\tau_A + \gamma_A) A, \\ \frac{dQ}{dt} = q E_2 - (\gamma_Q + \delta_Q) Q, \\ \frac{dL}{dt} = \tau_I I + \tau_A A - (\delta_L + \gamma_L) L, \\ \frac{dR}{dt} = \gamma_I I + \gamma_A A + \gamma_Q Q + \gamma_L L, \\ \frac{dD}{dt} = \delta_I I + \delta_L L + \delta_Q Q. \end{array} \right. \quad (1)$$

where the forces of infection are defined below

$$\left\{ \begin{array}{l} \lambda_{E_2} = \beta_{E_2} (1 - m\zeta) \frac{E_2}{N}, \\ \lambda_I = \beta_I (1 - m\zeta) \frac{I}{N}, \\ \lambda_A = \beta_A (1 - m\zeta) \frac{A}{N}, \\ \lambda_Q = \beta_Q (1 - m\zeta) \frac{Q}{N}, \\ \lambda_L = \beta_L (1 - m\zeta) \frac{L}{N}, \end{array} \right. \quad (2)$$

63 The parameters are described in Table 1.

64 We set $x = (S, E_1, E_2, I, A, Q, L, R, D)'$ the vector of state variable, Let $f : \mathbb{R}^9 \rightarrow \mathbb{R}^9$ the the right hand side
65 of system (1), which is a continuously differentiable function on $\mathbb{R}^9$. According to [23, Theorem III.10.VI], for

Parameter	Description
β_I (β_A) (β_Q) (β_L) (β_{E_2})	Effective contact rate (a measure of physical or social distancing)
m	Proportion of individuals who use face coverings or surgical masks
ζ	Efficacy of face coverings at reducing outward transmission by infected individuals as well as preventing acquisition
κ_1	Rate of progression from early-exposed class ($E_1(t)$) to pre-symptomatic class ($E_2(t)$)
$\rho\kappa_2$	Rate of progression from pre-symptomatic class ($E_2(t)$) to symptomatically-infectious class ($I(t)$)
$(1 - \rho)\kappa_2$	Rate of progression from pre-symptomatic class ($E_2(t)$) to asymptotically-infectious class ($A(t)$)
q	Confinement efficacy
τ_I	Rate of self or institutional isolation for symptomatically-infectious patients
τ_A	Rate of isolation for asymptotically-infectious patients
γ_I	Recovery rate for symptomatically-infectious patients
γ_A	Recovery rate for asymptotically-infectious individuals
γ_Q	Recovery rate for quarantined-infectious individuals
γ_L	Recovery rate for isolated or hospitalised individuals
δ_I	Disease-induced death rate for symptomatically-infectious individuals
δ_L	Disease-induced death rate for isolated individuals
δ_Q	Disease-induced death rate for quarantined-infectious

Table 1: Model parameters and meaning

any initial condition in Ω , a unique solution of (1) exists, at least locally, and remains in Ω for its maximal interval of existence [23, Theorem III.10.XVI]. Hence, model (1) is biologically well-defined.

2.2. Data sources

From the beginning of the COVID-19 outbreak, Center of Disease Control and Prevention (CDC) is providing authoritative and genuine data of daily confirmed COVID-19 cases. Daily confirmed data of five different countries named Argentina, Bangladesh, Brazil, Colombia and India have been compiled using that data repository. Johns Hopkins University Center for Systems Science and Engineering (JHU CSSE) is carefully maintaining the data repository supported by ESRI Living Atlas Team and the Johns Hopkins University Applied Physics Lab (JHU APL). The repository is really simple to use and publicly available [24].

3. Mathematical analysis

3.1. Asymptotic stability of disease-free equilibria

The disease-free equilibrium point denoted by x_0 can be defined as follows:

$$x_0 = (S_0, 0, 0, 0, 0, 0, 0, 0)' = (N_0, 0, 0, 0, 0, 0, 0, 0)'$$

Using notations in [25], matrices F and V for the new infection terms and the remaining transfer terms are, respectively, given by

$$F = \begin{pmatrix} 0 & \beta_{E_2}(1-m\zeta)\frac{S_0}{N_0} & \beta_I(1-m\zeta)\frac{S_0}{N_0} & \beta_A(1-m\zeta)\frac{S_0}{N_0} & \beta_Q(1-m\zeta)\frac{S_0}{N_0} & \beta_L(1-m\zeta)\frac{S_0}{N_0} \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix},$$

$$V = \begin{pmatrix} \kappa_1 & 0 & 0 & 0 & 0 & 0 \\ -\kappa_1 & \kappa_2 + q & 0 & 0 & 0 & 0 \\ 0 & -\rho\kappa_2 & \tau_I + \gamma_I + \delta_I & 0 & 0 & 0 \\ 0 & -(1-\rho)\kappa_2 & 0 & \tau_A + \gamma_A & 0 & 0 \\ 0 & -q & 0 & 0 & \gamma_Q + \delta_Q & 0 \\ 0 & 0 & -\tau_I & -\tau_A & 0 & \gamma_L + \delta_L \end{pmatrix}.$$

(3)

Then, the control reproduction ratio is defined, following [26, 25], as the spectral radius of the next generation matrix, FV^{-1} :

$$\mathcal{R}_c = \rho(FV^{-1}) = \mathcal{R}_{E_2} + \mathcal{R}_I + \mathcal{R}_A + \mathcal{R}_Q + \mathcal{R}_L$$

(4)

where,

$$\begin{aligned} \mathcal{R}_{E_2} &= \frac{\beta_{E_2}(1-m\zeta)}{\kappa_2 + q}, \\ \mathcal{R}_I &= \frac{\rho\kappa_2\beta_I(1-m\zeta)}{(\kappa_2 + q)(\gamma_I + \tau_I + \delta_I)}, \\ \mathcal{R}_A &= \frac{\kappa_2(1-\rho)\beta_A(1-m\zeta)}{(\kappa_2 + q)(\gamma_A + \tau_A)}, \\ \mathcal{R}_Q &= \frac{q\beta_Q(1-m\zeta)}{(\kappa_2 + q)(\gamma_Q + \delta_Q)}, \\ \mathcal{R}_L &= \frac{\kappa_2\beta_L(1-m\zeta) [\tau_A(\gamma_I + \delta_I)(1-\rho) + \tau_I(\rho\gamma_A + \tau_A)]}{(\kappa_2 + q)(\gamma_L + \delta_L)(\gamma_A + \tau_A)(\gamma_I + \tau_I + \delta_I)}, \end{aligned}$$

where $\rho(\cdot)$ represents the spectral radius operator.

The formula for control reproduction number has been formulated. Indeed, the insightful epidemic threshold, $\mathcal{R}_c$ calculates the average number of new secondary COVID-19 cases generated by a COVID-19 positive individual in a population a portion susceptible people are using effective face coverings. Different non-pharmaceutical measures are acting as control measures which lead to bring down $\mathcal{R}_c$ under unity [25]. Hence, we claim the following result followed by a direct consequence of the next generation operator method [25, Theorem 2]. where $\rho(\cdot)$ represents the spectral radius operator. The insightful epidemic threshold, $\mathcal{R}_0$ calculates the average number of new secondary COVID-19 infections generated by an COVID-19 positive patient in a completely susceptible population. The control of COVID-19 pandemic passes by the application of some control measures which contribute to decrease until $\mathcal{R}_0$ less than one [25]. Hence we claim the following result.

Theorem 1. The COVID-19 transmission dynamics is influenced by the basic reproduction number $\mathcal{R}_0$ as follows:

1. If $\mathcal{R}_0 < 1$, then a sufficiently small flow of infected individuals will not generate an outbreak of the COVID-19, i.e the disease-equilibrium $\mathcal{E}_0$ is locally asymptotically stable on ω .
2. If $\mathcal{R}_0 > 1$, then a sufficiently small flow of infected individuals will generate an outbreak of the COVID-19, and the disease-equilibrium $\mathcal{E}_0$ is unstable.

Lemma 1. If $\mathcal{R}_c < 1$, the disease-free equilibrium x_0 is locally asymptotically stable and unstable if $\mathcal{R}_c > 1$.

Remark 2. Lemma 1 implies that if $\mathcal{R}_c < 1$, then a sufficiently small flow of infected individuals will not generate an outbreak of COVID-19, whereas for $\mathcal{R}_c > 1$, epidemic curve reaches a peak by growing exponentially and then decreases to zero as $t \rightarrow \infty$.

The better control of the COVID-19 can be established by the fact that the DFE x_0 is globally asymptotically stable (GAS). In this context, we claim the following result.

Theorem 3. if $\mathcal{R}_c < 1$, then the manifold, $\mathcal{W}$, of disease-free equilibrium points of the model (1) is GAS in $\mathcal{D}$.

In the absence of use of face coverings, i.e. $m = 0$, $\mathcal{R}_c$ converges to the basic reproduction number, $\mathcal{R}_0$. Now, we will study the global stability of the disease-free equilibrium whenever the basic reproduction number is less than one ($\mathcal{R}_c < 1$). For this, we use the following Lyapunov function

$$\mathcal{L} = a_1 E_1 + a_2 E_2 + a_3 I + a_4 A + a_5 Q + a_6 L. \quad (5)$$

By deriving this function along the trajectories of the system (1), we obtain

$$\begin{aligned}\dot{\mathcal{L}} &= a_1 \dot{E}_1 + a_2 \dot{E}_2 + a_3 \dot{I} + a_4 \dot{A} + a_5 \dot{Q} + a_6 \dot{L} \\ &= a_1 [(\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2}) S - \kappa_1 E_1] \\ &\quad + a_2 [\kappa_1 E_1 - (\kappa_2 + q) E_2] + a_3 [\rho \kappa_2 E_2 - (\tau_I + \gamma_I + \delta_I) I] \\ &\quad + a_4 [(1 - \rho) \kappa_2 E_2 - (\tau_A + \gamma_A) A] + a_5 [q E_2 - (\gamma_Q + \delta_Q) Q] \\ &\quad + a_6 [\tau_I I + \tau_A A - (\delta_L + \gamma_L) L]\end{aligned}\tag{6}$$

Since $\frac{S}{N} \leq 1$, we obtain

$$\begin{aligned}\dot{\mathcal{L}} &\leq a_1 [\beta_I(1 - m\zeta)I + \beta_A(1 - m\zeta)A + \beta_Q(1 - m\zeta)Q + \beta_L(1 - m\zeta)L + \beta_{E_2}(1 - m\zeta)E_2 - \kappa_1 E_1] \\ &\quad + a_2 [\kappa_1 E_1 - (\kappa_2 + q) E_2] + a_3 [\rho \kappa_2 E_2 - (\tau_I + \gamma_I + \delta_I) I] \\ &\quad + a_4 [(1 - \rho) \kappa_2 E_2 - (\tau_A + \gamma_A) A] + a_5 [q E_2 - (\gamma_Q + \delta_Q) Q] \\ &\quad + a_6 [\tau_I I + \tau_A A - (\delta_L + \gamma_L) L] \\ &= (-a_1 \kappa_1 + a_2 \kappa_1) E_1 \\ &\quad + [a_1 \beta_{E_2}(1 - m\zeta) - a_2 (\kappa_2 + q) + a_3 \rho \kappa_2 + a_4 (1 - \rho) \kappa_2 + a_5 q] E_2 \\ &\quad + [a_1 \beta_I(1 - m\zeta) - a_3 (\tau_I + \gamma_I + \delta_I) + a_6 \tau_I] I \\ &\quad + [a_1 \beta_A(1 - m\zeta) - a_4 (\tau_A + \gamma_A) + a_6 \tau_A] A \\ &\quad + [a_1 \beta_Q(1 - m\zeta) - a_5 (\gamma_Q + \delta_Q)] Q + [a_1 \beta_L(1 - m\zeta) - a_6 (\delta_L + \gamma_L)] L\end{aligned}\tag{7}$$

We choose a_i , $i = 1, 2, \dots, 6$, such that coefficients of E_1 , I , A , Q , and L become zero. That is

$$\begin{cases} -a_1 \kappa_1 + a_2 \kappa_1 &= 0 \\ a_1 \beta_I(1 - m\zeta) - a_3 (\tau_I + \gamma_I + \delta_I) + a_6 \tau_I &= 0 \\ a_1 \beta_A(1 - m\zeta) - a_4 (\tau_A + \gamma_A) + a_6 \tau_A &= 0 \\ a_1 \beta_Q(1 - m\zeta) - a_5 (\gamma_Q + \delta_Q) &= 0 \\ a_1 \beta_L(1 - m\zeta) - a_6 (\delta_L + \gamma_L) &= 0 \end{cases}\tag{8}$$

which the non-zero solution is given by

$$\begin{cases} a_6 &= \frac{a_1 \beta_L(1 - m\zeta)}{\delta_L + \gamma_L} \\ a_5 &= \frac{a_1 \beta_Q(1 - m\zeta)}{\delta_Q + \gamma_Q} \\ a_4 &= \frac{a_1 \beta_A(1 - m\zeta) + a_6 \tau_A}{\tau_A + \gamma_A} \\ a_3 &= \frac{a_1 \beta_I(1 - m\zeta) + a_6 \tau_I}{\tau_I + \delta_I + \gamma_I} \\ a_2 &= a_1 \end{cases}\tag{9}$$

Plugging (9) into (7) gives

$$\begin{aligned} \dot{\mathcal{L}} \leq a_1(\kappa_2 + q) & \left[\frac{\beta_{E_2}(1 - m\zeta)}{\kappa_2 + q} + \frac{\rho\kappa_2\beta_I(1 - m\zeta)}{(\kappa_2 + q)(\gamma_I + \tau_I + \delta_I)} + \frac{\kappa_2(1 - \rho)\beta_A(1 - m\zeta)}{(\kappa_2 + q)(\gamma_A + \tau_A)} \right. \\ & \left. + \frac{q\beta_Q(1 - m\zeta)}{(\kappa_2 + q)(\gamma_Q + \delta_Q)} + \frac{\kappa_2\beta_L(1 - m\zeta) [\tau_A(\gamma_I + \delta_I)(1 - \rho) + \tau_I(\rho\gamma_A + \tau_A)]}{(\kappa_2 + q)(\gamma_L + \delta_L)(\gamma_A + \tau_A)(\gamma_I + \tau_I + \delta_I)} - 1 \right] E_2. \end{aligned} \quad (10)$$

Setting $a_1 = \frac{1}{\kappa_2 + q}$, we finally obtain

$$\dot{\mathcal{L}} \leq (\mathcal{R}_{E_2} + \mathcal{R}_I + \mathcal{R}_A + \mathcal{R}_Q + \mathcal{R}_L - 1) E_2 = (\mathcal{R}_c - 1) E_2. \quad (11)$$

From (11), it follows that $\dot{\mathcal{L}} < 0$ if $\mathcal{R}_c < 1$, and $\dot{\mathcal{L}} = 0$ if and only if $E_1 = E_2 = I = A = Q = L = 0$. Therefore, $\mathcal{L}$ is a Lyapunov function for system (1). Moreover, the maximal invariant set contained in $\{(S(t), E_1(t), E_2(t), I(t), A(t), Q(t), L(t), R(t), D(t)) \in \Omega : \dot{\mathcal{L}} = 0\}$ is the continuum of the disease-free equilibrium ($\mathcal{E}_0$). Thus, from Lyapunov theory, we deduce that the disease-free equilibrium $\mathcal{E}_0$ is GAS if $\mathcal{R}_c < 1$. Hence, it follows, by the LaSalle's Invariance Principal, that the continuum of disease-free equilibria of the model (1) is a stable global attractor in Ω whenever $\mathcal{R}_c \leq 1$. The previous analysis can be summarized as follows:

Theorem 4. If $\mathcal{R}_c \leq 1$, then the disease-free equilibrium $\mathcal{E}_0$ is globally asymptotically stable on Ω .

4. Model calibration and forecasting

The model (1) calibration has been performed using a newly developed optimization algorithm based on trust-region-reflective (TRR) algorithm, which can be regarded as an evolution of Levenberg-Marquardt algorithm [5]. This robust optimization procedure can be used effectively for solving nonlinear least-squares problems. This algorithm has been implemented using the **lsqcurvefit** function, which is available in the Optimization Toolbox in MATLAB. Necessary model parameters have been estimated using this optimization technique. Daily infected cases data have been collected from a trusted data repository, which is available online. A 7-day moving average of the daily reported cases has been used for our model calibration due to moderate volatile nature of real data. It has been observed that the number of daily testing in Argentina, Bangladesh, Brazil, Colombia and India have been really inconsistent. With an aim to capture the real outbreak scenario, the 7-day moving average has been used in this regard.

4.1. Argentina

On March 3, 2020, Argentina registered its first COVID-19 case. As a consequence, government officials immediately deploy strict lockdown, quarantine and isolation measures to curb the spread of the novel

Figure 2: Fitting performance of the model for daily infected cases in Argentina from March 04 to November 11, 2020.

Figure 3: Fitting performance of the model for cumulative infected cases in Argentina from March 04 to November 11, 2020.

coronavirus in the community. Due to unbearable economic crisis, inhabitants of the country have already started violating staying at home and quarantine orders. The country could face a second wave of infection in near future in the absence of non-pharmaceutical intervention measures. The robustness of our model fitting performance has been illustrated in Figs. 2 and 3 as of November 11, 2020. Moreover, projection results from mid November to early April for daily and cumulative cases in Argentina have been depicted in Figs. 4 and 5. According to our projection results, the daily could reach upto 2450K cases by the end of March 2021, if the current trend holds. Model parameters presented in the Table 2 have been calibrated using observed real data from March 4 to November 11, 2020.

As we can see the results from the proposed model responses complement the real data very well. The control reproduction number ($\mathcal{R}_c$) is estimated to be ~ 1.37 (95% CI : $1.03 - 1.59$) as of November 11 and prior established findings 1 – 5 for COVID-19 really match well with the estimation [30, 5]. The tally of cumulative infected cases is projected to reach 2450K and country's death toll could mount to 52.6K by the end of March 2021. Table 2 illustrates model calibration results and baseline parameter values.

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Figure 4: Projection results for daily new confirmed cases for Argentina from early March to late March 2021.

Figure 5: Projection results for cumulative cases for Argentina early March to late March 2021.

Parameter	Range (Unit)	Baseline value	TRR output	Reference
β_I	0.1–1.5 day^{-1}	0.55	0.3	[27, 5]
β_A	0.1–0.9 day^{-1}	0.3	0.15	[27, 5]
β_Q	0.1–0.9 day^{-1}	0.5	0.3	[27, 5]
β_L	0.1–0.9 day^{-1}	0.3	0.35	[5, 6]
β_{E_2}	0.05–0.3 day^{-1}	0.3	0.1	[6]
m	0.01–0.3 (dimensionless)	0.1	0.15	[6]
ζ	0.5 (dimensionless)	0.5	0.5	[6]
κ_1	$\frac{1}{4} day^{-1}$	$\frac{1}{4}$	$\frac{1}{4}$	[3, 6]
κ_2	1 day^{-1}	1	1	[3]
q	0.1–0.6 day^{-1}	0.3	0.25	Estimated
ρ	0.6–0.7 (dimensionless)	0.65	0.65	[6]
τ_I	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/10	[5, 6]
τ_A	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/10	[5, 6]
γ_I	$\frac{1}{14} - \frac{1}{7} day^{-1}$	1/7	0.143	[28, 29]
γ_A	$\frac{1}{10} - \frac{1}{7} day^{-1}$	1/7	0.143	[28, 29]
γ_Q	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/14	0.071	[28, 29]
γ_L	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/14	0.071	[28, 29]
δ_I	0.0001–0.01 day^{-1}	0.001	0.00033	[2, 5]
δ_L	0.0001–0.01 day^{-1}	0.001	0.001	[2, 5]
δ_Q	0.0001–0.01 day^{-1}	0.001	0.00033	[2, 5]

Table 2: Calibrated parameters of the proposed model (1) using trust-region-reflective algorithm and daily COVID-19 cases data of Argentina.

4.2. Bangladesh

Due to prolonged lockdown measures and severe economic recession, inhabitants of Bangladesh have started violating safety measures like wearing face coverings and maintaining physical distancing. Figs. 6 and 7 illustrate the model fitting performance with observed data from early March to mid November for

Figure 6: Fitting performance of the model for daily infected cases in Bangladesh from March 08 to November 11, 2020.

Figure 7: Fitting performance of the model for cumulative infected cases in Bangladesh from March 08 to November 11, 2020.

Bangladesh. The estimated error is found to be hovering around 10% for daily new cases. The actual outbreak scenario in Bangladesh is still a puzzle to be solved for the health officials due to scant COVID-19 testing program. The control reproduction number ($\mathcal{R}_c$) is estimated to be ~ 1.17 (95% CI : 0.95 – 1.39) as of November 11 and prior established findings for this metric go well with the estimation [30, 5]. The tally of cumulative infected cases is projected to reach 436K around March 31, 2021 and the estimated total death cases could reach 11,400 by the end of March, 2021. Table 3 illustrates the key features used to calibrate this scenario, which have been justified in prior clinical studies and relevant literature.

4.3. Brazil

COVID-19 pandemic emerged in Latin America a bit later than other continents and Brazil is among the hardest-hit countries in the world. When late-stage clinical trials of Chinese-developed CoronaVac vaccine are going on in Brazil, the inhabitants of Brazil are witnessing unprecedented spikes in new COVID-19 cases.

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Figure 8: Projection results for daily new confirmed cases for Bangladesh from early March to late March 2021.

Figure 9: Projection results for cumulative cases for Bangladesh early March to late March 2021.

Parameter	Range (Unit)	Baseline value	TRR output	Reference
β_I	0.1–1.5 day^{-1}	0.55	0.15	[27, 5]
β_A	0.1–0.9 day^{-1}	0.3	0.1	[27, 5]
β_Q	0.1–0.9 day^{-1}	0.5	0.1	[27, 5]
β_L	0.1–0.9 day^{-1}	0.3	0.12	[5, 6]
β_{E_2}	0.05–0.3 day^{-1}	0.3	0.11	[6]
m	0.01–0.3 (dimensionless)	0.1	0.3	[6]
ζ	0.5 (dimensionless)	0.5	0.5	[6]
κ_1	$\frac{1}{4} day^{-1}$	$\frac{1}{4}$	$\frac{1}{4}$	[3, 6]
κ_2	1 day^{-1}	1	1	[3]
q	0.1–0.6 day^{-1}	0.3	0.47	Estimated
ρ	0.6–0.7 (dimensionless)	0.65	0.65	[6]
τ_I	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/8	[5, 6]
τ_A	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/8	[5, 6]
γ_I	$\frac{1}{14} - \frac{1}{7} day^{-1}$	1/7	1/12	[28, 29]
γ_A	$\frac{1}{10} - \frac{1}{7} day^{-1}$	1/7	1/10	[28, 29]
γ_Q	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/21	0.071	[28, 29]
γ_L	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/21	0.071	[28, 29]
δ_I	0.0001–0.01 day^{-1}	0.001	0.0004	[2, 5]
δ_L	0.0001–0.01 day^{-1}	0.001	0.0009	[2, 5]
δ_Q	0.0001–0.01 day^{-1}	0.001	0.0007	[2, 5]

Table 3: Calibrated parameters of the proposed model (1) using trust-region-reflective algorithm and daily COVID-19 cases data of Bangladesh.

As of November 15, 2020, Brazil reported 5,863,093 cases and related 165,811 deaths. The model fitting performance for Brazil from late February to mid November are illustrated in Figs. 10 and 11. Historical data from late February to November 11, 2020 have been considered to calibrate the model parameters. It is clearly visible in the figures that the responses from the proposed model fit the real data very well.

158 The control reproduction number is estimated about ~ 2.12 as of November 11, which is in between the
 159 observed findings for COVID-19 [30, 5]. Fig. 13 illustrates that the tally of total infected cases could mount
 160 to 6368.6K by the end of March 2021 if current trend is held, and death toll could surpass 215K within this
 161 time period. Table 4 depicts the key features used to calibrate this scenario.

Parameter	Range (Unit)	Baseline value	TRR output	Reference
β_I	0.1–1.5 day^{-1}	0.55	0.15	[27, 5]
β_A	0.1–0.9 day^{-1}	0.3	0.1	[27, 5]
β_Q	0.1–0.9 day^{-1}	0.5	0.21	[27, 5]
β_L	0.1–0.9 day^{-1}	0.3	0.23	[5, 6]
β_{E_2}	0.05–0.3 day^{-1}	0.3	0.1	[6]
m	0.01–0.3 (dimension- less)	0.1	0.1	[6]
ζ	0.5 (dimensionless)	0.5	0.5	[6]
κ_1	$\frac{1}{4} day^{-1}$	$\frac{1}{4}$	$\frac{1}{4}$	[3, 6]
κ_2	1 day^{-1}	1	1	[3]
q	0.1–0.6 day^{-1}	0.3	0.2	Estimated
ρ	0.6–0.7 (dimension- less)	0.65	0.65	[6]
τ_I	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/12	[5, 6]
τ_A	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/12	[5, 6]
γ_I	$\frac{1}{14} - \frac{1}{7} day^{-1}$	1/10	0.143	[28, 29]
γ_A	$\frac{1}{10} - \frac{1}{7} day^{-1}$	1/10	0.143	[28, 29]
γ_Q	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/17	0.071	[28, 29]
γ_L	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/18	0.071	[28, 29]
δ_I	0.0001–0.01 day^{-1}	0.001	0.004	[2, 5]
δ_L	0.0001–0.01 day^{-1}	0.001	0.0099	[2, 5]
δ_Q	0.0001–0.01 day^{-1}	0.001	0.007	[2, 5]

Table 4: Calibrated parameters of the proposed model (1) using trust-region-reflective algorithm and daily COVID-19 cases data of Brazil.

Figure 10: Fitting performance of the model for daily infected cases in Brazil from February 25 to November 11, 2020.

Figure 11: Fitting performance of the model for cumulative infected cases in Brazil from February 25 to November 11, 2020.

4.4. Colombia

Colombia is one of the hardest-hit countries in Latin America which has already surpassed one million confirmed coronavirus cases. As of November 15, 2020, Colombia has seen 1,198,746 confirmed cases of COVID-19 and the country's death toll climbed to 34,031. Our model fitting performance with the historical data is quite outstanding depicted in Figs. 14 and 15. According to our projection results illustrated in Figs 16 and 17, the tally of cumulative cases could reach approximately 1730K cases and the number of daily cases could hover around 2000 cases by the end of March 2021. The control reproduction number ($\mathcal{R}_J$) is estimated to be ~ 1.19 as of November 11, 2020. According to our analysis, the epidemic threshold ($\mathcal{R}_J$) could be brought under unity by the end of December 2020 by following strict distance maintaining protocol and mass-level usage of efficacious face coverings.

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Figure 12: Projection results for daily new confirmed cases for Brazil from early March to late March 2021.

Figure 13: Projection results for cumulative cases for Brazil early March to late March 2021.

Figure 14: Fitting performance of the model for daily infected cases in Colombia from March 06 to November 11, 2020.

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Figure 15: Fitting performance of the model for cumulative infected cases in Colombia from March 06 to November 11, 2020.

Figure 16: Projection results for daily new confirmed cases for Colombia from early March to late March 2021.

Figure 17: Projection results for cumulative cases for Colombia early March to late March 2021.

Parameter	Range (Unit)	Baseline value	TRR output	Reference
β_I	0.1–1.5 day^{-1}	0.55	0.11	[27, 5]
β_A	0.1–0.9 day^{-1}	0.3	0.1	[27, 5]
β_Q	0.1–0.9 day^{-1}	0.5	0.1	[27, 5]
β_L	0.1–0.9 day^{-1}	0.3	0.12	[5, 6]
β_{E_2}	0.05–0.3 day^{-1}	0.3	0.11	[6]
m	0.01–0.3 (dimensionless)	0.1	0.3	[6]
ζ	0.5 (dimensionless)	0.5	0.5	[6]
κ_1	$\frac{1}{4} day^{-1}$	$\frac{1}{4}$	$\frac{1}{4}$	[3, 6]
κ_2	1 day^{-1}	1	1	[3]
q	0.1–0.6 day^{-1}	0.3	0.1	Estimated
ρ	0.6–0.7 (dimensionless)	0.65	0.65	[6]
τ_I	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/9	[5, 6]
τ_A	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/11	[5, 6]
γ_I	$\frac{1}{14} - \frac{1}{7} day^{-1}$	1/7	1/12	[28, 29]
γ_A	$\frac{1}{10} - \frac{1}{7} day^{-1}$	1/7	1/10	[28, 29]
γ_Q	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/14	1/20	[28, 29]
γ_L	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/14	1/21	[28, 29]
δ_I	0.0001–0.01 day^{-1}	0.001	0.0002	[2, 5]
δ_L	0.0001–0.01 day^{-1}	0.001	0.00058	[2, 5]
δ_Q	0.0001–0.01 day^{-1}	0.001	0.0005	[2, 5]

Table 5: Calibrated parameters of the proposed model (1) using trust-region-reflective algorithm and daily COVID-19 cases data of Colombia.

4.5. India

When India is celebrating a busy festival season, the tally of fresh COVID-19 cases continued to rise. Relaxation in protective and social-distancing measures could result in a significant upsurge in daily cases in upcoming days. Fig. 18 illustrates the fact that India is witnessing a downward trend after having

Figure 18: Fitting performance of the model for daily infected cases in India from January 30 to November 11, 2020.

Figure 19: Fitting performance of the model for cumulative infected cases in India from January 30 to November 11, 2020.

176 peak. According to our projection results, India could reel under the second wave of infection unless non-
177 pharmaceutical interventions strategies are followed comprehensively. As of November 15, the country's
178 caseload now stands at 8,850,338 and it's death toll has mounted to 130,187. Figs. 18 and 19 illustrate the
179 fitting performance of our proposed model for India from late January to mid November. Historical data
180 from January 30 to November 11 have been considered to calibrate the model parameters. As we can see
181 from the figures, model-fitting is exceptionally well for the historical observed data. Based on our projection
182 results from Fig. 20, the number of daily cases could be brought under 1000 cases if mass-level efficacious
183 face coverings is strictly maintained. The control reproduction number $\mathcal{R}_c$ is estimated to be ~ 1.41 as of
184 November 11, which complements the prior studied observations [30, 5]. The tally of cumulative infected
185 cases is projected to reach 11259K by the end of March 2021 if current trend continues. In addition, country's
186 death toll could mount to 187K over the period. Tab. 6 illustrates the key features used to calibrate this
187 scenario.

It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Figure 20: Projection results for daily new confirmed cases for India from late January 2020 to late March 2021.

Figure 21: Projection results of cumulative cases for India from late January 2020 to late March 2021.

Parameter	Range (Unit)	Baseline value	TRR output	Reference
β_I	0.1–1.5 day^{-1}	0.55	0.18	[27, 5]
β_A	0.1–0.9 day^{-1}	0.3	0.13	[27, 5]
β_Q	0.1–0.9 day^{-1}	0.5	0.2	[27, 5]
β_L	0.1–0.9 day^{-1}	0.3	0.22	[5, 6]
β_{E_2}	0.05–0.3 day^{-1}	0.3	0.15	[6]
m	0.01–0.3 (dimensionless)	0.1	0.23	[6]
ζ	0.5 (dimensionless)	0.5	0.5	[6]
κ_1	$\frac{1}{4} day^{-1}$	$\frac{1}{4}$	$\frac{1}{4}$	[3, 6]
κ_2	1 day^{-1}	1	1	[3]
q	0.1–0.6 day^{-1}	0.3	0.25	Estimated
ρ	0.6–0.7 (dimensionless)	0.65	0.65	[6]
τ_I	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/8	[5, 6]
τ_A	$\frac{1}{14} - \frac{1}{5} day^{-1}$	1/10	1/9	[5, 6]
γ_I	$\frac{1}{14} - \frac{1}{7} day^{-1}$	1/7	1/12	[28, 29]
γ_A	$\frac{1}{10} - \frac{1}{7} day^{-1}$	1/7	1/10	[28, 29]
γ_Q	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/14	1/16	[28, 29]
γ_L	$\frac{1}{21} - \frac{1}{10} day^{-1}$	1/14	1/18	[28, 29]
δ_I	0.0001–0.01 day^{-1}	0.001	0.0002	[2, 5]
δ_L	0.0001–0.01 day^{-1}	0.001	0.00032	[2, 5]
δ_Q	0.0001–0.01 day^{-1}	0.001	0.0003	[2, 5]

Table 6: Calibrated parameters of the proposed model (1) using trust-region-reflective algorithm and daily COVID-19 cases data of India.

5. Global sensitivity analysis

With an aim to quantify the most dominant mechanisms in the proposed model, a renowned global sensitivity analysis approach Partial Rank Correlation Coefficient (PRCC) method has been carried out. This

method can provide considerable insights about the relationship between model responses (state variables) and model parameters (sampled by Latin Hypercube Sampling method) in an outbreak setting [31]. PRCC values are generally bounded between -1 and 1. A monotonic relationship between the model input parameters and the model outputs is generally assumed in this method. Apart from qualitative relationship, precise quantitative impact of different parameters on model responses can be determined by calculating the PRCC values. Uniform distribution of all model parameters have been considered to generate LHS matrices. A positive PRCC value depicts that the model output can be increased by increasing the respective model input parameter. Similarly, model output can be decreased by forcing down the corresponding input parameter. In addition, a negative PRCC value indicates a negative correlation between the model input and output.

When it comes to analyse a complex model, it often get really challenging to control the parameters. In this context, sensitivity analysis can give considerable insights regarding the quantitative relationship between model responses and model input parameters. However, it is really challenging for complex models to determine the qualitative and quantitative relationship with sufficient accuracy. As we can see from Figure 22, nearly the same qualitative relationship has been found between the number of symptomatic infectious individuals (one of the crucial model responses) and three parameters which are effective contact rate with isolated infected individuals (β_L), efficacy of face coverings (ζ) and face coverings compliance for our studied five countries. The public health implications of these findings are the dynamics of COVID-19 could be controlled by encouraging mass-level usage of efficacious face coverings. In addition, the high significance of β_L indicates that immediate isolation of detected patients is highly required.

6. Solution of the model in Caputo-Fabrizio fractional derivative sense

6.1. Preliminaries

Here we recall the definitions of Caputo and Caputo-Fabrizio fractional derivatives.

Definition 1. [32] The Caputo definition of non-integer order derivative of order $\varrho > 0$ of a function $G : (0, \infty) \rightarrow \mathbb{R}$ is defined by

$$D_t^\varrho G(t) = \frac{1}{\Gamma(n - \varrho)} \int_0^t (t - \tau)^{n - \varrho - 1} G'(\tau) d\tau \quad (12)$$

where $n = [\varrho] + 1$ and $[\varrho]$ is the integer part of ϱ .

Definition 2. [33, 34] For $G \in H^1(c, d)$ and $0 < \varrho < 1$, the Caputo-Fabrizio (CF) fractional derivative (FD) of order ϱ is defined by

$${}_c^{CF} D_t^\varrho G(t) = \frac{1}{1 - \varrho} \int_c^t \frac{dG(\eta)}{d\eta} \exp[-\theta(t - \eta)] d\eta$$

where $\theta = \frac{\varrho}{1 - \varrho}$

The CF non-integer order integral is defined as

$${}_c^{CF} I_t^\varrho G(t) = (1 - \varrho)G(t) + \varrho \int_c^t G(\eta) d\eta.$$

(a) Sensitivity analysis for Argentina

(b) Sensitivity analysis for Bangladesh

(c) Sensitivity analysis for Brazil

214 6.2. Existence and uniqueness analysis

Now, we prove the existence of unique solution for the given COVID-19 model in the sense of Caputo-Fabrizio fractional derivative by the application of fixed-point theory. In this concern, the proposed system can be rewritten in the equivalent form as follows:

$$\begin{cases} {}^{CF}D_t^\varrho S(t) = \mathcal{G}_1(t, S(t)), \\ {}^{CF}D_t^\varrho E_1(t) = \mathcal{G}_2(t, E_1(t)), \\ {}^{CF}D_t^\varrho E_2(t) = \mathcal{G}_3(t, E_2(t)), \\ {}^{CF}D_t^\varrho I(t) = \mathcal{G}_4(t, I(t)), \\ {}^{CF}D_t^\varrho A(t) = \mathcal{G}_5(t, A(t)), \\ {}^{CF}D_t^\varrho Q(t) = \mathcal{G}_6(t, Q(t)), \\ {}^{CF}D_t^\varrho L(t) = \mathcal{G}_7(t, L(t)), \\ {}^{CF}D_t^\varrho R(t) = \mathcal{G}_8(t, R(t)), \\ {}^{CF}D_t^\varrho D(t) = \mathcal{G}_9(t, D(t)). \end{cases} \quad (13)$$

By applying the CF non-integer order integral operator, the above system (13), reduces to the following integral equation of Volterra type of order $0 < \varrho < 1$.

$$\begin{aligned} S(t) - S(0) &= (1 - \varrho)G_1(t, S) + \varrho \int_0^t \mathcal{G}_1(\chi, S)d\chi, \\ E_1(t) - E_1(0) &= (1 - \varrho)G_2(t, E_1) + \varrho \int_0^t \mathcal{G}_2(\chi, E_1)d\chi, \\ E_2(t) - E_2(0) &= (1 - \varrho)G_3(t, E_2) + \varrho \int_0^t \mathcal{G}_3(\chi, E_2)d\chi, \\ I(t) - I(0) &= (1 - \varrho)G_4(t, I) + \varrho \int_0^t \mathcal{G}_4(\chi, I)d\chi, \\ A(t) - A(0) &= (1 - \varrho)G_5(t, A) + \varrho \int_0^t \mathcal{G}_5(\chi, A)d\chi, \\ Q(t) - Q(0) &= (1 - \varrho)G_6(t, Q) + \varrho \int_0^t \mathcal{G}_6(\chi, Q)d\chi, \\ L(t) - L(0) &= (1 - \varrho)G_7(t, L) + \varrho \int_0^t \mathcal{G}_7(\chi, L)d\chi, \\ R(t) - R(0) &= (1 - \varrho)G_8(t, R) + \varrho \int_0^t \mathcal{G}_8(\chi, R)d\chi, \\ D(t) - D(0) &= (1 - \varrho)G_9(t, D) + \varrho \int_0^t \mathcal{G}_9(\chi, D)d\chi. \end{aligned} \quad (14)$$

Now, we get the subsequent iterative algorithm

$$\begin{aligned}
 S_{n+1}(t) &= (1 - \varrho)\mathcal{G}_1(t, S_n) + \varrho \int_0^t \mathcal{G}_1(\chi, S_n) d\chi, \\
 E_{1n+1}(t) &= (1 - \varrho)\mathcal{G}_2(t, E_{1n}) + \varrho \int_0^t \mathcal{G}_2(\chi, E_{1n}) d\chi, \\
 E_{2n+1}(t) &= (1 - \varrho)\mathcal{G}_3(t, E_{2n}) + \varrho \int_0^t \mathcal{G}_3(\chi, E_{2n}) d\chi, \\
 I_{n+1}(t) &= (1 - \varrho)\mathcal{G}_4(t, I_n) + \varrho \int_0^t \mathcal{G}_4(\chi, I_n) d\chi, \\
 A_{n+1}(t) &= (1 - \varrho)\mathcal{G}_5(t, A_n) + \varrho \int_0^t \mathcal{G}_5(\chi, A_n) d\chi, \\
 Q_{n+1}(t) &= (1 - \varrho)\mathcal{G}_6(t, Q_n) + \varrho \int_0^t \mathcal{G}_6(\chi, Q_n) d\chi, \\
 L_{n+1}(t) &= (1 - \varrho)\mathcal{G}_7(t, L_n) + \varrho \int_0^t \mathcal{G}_7(\chi, L_n) d\chi, \\
 R_{n+1}(t) &= (1 - \varrho)\mathcal{G}_8(t, R_n) + \varrho \int_0^t \mathcal{G}_8(\chi, R_n) d\chi, \\
 D_{n+1}(t) &= (1 - \varrho)\mathcal{G}_9(t, D_n) + \varrho \int_0^t \mathcal{G}_9(\chi, D_n) d\chi.
 \end{aligned} \tag{15}$$

Here we assume that we can get the exact solution by taking the limit as n tends to infinity.

6.2.1. Existence analysis by using Picard-Lindelof approach

Let us consider

$$\begin{aligned}
 F_1 &= \sup_{\mathcal{C}_{[c, z_1]}} \|\mathcal{G}_1(t, S)\|, F_2 = \sup_{\mathcal{C}_{[c, z_2]}} \|\mathcal{G}_2(t, E_1)\|, F_3 = \sup_{\mathcal{C}_{[c, z_3]}} \|\mathcal{G}_3(t, E_2)\|, F_4 = \sup_{\mathcal{C}_{[c, z_4]}} \|\mathcal{G}_4(t, I)\|, \\
 F_5 &= \sup_{\mathcal{C}_{[c, z_5]}} \|\mathcal{G}_5(t, A)\|, F_6 = \sup_{\mathcal{C}_{[c, z_6]}} \|\mathcal{G}_6(t, Q)\|, F_7 = \sup_{\mathcal{C}_{[c, z_7]}} \|\mathcal{G}_7(t, L)\|, F_8 = \sup_{\mathcal{C}_{[c, z_8]}} \|\mathcal{G}_8(t, R)\|, \\
 F_9 &= \sup_{\mathcal{C}_{[c, z_9]}} \|\mathcal{G}_9(t, D)\|,
 \end{aligned} \tag{16}$$

where

$$\begin{aligned}
 \mathcal{C}_{c, z_1} &= |t - c, t + c| \times [S - z_1, S + z_1] = D_1 \times B_1, \\
 \mathcal{C}_{c, z_2} &= |t - c, t + c| \times [E_1 - z_2, E_1 + z_2] = D_1 \times B_2, \\
 \mathcal{C}_{c, z_3} &= |t - c, t + c| \times [E_2 - z_3, E_2 + z_3] = D_1 \times B_3, \\
 \mathcal{C}_{c, z_4} &= |t - c, t + c| \times [I - z_4, I + z_4] = D_1 \times B_4, \\
 \mathcal{C}_{c, z_5} &= |t - c, t + c| \times [A - z_5, A + z_5] = D_1 \times B_5, \\
 \mathcal{C}_{c, z_6} &= |t - c, t + c| \times [Q - z_6, Q + z_6] = D_1 \times B_6, \\
 \mathcal{C}_{c, z_7} &= |t - c, t + c| \times [L - z_7, L + z_7] = D_1 \times B_7, \\
 \mathcal{C}_{c, z_8} &= |t - c, t + c| \times [R - z_8, R + z_8] = D_1 \times B_8, \\
 \mathcal{C}_{c, z_9} &= |t - c, t + c| \times [D - z_9, D + z_9] = D_1 \times B_9.
 \end{aligned} \tag{17}$$

considering the Picard operator as

$\phi : C(D_1, B_1, B_2, B_3, B_4, B_5, B_6, B_7, B_8, B_9) \rightarrow C(D_1, B_1, B_2, B_3, B_4, B_5, B_6, B_7, B_8, B_9)$, given as follows:

$$\phi\zeta(t) = \zeta_0(t) + \Delta(t, \zeta(t))(1 - \varrho) + \varrho \int_0^t \Delta(s, \zeta(s))ds,$$

where $\zeta(t) = \{S(t), E_1(t), E_2(t), I(t), A(t), Q(t), L(t), R(t), D(t)\}$, $\zeta_0(t) = \{S_0, E_{10}, E_{20}, I_0, A_0, Q_0, L_0, R_0, D_0\}$ and $\Delta(t, \zeta(t)) = \{\mathcal{G}_1(t, S(t)), \mathcal{G}_2(t, E_1(t)), \mathcal{G}_3(t, E_2(t)), \mathcal{G}_4(t, I(t)), \mathcal{G}_5(t, A(t)), \mathcal{G}_6(t, Q(t)), \mathcal{G}_7(t, L(t)), \mathcal{G}_8(t, R(t)), \mathcal{G}_9(t, D(t))\}$. Next we assume that the solution of the non-integer order model are bounded within a time period,

$$\|\zeta(t)\|_\infty \leq \max\{z_1, z_2, z_3, z_4, z_5, z_6, z_7, z_8, z_9\},$$

$$\|\zeta(t) - \zeta_0(t)\| = \|\Delta(t, \zeta(t))(1 - \varrho) + \varrho \int_0^t \Delta(s, \zeta(s))ds\|$$

$$\leq \|\Delta(t, \zeta(t))\| (1 - \varrho) + \varrho \int_0^t \|\Delta(s, \zeta(s))\| ds$$

$$\leq \left((1 - \varrho) + \varrho\zeta_0 \right) \max\{F_1, F_2, F_3, F_4, F_5, F_6, F_7, F_8, F_9\} < bF \leq d = \max\{z_1, z_2, z_3, z_4, z_5, z_6, z_7, z_8, z_9\},$$

where we demand that

$b < \frac{d}{F}$. Now by the application of fixed point theorem pertaining to Banach space along with the metric, we obtain

$\|\phi\zeta_1 - \phi\zeta_2\|_\infty = \sup_{t \rightarrow B} |\zeta_1 - \zeta_2|$. Now we have

$$\begin{aligned} \|\phi\zeta_1 - \phi\zeta_2\| &= \|\{\Delta(t, \zeta_1(t)) - \Delta(t, \zeta_2(t))\}(1 - \varrho) + \varrho \int_0^t \{\Delta(s, \zeta_1(s)) - \Delta(s, \zeta_2(s))\}ds\| \\ &\leq \|\Delta(t, \zeta_1(t)) - \Delta(t, \zeta_2(t))\| (1 - \varrho) + \varrho \int_0^t \|\Delta(s, \zeta_1(s)) - \Delta(s, \zeta_2(s))\| ds \\ &\leq (1 - \varrho)\beta \|\zeta_1(t) - \zeta_2(t)\| + \varrho\beta \int_0^t \|\zeta_1(s) - \zeta_2(s)\| ds \\ &\leq \{(1 - \varrho)\beta + \varrho\beta t_0\} \|\zeta_1(t) - \zeta_2(t)\| \\ &\leq \mathcal{U}\beta \|\zeta_1(t) - \zeta_2(t)\|, \end{aligned} \tag{18}$$

217 with $\beta < 1$. Since ζ is a contraction, we have $\mathcal{U}\beta < 1$, hence the given operator ϕ is also a contraction.

218 Therefore, the model involving C-F derivative given in Eq. (13) has a unique set of solution.

219 6.3. Solution method in Caputo-Fabrizio Operator

We now derive the solution method for $I(t)$ equation of the system (13) and for the rest of the equations it will be similar. The corresponding Volterra integral equation for $I(t)$ is as follows.

$$I(t) = I(0) + (1 - \varrho)\mathcal{G}_4(t, I(t)) + \varrho \int_0^t \mathcal{G}_4(s, I(s))ds \tag{19}$$

We have the following estimations at t_k

$$I(t_k) = I_0 + (1 - \varrho)\mathcal{G}_4(t_{k-1}, I(t_{k-1})) + \varrho \int_0^{t_k} \mathcal{G}_4(t, I(t))dt, \quad (20)$$

and at t_{k+1}

$$I(t_{k+1}) = I_0 + (1 - \varrho)\mathcal{G}_4(t_k, I(t_k)) + \varrho \int_0^{t_{k+1}} \mathcal{G}_4(t, I(t))dt. \quad (21)$$

Subtracting equation (21) from (20), we obtain

$$I(t_{k+1}) - I(t_k) = (1 - \varrho) (\mathcal{G}_4(t_k, I(t_k)) - \mathcal{G}_4(t_{k-1}, I(t_{k-1}))) + \varrho \int_{t_k}^{t_{k+1}} \mathcal{G}_4(t, I(t))dt. \quad (22)$$

Then by linear interpolation about $\mathcal{G}_4(t, I(t))$ and applying trapezoid rule for integration on the integral term, we can then write

$$\int_{t_k}^{t_{k+1}} \mathcal{G}_4(t, I(t))dt \cong \frac{3\Delta t}{2} \mathcal{G}_4(t_k, I(t_k)) - \frac{\Delta t}{2} \mathcal{G}_4(t_k, I(t_k)) \quad (23)$$

where $\Delta t = t_k - t_{k-1}$. Hence, we have the numerical approximation for equation of $I(t)$ as

$$I(t_{k+1}) = I(t_k) + \left(1 - \varrho + \frac{3\varrho\Delta t}{2}\right) \mathcal{G}_4(t_k, I(t_k)) - \left(1 - \varrho + \frac{\varrho\Delta t}{2}\right) \mathcal{G}_4(t_{k-1}, I(t_{k-1})). \quad (24)$$

Theorem 5. The numerical approximation (24) is unconditionally stable if

$$\|\mathcal{G}_4(t_{k+1}, I(t_{k+1})) - \mathcal{G}_4(t_k, I(t_k))\| \rightarrow 0.$$

Proof. Let $I(t)$ be the solution of a differential equation as shown in (19) under CF non-integer order derivative operator sense. Then we have to evaluate the norm

$$\begin{aligned} \|I(t_{k+1}) - I(t_k)\| &= \left\| (1 - \varrho) (\mathcal{G}_4(t_k, I(t_k)) - \mathcal{G}_4(t_{k-1}, I(t_{k-1}))) + \varrho \int_{t_k}^{t_{k+1}} \mathcal{G}_4(\eta, I(\eta))d\eta \right\| \\ &\leq (1 - \varrho) \|(\mathcal{G}_4(t_k, I(t_k)) - \mathcal{G}_4(t_{k-1}, I(t_{k-1})))\| + \varrho \left\| \int_{t_k}^{t_{k+1}} \mathcal{G}_4(\eta, I(\eta))d\eta \right\|. \end{aligned} \quad (25)$$

For $k \rightarrow \infty$, we have

$$\begin{aligned} \|I(t_{k+1}) - I(t_k)\|_\infty &\leq \lim_{k \rightarrow \infty} (1 - \varrho) \|(\mathcal{G}_4(t_k, I(t_k)) - \mathcal{G}_4(t_{k-1}, I(t_{k-1})))\| + \lim_{k \rightarrow \infty} \varrho \left\| \int_{t_k}^{t_{k+1}} \mathcal{G}_4(\eta, I(\eta))d\eta \right\| \\ &< \lim_{k \rightarrow \infty} (1 - \varrho) \|(\mathcal{G}_4(t_k, I(t_k)) - \mathcal{G}_4(t_{k-1}, I(t_{k-1})))\| \\ &\quad + \lim_{k \rightarrow \infty} \varrho \int_{t_k}^{t_{k+1}} \left| \sum_{j=0}^k \prod_{j=0}^k \frac{\eta - t_j}{\Delta t} \mathcal{G}_4(t_j, I(t_j)) \right| d\eta \end{aligned}$$

Clearly, the second term of the above inequality approaches zero when $k \rightarrow \infty$. Now, if $\|\mathcal{G}_4(t_{k+1}, I(t_{k+1})) - \mathcal{G}_4(t_k, I(t_k))\| \rightarrow 0$ as $k \rightarrow \infty$, we deduce that the numerical solution is stable. $\square$

6.4. Graphical simulations

After giving the all necessary theoretical concerns, now we do the graphical simulations via CF fractional derivative. In our paper, we are using the real numerical data of COVID-19 for five different countries named as Argentina, Bangladesh, Brazil, Colombia and India respectively. Here first we perform the graphs for Argentina COVID-19 cases. To perform numerical simulations, we use parameter values summarize in Table 2. In the family of Figure 23, we analysed the plots of $S(t)$, $E_1(t)$, $E_2(t)$, $I(t)$, $A(t)$ and $Q(t)$. We observed that for different fractional order values peaks are well defined and when we decrease the fractional order then the peaks sifted towards the later time period. In the group of Figs. 24- 25, first we show the nature of $L(t)$, $R(t)$, $D(t)$ and then analysed the plots of $I(t)$ versus $S(t)$, $A(t)$, $Q(t)$, $L(t)$ and $R(t)$. In the comparison of given classes with infected individuals, we see that when the infectious $I(t)$ increases then asymptomatic infectious $A(t)$ also increases with same nature. In sub-figs. 24d- 25b, we see that the fractional order does not play any big role because the nature of the classes is nearly same at all different fractional order values ρ . Initial values of given classes for Argentina are $S(0) = 45333107$, $E_1(0) = 10$, $E_2(0) = 4$, $I(0) = 2$, $A(0) = 1$, $Q(0) = 0$, $L(0) = 0$, $R(0) = 0$ and $D(0) = 0$. We have used the total population of the country for $S(0)$.

Figure 23: Plots of $S(t)$, $E_1(t)$, $E_2(t)$, $I(t)$, $A(t)$ and $Q(t)$ for Argentina data

Figure 24: Plots of $L(t)$, $R(t)$, $D(t)$ and relationship of $I(t)$ versus $S(t)$, $A(t)$, $Q(t)$, $L(t)$, $R(t)$ for Argentina data

Figure 25: $I(t)$ versus $L(t), R(t)$ for Argentina data

Now we perform the graphs for COVID-19 cases in Bangladesh. To perform numerical simulations, we use parameter values summarize in Table 3. In the family of Figure 26, we analysed the plots of $S(t), E_1(t), E_2(t), I(t), A(t)$ and $Q(t)$. We observed that for different fractional order values peaks are well defined and when we decrease the fractional order then the peaks sifted towards the later time period. In the group of Figures 27- 28, first we show the nature of $L(t), R(t), D(t)$ and then analysed the plots of $I(t)$ versus $S(t), A(t), Q(t), L(t)$ and $R(t)$. In the comparison of given classes with infected individuals, we see that the nature of infectious $I(t)$ is same as for Argentina, as when the population of infected individuals increases then asymptomatic infectious $A(t)$ also increases with same nature. In sub-figures 27d- 28b, we see that the fractional order does not play any big role because the nature of the classes is nearly same at all different fractional order values ρ . Initial values of given classes for Bangladesh are $S(0) = 164689383, E_1(0) = 10, E_2(0) = 4, I(0) = 2, A(0) = 1, Q(0) = 0, L(0) = 0, R(0) = 0$ and $D(0) = 0$. We have used the total population of the country for $S(0)$.

Figure 26: Plots of $S(t)$, $E_1(t)$, $E_2(t)$, $I(t)$, $A(t)$ and $Q(t)$ for Bangladesh data

Figure 27: Plots of $L(t)$, $R(t)$, $D(t)$ and relationship of $I(t)$ versus $S(t)$, $A(t)$, $Q(t)$, $L(t)$, $R(t)$ for Bangladesh data

Figure 28: $I(t)$ versus $L(t), R(t)$ for Bangladesh data

248 In the family of Figures 29- 31, we exemplified the graphs for COVID-19 cases in Brazil. To perform
 249 numerical simulations, we use parameter values summarize in Table 4. In the family of Figure 29, we
 250 analysed the plots of $S(t), E_1(t), E_2(t), I(t), A(t)$ and $Q(t)$. We observed that for different fractional order
 251 values peaks are well defined and when we decrease the fractional order then the peaks sifted towards the
 252 later time period. In the collection of Figures 30- 31, first we show the nature of $L(t), R(t), D(t)$ and then
 253 analysed the plots of $I(t)$ versus $S(t), A(t), Q(t), L(t)$ and $R(t)$. In the comparison of given classes with $I(t)$,
 254 we again observed that the nature of infectious $I(t)$ is same as for above countries, as when the population
 255 of infected individuals increases then asymptomatic infectious $A(t)$ also increases with same nature. In
 256 sub-figures 30d- 31b, we see that the fractional order does not play any big role because the nature of the
 257 classes is nearly same at all different fractional order values ρ . Initial values of given classes for Brazil are
 258 $S(0) = 212559417, E_1(0) = 10, E_2(0) = 4, I(0) = 2, A(0) = 1, Q(0) = 0, L(0) = 0, R(0) = 0$ and $D(0) = 0$.
 259 We have used the total population of the country for $S(0)$.

Figure 29: Plots of $S(t)$, $E_1(t)$, $E_2(t)$, $I(t)$, $A(t)$ and $Q(t)$ for Brazil data

Figure 30: Plots of $L(t)$, $R(t)$, $D(t)$ and relationship of $I(t)$ versus $S(t)$, $A(t)$, $Q(t)$, $L(t)$, $R(t)$ for Brazil data

Figure 31: $I(t)$ versus $L(t), R(t)$ for Brazil data

260 To continue the graphical simulations for the above mentioned countries to study the outbreaks of COVID-
261 19, in the family of Figures 32- 34, we exemplified the graphs for COVID-19 cases in Colombia. To perform
262 numerical simulations, we have taken the numerical values summarize in Table 5. In the family of Figure
263 32, we analysed the plots of $S(t), E_1(t), E_2(t), I(t), A(t)$ and $Q(t)$. We observed that the nature of peaks is
264 mostly same as for other above analysed data, for different fractional order values peaks are well defined and
265 when we decrease the fractional order then the peaks sifted towards the later time period. In the collection
266 of Figures 33- 34, first we show the nature of $L(t), R(t), D(t)$ and then analysed the plots of $I(t)$ versus
267 $S(t), A(t), Q(t), L(t)$ and $R(t)$. When we compare the given classes with $I(t)$, we again observed that when
268 the population of infected individuals increases then asymptomatic infectious $A(t)$ also increases with same
269 nature. In sub-figures 33d- 34b, we see that the fractional order does not play any big role because the
270 nature of the classes is nearly same at all different fractional order values ρ . Initial values of given classes
271 for Colombia are $S(0) = 50882891, E_1(0) = 10, E_2(0) = 4, I(0) = 2, A(0) = 1, Q(0) = 0, L(0) = 0, R(0) = 0$
272 and $D(0) = 0$. We have used the total population of the country for $S(0)$.

Figure 32: Plots of $S(t)$, $E_1(t)$, $E_2(t)$, $I(t)$, $A(t)$ and $Q(t)$ for Colombia data

Figure 33: Plots of $L(t)$, $R(t)$, $D(t)$ and relationship of $I(t)$ versus $S(t)$, $A(t)$, $Q(t)$, $L(t)$, $R(t)$ for Colombia data

Figure 34: $I(t)$ versus $L(t), R(t)$ for Colombia data

Eventually, we have done the graphical simulations for India which is the second highest populous country and also the second worst-hit country by COVID-19. To study the outbreak of COVID-19 in India, in the family of Figures 35- 37, we exemplified the all necessary graphs of given classes to observe the dynamics of COVID-19. To perform numerical simulations, we took the numerical values from the Table 6. In the family of Figure 35, we analysed the plots of $S(t), E_1(t), E_2(t), I(t), A(t)$ and $Q(t)$. We observed that the nature of peaks is mostly same as for other above analysed data, for different fractional order values peaks are well defined and when we decrease the fractional order then the peaks sifted towards the later time period. In the collection of Figures 36- 37, first we show the nature of $L(t), R(t), D(t)$ and then analysed the plots of $I(t)$ versus $S(t), A(t), Q(t), L(t)$ and $R(t)$. When we compare the given classes with $I(t)$, we again observed that when the population of infected individuals increases then asymptomatic infectious $A(t)$ also increases with same nature. In sub-figures 36d- 37b, we observed that at the different fractional order values the nature of the classes is nearly same. Initial values of given classes for India are $S(0) = 414001316, E_1(0) = 10, E_2(0) = 4, I(0) = 2, A(0) = 1, Q(0) = 0, L(0) = 0, R(0) = 0$ and $D(0) = 0$. We have used the 30% of the total population of India for $S(0)$.

Figure 35: Plots of $S(t)$, $E_1(t)$, $E_2(t)$, $I(t)$, $A(t)$ and $Q(t)$ for India data

Figure 36: Plots of $L(t)$, $R(t)$, $D(t)$ and relationship of $I(t)$ versus $S(t)$, $A(t)$, $Q(t)$, $L(t)$, $R(t)$ for India data

Figure 37: $I(t)$ versus $L(t), R(t)$ for India data

From the all above graphical observations we found that the Caputo-Fabrizio fractional derivative playing well to study the outbreaks of coronavirus in the aforesaid five countries.

7. Optimal control problem formulation

In this concern, our main aim is to decrease the number of infected individuals with COVID-19 at the same time decrease the cost $J(v)$ associated with their strategies. For this purpose, we use a control function $v = (v_1, v_2, v_3)$, where $v_1(t)$ is for introducing the public education or aware the public with health-care measures, $v_2(t)$ is the control function for enhancement of the strength of treatment for the infected individuals, $v_3(t)$ is the control function for the necessary suggestions of health care measures for those who are in asymptomatic infectious class and yet not admitted in the hospital.

$$\begin{cases} {}^{CF}D_t^\rho S(t) = -(1 - v_1)(\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2})S, \\ {}^{CF}D_t^\rho E_1(t) = (1 - v_1)(\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2})S - \kappa_1 E_1, \\ {}^{CF}D_t^\rho E_2(t) = \kappa_1 E_1 - (\kappa_2 + q)E_2, \\ {}^{CF}D_t^\rho I(t) = \rho \kappa_2 E_2 - (\tau_I + \gamma_I + \delta_I + v_2)I, \\ {}^{CF}D_t^\rho A(t) = (1 - \rho)\kappa_2 E_2 - (\tau_A + \gamma_A + v_3)A, \\ {}^{CF}D_t^\rho Q(t) = qE_2 - (\gamma_Q + \delta_Q)Q, \\ {}^{CF}D_t^\rho L(t) = \tau_I I + \tau_A A - (\delta_L + \gamma_L)L, \\ {}^{CF}D_t^\rho R(t) = \gamma_I I + \gamma_A A + \gamma_Q Q + \gamma_L L. \end{cases} \quad (26)$$

To define the optimal control problem (OCP), we are excluding the death equation $D(t)$, because there is no significance of deaths in optimal controls. Now consider the state system given in (26) in R^8 ,

with the set of admissible control function. $\Omega = \{(v_1(\cdot), v_2(\cdot), v_3(\cdot)) | v_i \text{ is Lebesgue measurable on } [0, 1]\}$
 $0 \leq (v_1(\cdot), v_2(\cdot), v_3(\cdot)) \leq 1$ So the objective functional is defined by

$$J((v_1(\cdot), v_2(\cdot), v_3(\cdot))) = \int_0^T [I(t) + \frac{1}{2}(k_1 v_1^2(t) + k_2 v_2^2(t) + k_3 v_3^2(t))] dt \quad (27)$$

where the constants k_1, k_2 and k_3 are a measure of associative cost with the controls v_1, v_2 and v_3 . Then we find the optimal controls v_1, v_2 and v_3 to minimize the cost function.

$$J((v_1, v_2, v_3)) = \int_0^T \mu(S, E_1, E_2, I, A, Q, L, R, v_1, v_2, v_3, t) dt \quad (28)$$

subject to constraint,

$${}_0^C D_t^\alpha S(t) = \phi_1, \quad {}_0^C D_t^\alpha E_1(t) = \phi_2, \dots \quad {}_0^C D_t^\alpha R(t) = \phi_8$$

where $\phi_j = \phi(S, E_1, E_2, I, A, Q, L, R, V_1, V_2, V_3, t)$, $j = 1, 2, \dots, 8$ and the given initial coordination are agreed $S(0) = S_0, E_1(0) = E_1, R(0) = R_0$

Now let us take the following modified cost function

$$\tilde{x} = \int_0^T [H(S_1, E_1, \dots, R, v_i, t) - \sum_{j=1}^8 (\theta_j \phi_j(S_1, E_1, \dots, R, v_i, t))] dt \quad (29)$$

where $i = 1, 2, 3$ and $j = 1, 2, 3, \dots, 8$ Hence the Hamiltonian is defined as follows:

$$H(S_1, E_1, \dots, R, v_i, t) = \mu(S_1, E_1, \dots, R, v_i, t) + \sum_{j=1}^8 (\theta_j \phi_j(S_1, E_1, \dots, R, v_i, t)) \quad (30)$$

where $i = 1, 2, 3$ and $j = 1, 2, 3, \dots, 8$ from Equation 29 and 30, the necessary and sufficient conditions for the functional optimal control problem (FOCP) are given as:

$${}_0^C D_t^\alpha \theta_1 = \frac{\partial H}{\partial S}, \quad {}_0^C D_t^\alpha \theta_2 = \frac{\partial H}{\partial E_1}, \dots \quad {}_0^C D_t^\alpha \theta_8 = \frac{\partial H}{\partial R} \quad (31)$$

$$0 = \frac{\partial H}{\partial v_i} \quad (32)$$

$${}_0^C D_t^\alpha S(t) = \frac{\partial H}{\partial \theta_1}, \dots \quad {}_0^C D_t^\alpha R(t) = \frac{\partial H}{\partial \theta_8} \quad (33)$$

Moreover, $\theta_j(T) = 0$, $j = 1, 2, \dots, 8$, are the lagseuges multipliers Eqn 31 and 32 express the necessary condition in terms of a Hamiltonian for the OCP defined above.

7.1. Optimality conditions for fractional order

Let us write the Hamiltonian function as follows:

$$H(S_1, E_1, \dots, R, v_i, \theta) = I + \frac{1}{2}(k_1 v_1^2 + k_2 v_2^2 + k_3 v_3^2) + \theta_1[-(1 - v_1)(\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2})S] + \theta_2[(1 - v_1)(\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2})S - \kappa_1 E_1] + \theta_3[\kappa_1 E_1 - (\kappa_2 - q)E_2] + \theta_4[\rho \kappa_2 E_2 - (\tau_I + \gamma_I + \delta_I + v_2)I] + \theta_5[(1 - \rho)\kappa_2 E_2 - (\tau_A + \gamma_A + v_3)A] + \theta_6[qE_2 - (\gamma_Q + \delta_Q)Q] + \theta_7[\tau_1 I + \tau_A A - (\delta_L + \gamma_L)L] + \theta_8[\gamma_I I + \gamma_A A + \gamma_Q Q + \gamma_L L]$$

where θ_j , $j = 1, 2, \dots, 8$, representing the lagrangars multipliers called co-states.

Theorem: If v_1^* , v_2^* and v_3^* are optimal controls of the given OCP if S^* , E_1^* , $\dots$, R^* are corresponding optimal paths, then there exists co-state variables θ_1^* , $\dots$, θ_8^* , such that besides the given control system is satisfied, the following conditions are satisfied:

Co-state equations:

$$\begin{aligned} {}^C D^e \theta_1^* &= (1 - v_1)(\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2})\theta_1^* - (1 - v_1)(\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2})\theta_2^*, \\ {}^C D^e \theta_2^* &= \kappa_1 \theta_2^* - \kappa_1 \theta_3^* \\ {}^C D^e \theta_3^* &= \beta_{E_2}(1 - m\zeta)\frac{S}{N}\theta_1^* - \beta_{E_2}(1 - m\zeta)\frac{S}{N}\theta_2^* + (\kappa_2 + q)\theta_3^* - \rho\kappa_2\theta_4^* - (1 - \rho)\kappa_2\theta_5^* - q\theta_6^* \\ {}^C D^e \theta_4^* &= \beta_I(1 - m\zeta)\frac{S}{N}\theta_1^* - \beta_I(1 - m\zeta)\frac{S}{N}\theta_2^* + (I_I + Y_I + \delta_I + v_2)\theta_4^* - \tau_I\theta_7^* - \gamma_I\theta_8^* \\ {}^C D^e \theta_5^* &= \beta_A(1 - m\zeta)\frac{S}{N}\theta_1^* - \beta_A(1 - m\zeta)\frac{S}{N}\theta_2^* + (\tau_A + \gamma_A + v_3)\theta_5^* - \tau_A\theta_7^* - \gamma_A\theta_8^* \\ {}^C D^e \theta_6^* &= \beta_Q(1 - m\zeta)\frac{S}{N}\theta_1^* - \beta_Q(1 - m\zeta)\frac{S}{N}\theta_2^* + (\gamma_Q + \delta_Q)\theta_6^* - \gamma_Q\theta_8^* \\ {}^C D^e \theta_7^* &= \beta_L(1 - m\zeta)\frac{S}{N}\theta_1^* - \beta_L(1 - m\zeta)\frac{S}{N}\theta_2^* + (\delta_L + \gamma_L)\theta_7^* - \gamma_L\theta_8^* \\ {}^C D^e \theta_8^* &= 0, \end{aligned} \tag{34}$$

with transversality conditions $\theta_j(T) = 0$, $j = 1, 2, \dots, 8$ and optimality conditions given by

$$H(S^*(t), \dots, R^*(t), \theta^*(t), v_i^*(t)) = \min_{0 \leq v_i \leq 1} H(S^*(t), \dots, R^*(t), \theta^*(t), v_i^*(t))$$

$$\begin{aligned} v_1^*(t) &= \min\{1, \max(0, \frac{(\theta_2^* - \theta_1^*)(\lambda_I + \lambda_A + \lambda_Q + \lambda_L + \lambda_{E_2})S^*}{D_1})\}, \\ v_2^*(t) &= \min\{1, \max(0, \frac{\theta_4^* I^*}{D_2})\}, \\ v_3^*(t) &= \min\{1, \max(0, \frac{\theta_5^* A^*}{D_3})\}. \end{aligned} \tag{35}$$

Proof: The adjoint system (34) i.e ${}^C D^e \theta_1^*, \dots, {}^C D^e \theta_8^*$ are obtained from the Hamiltonian H as

$$\frac{-d\theta_1}{dt} = \frac{\partial H}{\partial S}, \frac{-d\theta_2}{dt} = \frac{\partial H}{\partial E_1}, \dots, \frac{-d\theta_8}{dt} = \frac{\partial H}{\partial R}$$

with zero final time conditions or transversality conditions,

$\theta_1(T) = 0, \theta_2(T) = 0, \dots$ and $\theta_8(T) = 0$ and the characteristic of the fractional optimal control given by (35) is obtained by solving the Eqn $\frac{\partial H}{\partial v_1} = 0, \dots, \frac{\partial H}{\partial v_3} = 0$ on the interior of the control set and using the property of control space v .

8. Conclusions

Different mathematical paradigms can provide considerable insights and scientific evidences pertinent to any ongoing epidemic dynamics. Based on those valuable information, health officials and public health experts can set up potential control strategies to battle against any epidemic. From the emergence of the novel coronavirus in China, researchers and scientists are working relentlessly to develop various mathematical modeling approaches to gain a deeper understanding on the progression dynamics of COVID-19 in the world. In addition, in the absence of any safe, effective and widely available COVID-19 vaccine, different preventive measures are the most effective tool in combating against the virulent virus. On the basis of robust forecasting results of reliable epidemiological models, government officials can deploy different public health intervention strategies to control the rapid transmission of the virus. In this chapter, a compartmental mathematical has been designed to describe the transmission dynamics of the COVID-19 incorporating all possible real-life interactions and effective non-pharmaceutical interventions. Disease-free equilibrium (DFE) of the proposed model is found to be globally asymptotically stable (GAS), whenever control reproduction number ($\mathcal{R}_c$) less than unity. In addition, advanced forecasting techniques have also been applied for Argentina, Bangladesh, Brazil, Colombia and India to portray the future dynamics of the pandemic in near term. It has been enlightened in our study that mass-level using of highly effective face coverings could be a crucial factor in controlling the spread of coronavirus. Moreover, strict social-distancing measures and comprehensive contact-tracing are also effective strategies in battling against this pandemic. The public health implication of these insightful findings is government officials can undertake crucial clinical and public health decisions by analyzing all mathematical results and scientific evidences. Caputo-Fabrizio non-integer order derivative has been applied to solve the proposed mathematical model in fractional sense. We proved the existence of unique solution for the proposed fractional initial value problem. We proved the unconditional stability of the given technique. An important concern of fractional optimal control problem is given to suggest the health care measures for reducing the transmissibility of COVID-19 infection in the population.

Conflict of interest

This work does not have any conflicts of interest.

References

- [1] P. Folegatti, K. Ewer, P. Aley, B. Angus, S. e. a. Becker, Safety and immunogenicity of the chadox1 ncov-19 vaccine against sars-cov-2: a preliminary report of a phase 1/2, single-blind, randomised controlled trial, *The Lancet* 396 (10249) (2020) 447–512.

- [2] N. Ferguson, D. Laydon, G. Nedjati-Gilani, N. Imai, K. Ainslie, M. Baguelin, S. Bhatia, A. Boonyasiri, Z. M. Cucunubá, G. Cuomo-Dannenburg, A. Dighe, I. Dorigatti, H. Fu, K. Gaythorpe, W. Green, A. Hamlet, W. Hinsley, L. Okell, S. van Elsland, A. Ghani, Report 9: Impact of non-pharmaceutical interventions (npis) to reduce covid19 mortality and healthcare demand (2020). doi:10.25561/77482.
- [3] K. et. al., Effectiveness of isolation, testing, contact tracing, and physical distancing on reducing transmission of sars-cov-2 in different settings: a mathematical modelling study, *The Lancet Infectious Diseases* 20 (10) (2020) 1151–1160.
- [4] K. N. Nabi, H. Abboubakar, P. Kumar, Forecasting of covid-19 pandemic: From integer derivatives to fractional derivatives, *Chaos, Solitons & Fractals* 141 (2020) 110283.
- [5] K. N. Nabi, Forecasting covid-19 pandemic: A data-driven analysis, *Chaos, Solitons & Fractals* 139 (2020) 110046.
- [6] C. N. Ngonghala, E. Iboi, S. Eikenberry, M. Scotch, C. R. MacIntyre, M. H. Bonds, A. B. Gumel, Mathematical assessment of the impact of non-pharmaceutical interventions on curtailing the 2019 novel coronavirus, *Mathematical Biosciences* 325 (2020) 108364.
- [7] A. Babaei, H. Jafari, M. Ahmadi, A fractional order hiv/aids model based on the effect of screening of unaware infectives, *Mathematical Methods in the Applied Sciences* 42 (7) (2019) 2334–2343.
- [8] A. Babaei, H. Jafari, A. Liya, Mathematical models of hiv/aids and drug addiction in prisons, *The European Physical Journal Plus* 135 (5) (2020) 395.
- [9] K. Muhammad Altaf, A. Atangana, Dynamics of ebola disease in the framework of different fractional derivatives, *Entropy* 21 (3) (2019) 303.
- [10] S. Qureshi, A. Yusuf, Modeling chickenpox disease with fractional derivatives: From caputo to atangana-baleanu, *Chaos, Solitons & Fractals* 122 (2019) 111–118.
- [11] M. Rivero, J. J. Trujillo, L. Vázquez, M. P. Velasco, Fractional dynamics of populations, *Applied Mathematics and Computation* 218 (3) (2011) 1089–1095.
- [12] S. B. Bastos, D. O. Cajueiro, Modeling and forecasting the early evolution of the covid-19 pandemic in brazil, *arXiv preprint arXiv:2003.14288* (2020).
- [13] S. Çakan, Dynamic analysis of a mathematical model with health care capacity for pandemic covid-19, *Chaos, Solitons & Fractals* (2020) 110033.

- [14] C. Covid, C. COVID, C. COVID, S. Bialek, R. Gierke, M. Hughes, L. A. McNamara, T. Pilishvili, T. Skoff, Coronavirus disease 2019 in children—united states, february 12–april 2, 2020, Morbidity and Mortality Weekly Report 69 (14) (2020) 422.
- [15] D. Cucinotta, M. Vanelli, Who declares covid-19 a pandemic., Acta bio-medica: Atenei Parmensis 91 (1) (2020) 157–160.
- [16] V. S. Erturk, P. Kumar, Solution of a covid-19 model via new generalized caputo-type fractional derivatives, Chaos, Solitons & Fractals (2020) 110280.
- [17] W. Gao, P. Veeresha, H. M. Baskonus, D. Prakasha, P. Kumar, A new study of unreported cases of 2019-ncov epidemic outbreaks, Chaos, Solitons & Fractals (2020) 109929.
- [18] S. Jiang, S. Xia, T. Ying, L. Lu, A novel coronavirus (2019-ncov) causing pneumonia-associated respiratory syndrome, Cellular & molecular immunology 17 (5) (2020) 554–554.
- [19] P. Kumar, V. Suat Erturk, The analysis of a time delay fractional covid-19 model via caputo type fractional derivative, Mathematical Methods in the Applied Sciences (2020).
- [20] Q. Lin, S. Zhao, D. Gao, Y. Lou, S. Yang, S. S. Musa, M. H. Wang, Y. Cai, W. Wang, L. Yang, et al., A conceptual model for the outbreak of coronavirus disease 2019 (covid-19) in wuhan, china with individual reaction and governmental action, International journal of infectious diseases (2020).
- [21] N. Zhu, D. Zhang, W. Wang, X. Li, B. Yang, J. Song, X. Zhao, B. Huang, W. Shi, R. Lu, et al., A novel coronavirus from patients with pneumonia in china, 2019, New England Journal of Medicine (2020).
- [22] R. L. e. a. Tillett, Genomic evidence for reinfection with sars-cov-2: a case study, The Lancet Infect. Dis. (2020).
- [23] W. Walter, Ordinary Differential Equations, Springer, 1998.
- [24] COVID-19 Github Repository, Center for Systems Science and Engineering at Johns Hopkins University, <https://github.com/CSSEGISandData/COVID-19> (Date accessed November 16, 2020).
- [25] P. Riessche, J. Watmough, Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission, Math. Biosci. 180 (1-2) (2002) 29—48.
- [26] O. Diekmann, J. A. P. Heesterbeek, J. A. J. Metz, Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (sars-cov2), J. Math. Biol. 28 (4) (1990) 365–382.
- [27] R. Li, S. Pei, B. Chen, Y. Song, T. Zhang, W. Yang, J. Shaman, Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (sars-cov2), Science 368 (6490) (2020) 489–493.

- 406 [28] B. Tang, N. L. Bragazzi, Q. Li, S. Tang, Y. Xiao, J. Wu, An updated estimation of the risk of trans-
407 mission of the novel coronavirus (2019-ncov), *Infectious Disease Modelling* 5 (2020) 248—255.
- 408 [29] F. e. a. Zhou, Clinical course and risk factors for mortality of adult inpatients with covid-19 in wuhan,
409 china: a retrospective cohort study, *The Lancet* 395 (10229) (2020) 1054–1062.
- 410 [30] Y. Liu, A. A. Gayle, A. Wilder-Smith, J. Rocklöv, The reproductive number of covid-19 is higher
411 compared to sars coronavirus, *Journal of travel medicine* (2020).
- 412 [31] K. N. Nabi, C. N. Podder, Sensitivity analysis of chronic hepatitis c virus infection with immune response
413 and cell proliferation, *International Journal of Biomathematics* 13 (2020) 2050017.
- 414 [32] I. Podlubny, *Fractional differential equations: an introduction to fractional derivatives, fractional dif-*
415 *ferential equations, to methods of their solution and some of their applications*, Elsevier, 1998.
- 416 [33] A. Jajarmi, D. Baleanu, A new fractional analysis on the interaction of hiv with cd4+ t-cells, *Chaos,*
417 *Solitons & Fractals* 113 (2018) 221–229.
- 418 [34] J. Losada, J. J. Nieto, Properties of a new fractional derivative without singular kernel, *Progr. Fract.*
419 *Differ. Appl* 1 (2) (2015) 87–92.