1 Fatty acid profiles, cholesterol composition, and nutritional quality indices of 37 commonly
2 consumed local foods in Kuwait in relation to cardiovascular health

3

4 Hanan A. Al-Amiri^{1*}, Nisar Ahmed^{2¶}, Tahani Al-Sharrah^{2¶}

5 ¹ Food and Nutrition Program, Environment and Life Sciences Research Center, Kuwait
6 Institute for Scientific Research, Shuwaikh, Kuwait

7 ² Environment Pollution and Climate Program, Environment and Life Sciences Research
8 Center, Kuwait Institute for Scientific Research, Shuwaikh, Kuwait

9

10 * Corresponding author

11 E-mail: hamiri@kisr.edu.kw, hanan2303@yahoo.com

12 ORCID ID: <https://orcid.org/0000-0002-8338-5780>

13

14

15

16 ¶These authors contributed equally to this work

17

18

19

20

21

22

23

24

25

26

27

28 **Abstract**

29 Dietary fatty acids and cholesterol content are proved, by many research studies, to be
30 associated with various health conditions, including cardiovascular health. Knowledge of the
31 composition of these nutrients in food is essential for proper planning of health programs.
32 The present study aimed at assessing the fatty acid profile, fatty acid nutritional quality, and
33 cholesterol composition of 37 foods commonly consumed in the State of Kuwait and the
34 potential impact of these foods on cardiovascular risk. Fatty acid profile was determined by
35 gas chromatography-flame ionization detector into four types: saturated, monounsaturated,
36 polyunsaturated, and trans fatty acids. Nutritional quality was calculated using the
37 atherogenic index, thrombogenic index, hypocholesterolemic/hypercholesterolemic fatty acid
38 ratio, polyunsaturated fatty acid/ saturated fatty acid ratio, and n-3/n-6 fatty acids ratio.
39 Determination of cholesterol was performed by gas chromatography. Saturated fatty acid
40 levels ranged from 0.01–21.83, monounsaturated fatty acids 0.01–25.51, and 0.013–22.87
41 g/100 g edible portion of food. The predominant fatty acids identified in all studied foods
42 were C18:2c (n-3), C16:0, and C18:1c, with values 0.45–56.52, 10.12–44.90, and
43 16.99–42.56% of total fatty acids, respectively. The trans fatty acid content was low in all
44 foods. Cholesterol levels varied between traces (<0.05) and 454.79 mg/100 g edible portions
45 of food. Results show that seafood, rice-based, seed-based, and vegetable-based foods had
46 better nutritional quality in terms of the fatty acids content, as indicated by the
47 polyunsaturated fatty acid/saturated fatty acid and n-3/n-6 ratios, low thrombogenicity indices,
48 and high hypocholesterolemic/hypercholesterolemic fatty acid ratios. Cholesterol and fatty
49 acid data obtained in the present study will be of special interest for many studies, including
50 nutrition-related health research, and will help policymakers in proper strategies for health
51 programs.

52

53 Introduction

54 Dietary intake of fatty acids and cholesterol play an important role in the etiology of
55 cardiovascular disease (CVD) [1, 2], a leading cause of death in most countries [3].
56 According to the World Health Organization (WHO), CVD accounts for 17.9 million deaths
57 annually [3].

58 Several experimental and epidemiological studies have proven the strong association
59 between specific types of dietary fatty acids and blood cholesterol levels and therefore, the
60 risk of CVD [2, 4-8]. The degree of CVD risk may vary according to the type and level of
61 dietary fatty acids. Dietary saturated fatty acid (SFA), trans fatty acid (TFA), and/or improper
62 ratio of n-3/n-6 fatty acids are common dietary causes of CVD [6, 9]. Dietary SFAs elevate
63 total and low-density lipoprotein (LDL) blood cholesterol, whereas TFAs elevate total blood
64 cholesterol and LDL cholesterol, and lower blood high-density lipoprotein (HDL) cholesterol
65 [9, 10]. However, dietary polyunsaturated fatty acid (PUFA) and monounsaturated fatty acid
66 (MUFA) generally lower both total and LDL cholesterol levels, thus may cause reduction in
67 the incidence of CVD [11, 12], and possibly the prevention of some cancers, asthma, diabetes
68 and hypertension [13]. Additionally, a strong relationship has been revealed between cellular
69 cholesterol levels and Alzheimer's disease [14]. Nutritional strategies to achieve a better
70 balance of fatty acids in the diet by decreasing the intake of SFA and TFA as well as
71 cholesterol, appear to be significant and effective to reduce the potential risk of these diseases.
72 Thus, knowledge of the fatty acid and cholesterol composition of foods is essential before
73 any nutritional strategy can be implemented.

74 This article aims to provide detailed data on the fatty acid profiles and nutritional quality,
75 as well as cholesterol contents of 37 local foods commonly consumed in the State of Kuwait.
76 This data will contribute valuable information to fatty acid and cholesterol composition tables.
77 The enrichment of food composition tables with reliable analytical data on local foods is

78 significant for obtaining convincing results in future nutritional-health related research and in
79 the development and implementation of nutritional policies aimed at improving health.

80 **Materials and methods**

81 **Food sampling**

82 Top thirty seven most commonly consumed local foods among Kuwaitis were selected
83 from “Kuwait Total Diet Study” data. The selected foods included 21 foods commonly
84 cooked and prepared at home, and 16 ready-made foods often purchased by locals rather than
85 cooked at home. Each ready-made food was purchased from three different
86 restaurants/confectioneries based on its popularity among Kuwaitis. The most common
87 brands of cream cheese spread (four brands), triangle processed cheese (three brands), and
88 white cheese low fat (one brand “Al-Wafra”) were purchased from retailers.

89 As for cooked foods, data on the ingredients, their quantities, and cooking procedures
90 were collected based on a field survey of 250 Kuwaiti households. The food recipes were
91 standardized by identifying the foremost ingredients of each recipe and their weight ratio
92 from the total, ensuring that the coefficient of variation (CV) did not exceed 20%. Details on
93 the major ingredients of the 37 foods are given in S1 Table.

94 The cooked foods were prepared at the Ministry of Health (MOH) kitchen based on the
95 standardized recipes under the supervision of experienced Kuwaiti cooks. All ingredients
96 including water were weighed, and three identical preparations of each food were cooked
97 separately. Foods were cleared of inedible parts, if any, including bones and skin. Each
98 cooked and ready-made food was then homogenized thoroughly in a blender and sampled for
99 moisture analysis. The remaining food sample was dried in a freeze drier (VirTis, Unitop
100 800L, USA), ground, transferred into airtight bottle containers, labeled, and stored in a deep
101 freezer at -18°C until analysis.

Fatty acid analysis

Total fat content in g/100 g edible portion (EP) of food was determined gravimetrically following Soxhlet extraction (Tecator Soxtec system, HT 1043 extraction unit, Herndon, VA, USA) according to the method given in AOAC [15]. Fatty acids were extracted from each sample in triplicate according to Folch, Lees, and Sloane Stanley [16] with a modification by Segura and Lopez-Bote [17]. Briefly, 0.5 g of freeze dried food sample was homogenized with 20 ml chloroform:methanol (2:1 v/v) solution. The samples were ultrasonicated at room temperature for 15 min. After allowing to settle for a few minutes, the homogenate was filtered. The filtrate was collected in a tube and NaCl solution was added. The tube centrifuged for 20 min at approximately 3000 x rpm. The upper layer of methanol:water was removed, and the lower layer was filtered through anhydrous sodium sulphate and collected in a volumetric flask. The solvent was evaporated under nitrogen, the residual fat was esterified, and methyl esters of fatty acids (FAME) were prepared as described by the American Oil Chemists' Society (AOCS) Official Method Ce 2-66 [18]. A 0.5 µL esterified sample was injected into a gas chromatograph (GC) (Agilent 6890 Series, Agilent Technologies, Palo Alto, CA, USA) coupled with a flame ionization detector (FID). A cyano-polysiloxane (CP-Sil 88 for FAME, part number CP7489) capillary column (100 m × 0.25 mm, 0.20 µm; Agilent Technologies, Palo Alto, CA, USA) was used for the FAMEs separation under the following instrumental conditions: injector and FID detector temperatures were 260 and 270°C, respectively, with injector split ratio of 1:50, and carrier gas helium with a constant flow rate of 1.0 mL/min. The initial oven temperature was 80 °C, at which it was held for 5 min, followed by an increase to 220 °C at a rate of 4 °C/min then was held for 5 min, and finally the temperature was increased to 240 °C at a rate of 1 °C/min and was held for 10 min. A 37-component FAMEs standard mixture was used to identify the fatty acids, the mixture contained FAMEs ranging from C4:0 to C22:6. The FAMEs were

identified by comparing the retention times of standard peaks with the retention times of sample peaks. The quantification of the fatty acids was performed by the conversion of the peak areas in percentage of the extract using a GC Workstation software. The fatty acid content was reported as percentage of the total fatty acids. In order to compile data in food composition tables, fatty acid data need to be transformed to per 100 g EP on fresh weight basis, to create internationally interchangeable food composition data. The fatty acid composition in g/100 g EP was calculated from the corresponding fatty acid (percentage of total fatty acids), following the method described by Paul and Southgate [19].

Cholesterol analysis

Cholesterol was analyzed in triplicate via GC (Agilent 6890 Series, Agilent Technologies, Palo Alto, CA, USA) according to a method described by Naeemi et al. [20]. Each sample was spiked with the internal standard 5- α -cholestane after saponification with a saturated methanolic KOH solution; hexane was then added, centrifuged at 2000 rpm for 2 min, and finally 0.5 μ L of the upper layer was injected into the GC for cholesterol analysis.

The GC conditions used in this study were as follows: injector and detector temperatures were 300°C, with injector split ratio of 1:25, and carrier gas helium with flow rate of 1.0 mL/min. The initial oven temperature was 250 °C, the holding time was 2 min, and the temperature rise rate was 3 °C/min.

Fatty acid quality

Fatty acid quality was determined using the atherogenic index (AI) and thrombogenic index (TI) according to Ulbricht & Southgate [21]. The hypocholesterolemic to hypercholesterolemic (H/H) fatty acid ratio index which considers the specific effects of fatty acids on cholesterol metabolism, was determined as described by Testi et al. [22]. The indices were calculated using the following equations:

$$AI = [(C12:0 + (4 \times C14:0) + C16:0)] / (\sum MUFA + \sum n-6 + \sum n-3)$$

$$\begin{aligned}
 & TI = (C14:0 + C16:0 + C18:0) / [(0.5 \times \sum MUFA) + (0.5 \times \sum n-6) + (3 \times \sum n-3) + (\sum n-3 / \sum n-6)] \\
 & H/H = (C18:1c + C18:2n-6 + C20:4n-6 + C18:3n-3 + C20:5n-3 + C22:5n-3 + C22:6n-3) / (C14:0 + C16:0) \\
 & \text{where:} \\
 & \sum MUFA = \text{sum of monounsaturated fatty acids,} \\
 & \sum n-6 = \text{sum of n-6 fatty acids, and} \\
 & \sum n-3 = \text{sum of n-3 fatty acids.}
 \end{aligned}$$

Further indices were calculated as PUFA to SFA (P/S) ratio, and n-3/n-6 fatty acids ratio.

Statistical analysis

Each food sample was analyzed in triplicate for total lipid, fatty acid, and cholesterol composition. Fatty acid data were reported as the mean value on the wet weight of the edible portion of food in g/100 g, and as percentage of the total fatty acids. Cholesterol data were reported as the mean $\pm$ standard deviation (SD) in mg/100 g EP of food. The minimum and maximum values for each fatty acid and cholesterol content were interpreted as a measure of range.

Results and discussion

Fatty acid profile

Investigated foods in the current study were classified into nine categories: seafood (n=3), soup (n=3), cheese (n=3), rice-based (n=6), desserts (n=11), seed-based (n=2), meat-based (n=4), sandwiches (n=2), and vegetable-based (n=3) (Table 1). The classification of the foods is based on major ingredients in each recipe, as well as the common classification (i.e. cheese, soup, desserts, and sandwiches). All cheese, desserts (except *Khabeesa* “cooked semolina with oil and sugar” and *Aseeda* “cooked wheat flour with sugar”), sandwiches, one meat-based food (*Arayes* “pita bread stuffed with minced lamb”), and one seed-based food

(*Harda* “soft sesame butter, tahini”) were ready-made and purchased from the most popular restaurants and bakeries/confectioneries. The detected fatty acids were classified into four types: SFA, MUFA, PUFA, and TFA. The fatty acid compositions were expressed both as g/100 g EP and as % of the total fatty acids.

Saturated fatty acids

SFA was the most dominant type of fatty acids among analyzed foods (Fig 1). The SFA content was observed with values ranged from 0.01 to 21.85 g/100 g EP (S2 Table). The highest range of SFA content (8.32–21.85 g/100 g EP) and percentage of total fatty acids (62.27–67.33%) was found in cheese. Meat-based foods and sandwiches showed high levels of SFA ranged from 40.65 to 52.09, and 39.43 to 44.91% of total fatty acids, respectively. *Knafah* (shredded phyllo dough and cheese in sugar syrup) and *Gherayba* (shortbread cookie) showed high SFA concentrations of 10.96 and 15.95 g/100 g EP, respectively, and high SFA percentage of total fatty acids of 62.92 and 59.98%, respectively (S2 Table and Table 1). The high SFA content of the dessert *Gherayba* is attributed to the use of ghee in the recipe and to the use of cheese and butter in *Knafah*.

Fig 1. Concentrations in g/100 g edible portion of food: SFA (saturated fatty acid); MUFA (monounsaturated fatty acid); PUFA (polyunsaturated fatty acid); TFA (trans fatty acid); of 37 commonly consumed local foods in Kuwait. SFA was the most dominant type of fatty acids among foods, while TFA was found in very little amounts.

Table 1. Fatty acid composition profile on the wet weight of the edible portion (percentage of the total fatty acids) for 37 commonly consumed local foods in the State of Kuwait

Food	Saturated Fatty Acid (SFA)															ΣSFA
	4:0	6:0	10:0	12:0	13:0	14:0	15:0	16:0	17:0	18:0	20:0	21:0	22:0	23:0	24:0	
Seafood																
Marag Samak Maleh (Spanish mackerel broth)	tr	tr	tr	tr	tr	3.26	0.63	23.15	0.70	6.26	0.40	tr	tr	tr	tr	34.41
Kabab Rubyan (Shrimps kebab)	tr	tr	tr	tr	tr	1.20	0.37	18.40	0.77	5.84	0.40	tr	tr	tr	4.74	31.72
Marag Rubyan (Shrimps broth)	tr	tr	tr	tr	tr	0.26	0.17	13.25	0.42	3.49	0.44	0.05	0.06	tr	0.17	18.32
Soup																
Shorbat Shaereya (Vermicelli soup)	tr	tr	tr	tr	tr	1.77	tr	18.10	tr	3.52	tr	tr	tr	6.09	0.09	29.57
Shorbat Dajaj (Chicken soup)	tr	tr	tr	tr	0.08	6.71	0.67	30.55	0.35	9.87	0.12	tr	0.16	tr	tr	48.50
Shorbat Harees (Soup of peeled wheat with lamb)	tr	tr	tr	tr	tr	2.05	0.79	24.56	1.86	28.19	0.27	0.13	tr	tr	tr	57.86
Cheese																
Cream Cheese Spread, Regular	tr	tr	tr	tr	0.10	13.11	1.47	36.91	0.40	14.88	0.21	tr	tr	tr	tr	67.08
Triangle Processed Cheese	tr	tr	tr	tr	0.20	13.58	0.99	37.58	0.43	14.11	0.25	0.06	0.08	tr	0.05	67.33
“Al-Wafra” White Cheese, Low Fat	tr	tr	tr	tr	tr	10.70	1.10	36.90	1.13	12.43	tr	tr	tr	tr	tr	62.27
Rice-based																
Aish Muhammad (Sweet rice)	tr	tr	tr	tr	tr	tr	tr	12.71	tr	2.61	0.56	tr	tr	tr	tr	15.88
Aish Khothra (Rice with vegetables)	tr	tr	tr	tr	tr	0.09	tr	11.86	0.10	2.26	0.46	0.08	0.16	tr	0.09	15.09
Aish Fagae ma Laham (Rice, truffle with lamb)	tr	tr	tr	tr	tr	0.24	0.08	14.16	0.33	5.42	0.62	0.10	0.26	tr	0.17	21.39
Aish Bajella ma Shbent (Rice with green fava beans and dill)	tr	tr	tr	tr	tr	0.11	tr	13.31	0.12	2.47	0.59	tr	tr	tr	0.10	16.69
Aish Tomat (Rice with tomato)	tr	tr	tr	tr	tr	tr	3.30	28.96	tr	5.90	tr	tr	tr	tr	tr	38.15
Kushari (Rice with macaroni and mung beans)	tr	tr	tr	tr	tr	8.13	tr	12.10	tr	9.08	tr	tr	tr	tr	6.89	36.20
Desserts																
Gherayba (Shortbread cookie)	tr	tr	tr	tr	0.07	8.17	0.89	41.24	0.08	9.03	0.34	tr	0.04	tr	0.05	59.90
Knafah (Shredded phyllo dough and cheese in sugar syrup)	tr	tr	tr	tr	tr	11.77	1.27	38.38	0.77	10.74	tr	tr	tr	tr	tr	62.92
Baklava (Phyllo pastry with nuts, Baklava)	tr	tr	tr	tr	tr	6.63	0.77	32.68	tr	9.86	tr	tr	tr	tr	tr	49.93
Qatayef (Dumpling filled with mixed nuts)	tr	tr	tr	tr	tr	1.47	0.16	14.56	0.18	3.99	0.64	0.54	0.18	tr	0.08	21.79
Samsamia (Sesame bar)	tr	tr	tr	tr	tr	tr	tr	17.07	tr	6.73	tr	tr	tr	tr	tr	23.80

Table 1. (Continued)

Food	Saturated Fatty Acid (SFA)															ΣSFA
	4:0	6:0	10:0	12:0	13:0	14:0	15:0	16:0	17:0	18:0	20:0	21:0	22:0	23:0	24:0	
<i>Khabeesa</i> (Cooked semolina with oil and sugar)	tr	tr	tr	tr	tr	0.38	tr	12.17	0.44	2.72	0.48	tr	tr	1.94	0.13	18.26
<i>Darabeel</i> (Rolled dough stuffed with sugar and cinnamon)	tr	tr	tr	tr	tr	1.09	0.19	44.90	0.19	4.77	0.54	tr	tr	tr	tr	51.67
<i>Zalabia</i> (Jalebi)	tr	tr	tr	tr	tr	1.13	0.06	40.87	0.23	5.11	0.34	0.10	0.06	tr	tr	47.91
<i>Aseeda</i> (Cooked wheat flour with sugar)	tr	tr	tr	tr	tr	2.23	tr	22.53	tr	3.50	tr	tr	tr	tr	tr	28.26
<i>Halwa</i> (Sweet wheat starch)	tr	tr	tr	tr	0.05	6.33	0.61	22.27	0.33	9.30	0.31	0.05	0.05	tr	0.06	39.36
<i>Baksam</i> (Rusk)	tr	tr	tr	tr	tr	0.85	tr	32.48	0.46	4.22	0.61	1.82	1.12	tr	0.47	42.03
Seed-based																
<i>Heso</i> (Garden cress and fenugreek seeds stew)	tr	tr	tr	tr	tr	0.19	tr	10.88	0.09	2.51	tr	0.05	0.38	tr	0.15	14.25
<i>Harda</i> (Soft sesame butter, Tahini)	tr	tr	tr	tr	tr	tr	tr	10.12	tr	6.80	0.72	0.06	0.20	tr	tr	17.89
Meat-based																
<i>Madhrooba</i> (Crushed wheat with lamb)	tr	tr	tr	tr	tr	1.57	0.49	24.56	1.09	16.46	0.21	0.21	tr	tr	tr	44.58
<i>Arayes</i> (Pita bread stuffed with minced lamb)	tr	tr	tr	tr	tr	2.30	0.87	22.84	1.88	22.60	0.22	0.07	0.07	tr	tr	50.86
<i>Hameesat Kabdah</i> (Fried lamb liver in tomato sauce)	tr	tr	tr	tr	tr	0.53	0.44	22.07	1.78	26.62	0.32	0.12	0.13	tr	0.07	52.09
<i>Keema</i> (Minced lamb fried with peas)	tr	tr	tr	tr	tr	2.77	0.49	23.04	0.75	12.79	0.29	0.26	0.21	tr	0.05	40.65
Sandwich																
Cheeseburger	tr	tr	tr	tr	tr	3.60	0.67	28.68	0.87	11.10	tr	tr	tr	tr	tr	44.91
Chickenburger	tr	tr	tr	tr	tr	2.40	tr	27.62	tr	9.41	tr	tr	tr	tr	tr	39.43
Vegetable-based																
<i>Hameesat Mushroom</i> (Fried mushrooms in tomato sauce)	tr	tr	tr	tr	tr	0.07	tr	12.13	tr	2.03	0.43	0.07	0.21	tr	0.10	15.03
<i>Hameesat Fagae</i> (Fried truffles in tomato sauce)	tr	tr	tr	tr	tr	0.05	tr	12.59	tr	2.34	0.53	0.07	0.31	tr	0.16	16.06
<i>Marag Hawa</i> (Vegetable broth)	tr	tr	tr	tr	tr	0.07	tr	12.11	tr	2.28	0.51	0.08	0.17	tr	0.11	15.32

205 Table 1. (Continued)

206

Food	Monounsaturated Saturated Fatty Acid (MUFA)										Polyunsaturated Fatty Acid (PUFA)								Trans Fatty Acid
	14:1	15:1	16:1	17:1	18:1 _t	18:1 _c	20:1	22:1	24:1	ΣMUFA	18:2 _t	18:2 _c	18:3	20:4	20:5	22:2	22:6	ΣPUFA	ΣTFA
												n-6	n-3	n-6	n-3	n-3	n-3		
Seafood																			
<i>Marag Samak Maleh</i> (Spanish mackerel broth)	tr	tr	4.13	tr	tr	20.99	1.93	tr	tr	27.05	tr	29.38	tr	1.23	tr	tr	7.93	38.54	tr
<i>Kabab Rubyan</i> (Shrimps kebab)	tr	tr	2.44	0.40	tr	23.24	1.77	tr	tr	27.85	tr	33.49	tr	2.34	tr	tr	4.61	40.43	tr
<i>Marag Rubyan</i> (Shrimps broth)	tr	tr	0.76	0.20	tr	26.64	1.34	tr	0.07	29.01	0.08	49.34	0.06	0.07	0.09	1.72	1.32	52.68	0.08
Soup																			
<i>Shorbat Shaereya</i> (Vermicelli soup)	tr	tr	tr	tr	tr	21.57	3.09	tr	tr	24.66	tr	45.78	tr	tr	tr	tr	tr	45.78	tr
<i>Shorbat Dajaj</i> (Chicken soup)	0.49	0.19	3.22	tr	tr	30.62	1.62	0.18	0.15	36.48	0.17	14.63	0.12	tr	tr	tr	0.10	15.02	0.17
<i>Shorbat Harees</i> (Soup of peeled wheat with lamb)	0.41	0.34	1.16	tr	tr	36.31	0.89	tr	tr	39.10	0.24	2.81	tr	tr	tr	tr	tr	3.05	0.24
Cheese																			
Cream Cheese Spread, Regular	0.75	0.32	1.57	0.32	tr	26.32	0.98	tr	1.33	31.60	0.17	1.15	tr	tr	tr	tr	tr	1.32	0.17
Triangle Processed Cheese	0.90	0.33	1.83	0.35	tr	27.61	0.87	tr	tr	31.88	0.28	0.45	tr	0.06	tr	tr	tr	0.79	0.28
“Al-Wafra” White Cheese, Low Fat	0.90	tr	3.30	0.40	0.30	29.53	0.57	tr	tr	35.00	tr	2.73	tr	tr	tr	tr	tr	2.73	0.90
Rice-based																			
<i>Aish Muhammad</i> (Sweet rice)	tr	tr	tr	tr	tr	30.06	1.40	tr	tr	31.46	tr	52.66	tr	tr	tr	tr	tr	52.66	tr
<i>Aish Khothra</i> (Rice with vegetables)	tr	tr	0.11	tr	tr	27.89	1.28	tr	tr	29.29	tr	55.56	0.07	tr	tr	tr	tr	55.62	tr
<i>Aish Fagae ma Laham</i> (Rice, truffle, lamb)	tr	tr	0.17	0.18	tr	32.92	0.91	tr	0.16	34.34	0.15	43.95	tr	tr	0.17	tr	tr	44.28	0.15
<i>Aish Bajella ma Shbent</i> (Rice, fava beans and dill)	tr	tr	0.09	tr	0.38	28.01	1.23	tr	tr	29.71	tr	53.52	0.07	tr	tr	tr	tr	53.59	0.09
<i>Aish Tomat</i> (Rice with tomato)	tr	tr	tr	tr	tr	31.32	tr	tr	tr	31.32	tr	30.52	tr	tr	tr	tr	tr	30.52	tr
<i>Kushari</i> (Rice with macaroni and mung beans)	tr	tr	6.29	tr	tr	18.45	5.58	tr	tr	30.32	tr	33.48	tr	tr	tr	tr	tr	33.48	tr
Desserts																			
<i>Gherayba</i> (Shortbread cookie)	0.70	0.17	0.37	0.20	tr	30.43	0.85	tr	tr	32.72	0.09	7.15	tr	0.06	tr	0.05	0.05	7.39	0.09
<i>Knafah</i> (Shredded phyllo dough in sugar syrup)	0.83	tr	1.67	tr	1.53	27.24	0.60	tr	tr	31.87	tr	5.20	tr	tr	tr	tr	tr	5.20	1.53
<i>Baklava</i> (Phyllo pastry with nuts, Baklava)	tr	tr	1.20	tr	1.50	34.18	0.73	tr	tr	37.61	tr	12.46	tr	tr	tr	tr	tr	12.46	1.50
<i>Qatayef</i> (Dumpling filled with mixed nuts)	0.11	tr	0.26	tr	tr	28.53	1.02	tr	0.20	30.12	0.07	47.89	0.05	0.08	tr	tr	tr	48.09	0.07
<i>Samsamia</i> (Sesame bar)	tr	tr	tr	tr	tr	37.73	tr	tr	tr	37.73	tr	38.47	tr	tr	tr	tr	tr	38.47	tr

207 Table 1. (Continued)

Food	Monounsaturated Saturated Fatty Acid (MUFA)										Polyunsaturated Fatty Acid (PUFA)								Trans F
	14:1	15:1	16:1	17:1	18:1 _t	18:1 _c	20:1	22:1	24:1	ΣMUFA	18:2 _t	18:2 _c	18:3	20:4	20:5	22:2	22:6	ΣPUFA	ΣTFA
											n-6	n-3	n-6	n-3	n-3	n-3			
<i>Khabeesa</i> (Cooked semolina with oil and sugar)	tr	tr	0.30	tr	tr	27.48	1.55	0.57	0.31	30.21	tr	51.53	tr	tr	tr	tr	tr	51.53	tr
<i>Darabeel</i> (Rolled dough stuffed with sugar)	tr	tr	0.18	tr	0.27	35.88	0.80	tr	0.14	37.27	0.16	10.90	tr	tr	tr	tr	tr	11.06	0.44
<i>Zalabia</i> (Jalebi)	tr	tr	0.18	tr	0.25	39.35	0.34	tr	2.78	42.90	0.15	9.05	tr	tr	tr	tr	tr	9.20	0.40
<i>Aseeda</i> (Cooked wheat flour with sugar)	tr	tr	tr	tr	tr	16.99	5.13	tr	tr	22.13	tr	49.62	tr	tr	tr	tr	tr	49.62	tr
<i>Halwa</i> (Sweet wheat starch)	0.53	0.14	0.75	0.23	tr	34.04	3.02	tr	tr	38.70	0.15	21.50	0.33	tr	tr	tr	tr	21.97	0.15
<i>Baksam</i> (Rusk)	tr	tr	tr	tr	tr	30.20	2.26	0.96	tr	33.43	0.65	23.66	0.23	tr	tr	tr	tr	24.54	0.65
Seed-based																			
<i>Heso</i> (Garden cress and fenugreek seeds stew)	tr	tr	0.12	tr	tr	28.86	9.96	0.12	0.24	39.29	0.20	45.00	1.25	tr	tr	tr	tr	46.45	0.20
<i>Harda</i> (Soft sesame butter, Tahini)	0.05	tr	0.14	0.06	tr	42.56	0.52	tr	tr	43.33	tr	38.80	tr	tr	tr	tr	tr	38.80	tr
Meat-based																			
<i>Madhrooba</i> (Crushed wheat with lamb)	0.29	tr	1.00	0.39	tr	38.28	1.41	tr	tr	41.37	0.31	13.74	tr	tr	tr	tr	tr	14.05	0.31
<i>Arayes</i> (Pita bread stuffed with minced lamb)	0.37	0.29	1.00	0.85	4.47	35.79	1.13	tr	tr	43.89	0.31	4.93	tr	tr	tr	tr	tr	5.24	4.78
<i>Hameesat Kabdah</i> (Fried lamb liver in sauce)	0.18	0.19	1.13	0.90	tr	36.41	0.56	tr	tr	39.36	0.28	8.19	tr	0.06	tr	tr	tr	8.54	0.28
<i>Keema</i> (Minced lamb fried with peas)	0.75	0.32	2.33	0.49	1.47	35.68	1.05	tr	tr	42.10	0.12	16.93	0.12	0.05	tr	tr	tr	17.23	1.55
Sandwich																			
Cheeseburger	0.63	tr	2.37	0.53	2.33	29.18	2.03	tr	tr	37.07	tr	17.44	tr	0.57	tr	tr	tr	18.01	2.33
Chickenburger	tr	tr	0.80	tr	1.37	30.49	2.43	tr	tr	35.09	tr	25.48	tr	tr	tr	tr	tr	25.48	1.37
Vegetable-based																			
<i>Hameesat Mushroom</i> (Fried mushroom in sauce)	0.08	tr	0.16	tr	tr	26.88	1.17	tr	tr	28.28	tr	56.52	0.17	tr	tr	tr	tr	56.69	tr
<i>Hameesat Fagae</i> (Fried truffles in tomato sauce)	tr	tr	0.19	tr	tr	28.49	1.01	tr	tr	29.68	tr	53.97	0.09	0.05	0.16	tr	tr	54.28	tr
<i>Marag Hawa</i> (Vegetable broth)	tr	tr	0.11	tr	tr	27.84	1.60	tr	tr	29.54	tr	55.07	0.07	tr	tr	tr	tr	55.14	tr

tr: trace (values for peaks < 0.05%)

t: trans fatty acid; c: cis fatty acid

ΣTFA: 18:1_t+18:2_t

According to the joint Food and Agriculture Organization (FAO)/WHO [23] and FAO [24] reports, the SFAs with the most determinant effects on blood LDL cholesterol elevation are lauric (C12:0), myristic (C14:0), and to some extent palmitic (C16:0) acids, whereas stearic acid (C18:0) has been shown to reduce total blood cholesterol levels. Lauric acid was detected in trace amounts in all food samples (<0.05% of total fatty acids), whereas, C14:0 proportion was low with values ranging between trace amounts (<0.05%) and 8.2% of total fatty acids, except for cheese dishes (10.70–13.58%) and *Knafah* (11.77%) (Table 1). C16:0 was the most abundant SFA in 35 of 37 foods. The highest and lowest concentrations of C16:0, 12.02 and 0.01 g/100 g EP, were detected in cream cheese and *Shorbat Shaereya* (vermicelli soup), respectively (S2 Table). In contrast, the highest and lowest percentage of palmitic acid of the total fatty acids were identified in dessert *Darabeel* (rolled dough stuffed with sugar and cinnamon; 44.90%) and seed-based food *Harda* (10.12%), respectively. The C16:0 percent of the total fatty acids was predominant in cheese dishes (36.90–37.58%) and in the following desserts: *Baksam* (rusk; 32.48%), *Baklawaw* (baklava; 32.68%), *Knafah* (38.38%), *Zalabia* (jalebi; 40.87%), and *Gherayba* (41.24%) (Table 1). Thus, we recommend that these foods be consumed in moderation or the butter and ghee used in desserts preparation can be substituted with unsaturated fatty acids oil. The dominance of SFA in the analyzed foods and of palmitic acid in SFA is consistent with earlier studies on similar groups of foods from Saudi Arabia [25].

Monounsaturated fatty acids

The highest concentration of MUFA (25.66 g/100 g EP) was demonstrated by *Harda*, followed by *Samsamia* (sesame bar; 14.07 g/100 g EP) and cream cheese (10.29 g/100 g EP) (S2 Table). MUFA was the most dominant type of fatty acids in only two food samples *Keema* (minced lamb with peas) and *Harda* (42.10 and 43.33%, respectively). Oleic acid (C18:1c) was the most dominant MUFA across all 37 food samples. The oleic acid

percentage of total fatty acids was the highest in *Harda* (42.56%), the major ingredient of this food is sesame seeds, known to be high in oleic acid [26].

Polyunsaturated fatty acids

PUFA reduces blood LDL cholesterol, but has an undesirable effect of also reducing HDL cholesterol [11]. As presented in Table 1, PUFA was the most dominant type of fatty acids in vegetable-based foods (54.28–56.69% of total fatty acids), rice-based foods (44.28–55.62% of total fatty acids), and seafood dishes (38.54–52.68% of total fatty acids). Among PUFA, the concentration of linoleic acid (C18:2 *n*-6) was the highest in all foods, with *Harda* (22.98 g/100 g EP) and *Samsamia* (14.35 g/100 g EP) containing the highest levels which were attributed to sesame seeds constituting the major ingredient in *Harda* and *Samsamia* recipes. Linoleic acid was the dominant fatty acid in 17 of the 37 foods, which is nutritionally desirable owing to its metabolism at the tissue level, producing hormone-like prostaglandins that aid in reducing blood pressure, preventing gastric ulcers, and relieving nasal congestion and asthma, as well as diminishing platelet aggregation, thus preventing CVD [27]. Linolenic (C18:3 *n*-3), eicosapentaenoic (C20:5 *n*-3), and docosadienoic (C22:2 *n*-3) acids were detected in small quantities (<0.05 g/100 g EP) in all foods. Arachidonic (C20:4 *n*-6) and docosahexaenoic (DHA; C22:6 *n*-3) acids were detected in seafood dishes at concentrations of 0.001–0.09 and 0.02–0.51 g/100 g EP, respectively (S2 Table). The arachidonic acid and DHA percentage of total fatty acids detected in seafood dishes varied between traces–2.34 and traces–7.93%, respectively (Table 1). The availability of DHA in seafood dishes is in agreement with other studies [28, 29] pointed to the high levels of DHA in fish.

Trans fatty acids

Trans fats are commonly found in highly processed foods, although natural trans fats produced in animals and foods made from these animals (e.g. milk, meat products) may also

contain small amounts of these fats. TFA content investigated in the food samples ranged between traces and 0.45 g/100 g EP, all of which were in ready-made purchased foods and foods of animal origin. The highest TFA concentration (0.45 g/100 g EP) was in a meat-based food *Arayes*, followed by cheeseburger (0.36 g/100 g EP), and in desserts *Baklawah* (0.31 g/100 g EP) and *Knafah* (0.27 g/100 g EP) (S2 Table). The TFA percentage of total fatty acids for the same foods was the highest, with values of 4.78, 2.33, 1.50, and 1.53%, respectively (Table 1). The high TFA content in *Arayes* and cheeseburger could be attributed to the major ingredient of both of these foods being lamb and beef, respectively; whereas in *Baklawah* and *Knafah*, it could be attributed to the use of butter in *Baklawah*, and butter and cheese in *Knafah*.

In 2017, Saudi Arabia; a Middle East country with similar lifestyles, dietary patterns, eating habits, nutrition-related health concerns, and nature of foods as the State of Kuwait; implemented 2% and 5% TFA limits for fats and all other foods, respectively [30]. In light of this fact, it is interesting to note that none of the foods analyzed in the present study had TFA >5% of the total fatty acids, whereas the meat-based foods *Arayes* and cheeseburger had TFA proportion of total fatty acids exceeding 2%.

Overall, TFA has no known health benefits. There is a common understanding that high consumption of TFA has negative health effects, and has been strongly associated with a high risk of CVD [24]. TFA has worse effects on blood cholesterol levels than SFA as it not only raises LDL cholesterol but also lowers HDL cholesterol.

Cholesterol composition

The cholesterol content of the investigated foods varied between 4.42 and 454.79 mg/100 g EP of food (Table 2). Cholesterol was not detected in 19 foods, all of which contained no animal fat, the source of dietary cholesterol, detection limit was <0.05 mg/100 g.

283 Table 2. Cholesterol composition given in means $\pm$ SD (mg/100 g edible portion of food)

Food	
Seafood	
<i>Marag Samak Maleh</i> (Spanish mackerel broth)	86.75 $\pm$ 4.67
<i>Kabab Rubyan</i> (Shrimps kebab)	359.17 $\pm$ 8.04
<i>Marag Rubyan</i> (Shrimps broth)	212.58 $\pm$ 6.34
Soup	
<i>Shorbat Shaereya</i> (Vermicelli soup)	ND
<i>Shorbat Dajaj</i> (Chicken soup)	103.47 $\pm$ 5.56
<i>Shorbat Harees</i> (Soup of peeled wheat with lamb)	23.80 $\pm$ 0.64
Cheese	
Cream Cheese Spread, Regular	169.16 $\pm$ 5.67
Triangle Processed Cheese	145.16 $\pm$ 6.67
“Al-Wafra” White Cheese, Low Fat	97.13 $\pm$ 4.65
Rice-based	
<i>Aish Muhammar</i> (Sweet rice)	ND
<i>Aish Khothra</i> (Rice with vegetables)	ND
<i>Aish Fagae ma Laham</i> (Rice, truffle with lamb)	15.18 $\pm$ 2.46
<i>Aish Bajella ma Shbent</i> (Rice with green fava beans and dill)	ND
<i>Aish Tomat</i> (Rice with tomato)	ND
<i>Kushari</i> (Rice with macaroni and mung beans)	ND
Desserts	
<i>Gherayba</i> (Shortbread cookie)	10.10 $\pm$ 1.49
<i>Knafah</i> (Shredded phyllo dough and cheese in sugar syrup)	56.60 $\pm$ 4.63
<i>Baklava</i> (Phyllo pastry with nuts, Baklava)	4.42 $\pm$ 0.37
<i>Qatayef</i> (Dumpling filled with mixed nuts)	15.32 $\pm$ 1.75
<i>Samsamia</i> (Sesame bar)	ND
<i>Khabeesa</i> (Cooked semolina with oil and sugar)	ND
<i>Darabeel</i> (Rolled dough stuffed with sugar and cinnamon)	ND
<i>Zalabia</i> (Jalebi)	ND
<i>Aseeda</i> (Cooked wheat flour with sugar)	ND
<i>Halwa</i> (Sweet wheat starch)	33.21 $\pm$ 1.25
<i>Baksam</i> (Rusk)	ND
Seed-based	
<i>Heso</i> (Garden cress and fenugreek seeds stew)	ND
<i>Harda</i> (Soft sesame butter, Tahini)	ND
Meat-based	
<i>Madhrooba</i> (Crushed wheat with lamb)	19.78 $\pm$ 0.65
<i>Arayes</i> (Pita bread stuffed with minced lamb)	67.85 $\pm$ 2.46
<i>Hameesat Kabdah</i> (Fried lamb liver in tomato sauce)	454.79 $\pm$ 31.01
<i>Keema</i> (Minced lamb fried with peas)	72.42 $\pm$ 2.97
Sandwich	
Cheeseburger	123.42 $\pm$ 1.71
Chickenburger	68.00 $\pm$ 7.77
Vegetable-based	
<i>Hameesat Mushroom</i> (Fried mushrooms in tomato sauce)	ND
<i>Hameesat Fagae</i> (Fried truffles in tomato sauce)	ND
<i>Marag Hawa</i> (Vegetable broth)	ND

284 SD: standard deviation

285 ND: Not detected (detection limit <0.05 mg/100 g edible portion of food)

286

The highest cholesterol content of 454.79 mg/100 g was found in *Hameesat Kabdah*. This food is based on lamb liver that is known to be rich in cholesterol with 386 mg/100 g EP [31]. The high cholesterol concentrations in *Kabab Rubyan* (shrimp kebab; 359.17 mg/100 g) and *Marag Rubyan* (shrimp broth; 212.58 mg/100 g) are attributed to shrimp being the major ingredient in both food recipes. On a fresh weight basis, shrimp contain a high cholesterol concentration of 161 mg/100 g [32]. Cheese dishes (97.13–169.16 mg/100 g), cheeseburger based on beef (123.42 mg/100 g), *Shorbat Dajaj* based on chicken (103.47 mg/100 g), and *Samak Maleh* based on fish (86.75 mg/100 g) contained relatively high cholesterol levels. Among desserts, five dishes contained relatively low levels of cholesterol, *Baklawas* (4.42 mg/100 g), *Gherayba* (10.10 mg/100 g), *Qatayef* (dumpling filled with mixed nuts; 15.32 mg/100 g), *Halwa* (sweet wheat starch; 33.21 mg/100g) with the highest concentration of 56.60 mg/100 g in *Knafah*, that could be attributed to cheese and butter in the recipe. Other foods containing lower cholesterol levels were *Keema*, *Arayes*, *Madhrooba* (crushed wheat with lamb), and *Aish Fagae ma laham* (rice with truffles and lamb), all of which contained varying amounts of lamb. The cholesterol levels in meat-based foods analyzed in this study were higher than those reported for similar Saudi Arabian and Omani meat-based foods [25, 33].

Fatty acid quality

Foods with low PUFA to SFA (P/S) ratios have been considered as undesirable in the diet, as they induce elevated levels of blood cholesterol, which contribute to an increased risk of CVD and cancers. In order to maintain a healthy cardiovascular status, P/S values higher than 0.45 are desirable [13, 34]. As presented in Table 3, the P/S ratios for meat-based foods (0.10–0.42), cheese dishes (0.01–0.04), desserts (*Gherayba* 0.12, *Knafah* 0.08, *Baklawas* 0.25, *Darabeel* 0.21, and *Zalabia* 0.19), cheeseburger (0.40), soup (*Shorbat Harees* (soup of peeled wheat with lamb) 0.05, and *Shorbat Dajaj* (chicken soup) 0.31) were all <0.45, which is considered nutritionally unhealthy. The remaining foods had higher P/S ratios (>0.45) and

312

Table 3. Fatty acid n-3/n-6 and polyunsaturated/saturated fatty acid ratio

Food	n-3/n-6	P/S
Seafood		
<i>Marag Samak Maleh</i> (Spanish mackerel broth)	0.26	1.12
<i>Kabab Rubyan</i> (Shrimps kebab)	0.13	1.27
<i>Marag Rubyan</i> (Shrimps broth)	0.06	2.88
Soup		
<i>Shorbat Shaereya</i> (Vermicelli Soup)	0.00	1.55
<i>Shorbat Dajaj</i> (Chicken soup)	0.02	0.31
<i>Shorbat Harees</i> (Soup of peeled wheat with lamb)	0.00	0.05
Cheese		
Cream Cheese Spread, Regular	0.00	0.02
Triangle Processed Cheese	0.00	0.01
“Al-Wafra” White Cheese, Low Fat	0.00	0.04
Rice-based		
<i>Aish Muhammar</i> (Sweet rice)	0.00	3.32
<i>Aish Khothra</i> (Rice with vegetables)	0.00	3.68
<i>Aish Fagae ma Laham</i> (Rice, truffle with lamb)	0.00	2.07
<i>Aish Bajella ma Shbent</i> (Rice with green fava beans and dill)	0.00	3.21
<i>Aish Tomat</i> (Rice with tomato)	0.00	0.80
<i>Kushari</i> (Rice with macaroni and mung beans)	0.00	0.92
Desserts		
<i>Gherayba</i> (Shortbread cookie)	0.01	0.12
<i>Knafah</i> (Shredded phyllo dough and cheese in sugar syrup)	0.00	0.08
<i>Baklava</i> (Phyllo pastry with nuts, Baklava)	0.00	0.25
<i>Qatayef</i> (Dumpling filled with mixed nuts)	0.00	2.21
<i>Samsamia</i> (Sesame bar)	0.00	1.62
<i>Khabeesa</i> (Cooked semolina with oil and sugar)	0.00	2.82
<i>Darabeel</i> (Rolled dough stuffed with sugar and cinnamon)	0.00	0.21
<i>Zalabia</i> (Jalebi)	0.00	0.19
<i>Aseeda</i> (Cooked wheat flour with sugar)	0.00	1.76
<i>Halwa</i> (Sweet wheat starch)	0.02	0.56
<i>Baksam</i> (Rusk)	0.01	0.58
Seed-based		
<i>Heso</i> (Garden cress and fenugreek seeds stew)	0.03	3.26
<i>Harda</i> (Soft sesame butter, Tahini)	0.00	2.17
Meat-based		
<i>Madhrooba</i> (Crushed wheat with lamb)	0.00	0.32
<i>Arayes</i> (Pita bread stuffed with minced lamb)	0.00	0.10
<i>Hameesat Kabdah</i> (Fried lamb liver in tomato sauce)	0.00	0.16
<i>Keema</i> (Minced lamb fried with peas)	0.01	0.42
Sandwich		
Cheeseburger	0.00	0.40
Chickenburger	0.00	0.65
Vegetable-based		
<i>Hameesat Mushroom</i> (Fried mushrooms in tomato sauce)	0.00	3.77
<i>Hameesat Fagae</i> (Fried truffles in tomato sauce)	0.00	3.38
<i>Marag Hawa</i> (Vegetable broth)	0.00	3.60

313 n-3 fatty acid: 18:3, 20:5, 22:2, and 22:6

314 n-6 fatty acid: 18:2 and 20:4

315 P/S: polyunsaturated/saturated fatty acid ratio

can thus be said to have good fat content for human nutrition concerning the balance between PUFA and SFA.

A balanced n-3/n-6 fatty acid ratio has positive effects on decreasing the risk of CVD and cancers [35]. The n-3/n-6 fatty acid ratios below 4.0 were recommended by the Department of Health of the United Kingdom [36]. The n-3/n-6 ratio of all foods examined in this study (0.00 and 0.26) were within the recommendations (Table 3). Although, recent studies by Marventano et al. and FAO/WHO [23, 37] have suggested that there might not be any impact of n-3/n-6 fatty acid ratio on human health.

An atherogenic and/or thrombogenic food is one that has the potential to induce platelet aggregation, which might stimulate CVD, and is thus considered less nutritionally beneficial. The SFAs, C12:0, C14:0, and C16:0 are considered atherogenic and thrombogenic as many studies have shown that they have a cholesterol-raising effect that may induce platelets aggregation [38]. Thus, foods with higher concentrations of these fatty acids have higher atherogenic (AI) and thrombogenic indices (TI).

AI and TI values were calculated in the present study, these values indicate the potency of foods to stimulate platelet aggregation, and lower values indicate better nutritional quality of food. AI and TI values were the highest for cheese dishes, and ranged from 2.11 to 2.84, and 3.18 to 4.03; respectively (Table 4). The AI and TI values were variable for desserts, AI values ranged from 0.17 in *Khabeesa* and 2.31 in *Knafah*, while TI values ranged from 0.37 in *Khabeesa* and 3.29 in *Knafah*. The high AI and TI values for *Knafah* were attributed to the presence of around 50% C14:0 and C16:0 SFA. Soup was the third highest food group with AI ranging from 0.36–1.12, and TI from 0.66–2.61. AI ranged between 0.61–0.78 and TI between 1.30–1.58 for sandwiches, 0.51–0.66 and 1.29–2.07 for meat-based food, 0.18–0.55 and 0.32–0.58 for seafood, and 0.14–0.70 and 0.33–1.13 for rice-based food. A low AI of 0.15 was recorded for all vegetable-based foods, as well as low TI values varied between 0.33

341 Table 4. Fatty acid quality of 37 local foods of the State of Kuwait using atherogenic, thrombogenic and
342 hypocholesterolemic/hypercholesterolemic indices.

Food	AI	TI	H/H
Seafood			
<i>Marag Samak Maleh</i> (Spanish mackerel broth)	0.55	0.58	2.25
<i>Kabab Rubyan</i> (Shrimps kebab)	0.34	0.52	3.25
<i>Marag Rubyan</i> (Shrimps broth)	0.18	0.32	5.87
Soup			
<i>Shorbat Shaereya</i> (Vermicelli soup)	0.36	0.66	3.39
<i>Shorbat Dajaj</i> (Chicken soup)	1.12	1.77	1.22
<i>Shorbat Harees</i> (Soup of peeled wheat with lamb)	0.78	2.61	1.47
Cheese			
Cream Cheese Spread, Regular	2.73	3.96	0.55
Triangle Processed Cheese	2.84	4.03	0.55
“Al-Wafra” White Cheese, Low Fat	2.11	3.18	0.68
Rice-based			
<i>Aish Muhammar</i> (Sweet rice)	0.15	0.36	6.51
<i>Aish Khothra</i> (Rice with vegetables)	0.14	0.33	6.99
<i>Aish Fagae ma Laham</i> (Rice, truffle with lamb)	0.19	0.50	5.35
<i>Aish Bajella ma Shbent</i> (Rice with green fava beans and dill)	0.17	0.38	6.08
<i>Aish Tomat</i> (Rice with tomato)	0.47	1.13	2.14
<i>Kushari</i> (Rice with macaroni and mung beans)	0.70	0.92	2.57
Desserts			
<i>Gherayba</i> (Shortbread cookie)	1.85	2.88	0.76
<i>Knafah</i> (Shredded phyllo dough and cheese in sugar syrup)	2.31	3.29	0.65
<i>Baklava</i> (Phyllo pastry with nuts, Baklava)	1.18	1.96	1.19
<i>Qatayef</i> (Dumpling filled with mixed nuts)	0.26	0.51	4.78
<i>Samsamia</i> (Sesame bar)	0.22	0.62	4.46
<i>Khabeesa</i> (Cooked semolina with oil and sugar)	0.17	0.37	6.30
<i>Darabeel</i> (Rolled dough stuffed with sugar and cinnamon)	1.02	2.11	1.02
<i>Zalabia</i> (Jalebi)	0.87	1.81	1.15
<i>Aseeda</i> (Cooked wheat flour with sugar)	0.44	0.79	2.69
<i>Halwa</i> (Sweet wheat starch)	0.79	1.21	1.95
<i>Baksam</i> (Rusk)	0.63	1.28	1.62
Seed-based			
<i>Heso</i> (Garden cress and fenugreek seeds stew)	0.14	0.29	6.79
<i>Harda</i> (Soft sesame butter, Tahini)	0.12	0.41	8.04
Meat-based			
<i>Madhrooba</i> (Crushed wheat with lamb)	0.56	1.55	1.99
<i>Arayes</i> (Pita bread stuffed with minced lamb)	0.66	1.96	1.62
<i>Hameesat Kabdah</i> (Fried lamb liver in tomato sauce)	0.51	2.07	1.98
<i>Keema</i> (Minced lamb fried with peas)	0.58	1.29	2.04
Sandwich			
Cheeseburger	0.78	1.58	1.46
Chickenburger	0.61	1.30	1.86
Vegetable-based			
<i>Hameesat Mushroom</i> (Fried mushrooms in tomato sauce)	0.15	0.33	6.85
<i>Hameesat Fagae</i> (Fried truffles in tomato sauce)	0.15	0.35	6.55
<i>Marag Hawa</i> (Vegetable broth)	0.15	0.34	6.81

343 AI: Atherogenic Index, TI: Thrombogenic Index, H/H: Hypocholesterolemic/Hypercholesterolemic Index.

344

and 0.35. AI for seed-based *Harda* and *Heso* (garden cress and fenugreek seeds stew) was the lowest among all foods investigated in this study, with values of 0.12 and 0.14, respectively; and the lowest TI among all foods was 0.29 for *Heso*. This is attributed to low C14:0 and C16:0 SFA and high MUFA and PUFA levels in *Harda* and *Heso*.

Contrary to AI and TI, foods with higher H/H indices are more nutritionally desirable [39]. The highest H/H (8.04) was recorded for *Harda*. The H/H range for the different foods examined in this study from the highest to lowest values were: rice-based foods (2.14–6.99), vegetable-based foods (6.55–6.85), *Heso* (6.79), seafood (2.25–5.87), desserts (0.65–4.78, except *Khabeesa* with H/H 6.29), soup (1.22–3.39), meat-based foods (1.62–2.04), sandwiches (1.46–1.86), and cheese dishes with the lowest H/H values (0.55–0.68). The AI, TI, and H/H values may indicate that the consumption of seed-based, vegetable-based, rice-based, and seafood dishes analyzed in the present study is potentially healthier, in terms of the fatty acid quality, for humans than that of cheese, desserts, sandwiches, and meat-based foods.

Conclusions

The results of this study revealed a high percentage of unsaturated fatty acids, predominantly oleic and linoleic acids, and favorable indices such as P/S, AI, TI, and H/H for seafood dishes, rice-based, seed-based, and vegetable-based foods, suggesting that these foods can be consumed as part of a healthy diet. The cholesterol content was high in two shrimp-based seafood dishes; hence, it is preferable that these foods be consumed with caution by people with high blood cholesterol levels. The TFA content was low in all foods examined, at levels that do not represent any known significant health risk. Interestingly, all foods showed n-3/n-6 fatty acid ratios within the nutritional recommendations. The highest SFA content (predominantly C16:0 and C14:0) was recorded in cheese and some ready-made desserts including *Knafah*, *Gherayba*, *Baklawa*, and *Zalabia*. It is worth mentioning that

these desserts are also popular among the Middle East and Mediterranean populations. The same desserts showed low fatty acid quality based on their P/S, AI, TI, and H/H values; thus, they should be consumed in moderation or other strategies should be adopted, such as modifying the composition of these foods, by for example substituting ghee or butter with vegetable oil. Published data on foods with similar structure could not be compared with our data, since other studies reported fatty acid data on a dry weight basis. The fatty acid and cholesterol data obtained in the present study can be compiled to national food composition tables and incorporated in international databases such the United States Department of Agriculture (USDA) Nutrient Database for Standard Reference and EuroFIR FoodEXplorer to fill in missing data on fatty acids and cholesterol composition of local foods. Furthermore, data of the present study could be used in epidemiological and nutrition-health related studies aim at assessing population fatty acid nutritional status, and hence studying the associations between dietary fatty acids and the etiology of CVD and/or other nutrition-related diseases in humans.

Supplementary material

S1 Table. Detailed ingredients for 37 local Kuwaiti foods.

S2 Table. Fatty acid composition (g/100 g EP) for 37 local Kuwaiti foods.

Acknowledgments

The authors gratefully acknowledge the Ministry of Health of the State of Kuwait (MOH) for their financial support, and the Management of Kuwait Institute for Scientific Research (KISR) for their support and encouragement.

Author contributions

Hanan A. Al-Amiri: Conceptualization, Validation, Data Analysis, Visualization,
Supervision, Project Administration, Funding acquisition, Writing-Original Draft,
Writing-Reviewing & Editing

Nisar Ahmed, Tahani Al-Sharrah: Methodology, Investigation, Resources.

References

1. Michas G, Micha R, Zampelas A. Dietary fats and cardiovascular disease: putting together the pieces of a complicated puzzle. *Atherosclerosis*. 2014;234:320-8. <https://doi.org/10.1016/j.atherosclerosis.2014.03.013>
2. Jakobsen MU, O'Reilly EJ, Heitmann BL, Pereira MA, Bälter K, Fraser GE, et al. Major types of dietary fat and risk of coronary heart disease: a pooled analysis of 11 cohort studies. *Am J Clin Nutr*. 2009; 89:1425-32. <https://doi.org/10.3945/ajcn.2008.27124>
3. World Health Organization [Internet]. Noncommunicable diseases [cited 2018]. Available from: <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases/>
4. He K. Fish, long-chain omega-3 polyunsaturated fatty acids and prevention of cardiovascular disease-eat fish or take fish oil supplement? *Prog Cardiovasc Dis*. 2009; 52(2): 95-114. <https://doi.org/10.1016/j.pcad.2009.06.003>
5. Assmann G, Schulte H. Relation of high-density lipoprotein cholesterol and triglycerides to incidence of atherosclerotic coronary artery disease (the PROCAM experience). *Am J Cardiol*. 1992;70(7):7733-7. [https://doi.org/10.1016/0002-9149\(92\)90550-1](https://doi.org/10.1016/0002-9149(92)90550-1)
6. De Lorgeril M, Salen P. New insights into the health effects of dietary saturated and omega-6 and omega-3 polyunsaturated fatty acids. *BMC Med*. 2012;10, 50. <https://doi.org/10.1186/1741-7015-10-50>
7. Astrup A, Dyerberg J, Elwood P, Hermansen K, Hu FB, Jakobsen MU, et al. The role of reducing intakes of saturated fat in the prevention of cardiovascular disease: where does

- the evidence stand in 2010? *Am J Clin Nutr.* 2011;93:684-8.
<https://doi.org/10.3945/ajcn.110.004622>
8. Murry CE, Jennings RB, Reimer KA. New insights into potential mechanisms of
ischemic preconditioning. *Circulation.* 1991;84:442-5. [https://doi:](https://doi.org/10.1161/01.cir.84.1.442)
[10.1161/01.cir.84.1.442](https://doi.org/10.1161/01.cir.84.1.442)
9. Mozaffarian D, Katan MB, Ascherio A, Stampfer MJ, Willett WC. Trans fatty acids and
cardiovascular disease. *New Eng J Med.* 2006;354(15):1601-13. [http://doi:](http://doi.org/10.1056/NEJMra054035)
[10.1056/NEJMra054035](http://doi.org/10.1056/NEJMra054035)
10. Brouwer IA, Wanders AJ, Katan MB. Trans fatty acids and cardiovascular health:
Research completed? *Eur J Clin Nutr.* 2013;67(5):541-7.
<https://doi.org/10.1038/ejcn.2013.43>
11. Ramsden CE, Hibbeln JR, Majchrzak SF, Davis JM. N-6 fatty acid-specific and mixed
polyunsaturated dietary interventions have different effects on CHD risk: a meta-analysis
of randomized controlled trials. *Br J Nutr.* 2010;104:1586-600.
<https://doi.org/10.1017/S0007114510004010>
12. Grundy SM. Monounsaturated fatty acids and cholesterol metabolism: Implications for
dietary recommendations. *J Nutr.* 1989;119:529-33. <https://doi.org/10.1093/jn/119.4.529>
13. Wood JD, Richardson RI, Nute GR, Fisher AV, Campo MM, Kasapidou E, et al. Effect
of fatty acids on meat quality: a review. *Meat Sci.* 2003;66:21-32.
[https://doi.org/10.1016/S0309-1740\(03\)00022-6](https://doi.org/10.1016/S0309-1740(03)00022-6)
14. Michikawa M. The role of cholesterol in pathogenesis of Alzheimer's disease: dual
metabolic interaction between amyloid beta-protein and cholesterol. *Mol Neurobiol.* 2003;
27:1-12. <https://doi.org/10.1385/MN:27:1:1>
15. Association of Official Analytical Chemists. Official Methods of Analysis, 17th Ed.
Washington (DC): The Association; 2002.

16. Folch J, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem.* 1957;226:497-509.
<https://www.jbc.org/content/226/1/497.long>
17. Segura J, Lopez-Bote C J. A laboratory efficient method for intramuscular fat analysis. *Food Chem.* 2014;145:821-5.
18. American Oil Chemists' Society. Official Method Ce 2–66, Preparation of Methyl Esters of Fatty Acids. Champaign, IL: The Society; 2000.
19. Paul AA, Southgate DAT. McCance and Windowson's, The composition of Foods. London: Her Majesty's Stationary Office; 1978.
20. Naeemi ED, Ahmad N, Al-Sharrah TK, Behbahani M. Rapid and simple methods for determination of cholesterol in processed food. *J AOAC Int.* 1995;78(6):1522-4.
<https://doi.org/10.1093/jaoac/78.6.1522>
21. Ulbricht TLV, Southgate DAT. Coronary Heart Disease: Seven Dietary Factors. *Lancet.* 1991;338:985-92. [https://doi.org/10.1016/0140-6736\(91\)91846-M](https://doi.org/10.1016/0140-6736(91)91846-M)
22. Testi S, Bonaldo A, Gatta P, Badiani A. Nutritional traits of dorsal and ventral fillets from three farmed fish species. *Food Chem.* 2006;98:104-11.
<https://doi.org/10.1016/j.foodchem.2005.05.053>
23. Joint FAO/WHO Expert Consultation on Fats and Fatty Acids in Human Nutrition. Interim Summary of Conclusions and Dietary Recommendations on Total Fat & Fatty Acids. Geneva: WHO; 2008.
https://www.who.int/nutrition/topics/FFA_interim_recommendations/en/
24. Food and Agriculture Organization of the United Nations (FAO). Fats and fatty acids in human nutrition: Report of an expert consultation. Geneva: FAO; 2010;91:1-166.
<http://www.fao.org/3/i1953e/i1953e00.pdf>

25. Sawaya WN, Al-Jebrin A, Salji JP, Ayaz M, Khalil J. Cholesterol contents and fatty acid composition of selected Saudi Arabian dishes. *Nutr Rep Int.* 1985;31(3):593-600.
26. United States Department of Agriculture, Agricultural Research Service [Internet]. USDA Nutrient Database for Standard Reference, Release 18. Available from: <https://fdc.nal.usda.gov/fdc-app.html#/food-details/170152/nutrients>
27. Bhattacharya A, Banu J, Rahman M, Causey J, Fernandes G. Biological effects of conjugated linoleic acids in health and disease. *J Nutr Biochem.* 2006;17:789-810. <https://doi.org/10.1016/j.jnutbio.2006.02.009>
28. Riera-Heredia N, Lutfi E, Gutiérrez J, Navarro I, Capilla E. Fatty acids from fish or vegetable oils promote the adipogenic fate of mesenchymal stem cells derived from gilthead sea bream bone potentially through different pathways. *PLoS ONE.* 2019;14:e0215926. <https://doi.org/10.1371/journal.pone.0215926>
29. Pike IH, Andrew J. Fish oil: production and use now and in the future. *Lipid Technol.* 2010;22:59-61. <https://doi.org/10.1002/lite.201000003>
30. World Health Organization [Internet]. Policies to eliminate industrially produced trans fat [cited 2019]. Available from: <https://apps.who.int/iris/bitstream/handle/10665/331302/WHO-NMH-NHD-18.5-eng.pdf/>
31. U.S. Department of Agriculture, Agricultural Research Service [Internet]. USDA Nutrient Database for Standard Reference, Release 18. Available from: <https://fdc.nal.usda.gov/fdc-app.html#/food-details/172615/nutrients>
32. U.S. Department of Agriculture, Agricultural Research Service [Internet]. USDA Nutrient Database for Standard Reference, Release 18. Available from: <https://fdc.nal.usda.gov/fdc-app.html#/food-details/175179/nutrients/>
33. Musaiger AO, Ahmed MA, Rao MV. Chemical composition of some local dishes of Oman. *Food Chem.* 1989;61:17-22. [https://doi.org/10.1016/S0308-8146\(97\)00148-9](https://doi.org/10.1016/S0308-8146(97)00148-9)

- 491 34. Ospina-E JC, Sierra-C A, Ochoa O, Pérez-Álvarez J, Fernández-López J. Substitution of
492 saturated fat in processed meat products: A review. *Crit Rev Food Sci Nutr*.
493 2012;52(2):113-22. <https://doi.org/10.1080/10408398.2010.493978>
- 494 35. Simopoulos AP. The importance of the ratio of omega-6/omega- 3 essential fatty acids.
495 *Biomed Pharmacother*. 2002;56:365-79. [https://doi.org/10.1016/S0753-3322\(02\)00253-6](https://doi.org/10.1016/S0753-3322(02)00253-6)
- 496 36. Department of Health of the United Kingdom. Nutritional Aspects of Cardiovascular
497 Disease: Report of the Health and Social Subjects. London: Her Majesty's Stationery
498 Office; 1994;46:37-46.
- 499 37. Marventano S, Kolacz P, Castellano S, Galvano F, Buscemi S, Mistretta A, et al. A
500 review of recent evidence in human studies of n-3 and n-6 PUFA intake on
501 cardiovascular disease, cancer, and depressive disorders: does the ratio really matter? *Int*
502 *J Food Sci Nutr*. 2015;66:611-22. <https://doi.org/10.3109/09637486.2015.1077790>
- 503 38. Krauss RM, Deckelbaum RJ, Ernst N, Fisher E, Howard BV, Knopp RH, et al. Dietary
504 guidelines for healthy American adults: A statement for health professionals from the
505 Nutrition Committee, American Heart Association. *Circulation*. 1996;94:1795-1800.
506 <https://doi.org/10.1161/01.CIR.94.7.1795>
- 507 39. Sasyte V, Grashorn MA, Klementaviciute J, Viliene V, Raceviciute-Stupeliene
508 A, Gruzauskas R, et al. Effect of extruded full-fat rapeseed on egg quality in laying hens.
509 *Eur Poult Sci*. 2017;81:1-15.
510 [https://www.european-poultry-science.com/Effect-of-extruded-full-fat-rapeseed-on-egg-q](https://www.european-poultry-science.com/Effect-of-extruded-full-fat-rapeseed-on-egg-quality-in-laying-hens,QUIEPTU1NzU5OTgmTUIEPTE2MTAxNA.html)
511 [uality-in-laying-hens,QUIEPTU1NzU5OTgmTUIEPTE2MTAxNA.html](https://www.european-poultry-science.com/Effect-of-extruded-full-fat-rapeseed-on-egg-quality-in-laying-hens,QUIEPTU1NzU5OTgmTUIEPTE2MTAxNA.html)