Psychosocial Health of School-aged Children during the Initial COVID-19 Safer-at-Home School mandates in Florida: A cross-sectional study

Sarah McKune¹, Daniel Acosta¹, Nick Diaz¹, Kaitlin Brittain¹, Diana Joyce- Beaulieu², Anthony T. Maurelli^{1,3}, Eric J. Nelson^{1,3,4}

¹ Departments of Environmental and Global Health, College of Public Health and Health Professions, University of Florida, Gainesville, FL, USA

² Department of Special Education, School Psychology, & Early Childhood Studies, College of Education, University of Florida, Gainesville, FL, USA

³ Emerging Pathogens Institute, University of Florida, Gainesville, FL, USA

⁴ Department of Pediatrics, College of Medicine, University of Florida, Gainesville, FL, USA

Corresponding Author: Sarah McKune, smckune@ufl.edu

Abstract

Background: Given the emerging literature regarding the impacts of lockdown measures on mental health, this study aims to identify risk factors in school-aged children for being at risk for psychosocial disorders during the COVID-19 Safer-at-Home School mandates in Florida

Methods: A cross-sectional study was conducted in April 2020 (n=280). Bivariate analysis and logistic and multinomial logistic regression models are used to examine socio-demographic and knowledge, attitude, and practice (KAP) predictors of anxiety, depression, and obsessive-compulsive disorder (OCD).

Results: Loss of household income was associated with being at risk for depression [aOR=3.130, 95% CI= (1.41-6.97)], anxiety [aOR=2.531, 95%CI= (1.154-5.551)], and OCD [aOR=2.90, 95%CI= (1.32-6.36)]. Being female was associated with risk for depression [aOR=1.72, 95% CI=(1.02-2.93)], anxiety [aOR=1.75, 95% CI=(1.04-2.97)], and OCD[aOR=1.764, 95%CI= (1.027-3.028)]. Parental practices that are protective against COVID-19 were associated with children being at risk of depression [aOR=1.55, 95% CI= (1.04-2.31)]. Being at a lower school level was risk factor for anxiety and OCD.

Conclusions: Efforts to address mental health risk in children, as a result schools should prioritize girls, younger children, and children of families who lose income. Limiting the spread of COVID-19 through school closure may exacerbate the risk of psychosocial disorders in children, thus school administrators should move quickly to target those at greatest risk.

Keywords:

COVID-19, Mental Health, Psychosocial impacts, Vulnerable Population, Pandemic

Background

Social distancing is the primary public health intervention to reduce the spread of SARS-CoV-2, the virus that causes COVID-19. In the Spring of 2020, school closures, limits on the number of individuals allowed at social gatherings, and closing restaurants, bars, and businesses rapidly reduced transmission and flattened the epidemiologic curve in many parts of the US, including Florida. Though children appear to be at less risk of severe illness and mortality associated with COVID-19¹, these interventions significantly disrupted the lives of children and families – and not equally. Disparities in health care, education, and wealth, which are prevalent across the United States, put minorities and disadvantaged groups at higher risk for negative outcomes associated with the pandemic.

The effects of COVID-19 are far reaching, and scientists urged the research community early in the pandemic to prioritize high quality data on mental health and psychosocial effects of the COVID-19 pandemic^{2,3}. Past evidence has shown significant psychological effects on children during disasters⁴, and in the context of the COVID-19 pandemic, reports of psychosocial distress in children and adolescents have increased compared to the pre-pandemic baseline^{5,6}. COVID-19 related hardships have also affected psychological wellbeing of children, with greater impact on those who have pre-existing mental illness or who live in households facing larger economic distress^{6,7}. Findings from early studies in China found an increase in the prevalence of children reporting symptoms of depression and anxiety during home confinement⁸. A similar study of the general population in China found that some preventive measures were associated with reporting mental health symptoms⁹. COVID-19 psychosocial data are limited, especially data representative of racial and ethnically diverse populations of children in the US. Given that minority communities and communities of color in the US have experienced much higher rates of infection and death¹⁰, the distribution of the psychological toll of COVID-19, including fear, loss, and trauma, is disproportionately affecting children of these communities^{11,12}. Unequal access to healthcare and educational options during the pandemic, along with any

discrimination that comes from being a member of a community with high rates of transmission^{5,13}, will only serve to amplify these psychosocial ramifications.

Families have varying degrees of knowledge, and myriad attitudes and practices surrounding COVID-19. If/how these knowledge, attitudes, and practices (KAP) are associated with risk of poor psychosocial outcomes in children is unknown. KAP studies, often used to develop appropriate public health interventions, are built on the well-established Theory of Planned Behavior and assume a linear relationship, culminating in behaviors that improve health¹⁴⁻¹⁶. Given that the situation with COVID-19 has had high levels of uncertainty, there might be a relationship with KAP and psychosocial distress.

Understanding the effects of COVID-19 on the psychosocial health of school-aged children and identifying those groups of children most at risk is necessary for the development of appropriate, targeted mental health interventions. The aim of this study is to describe the psychosocial health of a population of school-aged children during the early phase of the COVID-19 pandemic. We examine parental KAP related to COVID-19 by race/ethnicity. We then examine these, and other socio-demographic risk factors associated with poor psychosocial health in school-aged children. This information will assist school administrators, public health practitioners, and policy makers in designing evidence-based, targeted interventions to address the psychosocial needs of the school-aged population during the ongoing COVID-19 pandemic. This study examines social groups (sex, age, race, and ethnicity) and COVID-19 related KAP as risk factors for indicators of poor psychosocial outcomes, including anxiety, depression, and obsessive compulsive disorder (OCD), among school-aged (K-12) children in Florida at the beginning of the pandemic.

Methods

Study Design

A cross-sectional study was conducted in April 2020 to describe the effects of COVID-19 on school-aged children and their parents. We present results from an online survey that collected information on socio-demographics, clinical risk factors, parental COVID-19 related KAP, and indicators of psychosocial risk in children. The survey was coupled with the collection of

oropharyngeal swabs (PCR for viral detection) and finger sticks (ELISA for antibodies) from student participants at a drive-thru testing site. Laboratory data associated with this study will be published independently and are not included herein.

Data were collected at a public, K-12 school in Florida. The school is mandated to have a student body that reflects the demographics of the State of Florida. Students are characterized by gender, race/ethnicity, family income, and academic needs. Parents/guardians of all current students (N=1178) were invited by email to have their child participate in the study. Those who were interested were able to consent/assent to participation online via a HIPAA compliant interface connected to RedCap (Vanderbilt University). Parental consent was required for all children under the age of 18; and, all children over the age of eight assented for themselves. All data were deidentified prior to analysis and were stored on secured servers to ensure the protection of participants (RedCap). The study was approved by the University of Florida Institutional Review Board, protocol IRB202001345

Variables and statistical analyses

The demographics section of the survey included questions about child's age, sex, race, ethnicity, and the loss of household income. Other risk factors evaluated in the survey included, clinical symptoms and parental occupation. The COVID-19 related KAP section consisted of fourteen knowledge, eight attitude, and five practice questions. Knowledge scores were created by assigning one (1) to each correctly answered knowledge question about COVID-19 (transmission, prevention, and/or general information) and summing them for a total possible score of 14. Similarly, attitudes and practices were categorized as protective and unprotective, with protective attitudes and practices coded as one (1) and all others as zero (0), and summed, for respective possible scores of 8 and 5.

The questions used to measure psychosocial risk evaluate three internalizing disorders: depression, anxiety, and OCD. These emotional and behavioral disorders were chosen because of their critical importance and generalizability to the pediatric COVID-19 literature^{6,8,17}. Importantly, externalizing disorders are not evaluated in this study. Each psychosocial disorder (depression, anxiety, and OCD) was evaluated using a set of categorical questions (5-point

Likert scale). All questions used age-appropriate language and response options (e.g. 1-Never, 2-A Little, 3-Sometimes, 4-A Lot, or 5-Always/Constantly). Within each group of questions, if a participant's responses were all 1s or 2s, the child was categorized *Not at Risk*; everyone else was considered *At Risk* and further categorized as *High*, *Medium*, or *Low Risk*, using the frequency of their highest response options (see Likert scale above). Participants who answered "a lot" (4) to two or more questions in a group or "always/constantly" (5) to any one of the questions in the group were considered *High Risk*; participants who answered "a lot" (4) only once, with all other questions at a 3 or below, were considered *Medium Risk*; and any participant who answered "sometimes" (3) to one or more question was considered *Low Risk*. Each group of questions was designed by a team of psychologists working with the research team to identify students at risk of developing one of the specific internalized disorders mentioned above. The six primary outcome variables in this study are *At Risk* (Using *No Risk as reference (REF)*) and *High Risk* (REF *No Risk*) of anxiety, depression, and OCD. A summary outcome variable, *Any Risk* indicates if a child presents as *At Risk* for any of the three psychosocial outcomes assessed here.

Bivariate analysis of all outcome variables was conducted to test for association with knowledge score, attitude score, and practice score, demographic variables, and parental occupation. An additional bivariate analysis of each individual KAP question was used to identify association between any KAP item (question) and race/ethnicity using Fisher's Exact Test and logistic regression. Further analysis was conducted using logistic and multinomial logistic regression models to examine predictors of *Any Risk* (REF *No Risk*) and *High Risk* (REF *No Risk*), respectively, for each depression, anxiety, and OCD. The covariates included in the models were race/ethnicity, sex, school level, household loss of income during the pandemic, parent working in a medical setting, knowledge score, attitude score, and practice score. Survey data were collected in RedCap and analyzed in R Software (version 4.0.0) and SPSS (version 26).

Results

A total of 280 students were enrolled out of student body of 1178. Table 1 presents the demographic characteristic of the sample and the adjusted odds ratio (aOR) for being at *Any*

Risk for one or more of three internalizing syndromes considered in this study (depression, anxiety, or OCD). The study population was nearly 2/3 White, a fifth Hispanic (regardless of race), and 9% Black. Participants were 48% male, and distributed across high school (40%), middle school (29%) and primary school (31%).

Parental knowledge about COVID-19 was high, with 31.4% answering all questions correctly, and only 13.6% answering three or more questions incorrectly (Table 2). Attitudes had similar results, with most parents expressing agreement with protective attitudes. Preventative practices were also high, with at least 90% of respondents reporting increased hand washing, avoiding physical contact with those outside their home, and adhering to social distancing guidelines (Table 3).

There was a statistically significant association between race/ethnicity and knowledge questions regarding the existence of an effective cure of the virus, where Multiracial/Other respondents had a higher rate of incorrect responses compared to White respondents. The use of masks as a protective factor against COVID-19 was also significant, with White respondents answering that masks do not protect against the virus at a higher proportion than African Americans and Hispanics (Table 2).

There were race/ethnic differences in who was worried about getting infected by the virus and, correspondingly, with the knowledge question about mask usage; White respondents were less worried overall about getting infected with the virus compared to African Americans and Hispanics (Table 3). There were no significant differences between Multiracial/Other and White respondents in mask usage or being worried about infection with the virus. Among those practices assessed, no significant difference by race/ethnicity was found for social distancing, handwashing, or general hygiene; however, Hispanic respondents were less likely to report purchasing larger amounts of staple foods than normal.

Loss of household income was significantly associated with students being *At Risk* of depression [aOR=3.130, 95% CI= (1.41-6.97)], anxiety [aOR=2.531, 95%CI= (1.154-5.551)], and OCD [aOR=2.90, 95%CI= (1.32-6.36)], see Table 4. Statistical results also indicate a marginally significant association (where $p < 0.1$) between loss of household income and a student being at *High Risk* for depression [p value 0.05, aOR=3.74, 95% CI=(1.00-14.01)] and OCD [p value 0.069, aOR=3.19, 95% CI=(0.91-11.12)], see Table 4. Being female was significantly associated with being *At Risk* for depression [aOR=1.72, 95% CI=(1.02-2.93)], anxiety [aOR=1.75, 95% CI=(1.04-2.97)], and OCD[aOR=1.764, 95%CI= (1.027-3.028)]. Girls may also be more likely to present as *High Risk* for depression [p value=0.057, aOR=2.93, 95% CI= (0.970-8.84), Table 3] and OCD [p value= 0.050 aOR=2.58, 95%CI= (0.99-6.68), table 4], as a marginally significant association was identified.

School level was significantly associated with being *At Risk* and *High Risk* of both anxiety and OCD (Table 4). Those in primary school and middle school are more likely to be *At Risk* than students in high school, and those in primary school being more likely to be at *High Risk*. A family's COVID-19 Practice score was significantly associated with a child being *At Risk* for depression [aOR=1.55, 95% CI= (1.04-2.31), Table 4]; those families who followed stricter protective practices were more likely to have a child who presented as *At Risk* for depression. A family's COVID-19 Attitude score was marginally significant in the multinomial regression model for *High Risk* of anxiety [p value=0.062, aOR 1.82, 95% CI (0.971-3.43)]. Inversely, increased parental COVID-19 Knowledge was protective against children presenting as *High Risk* for OCD. Parents working in a frontline medical setting was protective against children being *At Risk* for depression [p value=0.084, aOR=0.59, 95% CI= (0.33-1.07)] or OCD [p value= 0.091 aOR=0.592, 95%CI= (0.322-1.088)].

Discussion

A number of items present in this analysis are risk factors for psychosocial distress among school-aged children. These include a loss of income in their household, being female, and being in primary school. In addition, families who report more protective practices were more likely to have children who present as *At Risk* for depression, and families who indicate

attitudes consistent with protection from COVID-19 are more likely to have children who present as high risk for anxiety. These results merit further discussion.

Loss of household income was a clear indicator of mental distress across the sample population, which is congruent with other studies, suggesting that the economic impacts of the lockdown are an important trigger for mental distress in children.^{3,7,9} Economic hardship alone might be associated with increased risk of socioemotional problems in children, exacerbated by their parents' response to the situation¹⁸. This may also be a partial explanation for why having a parent working on the medical frontline was marginally significant as a protective factor for depression and OCD, as those may be households that are economically advantaged (doctors, nurses), and did not lose income compared to others.

Sex was also an important risk factor, as girls were more likely to be at risk for depression, anxiety, and OCD. This finding echoes results from post-disaster related studies, where being female was associated with increased psychosocial risk¹⁹. On the other hand, this result might be related to reporting, as girls are more likely than boys to report their risk and/or clinical symptoms of depression²⁰⁻²². The results also suggest that children in primary school are more likely to present as at high risk for anxiety and OCD, making them priority groups for interventions at schools.

Reporting protective practices was associated with children presenting as *At Risk* for depression. Distancing and isolation can increase stress, which can aggravate feelings of loneliness and impact long term health³. Protective attitudes in parents was also associated with children presenting as high risk for anxiety. Higher practice scores were associated with hand washing, avoiding physical contact, and adhering to social distancing guidelines, while higher attitudes scores are related to supporting closures and limiting social gatherings. Even though these attitudes and practices are crucial to slow the spread of viral diseases, to continue implementing them might increase risk of psychosocial distress.

Thus, the pandemic paradox arises: do we limit the spread of disease at the cost of higher risk of psychosocial distress, or do we risk overwhelming the health system and witness a spike in deaths, as we seek to protect children from risk of psychosocial distress? There is no simple

answer, as both options have important negative impacts and implications. Extended quarantine can expose children to domestic violence²³⁻²⁵, as well as aggravating loss of household income to those who are unable to hire private childcare, thus are unable to return to work. On the other hand, reopening schools could expose children to deaths of teachers, staff, and fellow students, which would only exacerbate negative psychosocial impacts. Additionally, there are important disparities that will influence which children become infected: a recent study of pediatric cases showed that 51% of the children infected came from low-income communities, while only 2% came from high-income communities²⁶. Yonker et al. also suggest that children from lower income settings pose a larger threat to their families as household size may be larger with multi-generational co-habitation and higher household density²⁶.

Conclusion

Many studies aim to predict how lockdown measures will flatten the pandemic curve, but few studies focus on the psychosocial impacts of these interventions on children and their families. As public health experts focus on reducing the spread of COVID-19 infections, it is imperative that they also focus on addressing the psychosocial needs of children. Additional research is needed to better understand and address the impacts of the pandemic and its societal responses on children, but doing so must be inclusive of vulnerable populations, including those whose households have lost income. Future research must include economic status and ensure diversity and inclusion of minorities in order to understand how the impacts affect vulnerable groups differently. Given that loss of household income was clearly an important risk factor for depression, anxiety, and OCD, understanding a household's baseline economic status becomes important in considering vulnerability and possible mitigating conditions. These findings have important implications for policy makers as they negotiate the continuation of funding for unemployment benefits and other safety nets for those economically disadvantaged, as losing these could increase the number of children at risk of psychosocial distress. There should be a national, coordinated effort to continue collecting psychosocial data on children, as well as to design strategies and coping mechanisms for these children as societies across the country strive to balance the risks of COVID-19 infection and the risk of lockdown on psychosocial

wellbeing. Previous disasters have shown that adverse psychological effects are not only present during the event, but also remain long after the incident¹⁹. Efforts to address mental health cannot wait until the pandemic is over, neither for school age population as the present study suggests, nor for adults, as other studies indicate^{7,9,11,27-29}. At the very minimum, high risk groups must be identified across the US. These data contribute to evidence that lay bare the urgent need for primary and secondary school administrators, in collaboration with public health practitioners and medical professionals, to roll out targeted and group interventions as early as possible to respond to the emotional and psychosocial needs of children after a disaster occurs.

Limitations

Data were not collected on household income; only loss of income due to COVID-19 was captured, thus limiting interpretation of some of these analyses. The survey instrument was designed in March 2020, when the use of masks was not widely recommended³⁰ and there were many uncertainties about the virus and its spread. As a result, only one question about mask usage was included in the knowledge section of the questionnaire. The survey was completed in mid-April when schools had already moved to remote learning and prevention campaigns were already in effect, which could explain the high rate at which knowledge questions were answered correctly.

List of abbreviations

aOR: Adjusted odds ratio

CI: Confidence Interval

HIPPA: Health Insurance Portability and Accountability Act

KAP: Knowledge, attitudes, and practices

OCD: Obsessive Compulsive Disorder

Declarations

303 **Ethics approval and consent to participate**

304 The study was approved by the University of Florida Institutional Review Board, protocol
305 IRB202001345

306 **Consent for publication**

307 Not applicable

308 **Availability of data and materials**

309 All data will be de-identified and made publicly available 12 months after the end of the cohort
310 study.

311 **Competing interests**

312 The authors declare that they have no competing interests

313 **Funding**

314 This research was funded and supported by the University of Florida College of Public Health
315 and Health Professions, the College of Medicine, the Clinical and Translational Science Institute,
316 and the Emerging Pathogens Institute at the University of Florida.

317 **Authors' Contributions**

318 SM designed and supervised the study, supervised the analyses, and led the writing. DA
319 completed the analyses and significantly contributed to the writing. ND assisted with the
320 analyses and assisted with the writing. KB assisted with the analyses and assisted with the
321 writing. DJB assisted with the design, with the analyses, and assisted with the writing. ATM
322 designed and supervised the study, assisted with the writing. EJM designed and supervised the
323 study, contributed to the writing

324 **Acknowledgements**

325 A special thanks to the school personnel and staff; the medical, public health, and nursing
326 student volunteers; Drs. Tara Sabo-Atwood, John Lednický, and the entire research testing lab

staff in Environmental and Global Health; and Dr. Glenn Morris, and Dr. Michael Lauzardo for their support and guidance.

References

1. Dong Y, Mo X, Hu Y, et al. Epidemiology of COVID-19 Among Children in China. *Pediatrics*. 2020;145(6). doi:10.1542/peds.2020-0702
2. Holmes* EA, O'Connor* RC, Perry VH, et al. Multidisciplinary research priorities for the COVID-19 pandemic: *Lancet Psychiatry*. 2020;(January):19-21.
3. Bavel JJV, Baicker K, Boggio PS, et al. Using social and behavioural science to support COVID-19 pandemic response. *Nat Hum Behav*. 2020;4(May). doi:10.1038/s41562-020-0884-z
4. Rolfsnes ES, Idsoe T. School-Based Intervention Programs for PTSD Symptoms: A Review and Meta-Analysis. 2007;20(3):251-262. doi:10.1002/jts.20622
5. Duan L, Shao X, Wang Y, et al. An investigation of mental health status of children and adolescents in china during the outbreak of COVID-19. *J Affect Disord*. 2020;275(April):112-118. doi:10.1016/j.jad.2020.06.029
6. Taquet M, Quoidbach J, Fried EI, Goodwin GM. Mood Homeostasis before and during the Coronavirus Disease 2019 (COVID-19) Lockdown among Students in the Netherlands. *JAMA Psychiatry*. 2020;2019:2019-2021. doi:10.1001/jamapsychiatry.2020.2389
7. Gassman-Pines A, Ananat EO, Fitz-Henley J. COVID-19 Crisis Impacts on Parent and Child Psychological Well-being. *Pediatrics*. 2020;146(4):e2020007294. doi:10.1542/peds.2020-007294
8. Xinyan X, Qi X, Yu Z, Qi L, Jiajia Z, Ranran S. Mental Health Status Among Children in Home Confinement During the Coronavirus Disease 2019 Outbreak in Hubei Province, China. *Jama Pediatr*. 2020;7:2-4.
9. Shi L, Lu ZA, Que JY, et al. Prevalence of and Risk Factors Associated With Mental Health Symptoms Among the General Population in China During the Coronavirus Disease 2019

- 353 Pandemic. *JAMA Netw open*. 2020;3(7):e2014053.
354 doi:10.1001/jamanetworkopen.2020.14053
- 355 10. Laurencin CT, McClinton A. The COVID-19 Pandemic: a Call to Action to Identify and
356 Address Racial and Ethnic Disparities. *J Racial Ethn Heal Disparities*. 2020;7(3):398-402.
357 doi:10.1007/s40615-020-00756-0
- 358 11. Shim RS, Compton MT. The Social Determinants of Mental Health: Psychiatrists' Roles in
359 Addressing Discrimination and Food Insecurity. *Focus (Madison)*. 2020;18(1):25-30.
360 doi:10.1176/appi.focus.20190035
- 361 12. Ambrose AJH. Inequities during COVID-19. *Pediatrics*. 2020;146(2).
362 doi:10.1542/peds.2020-1501
- 363 13. Zheng G, Jimba M, Wakai S. Exploratory study on psychosocial impact of the severe acute
364 respiratory syndrome (SARS) outbreak on Chinese students living in Japan. *Asia-Pacific J*
365 *Public Heal*. 2005;17(2):124-129. doi:10.1177/101053950501700211
- 366 14. Werner PD. Implications of Attitude-Behavior Studies for Population Research and Action
367 Author (s): Paul D . Werner Published by: Population Council Stable URL:
368 <https://www.jstor.org/stable/1966280> Implications of Attitude-Behavior Studies for
369 Population Researc. 2020;8(11):294-299.
- 370 15. Ajzen I. Perceived behavioral control, self-efficacy, locus of control, and the theory of
371 planned behavior. *J Appl Soc Psychol*. 2002;32(4):665-683. doi:10.1111/j.1559-
372 1816.2002.tb00236.x
- 373 16. Warwick D. *The KAP Survey: Dictates of Mission Versus Demands of Science*. London:
374 Harvard Institute for International Development; 1983.
- 375 17. Seğer İ, Ulaş S. An Investigation of the Effect of COVID-19 on OCD in Youth in the Context
376 of Emotional Reactivity, Experiential Avoidance, Depression and Anxiety. *Int J Ment*
377 *Health Addict*. 2020. doi:10.1007/s11469-020-00322-z
- 378 18. McLoyd VC. Socialization and development in a changing economy. *Am Psychol*.

- 1989;44(2):293-302.
<http://eds.a.ebscohost.com.libproxy.txstate.edu/eds/pdfviewer/pdfviewer?vid=80&sid=d35e1ab5-fcdd-4a23-8336-2c0a867a8837@sessionmgr4005&hid=4202>.
19. Goldmann E, Galea S. Mental health consequences of disasters. *Annu Rev Public Health*. 2014;35:169-183. doi:10.1146/annurev-publhealth-032013-182435
20. Kaczynski KJ, Claar RL, Logan DE. Testing gender as a moderator of associations between psychosocial variables and functional disability in children and adolescents with chronic pain. *J Pediatr Psychol*. 2009;34(7):738-748. doi:10.1093/jpepsy/jsn113
21. Nolen-Hoeksema S, Girgus JS. The Emergence of Gender Differences in Depression During Adolescence. *Psychol Bull*. 1994;115(3):424-443. doi:10.1037//0033-2909.115.3.424
22. Lewinsohn PM, Lewinsohn M, Gotlib IH, Seeley JR, Allen NB. Gender differences in anxiety disorders and anxiety symptoms in adolescents. *J Abnorm Psychol*. 1998;107(1):109-117. doi:10.1037/0021-843X.107.1.109
23. Bradbury-Jones C, Isham L. The pandemic paradox: The consequences of COVID-19 on domestic violence. *J Clin Nurs*. 2020;29(13-14):2047-2049. doi:10.1111/jocn.15296
24. Campbell AM. An increasing risk of family violence during the Covid-19 pandemic: Strengthening community collaborations to save lives. *Forensic Sci Int Reports*. 2020;2(April):100089. doi:10.1016/j.fsir.2020.100089
25. Humphreys KL, Myint MT, Zeanah CH. Increased risk for family violence during the COVID-19 pandemic. *Pediatrics*. 2020;146(1). doi:10.1542/peds.2020-0982
26. Yonker LM, Neilan AM, Bartsch Y, et al. Pediatric SARS-CoV-2: Clinical Presentation, Infectivity, and Immune Responses. *J Pediatr*. 2020. doi:10.1016/j.jpeds.2020.08.037
27. Ettman CK, Abdalla SM, Cohen GH, Sampson L, Vivier PM, Galea S. Prevalence of Depression Symptoms in U.S. Adults Before and During the COVID-19 Pandemic. *JAMA Netw Open*. 2020;In Press(9). doi:10.1001/jamanetworkopen.2020.19686

28. Islam MA, Barna SD, Raihan H, Khan MNA, Hossain MT. Depression and anxiety among university students during the COVID-19 pandemic in Bangladesh: A web-based cross-sectional survey. *PLoS One*. 2020;15(8):e0238162. doi:10.1371/journal.pone.0238162
29. Math S, Nirmala M, Moirangthem S, Kumar N. Disaster management: Mental health perspective. *Indian J Psychol Med*. 2015;37(3):261-271. doi:10.4103/0253-7176.162915
30. CDC. CDC's recommendations for implementation of mitigation strategies for Florida , based on current situation with COVID-19 transmission and consideration of the state's large older adult population. 2020:1-5. https://www.cdc.gov/coronavirus/2019-ncov/downloads/FL_Community_Mitigation.pdf.

Tables

Table 1: Demographic characteristics of the study population and Adjusted Odds Ratio (aOR) for being at Any Risk of anxiety, depression, or OCD (n=280)

		Study sample % (school %)	At Any Risk of anxiety, depression, or OCD	95% CI
School Level	High School	40% (42%)	Ref	NA
	Middle School	29% (28%)	1.72	0.928-3.19
	Primary School	31% (30%)	2.1	1.14-3.88
Sex	Male	48% (50%)	Ref	NA
	Female	52% (50%)	1.745	1.047-3.88
Race/Ethnicity	White	62% (48%)	Ref	NA
	Latinx/Hispanic	19% (24%)	0.532	0.266-1.066
	Black	9% (26%)	0.591	0.234-1.492
	Multiracial/Other	10% (7%)	0.603	0.251-1.449

419 **Table 2: Knowledge Results. P-values are for Fisher's Exact Test comparing knowledge answers and**
420 **race/ethnicity.**

Knowledge Results			
Questions		n=280	
	Correct	Incorrect	p-value
	No (%)	No (%)	
The main clinical symptoms of COVID-19 are fever, fatigue, dry cough, and muscle aches.	265(94.6%)	15(5.4%)	0.589
Unlike the common cold, stuffy nose, runny nose, and sneezing are less common in persons infected with the COVID-19 virus.	220(78.6%)	60(21.4%)	0.747
There currently is no effective cure for COVID-19, but early symptomatic and supportive treatment can help most patients recover from the infection.	265(94.6%)	15(5.4%)	0.016*
Antibiotics can be used to treat COVID-19 [^]	210 (75.0%)	70(25%)	0.732
Not all persons with COVID-19 will develop to severe cases. Those who are elderly and have chronic illnesses are more likely to be severe cases.	277(98.9%)	3(1.1%)	1.000
People of all racial and ethnic groups can become infected with the COVID-19 virus.	276(98.6%)	4(1.4%)	0.818
Most people who are infected with the COVID-19 virus recover from it	258(92.1%)	22(7.9%)	0.609
Handwashing can help reduce transmission of the COVID-19 virus.	279(99.6%)	1(0.4%)	1.000
Persons with COVID-19 cannot pass the virus to others if they do not have symptoms.	261(93.2%)	19(6.8%)	0.823
The COVID-19 virus spreads via respiratory droplets of infected individuals	269(96.1%)	11(3.9%)	0.657
Ordinary residents can wear general medical masks to prevent infection by the COVID-19 virus.	173(61.8%)	107(38.2%)	0.015*
It is not necessary for children and young adults to take measures to prevent infection by the COVID-19 virus.	272(97.1%)	8(2.9%)	0.533
Isolation and treatment of people who are infected with the COVID-19 virus are effective ways to reduce the spread of the virus.	278(99.3%)	2(0.7%)	1.000
People who have contact with someone infected with the COVID-19 virus should be immediately isolated in a proper place. In general, the observation period is 14 days.	278(99.3%)	2(0.7%)	0.379
Cumulative Knowledge Score		n=280	
Scores out of 14		No.	%
<11		8	2.9%
11		30	10.7%

12	55	19.6%
13	99	35.4%

*p-value<0.05

^This item is false. The survey was developed and implemented prior to the use of hydroxychloroquine, azithromycin, or remdesivir in experimental trials.

Table 3: Attitudes and Practices Results. P-values are for Fisher's Exact Test comparing protective and un-protective attitudes and practices with race/ethnicity.

Attitudes	n=280			p-value
	Agree No (%)	Disagree No (%)	Don't know No (%)	
I am worried about getting infected with the COVID-19 virus.	176(63%)	82(29.3%)	22(7.9%)	0.030*
I feel confident I can prevent myself and my family from becoming infected with the COVID-19 virus if it becomes more widespread in Florida.	151(53.9%)	64(22.9%)	65(23.2%)	0.275
I know what actions to take to prevent myself and my family from becoming infected with the COVID-19 virus.	256(92.5%)	9(3.2%)	12(4.3%)	1.000
I support a government-imposed mandatory quarantine for those who are infected with the COVID-19 virus.	266(95.0%)	4(1.4%)	10(3.6%)	0.697
I support voluntary home quarantine for up to 2 weeks for people who have been in contact with someone who has COVID-19.	275(98.2%)	1(0.4%)	4(1.8%)	0.064
I support postponing or canceling mass gatherings such as concerts, festivals, and sporting events.	268(95.7%)	4(1.4%)	8(2.9%)	0.325
I support closure of K-12 schools if any student, staff member, or teacher is found to have COVID-19	251(89.6%)	11(3.9%)	18(6.4%)	0.184
If I were exposed to and could possibly be infected with the COVID-19 virus, I would be willing to quarantine myself at home for 2 weeks until I was sure I was not infected in order to prevent others from getting COVID-19 from me.	278(99.3%)	0(0%)	2(0.7%)	0.103
Practice	n=280			p-value
	Yes No (%)	No No (%)		
In recent days, are you washing your hands with soap and water more often than normal?	262(93.6%)	18(6.4%)		1.000
In recent days, are you using more disinfectants, such as hand sanitizers and cloth wipes?	252(90.0%)	28(10.0%)		0.878

In recent days, are you avoiding shaking hands or other physical contact with others outside your home?	277(98.9%)	3(1.1%)	1.000
In recent days, have you adhered to other social distancing guidelines, such as avoiding meetings of more than 10 people and keeping a distance of 6 feet apart?	279(99.6%)	1(0.4%)	1.000
In recent days, have you bought larger amounts of staple foods (flour, sugar, pasta, rice, canned food) than normal?	148(52.9%)	132(47.1%)	0.042 *

*p-value<0.05

Table 4 Adjusted Odds Ratios Children At Risk and at High Risk for depression, anxiety, and OCD, by race/ethnicity, sex, school level, household loss of income, parental employment, and COVID-19 KAP.

	Anxiety		Depression		OCD	
	aOR At Risk (95% C.I.) ¹	aOR High Risk (95% C.I.) ²	aOR At Risk (95% C.I.) ¹	aOR High Risk (95% C.I.) ²	aOR At Risk (95% C.I.) ¹	aOR High Risk (95% C.I.) ²
Covariates						
Race/Ethnicity						
White	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)
Black	0.758(0.29-1.97)	0.34(0.04-2.96)	0.97(0.39-2.48)	0.74(0.082-6.78)	1.01(0.39-2.66)	2.17(0.57(8.37)
Hispanic	0.589(0.28-1.22)	0.82(0.26-2.64)	0.68(0.33-1.42)	1.83(0.56-5.99)	1.02(0.49-2.08)	0.79(0.22-2.82)
Multiracial	1.034(0.43-2.49)	1.47(0.34-6.36)	0.78 (0.31-1.96)	2.53(0.49-11.41)	1.41(0.58-3.43)	1.55(0.35-6.87)
Sex						
Male	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)
Female	1.753(1.04-2.97)	1.93(0.76-4.88)	1.72 (1.02-2.93)	2.93(0.970-8.84)	1.76(1.03-3.03)	2.58(0.99-6.68)
School Level						
High School	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)
Middle School	1.82(0.95-3.56)	1.26(0.37-4.33)	1.67(0.88-3.18)	0.83(0.23-2.99)	1.16(0.59-2.26)	1.46(0.44-4.87)
Primary School	2.32(1.23-4.36)	2.94(1.04-8.30)	1.63 (0.88-3.07)	1.03(0.33-3.26)	2.01(1.07-3.77)	3.10(1.05-9.12)
Household loss of income due to COVID-19						
No	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)	Ref (NA)
Yes	2.53(1.15-5.55)	1.17(0.23-5.8)	3.13(1.41-6.97)	3.74(1.00-14.01)	2.90(1.32-6.36)	3.19(0.91-11.12)
Parent working on frontline						
No	Ref (NA)	NA	Ref (NA)	Ref (NA)	Ref	NA
Yes	0.66(0.37-1.18)	0.41(0.13-1.25)	0.59(0.33-1.07)	0.43(0.13-1.45)	0.59(0.32-1.09)	0.42(0.14-1.31)
KAP Scores						
Knowledge Score	1.05(0.83-1.32)	0.96(0.64-1.43)	0.96(0.77-1.21)	1.11(0.69-1.76)	0.89(0.77-1.25)	0.72(0.50-1.04)
Attitude Score	0.97(0.75-1.25)	1.82(0.97-3.43)	0.90(0.71-1.16)	1.51(0.79-2.89)	0.25(0.89-1.56)	1.46(0.87-2.44)
Practice Score	1.08(0.75-1.58)	1.19(0.57-2.52)	1.55(1.04-2.31)	1.34(0.63-2.86)	0.42(0.79-1.74)	1.13(0.57-2.22)

¹ Binary Logistic Regression was used to model At Risk, using No Risk as the reference category

² Multinomial Logistic Regression was used to model High Risk using No Risk as the reference category