

## Supplementary File

### Frailty and comorbidity in predicting community COVID-19 mortality in the UK Biobank: the effect of sampling

Jonathan K. L. Mak<sup>1\*</sup>, Ralf Kuja-Halkola<sup>1</sup>, Yunzhang Wang<sup>1</sup>, Sara Hägg<sup>1</sup>, Juulia Jylhävä<sup>1</sup>

<sup>1</sup> Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden


\*Corresponding author: Jonathan K. L. Mak

Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Nobels väg 12A, 171 65 Stockholm, Sweden

Email: [jonathan.mak@ki.se](mailto:jonathan.mak@ki.se)

#### Table of Contents

Supplementary Fig. 1. Flowchart of selection of study samples from the UK Biobank.....	2
Supplementary Table 1. List of the five frailty criteria for construction of the frailty phenotype.....	3
Supplementary Table 2. List of the 49 frailty items and scoring for construction of the frailty index.....	4
Supplementary Table 3. List of the 109 ICD-10 codes and assigned weights for construction of the Hospital Frailty Risk Score.....	5
Supplementary Table 4. List of the 17 comorbidities, assigned weights and associated ICD-10 codes for construction of Charlson Comorbidity Index.....	7
Supplementary Table 5. Characteristics of individuals with and without hospital data .....	8
Supplementary Table 6. Characteristics of the COVID-19 positive subsample (n=2,287) by the source of the disease status .....	9
Supplementary Table 7. Spearman's rank correlation coefficients among the frailty measures and comorbidity .....	10
Supplementary Table 8. Associations between different frailty and comorbidity measures, and being COVID-19 positive (n=428,754) .....	11
Supplementary Table 9. Associations between the Hospital Frailty Risk Score and COVID-19 mortality stratified by age in the full sample and COVID-19 positive subsample .....	12
Supplementary Table 10. Associations between different frailty and comorbidity measures, and mortality due to COVID-19 and other causes in the full sample and COVID-19 positive subsample .....	13



**Supplementary Fig. 1** Flowchart of selection of study samples from the UK Biobank

**Supplementary Table 1** List of the five frailty criteria for construction of the frailty phenotype

Frailty criteria	Measurement	Scoring
Weight loss	Self-reported question: “Compared with one year ago, has your weight changed?”	Yes, lost weight=1; other=0
Exhaustion	Self-reported question: “Over the past two weeks, how often have you felt tired or had little energy?”	More than half the days or nearly every day=1; other=0
Slowness	Self-reported question: “How would you describe your usual walking pace?”	Slow=1; other=0
Low physical activity	Self-reported physical activity questionnaire, categorized into 4 levels: (i) none – no physical activity in the last 4 weeks (ii) low – light DIY activity (e.g., pruning, watering the lawn) only in the last 4 weeks (iii) medium – heavy DIY activity (e.g., weeding, lawn mowing, carpentry and digging), walking for pleasure, or other exercises in the last 4 weeks (iv) high – strenuous sports in the last 4 weeks	None or light activity with a frequency of once per week or less=1; medium or heavy activity, or light activity more than once per week=0
Weakness	Measured grip strength	Sex and body mass index (BMI) adjusted cutoffs: Men: ≤29 kg for BMI ≤24; ≤30 kg for BMI 24.1–26; ≤30 kg for BMI 26.1–28; or ≤32 kg for BMI >28 Women: ≤17 kg for BMI ≤23; ≤17.3 kg for BMI 23.1–26; ≤18 kg for BMI 26.1–29; or ≤21 kg for BMI >29

**Supplementary Table 2** List of the 49 frailty items and scoring for construction of the frailty index

#	Frailty item	Scoring
1	Glaucoma	No=0; Yes=1
2	Cataracts	No=0; Yes=1
3	Hearing difficulty	No=0; Yes/completely deaf=1
4	Migraine	No=0; Yes=1
5	Dental problems (ulcers, painful gums, bleeding gums, loose teeth, toothache, dentures)	None=0; Any=1
6	Self-rated health	Excellent=0; Good=0.25; Fair=0.5; Poor=1
7	Fatigue: frequency of tiredness/lethargy in last two weeks	Not at all=0; Several days=0.25; More than half=0.5; Nearly every day=1
8	Sleep: experience of sleeplessness/insomnia	Never/rarely=0; Sometimes=0.5; Usually=1
9	Depressed feelings: frequency in last two weeks	Not at all=0; Several days=0.5; More than half=0.75; Nearly every day=1
10	Self-described nervous personality	No=0; Yes=1
11	Severe anxiety/panic attacks	No=0; Yes=1
12	Common to feel loneliness	No=0; Yes=1
13	Sense of misery (ever/never)	No=0; Yes=1
14	Infirmity: long-standing illness or disability	No=0; Yes=1
15	Falls in last year	No falls=0; one fall=0.5; more than one=1
16	Fractures/broken bones in last five years	No=0; Yes=1
17	Diabetes	No=0; Yes=1
18	Myocardial infarction	No=0; Yes=1
19	Angina	No=0; Yes=1
20	Stroke	No=0; Yes=1
21	High blood pressure	No=0; Yes=1
22	Hypothyroidism	No=0; Yes=1
23	Deep-vein thrombosis	No=0; Yes=1
24	High cholesterol	No=0; Yes=1
25	Breathing: wheeze in last year	No=0; Yes=1
26	Pneumonia	No=0; Yes=1
27	Chronic bronchitis/emphysema	No=0; Yes=1
28	Asthma	No=0; Yes=1
29	Rheumatoid arthritis	No=0; Yes=1
30	Osteoarthritis	No=0; Yes=1
31	Gout	No=0; Yes=1
32	Osteoporosis	No=0; Yes=1
33	Hay fever, allergic rhinitis or eczema	No=0; Yes=1
34	Psoriasis	No=0; Yes=1
35	Any cancer diagnosis	No=0; Yes=1
36	Multiple cancers diagnosed	No/single cancer=0; Multiple cancers=1
37	Chest pain	No=0; Yes=1
38	Head and/or neck pain	No=0; Yes=1
39	Back pain	No=0; Yes=1
40	Stomach/abdominal pain	No=0; Yes=1
41	Hip pain	No=0; Yes=1
42	Knee pain	No=0; Yes=1
43	Whole-body pain	No=0; Yes=1
44	Facial pain	No=0; Yes=1
45	Sciatica	No=0; Yes=1
46	Gastric reflux	No=0; Yes=1
47	Hiatus hernia	No=0; Yes=1
48	Gall stones	No=0; Yes=1
49	Diverticulitis	No=0; Yes=1

**Supplementary Table 3** List of the 109 ICD-10 codes and assigned weights for construction of the Hospital Frailty Risk Score

#	ICD-10 code	Description	Weight
1	F00	Dementia in Alzheimer's disease	7.1
2	G81	Hemiplegia	4.4
3	G30	Alzheimer's disease	4.0
4	I69	Sequelae of cerebrovascular disease (secondary codes)	3.7
5	R29	Other symptoms and signs involving the nervous and musculoskeletal systems (R29.6 Tendency to fall)	3.6
6	N39	Other disorders of urinary system (includes urinary tract infection and urinary incontinence)	3.2
7	F05	Delirium, not induced by alcohol and other psychoactive substances	3.2
8	W19	Unspecified fall	3.2
9	S00	Superficial injury of head	3.2
10	R31	Unspecified hematuria	3.0
11	B96	Other bacterial agents as the cause of diseases classified to other chapters (secondary code)	2.9
12	R41	Other symptoms and signs involving cognitive functions and awareness	2.7
13	R26	Abnormalities of gait and mobility	2.6
14	I67	Other cerebrovascular diseases	2.6
15	R56	Convulsions, not elsewhere classified	2.6
16	R40	Somnolence, stupor and coma	2.5
17	T83	Complications of genitourinary prosthetic devices, implants and grafts	2.4
18	S06	Intracranial injury	2.4
19	S42	Fracture of shoulder and upper arm	2.3
20	E87	Other disorders of fluid, electrolyte and acid- base balance	2.3
21	M25	Other joint disorders, not elsewhere classified	2.3
22	E86	Volume depletion	2.3
23	R54	Senility	2.2
24	Z50	Care involving use of rehabilitation procedures	2.1
25	F03	Unspecified dementia	2.1
26	W18	Other fall on same level	2.1
27	Z75	Problems related to medical facilities and other health care	2.0
28	F01	Vascular dementia	2.0
29	S80	Superficial injury of lower leg	2.0
30	L03	Cellulitis	2.0
31	H54	Blindness and low vision	1.9
32	E53	Deficiency of other B group vitamins	1.9
33	Z60	Problems related to social environment	1.8
34	G20	Parkinson's disease	1.8
35	R55	Syncope and collapse	1.8
36	S22	Fracture of rib(s), sternum and thoracic spine	1.8
37	K59	Other functional intestinal disorders	1.8
38	N17	Acute renal failure	1.8
39	L89	Decubitus ulcer	1.7
40	Z22	Carrier of infectious disease	1.7
41	B95	Streptococcus and staphylococcus as the cause of diseases classified to other chapters	1.7
42	L97	Ulcer of lower limb, not elsewhere classified	1.6
43	R44	Other symptoms and signs involving general sensations and perceptions	1.6
44	K26	Duodenal ulcer	1.6
45	I95	Hypotension	1.6
46	N19	Unspecified renal failure	1.6
47	A41	Other septicemia	1.6
48	Z87	Personal history of other diseases and conditions	1.5
49	J96	Respiratory failure, not elsewhere classified	1.5
50	X59	Exposure to unspecified factor	1.5
51	M19	Other arthrosis	1.5
52	G40	Epilepsy	1.5
53	M81	Osteoporosis without pathological fracture	1.4

**Supplementary Table 3 (continued)**

#	ICD-10 code	Description	Weight
54	S72	Fracture of femur	1.4
55	S32	Fracture of lumbar spine and pelvis	1.4
56	E16	Other disorders of pancreatic internal secretion	1.4
57	R94	Abnormal results of function studies	1.4
58	N18	Chronic renal failure	1.4
59	R33	Retention of urine	1.3
60	R69	Unknown and unspecified causes of morbidity	1.3
61	N28	Other disorders of kidney and ureter, not elsewhere classified	1.3
62	R32	Unspecified urinary incontinence	1.2
63	G31	Other degenerative diseases of nervous system, not elsewhere classified	1.2
64	Y95	Nosocomial condition	1.2
65	S09	Other and unspecified injuries of head	1.2
66	R45	Symptoms and signs involving emotional state	1.2
67	G45	Transient cerebral ischemic attacks and related syndromes	1.2
68	Z74	Problems related to care-provider dependency	1.1
69	M79	Other soft tissue disorders, not elsewhere classified	1.1
70	W06	Fall involving bed	1.1
71	S01	Open wound of head	1.1
72	A04	Other bacterial intestinal infections	1.1
73	A09	Diarrhea and gastroenteritis of presumed infectious origin	1.1
74	J18	Pneumonia, organism unspecified	1.1
75	J69	Pneumonitis due to solids and liquids	1.0
76	R47	Speech disturbances, not elsewhere classified	1.0
77	E55	Vitamin D deficiency	1.0
78	Z93	Artificial opening status	1.0
79	R02	Gangrene, not elsewhere classified	1.0
80	R63	Symptoms and signs concerning food and fluid intake	0.9
81	H91	Other hearing loss	0.9
82	W10	Fall on and from stairs and steps	0.9
83	W01	Fall on same level from slipping, tripping and stumbling	0.9
84	E05	Thyrotoxicosis [hyperthyroidism]	0.9
85	M41	Scoliosis	0.9
86	R13	Dysphagia	0.8
87	Z99	Dependence on enabling machines and devices	0.8
88	U80	Agent resistant to penicillin and related antibiotics	0.8
89	M80	Osteoporosis with pathological fracture	0.8
90	K92	Other diseases of digestive system	0.8
91	I63	Cerebral Infarction	0.8
92	N20	Calculus of kidney and ureter	0.7
93	F10	Mental and behavioral disorders due to use of alcohol	0.7
94	Y84	Other medical procedures as the cause of abnormal reaction of the patient	0.7
95	R00	Abnormalities of heartbeat	0.7
96	J22	Unspecified acute lower respiratory infection	0.7
97	Z73	Problems related to life-management difficulty	0.6
98	R79	Other abnormal findings of blood chemistry	0.6
99	Z91	Personal history of risk-factors, not elsewhere classified	0.5
100	S51	Open wound of forearm	0.5
101	F32	Depressive episode	0.5
102	M48	Spinal stenosis (secondary code only)	0.5
103	E83	Disorders of mineral metabolism	0.4
104	M15	Polyarthritis	0.4
105	D64	Other anemias	0.4
106	L08	Other local infections of skin and subcutaneous tissue	0.4
107	R11	Nausea and vomiting	0.3
108	K52	Other noninfective gastroenteritis and colitis	0.3
109	R50	Fever of unknown origin	0.1

Abbreviation: *ICD-10*, International Statistical Classification of Diseases and Related Health Problems, Tenth Revision

**Supplementary Table 4** List of the 17 comorbidities, assigned weights and associated ICD-10 codes for construction of Charlson Comorbidity Index

#	Condition	Weight	ICD-10 code
1	Myocardial infarction	1	I21, I22, I25.2
2	Congestive heart failure	1	I50
3	Peripheral vascular disease	1	I71, I73.9, I79.0, R02, Z95.8, Z95.9
4	Cerebral vascular disease	1	G45, G46, I60–I64, I67–I69
5	Dementia	1	F00–F02, F05.1
6	Pulmonary disease	1	J40–J47, J60–J67
7	Connective tissue disorder	1	M05, M06, M32–M34, M35.3
8	Peptic ulcer	1	K25–K28
9	Liver disease	1	K70.2, K70.3, K71.7, K73, K74
10	Diabetes	1	E10–E14, excluding subdivisions 2–5
11	Diabetes complications	2	E10–E14, subdivisions 2–5
12	Paraplegia	2	G04.1, G81, G82
13	Renal disease	2	N01, N03, N18, N19, N25, N05.2–N05.6, N07.2–N07.4,
14	Cancer	2	C00–C41, C43, C45–C76, C81–C85, C88, C90–C97
15	Metastatic cancer	3	C77–C80
16	Severe liver disease	3	K72.1, K72.9, K76.6, K76.7
17	AIDS	6	B20–B24

Abbreviations: *AIDS*, Acquired immune deficiency syndrome; *ICD-10*, International Statistical Classification of Diseases and Related Health Problems, Tenth Revision

**Supplementary Table 5** Characteristics of individuals with and without hospital data

Characteristic	With hospital data (n=370,660)	Without hospital data <sup>a</sup> (n=58,094)	<i>p</i> <sup>b</sup>
Deaths, n (%)	2,291 (0.6)	60 (0.1)	<0.001
Age, mean (SD), years	68.5 (8.1)	65.6 (7.7)	<0.001
Age category, n (%)			<0.001
<65	119,127 (32.1)	26,173 (45.1)	
65–74	149,822 (40.4)	23,685 (40.8)	
≥75	101,711 (27.4)	8,236 (14.2)	
Sex, n (%)			<0.001
Female	207,044 (55.9)	29,600 (51.0)	
Male	163,616 (44.1)	28,494 (49.0)	
Ethnicity, n (%)			<0.001
White	349,300 (95.6)	54,110 (93.5)	
Asian	8,313 (2.3)	1,593 (2.8)	
Black	6,229 (1.7)	1,122 (1.9)	
Others	5,576 (1.5)	1,065 (1.8)	
Smoking status, n (%)			<0.001
Never	201,756 (54.6)	35,951 (62.0)	
Previous	130,554 (35.4)	16,923 (29.2)	
Current	37,038 (10.0)	5,073 (8.8)	
Education, n (%)			<0.001
Low	63,227 (17.2)	5,404 (9.4)	
Intermediate	187,950 (51.2)	28,385 (49.2)	
High	115,753 (31.6)	23,868 (41.4)	
Income, n (%)			<0.001
<£18,000	72,821 (23.0)	7,096 (13.9)	
£18,000–30,999	82,596 (26.1)	10,807 (21.1)	
£31,000–51,999	83,099 (26.3)	14,304 (28.0)	
≥£52,000	77,998 (24.6)	18,964 (37.1)	
Townsend deprivation quintile, n (%)			<0.001
1 (least deprived)	74,039 (20.0)	12,405 (21.4)	
2	75,329 (20.4)	12,079 (20.8)	
3	75,128 (20.3)	11,677 (20.1)	
4	74,216 (20.1)	11,596 (20.0)	
5 (most deprived)	71,511 (19.3)	10,262 (17.7)	
Frailty phenotype category, n (%)			<0.001
Non-frail	209,784 (56.6)	39,036 (67.2)	
Pre-frail	146,773 (39.6)	18,419 (31.7)	
Frail	14,103 (3.8)	639 (1.1)	
Frailty index category, n (%)			<0.001
Relatively fit	18,956 (5.1)	6,942 (12.0)	
Less fit	132,051 (35.6)	30,098 (51.8)	
Least fit	170,396 (46.0)	19,340 (33.3)	
Frail	49,257 (13.3)	1,714 (3.0)	

<sup>a</sup> Resided in England at baseline and without hospital data between 1980 and 2020

<sup>b</sup> Based on t-tests or Mann–Whitney *U* tests for continuous variables and  $\chi^2$ -tests for categorical variables

Abbreviations: *IQR*, interquartile range; *SD*, standard deviation



**Supplementary Table 6** Characteristics of the COVID-19 positive subsample (n=2,287) by the source of the disease status

	Confirmed COVID-19 cases (n=2,201; deaths=383) <sup>a</sup>	Died of COVID-19 without positive test record (n=86) <sup>b</sup>	<i>p</i> <sup>c</sup>
Age (year), mean (SD)	67.9 (9.1)	75.4 (6.0)	<0.001
Age category, n (%)			<0.001
<65	857 (38.9)	6 (7.0)	
65–74	645 (29.3)	20 (23.3)	
≥75	699 (31.8)	60 (69.8)	
Sex, n (%)			0.17
Female	1,062 (48.3)	35 (40.7)	
Male	1,139 (51.8)	51 (59.3)	
Ethnicity, n (%)			0.20
White	1,925 (87.8)	82 (95.4)	
Asian	110 (5.0)	2 (2.3)	
Black	103 (4.7)	1 (1.2)	
Others	55 (2.5)	1 (1.2)	
Smoking status, n (%)			0.23
Never	1,070 (49.0)	34 (39.5)	
Previous	847 (38.8)	40 (46.5)	
Current	268 (12.3)	12 (14.0)	
Education, n (%)			0.76
Low	530 (24.4)	23 (27.7)	
Intermediate	1,097 (50.6)	39 (47.0)	
High	543 (25.0)	21 (25.3)	
Income, n (%)			0.015
<£18,000	580 (31.5)	31 (40.8)	
£18,000–30,999	459 (24.9)	25 (32.9)	
£31,000–51,999	417 (22.7)	14 (18.4)	
≥£52,000	385 (20.9)	6 (7.9)	
Townsend deprivation quintile, n (%)			0.36
1 (least deprived)	329 (15.0)	14 (16.3)	
2	366 (16.6)	13 (15.1)	
3	394 (17.9)	22 (25.6)	
4	465 (21.1)	18 (20.9)	
5 (most deprived)	646 (29.4)	19 (22.1)	
Frailty phenotype category, n (%)			0.07
Non-frail	1,097 (49.8)	34 (39.5)	
Pre-frail	925 (42.0)	47 (54.7)	
Frail	179 (8.1)	5 (5.8)	
Frailty index category, n (%)			0.30
Relatively fit	96 (4.4)	2 (2.3)	
Less fit	678 (30.8)	20 (23.3)	
Least fit	996 (45.3)	46 (53.5)	
Frail	431 (19.6)	18 (20.9)	
Hospital Frailty Risk Score category, n (%)			0.001
Low risk	1,730 (78.6)	55 (64.0)	
Intermediate risk	220 (10.0)	10 (11.6)	
High risk	251 (11.4)	21 (24.4)	
Charlson Comorbidity Index score, mean (SD)	0.93 (1.73)	1.09 (1.64)	0.16

<sup>a</sup> Positive in at least one of the COVID-19 tests or recorded as a COVID-19 inpatient in hospital data

<sup>b</sup> With COVID-19 as the primary or contributory cause of death but were not recorded as tested positive or was a COVID-19 inpatient

<sup>c</sup> Based on t-tests or Mann–Whitney *U* tests for continuous variables and  $\chi^2$ -tests for categorical variables

Abbreviations: *IQR*, interquartile range; *SD*, standard deviation

**Supplementary Table 7** Spearman's rank correlation coefficients among the frailty measures and comorbidity

Variable	FP score	FI score	HFRS score	CCI score
FP score <sup>a</sup>	1			
FI score	0.341*	1		
HFRS score	0.115*	0.189*	1	
CCI score	0.123*	0.216*	0.554*	1

<sup>a</sup> FP is considered as a continuous variable using the score across the five criteria (ranging from 0 to 5)

\*  $p < 0.001$

Abbreviations: *CCI*, Charlson Comorbidity Index; *FI*, frailty index; *FP*, frailty phenotype; *HFRS*, hospital frailty risk score

**Supplementary Table 8** Associations between different frailty and comorbidity measures, and being COVID-19 positive (n=428,754)

Variable	Model 1 <sup>a</sup>	Model 2 <sup>a</sup>	Model 3 <sup>a</sup>	Model 4 <sup>a</sup>	Model 5 <sup>a</sup>	Model 6 <sup>a</sup>
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
Age	1.00 (1.00–1.01)	1.00 (0.99–1.00)	1.00 (0.99–1.00)	0.98 (0.98–0.99)*	0.99 (0.98–0.99)*	0.98 (0.98–0.99)*
Male sex	1.34 (1.23–1.45)*	1.38 (1.27–1.50)*	1.38 (1.27–1.50)*	1.30 (1.19–1.41)*	1.26 (1.16–1.36)*	1.30 (1.20–1.41)*
FP category (ref. non-frail)						
Pre-frail	-	1.32 (1.21–1.44)*	-	-	-	1.11 (1.01–1.21)*
Frail	-	2.89 (2.47–3.38)*	-	-	-	1.48 (1.24–1.77)*
FI category (ref. relatively fit)						
Less fit	-	-	1.17 (0.94–1.44)	-	-	1.14 (0.92–1.41)
Least fit	-	-	1.51 (1.23–1.86)*	-	-	1.30 (1.06–1.61)*
Frail	-	-	2.48 (1.99–3.09)*	-	-	1.44 (1.14–1.81)*
HFRS category (ref. low risk)						
Intermediate risk	-	-	-	3.66 (3.18–4.21)*	-	2.55 (2.18–2.97)*
High risk	-	-	-	19.29 (16.82–22.12)*	-	10.72 (9.02–12.72)*
CCI score	-	-	-	-	1.45 (1.42–1.49)*	1.17 (1.13–1.21)*
AUC (95% CI)	0.54 (0.53–0.55)	0.57 (0.56–0.58)	0.58 (0.57–0.59)	0.64 (0.62–0.65)	0.64 (0.63–0.65)	0.67 (0.66–0.68)

<sup>a</sup> Model 1: age (continuous) and sex; models 2–5: added FP, FI, HFRS, and CCI as independent variables respectively; model 6: all listed variables were mutually adjusted for

\* Significant with a false discovery rate corrected significance level at 0.041

Abbreviations: *AUC*, area under the receiver operating characteristic curves; *CCI*, Charlson Comorbidity Index; *CI*, confidence interval; *FI*, frailty index; *FP*, frailty phenotype; *HFRS*, Hospital Frailty Risk Score; *OR*, odds ratio.

**Supplementary Table 9** Associations between the Hospital Frailty Risk Score and COVID-19 mortality stratified by age in the full sample and COVID-19 positive subsample

Variable	Age category			$P_{\text{interaction}}^a$
	<65 years OR (95% CI)	65–74 years OR (95% CI)	$\geq 75$ years OR (95% CI)	
<b>(a) Full sample (n=428,754)</b>				
HFRS category (ref. low risk)				
Intermediate risk	10.60 (3.66–30.68)*	6.53 (3.90–10.93)*	4.70 (3.38–6.54)*	0.003
High risk	77.71 (26.68–226.29)*	34.19 (20.59–56.76)*	18.71 (13.78–25.41)*	
Male sex	1.14 (0.56–2.31)	1.89 (1.29–2.76)*	2.13 (1.63–2.78)*	
AUC (95% CI)	0.62 (0.51–0.72)	0.71 (0.65–0.76)	0.73 (0.69–0.76)	
<b>(b) COVID-19 positive sample<sup>b</sup> (n=2,287)</b>				
HFRS category (ref. low risk)				
Intermediate risk	5.27 (1.67–16.62)*	2.23 (1.23–4.06)*	1.28 (0.85–1.93)	<0.001
High risk	15.39 (4.41–53.64)*	1.70 (0.96–3.00)	1.15 (0.79–1.67)	
Male sex	1.04 (0.49–2.19)	1.54 (1.01–2.34)	1.54 (1.12–2.12)	
AUC (95% CI)	0.61 (0.50–0.72)	0.60 (0.54–0.65)	0.57 (0.52–0.61)	

<sup>a</sup>  $P_{\text{interaction}}$  represents the  $p$ -values of the interaction terms fitted between the HFRS and age as continuous variables

<sup>b</sup> The COVID-19 positive subsample is nested within the full sample

\* Significant with a false discovery rate corrected significance level at 0.033

Abbreviations: *AUC*, area under the receiver operating characteristic curves; *CI*, confidence interval; *HFRS*, Hospital Frailty Risk Score; *OR*, odds ratio.

**Supplementary Table 10** Associations between different frailty and comorbidity measures, and mortality due to COVID-19 and other causes in the full sample and COVID-19 positive subsample

Variable	Model 1 <sup>a</sup>		Model 2 <sup>a</sup>		Model 3 <sup>a</sup>		Model 4 <sup>a</sup>		Model 5 <sup>a</sup>	
	COVID-19 mortality	Other deaths <sup>b</sup>	COVID-19 mortality	Other deaths <sup>b</sup>	COVID-19 mortality	Other deaths <sup>b</sup>	COVID-19 mortality	Other deaths <sup>b</sup>	COVID-19 mortality	Other deaths <sup>b</sup>
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
<b>(a) Full sample (n=428,754)</b>										
Age	1.14 (1.12–1.16)*	1.12 (1.11–1.13)*	1.14 (1.12–1.16)*	1.12 (1.11–1.13)*	1.11 (1.09–1.13)*	1.09 (1.08–1.10)*	1.12 (1.10–1.14)*	1.09 (1.08–1.10)*	1.11 (1.09–1.13)*	1.08 (1.07–1.09)*
Male sex	2.16 (1.76–2.66)*	1.62 (1.48–1.78)*	2.15 (1.75–2.65)*	1.62 (1.48–1.77)*	1.98 (1.61–2.44)*	1.51 (1.38–1.65)*	1.80 (1.46–2.21)*	1.26 (1.15–1.38)*	1.93 (1.57–2.38)*	1.26 (1.14–1.38)*
FP category (ref. non-frail)										
Pre-frail	1.55 (1.26–1.91)*	1.34 (1.22–1.47)*	-	-	-	-	-	-	1.19 (0.95–1.48)	1.01 (0.91–1.11)
Frail	3.82 (2.69–5.41)*	2.94 (2.49–3.47)*	-	-	-	-	-	-	1.54 (1.04–2.28)	1.10 (0.90–1.34)
FI category (ref. relatively fit)										
Less fit	-	-	1.41 (0.74–2.71)	1.10 (0.86–1.41)	-	-	-	-	1.32 (0.69–2.54)	0.99 (0.77–1.27)
Least fit	-	-	2.00 (1.06–3.79)	1.40 (1.09–1.78)*	-	-	-	-	1.49 (0.78–2.84)	0.91 (0.71–1.16)
Frail	-	-	3.75 (1.05–7.21)*	2.36 (1.83–3.04)*	-	-	-	-	1.63 (0.83–3.19)	0.75 (0.57–0.98)
HFRS category (ref. low risk)										
Intermediate risk	-	-	-	-	5.42 (4.13–7.11)*	8.47 (7.58–9.46)*	-	-	3.22 (2.38–4.35)*	2.91 (2.55–3.32)*
High risk	-	-	-	-	24.57 (18.94–31.87)*	25.47 (22.43–28.92)*	-	-	11.41 (8.27–15.74)*	5.58 (4.77–6.53)*
CCI score	-	-	-	-	-	-	1.59 (1.52–1.66)*	1.86 (1.82–1.89)*	1.26 (1.19–1.34)*	1.64 (1.60–1.68)*

**Supplementary Table 10** (*continued*)

Variable	Model 1 <sup>a</sup>		Model 2 <sup>a</sup>		Model 3 <sup>a</sup>		Model 4 <sup>a</sup>		Model 5 <sup>a</sup>	
	COVID-19 mortality	Other deaths <sup>b</sup>	COVID-19 mortality	Other deaths <sup>b</sup>	COVID-19 mortality	Other deaths <sup>b</sup>	COVID-19 mortality	Other deaths <sup>b</sup>	COVID-19 mortality	Other deaths <sup>b</sup>
	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)	OR (95% CI)
<b>(b) COVID-19 positive subsample<sup>c</sup> (n=2,287)</b>										
Age	1.13 (1.11–1.15)*	1.10 (1.07–1.14)*	1.13 (1.11–1.15)*	1.10 (1.06–1.14)*	1.13 (1.11–1.15)*	1.08 (1.05–1.12)*	1.13 (1.11–1.15)*	1.08 (1.05–1.12)*	1.13 (1.11–1.15)*	1.09 (1.05–1.13)*
Male sex	1.55 (1.22–1.98)*	1.74 (1.07–2.83)*	1.55 (1.22–1.97)*	1.77 (1.09–2.87)*	1.54 (1.21–1.96)*	1.74 (1.07–2.82)*	1.47 (1.16–1.88)*	1.53 (0.93–2.50)	1.50 (1.17–1.91)*	1.52 (0.92–2.51)
FP category (ref. non-frail)										
Pre-frail	1.19 (0.93–1.52)	1.15 (0.70–1.87)	-	-	-	-	-	-	1.11 (0.85–1.44)	0.96 (0.57–1.63)
Frail	1.29 (0.85–1.96)	1.53 (0.71–3.31)	-	-	-	-	-	-	1.03 (0.65–1.63)	0.84 (0.34–2.07)
FI category (ref. relatively fit)										
Less fit	-	-	1.07 (0.51–2.24)	0.45 (0.14–1.41)	-	-	-	-	1.04 (0.49–2.17)	0.40 (0.12–1.27)
Least fit	-	-	1.19 (0.58–2.45)	0.65 (0.22–1.91)	-	-	-	-	1.08 (0.52–2.24)	0.52 (0.17–1.56)
Frail	-	-	1.35 (0.65–2.84)	0.82 (0.27–2.52)	-	-	-	-	1.05 (0.49–2.24)	0.45 (0.14–1.50)
HFRS category (ref. low risk)										
Intermediate risk	-	-	-	-	1.79 (1.28–2.52)*	3.07 (1.68–5.60)*	-	-	1.29 (0.89–1.86)	1.36 (0.70–2.65)
High risk	-	-	-	-	1.41 (1.03–1.93)	2.23 (1.23–4.05)*	-	-	0.89 (0.61–1.29)	0.71 (0.34–1.45)
CCI score	-	-	-	-	-	-	1.20 (1.12–1.27)*	1.44 (1.32–1.58)*	1.19 (1.11–1.28)*	1.47 (1.32–1.65)*

<sup>a</sup> Associations were assessed using multinomial logistic regression models, in which “COVID-19 mortality” and “other deaths” were compared to those who were alive as of. Model 1–4: FP, FI, HFRS, and CCI as independent variables respectively, adjusted for age and sex; model 6: all listed variables were mutually adjusted for

<sup>b</sup> Other deaths are individuals died between March 1 and September 21, 2020 whose primary and contributory causes of death do not include COVID-19

<sup>c</sup> The COVID-19 positive subsample is nested within the full sample

\* Significant with a false discovery rate corrected significance level at 0.031

Abbreviations: *CCI*, Charlson Comorbidity Index; *CI*, confidence interval; *FI*, frailty index; *FP*, frailty phenotype; *HFRS*, Hospital Frailty Risk Score; *OR*, odds ratio.