Figure S1


Nucleoprotein: IgG
Number of outlier samples
0
20
40
60
0
100
200
300
400
Spike: IgG
Number of outlier samples
0
10
20
30
0
500
1000
Protein 3a: IgG
Number of outlier samples
0
5
10
15
20
25
0
100
200
Non-structural protein 8: IgG
Number of outlier samples
0
5
10
15
0
25
50
75
100
125
Membrane protein: IgG
Number of outlier samples
0.0
2.5
5.0
7.5
0
50
100
150
200
Membrane protein: IgM
Number of outlier samples
0
25
50
75
0
50
100
150
200

Figure S2

YWXYFXK



PDB ID: 7A97

YXXYFXK: site #2



PDB ID: 7A95

YXXYFXK: site #3



PDB ID: 7A95

Figure S3


Training
Testing
Model prediction
Mild
Moderate/Severe


Mild vs. Moderate/Severe
Sensitivity
Specificity
Training AUC =0.92 [0.88, 0.96]
Testing AUC =0.9 [0.85, 0.96]


SERA IgG Score
Days since symptom onset
severity
Severe
Moderate
Mild


IWAS S1 Furin Cleavage
IWAS ENAC ~205
severity
Mild
Moderate
Severe
Pearson's r = 0.88


Max. IWAS, SPIKE Furin Cleavage Site
Days since Symptoms Onset
severity
Mild_Asymp
Moderate
Severe


Max. IWAS, ENAC ~205
Days since Symptoms Onset
severity
Mild_Asymp
Moderate
Severe

Figure S4


Observed
WT > Mut
Mut > WT
Count
1000
100
10
0
-10
-100
-1000
0
100
200
Number of samples with significant change in PIWAS value
Random
WT > Mut
Mut > WT
Count
1000
100
10
0
-10
-100
-1000
0
100
200
Number of samples with significant change in PIWAS value