Title

Modelling the impact of health-related variables, age, migration, and socio-economic factors in the geographical distribution of early tested case-fatality risks associated with COVID-19 in Mexico

Authors

*Ricardo Ramírez-Aldana¹, Juan Carlos Gomez-Verjan¹, Omar Yaxmehen Bello-Chavolla¹,
Carmen García-Peña¹

Institutions

¹Research Division, Instituto Nacional de Geriatria, Mexico City, Mexico

***Corresponding Author:** Ricardo Ramirez-Aldana **e-mail:**
ricardoramirezaldana@gmail.com, Research Division, Instituto Nacional de Geriatria
(INGER), Anillo Perif. 2767, San Jerónimo Lídice, La Magdalena Contreras, 10200, Mexico
City, Mexico.

ABSTRACT

COVID-19 is a respiratory disease caused by SARS-CoV-2, which has significantly impacted economic and public healthcare systems world-wide. SARS-CoV-2 is highly lethal in older adults (>65 years old) and in cases with underlying medical conditions including chronic respiratory diseases, immunosuppression, and cardio-metabolic diseases including severe obesity, diabetes, and hypertension. The course of the COVID-19 pandemic in Mexico has led to many fatal cases in younger patients attributable to cardio-metabolic conditions. Here, we aimed to perform an early spatial epidemiological analysis for the COVID-19 outbreak in Mexico to evaluate how tested case-fatality risks (t-CFRs) are geographically distributed and to explore spatial predictors of early t-CFRs considering the variation of their impact on COVID-19 fatality across different states in Mexico, controlling for the severity of the disease. As results, considering health related variables; diabetes and obesity were highly associated with COVID-19 fatality. We identified that both external and internal migration had an important impact over early COVID-19 risks in Mexico, with external migration having the second highest impact when analyzing Mexico as a whole. Physicians-to-population ratio, as a representation of urbanity, population density, and overcrowding households, has the highest impact on t-CFRs, whereas the age group of 10 to 39 years was associated with lower risks. Geographically, the states of Quintana Roo, Baja California, Chihuahua, and Tabasco had higher t-CFRs and relative risks comparing with a national standard, suggesting that risks in these states were above of what was nationally expected; additionally, the strength of the association between some spatial predictors and the COVID-19 fatality risks variates by zone depending on the predictor.

Keywords: COVID-19, generalized geographically weighted regression, Mexico, SARS-CoV-2, spatial clusters, spatial epidemiology, spatial statistics.

49

50 INTRODUCTION

51 COVID-19 is a respiratory disease caused by SARS-CoV-2 (severe acute respiratory
52 syndrome coronavirus 2), which has caused almost twenty million cases around the world
53 and caused 790,000 deaths as of August 20th, 2020 (1). SARS-CoV-2 is a highly
54 contagious RNA virus from the Coronaviridae family, with a small genome of about 30,000
55 nucleotides closely related to the bat coronavirus (RaTG13) (2). Given its global spread,
56 there is an urgent need for scientific and health systems around the world to understand
57 the epidemiology, pathogenicity, and mechanisms of immunological defenses to develop
58 possible therapeutic and public health alternatives to fight against one of the most
59 outstanding threats to public health since *Spanish Influenza* over a hundred years ago (3).
60 According to the World Health Organization (WHO), the groups most susceptible to
61 acquire infection and develop adverse outcomes are people with underlying medical
62 conditions and older adults (>65 years old), particularly those living at nursing homes.
63 Medical conditions which have been associated with increased susceptibility for adverse
64 outcomes related to SARS-CoV-2 infection include chronic obstructive pulmonary disease
65 (COPD), chronic kidney disease (CKD), cardiovascular diseases, liver diseases, moderate
66 asthma, immunosuppression (HIV/AIDS, bone marrow transplantation, cancer treatment,
67 and genetic immune deficiencies), and particularly, severe obesity, diabetes, and
68 hypertension (4). These associations may be related to the strong link between pro-
69 inflammatory cytokines in response to infection and the pathogenesis of SARS-CoV-2,
70 which can be seen in pneumonia patients with severe COVID-19 disease exhibiting
71 systemic hyper-inflammation known as cytokine storm or as a secondary hemophagocytic
72 lymphohistocytosis (5).
73 On the other hand, accordingly to the OCDE, in the last ten years the countries with the
74 highest prevalence of cardio-metabolic conditions linked to increased risk of severe

COVID-19, including obesity, type 2 diabetes, and hypertension, are Mexico and the United States of America (USA), when considering adults from 15 to 74 years old (6). Thus, considering that the population pyramid is flattening more and more, indicating an aging population structure; the high rates of diabetes and obesity in Mexico are likely to increase susceptibility to higher rates of mortality attributable to COVID-19 even in younger populations (7).

In response to SARS-CoV-2 spread worldwide, mobility restrictions have been imposed to reduce community-level transmission; nevertheless, early influence of human mobility particularly internal and external migration has largely driven the spread of COVID-19 (8). When considering the high transmissibility of SARS-CoV-2, human mobility gains relevance in early stages of spread, in which people travelling from other countries can drive increased rates of transmission (9). SARS-CoV-2 spread related to human mobility is relevant, particularly considering that the risk of transmission and adverse outcomes are related to inequalities and suboptimal socio-economic conditions(10). In this sense, the risk of becoming infected and dying might increase in areas without optimum socio-economic conditions, for instance, spatial units in which people live in overcrowding households or without access to potable water or drainage.

Spatial analyses allow us to understand how the fatality risks are distributed along a territory, the presence of spatial clusters, and how the effects of the variables associated with the risks variate along any given territory. In this sense, several examples of geoeidemiology studies in autoinflammatory diseases, specific syndromes, infectious diseases, among others, have been helpful for the development of public health policies and to understand the structure of disease spread (11–13). In the present work, we performed spatial analyses in order to derive spatial relationships corresponding to the geographical phenomenon of the COVID-19 outbreak in Mexico and its associated fatality risks, using for this task statistical methods including spatial clustering through local

indicators of spatial autocorrelation and generalized geographically weighted regression. Additionally, we wanted to identify spatial units or regions in Mexico that should be considered and analyzed with care to better understand the propagation of the disease and its associated fatality risks in the early stages of COVID-19 spread.

MATERIAL AND METHODS

Data sources

We obtained state-level variables considering the 32 states in Mexico (**Table 1**). A first group of variables were obtained from the epidemiological surveillance entity of Mexico (Dirección General de Vigilancia Epidemiológica, Secretaría de Salud) at an individual level, the data set corresponded to observations until April 21st, 2020 (14). These variables concern health and very general socioeconomic information associated with people who were suspected for COVID-19 in Mexico and underwent real time RT-PCR for SARS-CoV-2 confirmation. Available variables include the presence of diabetes, obesity, chronic renal problems (CKD), chronic obstructive pulmonary disease (COPD), pregnancy, hypertension, immunosuppression, cardiovascular disease, pneumonia, as well as age, and whether the patient was hospitalized, admitted to an intensive care unit (ICU), or required intubation. We grouped age into groups as explained in the model selection process below and we finally used three age groups: 10-39, 40-69, and 70 years old and over. We also computed the risk of death due to the disease on individuals who were tested for SARS-CoV-2, or tested case-fatality risks (t-CFRs), by considering as a death that which was recorded after having a positive test (the information concerning positive tests and death is available in the epidemiological data set), method consistent with the official numbers. We aggregated all variables at a state level as counts and used them as relative frequencies (rates) in all analyses.

Additional variables concerning socio-demographic, economic, mobility, and climatic features were obtained at a state-level. In terms of socio-demographic variables, from the

National Institute of Geography and Statistics (INEGI), we extracted information concerning clustering of individuals: population density (people per km^2) in 2015 and the proportion of people in a household living in an overcrowded place in 2017; literacy rate of population aged ≥ 15 years in 2015; people settled in rural areas in 2010 (%) (a location was considered as rural when there are $< 2,500$ habitants); and the number of physicians available by every 1000 people in 2015, which was obtained from the National Institute of Public Health (INSP). In terms of economic variables, we obtained from INEGI the state contribution to gross domestic product (GDP) in 2018, which we modified by considering only those values associated with states not containing the biggest cities in Mexico to improve the linearity assumption with the transformed response. We also obtained information concerning people living in poverty in 2018 (%), as it is defined and calculated by CONEVAL according to a multidimensional index obtained from *per capita* income and an index of social deprivation (15). In terms of mobility, we extracted from INEGI information concerning internal migration, as the rate of people aged 5 years and over living in another state five years before 2014, and external migration, as the rate of people aged ≥ 5 years living in another country five years before 2014, both proxies of internal and external mobility, respectively; and, the number of flights in 2019 by state, which we calculated from information associated with the number of flights by airport in Mexico (Ministry of Communication and Transport)(16). Finally, information concerning average temperature ($^{\circ}C$) in March 2020 was obtained from the National Council of Water (CONAGUA).

The risk of dying in tested individuals (t-CFRs) was chosen as the dependent variable on the basis of relevant indicators of COVID-19 epidemiology in Mexico; notably, the number of tests per 100,00 individuals is limited compared to other countries, which decreases detection rates and given the likely under-detection of mild SARS-CoV-2 cases in this setting, standardizing deaths by tested cases considers the extent of detection, which

could similarly be influenced by structural factors (17). The remaining variables obtained and calculated from the different data sources are treated as explanatory, except for hospitalization, ICU, and intubation, which we considered as control variables, being an approximate measure of the presence of severe COVID-19 cases and possibly access to services attending COVID-19 in a region.

162 **Table 1.** Features extracted for all the analyses by state used to predict tested case-fatality risks of COVID-19 in Mexico.
 163 Abbreviations: GDP, Gross Domestic Product; COPD, Chronic Obstructive Pulmonary Disease; and ICU, Intensive Care Unit.

State	Deaths caused by COVID-19*	Number of tested individuals (offset)*	Rural 2006 (%)**	Poverty 2018 (%)***	Density 2015+	Literacy 2015 (%)+	GDP 2018 (%)+	Temperature March 2020 (°C)++	Internal migration 2014 (%)+	External migration 2014 (%)+	Physicians by 1000 people 2015+++	Overcrowding 2017 (%)+
AGUASCALIENTES	2	1070	18.84	26.18	233.70	97.00	1.31	19.30	3.60	0.70	1.33	3.90
BAJA CALIFORNIA	100	2353	6.99	23.26	46.40	97.60	3.14	14.40	3.80	1.90	1.05	6.40
BAJA CALIFORNIA SUR	9	978	15.22	18.07	9.60	96.80	0.98	19.80	8.20	0.50	1.50	10.40
CAMPECHE	4	229	26.00	46.25	15.60	92.90	2.98	28.00	4.10	0.20	1.16	16.50
COAHUILA DE ZARAGOZA	23	2472	9.95	22.49	19.50	97.10	3.50	22.10	2.10	0.40	1.44	4.50
COLIMA	2	133	12.41	30.87	126.40	95.60	0.61	26.60	5.60	1.50	1.56	10.70
CHIAPAS	4	440	52.26	76.41	71.20	84.20	1.56	25.60	1.50	0.50	0.53	20.50
CHIHUAHUA	36	705	15.51	26.28	14.40	95.00	3.20	16.70	1.20	1.20	1.11	5.60
CIUDAD DE MÉXICO	224	10183	0.35	30.55	5967.30	97.70	17.64	19.40	2.10	0.40	2.99	7.10
DURANGO	4	407	32.77	37.35	14.20	96.20	1.13	18.40	2.60	0.80	1.17	5.40
GUANAJUATO	9	2482	30.29	43.38	191.30	93.00	4.11	20.60	1.50	1.00	0.84	5.50
GUERRERO	16	583	42.44	66.47	55.60	85.50	1.37	26.20	1.90	1.10	0.64	27.60
HIDALGO	16	608	47.69	43.85	137.30	91.00	1.54	20.50	3.90	0.90	0.66	8.20
JALISCO	15	3440	13.85	28.43	99.80	95.80	6.91	22.40	2.20	1.10	1.26	5.30
MÉXICO	71	5759	12.74	42.72	724.20	95.80	8.93	16.30	2.80	0.40	0.81	8.60
MICHOACÁN	19	1124	32.06	46.04	78.20	90.80	2.43	21.70	1.70	1.20	0.72	10.10
MORELOS	14	525	13.91	50.82	390.20	94.30	1.13	24.70	3.30	0.90	1.04	10.80
NAYARIT	5	299	33.48	34.77	42.40	94.30	0.68	24.70	4.30	1.40	1.06	8.10
NUEVO LEÓN	6	4482	5.64	14.53	79.80	97.40	7.47	22.90	2.70	0.40	1.30	3.10
OAXACA	7	522	52.89	66.35	42.30	84.20	1.48	24.60	2.00	1.20	0.61	16.90
PUEBLA	53	1651	29.38	58.92	179.80	90.80	3.39	20.00	2.30	0.60	0.83	9.70

QUERÉTARO	6	630	30.13	27.58	174.40	94.70	2.33	22.00	5.60	0.90	0.85	6.90
QUINTANA ROO	47	982	14.42	27.57	33.60	95.50	1.63	26.80	8.00	0.70	0.96	17.50
SAN LUIS POTOSÍ	5	1167	37.31	43.40	44.50	92.90	2.11	24.60	2.70	0.80	0.85	6.20
SINALOA	52	1580	29.17	30.88	51.70	95.20	2.23	22.90	2.50	0.70	1.16	9.20
SONORA	13	869	14.23	28.19	15.90	96.70	3.26	17.70	2.90	1.10	1.39	7.00
TABASCO	55	1266	44.98	53.65	96.90	93.80	2.61	27.60	2.30	0.30	0.92	12.00
TAMAULIPAS	7	1210	12.74	35.07	42.90	96.00	2.82	25.20	1.80	0.30	1.20	6.90
TLAXCALA	5	587	21.77	48.38	318.40	95.20	0.56	16.30	3.40	0.60	0.81	9.50
VERACRUZ	13	1520	39.33	61.78	113.00	89.80	4.59	23.70	3.00	0.60	0.87	12.60
YUCATÁN	12	959	17.01	40.80	53.10	91.90	1.46	26.90	3.60	0.40	1.39	13.90
ZACATECAS	3	419	42.73	46.76	21.00	94.90	0.88	19.10	2.10	1.40	0.89	5.00

* Dirección General de Epidemiología, Secretaría de Salud (General Direction of Epidemiology, Health Ministry)

** Instituto Nacional de Salud Pública, Encuesta Nacional de la Dinámica Demográfica (National Public Health Institute, National Survey of Demographic Dynamic)

*** Estimations obtained by CONEVAL (Consejo Nacional de Evaluación de la Política de Desarrollo Social, National Council for the Evaluation of Social Development Policy) based on the ENIGH (National Survey of Household Income and Expenditure) 2008, 2010, 2012, 2014, 2016, and 2018.

+ Instituto Nacional de Estadística y Geografía (National Institute of Statistics and Geography)

++ Consejo Nacional del Agua (CONAGUA) (National Water Commission)

+++ Instituto Nacional de Salud Pública (INSP) (The National Institute of Public Health of Mexico)

Table 1(continued)

State	Diabetes (%)*	Obesity (%)*	Chronic renal problems (%)*	COPD (%)*	Pregnancy (%)*	Hypertension (%)*	Immuno-deficiency (%)*	Cardio-vascular (%)*	Pneumonia (%)*	ICU (%)*	Intubated (%)*	Hospitalized (%)*	Age group 10-39(%)*	Age group 40-69(%)*	Age group 70 years and over(%)*
AGUASCALIENTES	7.216	14.204	1.947	1.833	0.573	13.402	1.833	2.864	7.446	0.458	0.916	12.257	54.296	33.333	4.582
BAJA CALIFORNIA	12.420	18.059	1.570	1.713	1.071	22.484	2.355	3.854	21.485	1.213	2.784	27.195	49.179	43.683	3.854
BAJA CALIFORNIA SUR	11.350	10.763	0.978	0.978	2.544	14.677	2.153	3.131	13.503	1.566	1.370	17.613	53.816	35.225	4.892

CAMPECHE	11.111	16.239	0.000	0.855	0.000	12.821	2.137	2.137	16.667	2.137	1.282	26.923	46.581	45.299	5.556
COAHUILA DE ZARAGOZA	11.965	16.079	2.015	1.847	0.756	16.037	2.603	2.981	8.942	1.847	0.672	14.484	47.397	44.249	5.584
COLIMA	14.388	19.424	5.036	2.878	2.158	15.827	2.158	3.597	25.899	2.878	2.878	33.094	52.518	34.532	7.194
CHIAPAS	11.836	13.050	1.973	2.883	1.669	15.630	2.276	2.731	17.299	3.642	3.035	27.314	49.469	39.150	4.856
CHIHUAHUA	15.385	13.501	2.826	4.710	1.413	21.821	3.297	5.181	33.438	9.733	3.925	45.997	45.997	40.816	7.849
CIUDAD DE MÉXICO	10.796	15.511	1.658	1.872	0.618	14.970	2.688	3.049	14.670	2.044	2.113	20.356	44.619	47.926	4.767
DURANGO	15.615	13.621	2.990	4.983	0.831	24.917	4.485	5.980	19.767	2.326	2.159	33.389	43.355	39.203	13.787
GUANAJUATO	10.131	11.997	1.908	3.942	1.399	14.455	2.755	2.671	10.428	1.696	1.526	17.338	50.996	34.718	6.867
GUERRERO	15.456	17.531	2.801	3.216	1.245	17.635	1.763	3.734	26.349	5.394	4.253	33.091	43.154	44.813	7.573
HIDALGO	18.491	15.976	4.142	4.734	1.479	19.970	4.438	2.071	21.893	1.627	2.959	31.953	42.751	44.231	9.172
JALISCO	10.876	13.425	3.533	3.625	1.475	16.559	2.888	3.840	12.442	1.874	1.260	20.399	49.124	37.665	7.957
MÉXICO	12.899	15.374	2.358	2.358	0.838	15.140	2.747	3.079	24.318	2.864	3.254	33.164	45.733	44.564	5.943
MICHOACÁN	13.541	16.003	2.752	4.272	1.376	19.406	2.462	4.345	22.230	2.172	2.462	29.616	41.999	41.202	11.369
MORELOS	14.397	14.786	2.918	4.086	1.167	17.899	2.724	2.529	24.708	4.086	3.502	34.825	43.774	42.218	9.728
NAYARIT	16.188	16.449	4.178	4.700	0.261	23.760	3.394	4.439	20.888	2.350	2.350	30.548	41.253	45.431	9.922
NUEVO LEÓN	10.250	11.597	1.594	1.374	0.879	14.702	2.198	3.023	7.227	0.989	0.632	11.761	49.217	40.918	5.001
OAXACA	15.768	12.656	3.631	5.394	1.141	16.183	3.423	3.008	25.519	1.971	2.697	33.299	43.050	44.295	9.232
PUEBLA	14.850	16.630	2.948	3.281	1.001	16.407	2.169	3.504	23.637	2.503	3.003	31.535	40.712	48.498	8.343
QUERÉTARO	11.736	14.425	3.423	2.689	0.978	17.359	4.890	3.178	17.604	1.711	2.200	28.362	47.677	38.631	6.846
QUINTANA ROO	11.429	15.238	1.270	0.635	1.905	9.206	2.857	2.222	23.492	6.032	4.762	38.413	51.111	30.476	4.127
SAN LUIS POTOSÍ	12.211	11.469	2.723	4.785	1.650	16.007	2.805	3.548	16.089	2.063	1.815	24.340	48.102	35.066	9.901
SINALOA	12.863	19.911	2.218	3.549	0.641	21.883	2.661	3.401	19.961	2.366	1.676	32.134	40.069	48.053	9.463
SONORA	13.734	18.842	2.043	2.838	2.270	23.156	2.724	5.675	20.318	1.476	1.022	29.512	48.354	41.317	7.151
TABASCO	15.422	21.687	1.205	2.088	1.526	20.321	1.124	2.972	13.976	3.614	1.687	21.365	42.570	50.361	5.221
TAMAULIPAS	13.875	15.107	2.381	1.888	1.888	18.062	2.545	3.530	9.278	1.642	0.903	16.256	49.015	42.529	5.008
TLAXCALA	14.211	10.702	2.456	2.807	0.877	15.088	3.684	2.632	18.947	4.211	1.930	22.982	43.333	45.789	7.544
VERACRUZ	13.582	15.734	2.965	3.252	1.387	18.747	3.396	2.965	18.890	2.726	1.435	31.707	46.389	42.037	8.082
YUCATÁN	10.733	16.718	1.651	2.786	1.342	18.885	1.238	2.683	13.829	3.096	2.167	22.910	47.678	40.454	7.637

ZACATECAS	14.390	16.211	4.007	5.100	1.821	25.137	4.736	4.918	23.133	2.186	1.457	27.687	41.712	41.894	14.026
-----------	--------	--------	-------	-------	-------	--------	-------	-------	--------	-------	-------	--------	--------	--------	--------

* Dirección General de Epidemiología, Secretaría de Salud (General Direction of Epidemiology, Health Ministry)

** Instituto Nacional de Salud Pública, Encuesta Nacional de la Dinámica Demográfica (National Public Health Institute, National Survey of Demographic Dynamic)

*** Estimations obtained by CONEVAL (Consejo Nacional de Evaluación de la Política de Desarrollo Social, National Council for the Evaluation of Social Development Policy) based on the ENIGH (National Survey of Household Income and Expenditure) 2008, 2010, 2012, 2014, 2016, and 2018.

+ Instituto Nacional de Estadística y Geografía (National Institute of Statistics and Geography)

++ Consejo Nacional del Agua (CONAGUA) (National Water Commission)

+++ Instituto Nacional de Salud Pública (INSP) (The National Institute of Public Health of Mexico)

COVID-19 t-CFRs estimation by state

We obtained quantile maps associated with raw and smoothed t-CFRs of COVID-19 cases. The risks by state were smoothed by using an empirical Bayes estimator, which is a biased estimator that improves variance instability proper of risks estimated in small-sized spatial units (18); however, the analyses were performed with both raw and smoothed risks to compare results. We also obtained maps concerning relative risks, understanding them as a comparison of the observed number of events by state to a national standard, the latter using the expected number of events considering as if risks in a state were the same as those at a national level.

Spatial weight estimation and spatial autocorrelation

We obtained queen contiguity weights (19), which consider as neighbors those states sharing at least a point in common, obtaining a squared matrix of dimension 32 with all entries equal to zero or one, where a one indicates that two states are neighbors. From these neighbors, weights are calculated by integrating a matrix in a row-standardized form. Moran's I statistic (20) was obtained as a measure of global spatial autocorrelation and its significance was assessed through a random permutation inference technique based on simulations. Local indicators of spatial autocorrelation (LISA) were obtained (21) and used to derive significant spatial clustering through four cluster types: High-High, Low-Low, High-Low, and Low-High. For instance, the Low-Low cluster indicates states with low values of a variable that are significantly surrounded by regions with similarly low values.

Spatial multivariable linear model

A multivariable Generalized Geographically Weighted Regression (GGWR) was fitted (22). In this model, a dependent variable is measured for each spatial unit and independent variables (inputs) are simultaneously considered as well, such that the corresponding parameters depend on the coordinates in which the state is spatially located (centroids in the case of polygons); therefore, a parameter is associated with each state and

independent variable. To be able to estimate such model, a weighting diagonal matrix is considered, we used Gaussian spatial weighting to generate it. These weights determine the relationship from any state to another in terms of the distance (Euclidean) between states and a bandwidth. The bandwidth determines which spatial units are similar under the GGWR and can be selected using automatized methods, we used a cross-validation (CV) method with an adaptive scheme, i.e. a different bandwidth was used for each unit. Since the response variable is a count (number of deaths), a GGWR with a Poisson distribution and logarithm link function was used, including as offset term the number of people tested with COVID-19 in a logarithmic scale, thus modelling the t-CFRs instead of just the number of deaths.

A global multivariable model for all states, a Poisson multivariable linear model (generalized linear model or GLM) with offset and a logarithmic link function, was also fitted and significant variables were identified (23). To obtain the best possible model, including variables with the greatest effect on the risks and satisfying as much as possible all statistical assumptions, we used the following selection process: 1) We fitted univariable Poisson models with offset and logarithmic link function, identified significant effects, and ordered them in absolute value from highest to lowest, 2) We identified variables with good and acceptable linear association with the log-transformed mortality risk by obtaining scatter plots between variables and the transformed risk, including a smoothed LOESS (locally estimated scatterplot smoothing) curve; and, when possible, variables were transformed to improve this assumption, as for GDP as explained above, 3) VIF was used to assess multicollinearity; thus, we fitted a model including variables with acceptable and good linear association, and eliminated any variable with a $VIF > 10$, 4) We added to the resulting model one by one significant variables in the univariable models. First, we added the variable with the highest effect and fitted the associated model identifying whether $VIF > 10$. If not, we added the variable, and if $VIF > 10$, the model was

not modified. We proceeded with the resulting model repeating the same process with the second highest effect; and so on, 5) From this process, we ended with a model consisting on all variables with the greatest univariable effects (0.08 and above in absolute value), better linear behavior, and without multicollinearity ($VIF \leq 10$), 6) We used residuals associated with other three Poisson models concerning ICU, intubated, and hospitalization, including a subset of variables pertaining to the model for fatality risk as inputs, thus obtaining the effects each confounder has over the risks without the effects that explanatory variables have over these confounders. These residuals and the corresponding estimated parameters do not have a meaning, 7) We evaluated goodness-of-fit and validated all model assumptions. For age, we obtained age groups: 0-1, 2-9, 10-19, ..., 80-89, and 90 years old and over, fitted the corresponding univariable Poisson models with offset, and identified significant age groups and the direction of the association to obtain a new set of age groups: 10-39, 40-69, and 70 and more; and proceeded with these variables as with the others.

The GGWR with the same variables as in the GLM was fitted and multiplicative effects over the t-CFRs (i.e. the exponentiated estimated parameters) associated with each variable were calculated. Maps associated with these effects were obtained, presenting only those associated with the explanatory variables that were significant in the global model. All statistical analyses were conducted using R version 3.6.2 through the *spdep*, *rgdal*, and *spgwr* packages for spatial analyses and *car* package for the correlation analysis and GeoDa 1.14.0 was used also for some spatial analyses. The significance level for all analyses was 5% (i.e., $\alpha=0.05$).

RESULTS

t-CFR description and spatial autocorrelation of COVID-19 t-CFRs between states

Maps for quartiles corresponding to the raw and smoothed t-CFRs are similar, except for two states, Tlaxcala, which from a category using the raw risks moves to the next superior

one in the smoothed risks, the opposite occurring for Veracruz. Thus, only the map concerning the smoothed values is shown (**Figure 1A**). We observed the largest t-CFRs (above 4%) in Quintana Roo, Baja California, Chihuahua, and Tabasco, and the greatest relative risks (2.00-4.00) in Baja California, Chihuahua, Quintana Roo, and Tabasco (**Figure 1B**). Globally, there is a non-significant spatial autocorrelation (Moran's $I=-0.079$, $p=0.390$); however, there is a noticeable Low-Low cluster on the northeast around San Luis Potosi and two Low-High clusters around Yucatan and Sonora (**Figure 2**). These Low-High clusters make sense since the associated states, especially Sonora, are surrounded by the states with the highest t-CFRs in all the country. We verified that the spatial autocorrelation value and clustering were exactly the same using both the raw values or those obtained with the Bayes spatial technique.

Figure 1. Maps associated with COVID-19 deaths in Mexico by state until April 21st, 2020. A) Quartiles corresponding to tested case-fatality risks smoothed through an empirical Bayes procedure. B) Relative risk.

Figure 2. Spatial clustering associated with tested case-fatality risks of COVID-19 in Mexico by state until April 21st, 2020, considering queen contiguity. A) Significant spatial clustering obtained through Local Indicators of Spatial Autocorrelation (LISA) comparisons. There are four types of clusters: High-High, Low-Low, High-Low, and Low-High, e.g. a Low-Low cluster (blue) indicates states with low values of a variable significantly surrounded by regions with similarly low values. B) P-values associated with the spatial clustering in A), C) Scatter plot associated with the smoothed risks vs. their corresponding spatially lagged values, including the associated linear regression fitting, whose slope is the Moran's I statistic.

Fit of multivariable generalized global and geographical linear models for COVID-19 tested case-fatality risks

Through a preliminary analysis obtained by fitting a Poisson multivariable linear model with offset and including all variables, we found the presence of serious multicollinearity problems, since we obtained Variance Inflation Factors (VIF) with values above 50 for some variables. A correlation analysis between all variables was performed (**Figure 3A**), including the response (raw t-CFRs) identifying very correlated variables. Thus, we followed the model selection process as explained above. Our final global model included as inputs: diabetes, obesity, GDP, internal and external migration, age group of 10 to 39 years, physicians-to-population ratio, cardiovascular disease, ICU, hospitalization, and intubated. The latter three variables are confounders and to eliminate effects of other variables on them, they are used as residuals associated with appropriate Poisson models with independent variables: diabetes, obesity, age, physicians-to-population ratio, cardiovascular disease, plus ICU for the model associated with the intubated variable. Goodness-of-fit in our final model was assessed, finding that all variables were jointly significant (LR=472.19, p-value<0.001). Additionally, through a PP-plot associated with the standardized residuals and associated Anderson-Darling test (A=0.260, p-value = 0.689) and scatter plots between the fitted values and standardized residuals, and a similar plot using the root of the residuals instead, we determined that the link function and the way the explanatory variables are related with the response seems correct. However, we found some overdispersion (Chi-squared statistic divided by its degrees of freedom of 1.972). There were not any significant pairwise interaction effects.

A multivariable GGWR with a Poisson distribution, adaptive kernel, and the same input variables was also fitted. In **Table 2**, we present a summary of the multiplicative effects over the risks, exponentiated parameters, i.e. minimum, quartiles, and maximum, associated with each variable for all states in the GGWR and the global values

corresponding to the GLM. Pearson residuals associated with the GGWR are shown in **Figure 3B**, showing that the worst fit corresponded to two states: Veracruz and Yucatan.

Figure 3. Figures associated with the selection and goodness-of-fit of a multivariable generalized geographically weighted model (GGWR); with a Poisson distribution, offset, and a logarithm link function, used to explain tested case-fatality risks. A) Correlation plot including the raw risks, and B) Representation of Pearson residuals by state.

Table 2: Statistics by variable (minimum, maximum, and quartiles) associated with the multiplicative effects by state over the tested case-fatality risks under the GGWR and analogous effects and p-values associated with a global model (all models consider a Poisson distribution, offset term, and logarithmic link function).

Variable	Min	1st quartile	Median	3rd quartile	Max	Global	
						Effect	p-value
Intercept	0.004	0.005	0.005	0.005	0.005	0.005	<0.001
Diabetes	1.147	1.148	1.150	1.151	1.154	1.152	<0.001
Obesity	1.121	1.122	1.123	1.125	1.125	1.122	<0.001
ICU (residual)	1.219	1.220	1.220	1.222	1.225	1.223	<0.001
GDP (modified)	1.298	1.298	1.298	1.300	1.307	1.304	<0.001
Internal migration	1.250	1.252	1.262	1.268	1.276	1.267	<0.001
External migration	1.322	1.323	1.325	1.332	1.361	1.349	0.018
Age group 10-39	0.907	0.907	0.908	0.909	0.910	0.908	<0.001
Intubated (residual)	1.214	1.216	1.219	1.221	1.222	1.219	<0.001
Physicians' ratio	1.845	1.853	1.856	1.859	1.865	1.866	<0.001
Cardiovascular	0.948	0.951	0.953	0.954	0.955	0.951	0.478
Hospitalization (residual)	0.957	0.957	0.957	0.959	0.960	0.958	0.009

Predictors of COVID-19 spatial lethality in Mexico

The exponentiated estimated parameters under the GGWR for each state were obtained for each variable (not shown) and maps were obtained based on the .shp file provided in

(24). Nevertheless, we only present the maps for those explanatory variables that significantly impacted over the log-transformed t-CFRs in the global model (GLM) (**Figure 4**). These significant variables were diabetes, obesity, GDP, internal and external migration, age group of 10 to 39 years, and physicians-to-population ratio. However, care should be taken when the map for GDP is interpreted considering this variable corresponds to GDP on states not having the biggest cities, as an interaction term between GDP and a binary variable, thus having a fixed value of zero in four states, which are represented as blank spaces in the map. All estimated terms were interpreted by considering fixed values for all variables except the one being interpreted.

Figure 4. Multiplicative estimated effects over the tested case-fatality risks due to COVID-19 under a GGWR for those variables that significantly impact the response under a global model (Poisson models with offset and a logarithmic link function).

Age and metabolic predictors of t-CFRs in Mexico

Prevalence of obesity has a global (for all states) significant positive association (multiplicative effect) with the COVID-19 t-CFRs (1.122; 95%CI 1.081-1.166), locally the effect is between 1.120 in Quintana Roo to 1.125 in Sinaloa, having a similar effect on all Mexico, though slightly larger on the north and center. For diabetes, there is also a significant positive association with the t-CFRs (1.152; 95%CI 1.079-1.230) with local effects between 1.147 in Colima to 1.154 in Yucatan, having a positive effect in all Mexico; but specially in the center, south, and Yucatan peninsula. On the other hand, the proportion of individuals between 10 and 39 years old has a significant negative association with the COVID-19 t-CFRs (0.907; 95%CI 0.873-0.944), locally the effect is between 0.907 in Chiapas to 0.910, thus having a similar association in all the country.

Mobility and socio-economic predictors of t-CFRs in Mexico

The percentage of internal migration in the spatial unit a patient comes from has a significant positive association with the COVID-19 t-CFRs (1.267; 95%CI 1.183-1.356), locally the effect is between 1.250 in Durango to 1.276 in Quintana Roo having more effect in the south, center, and Yucatan peninsula. The percentage of external migration in the spatial unit a patient comes from is also significantly positively associated with the COVID-19 t-CFRs (1.349; 95%CI 1.052-1.728), locally the effect is between 1.322 in Morelos to 1.361 in Baja California, this effect exists over all Mexico, but it is stronger in the North and Baja California peninsula. Physicians-to-population ratio is also significantly positively associated with the t-CFRs (1.866; 95%CI 1.625-2.142), with local effects between 1.845 in Colima to 1.865 in Yucatan, having a slightly larger effect on the north and Yucatan peninsula. Finally, GDP excluding the states having the biggest cities in the country (Nuevo León, México, Ciudad de México, and Jalisco) is also significantly positively associated with the t-CFRs (1.304; 95%CI 1.198-1.429), locally the effects are between 1.298 in Oaxaca and 1.307 in Baja California.

DISCUSSION

Here, we show the relevance of spatial analyses to allow us not only to understand how t-CFRs are distributed along Mexico and the presence of spatial clusters related to COVID-19 t-CFRs, but also how some variables are associated with these risks, with an association that varies along all territory. Through our analysis, we were able to identify spatial units or regions in which care could have been considered to avoid SARS-CoV-2 spread and its related adverse outcomes. Identifying these areas, might allow to understand the propagation of the disease in other possible waves of SARS-CoV-2 spread or similar infectious pathogens.

Considering the global results, variables that are significantly associated with an increase on the COVID-19 t-CFRs include percentages associated with diabetes, obesity, external and internal migration, physicians-to-population ratio, and GDP in states that do not

include one of the greatest four cities in Mexico, whereas the proportion of individuals between 10 to 39 years old is significantly associated with a decrease on the risks. The association of cardio-metabolic diseases with adverse COVID-19 outcomes has been well documented and has been linked to mild, but sustained chronic inflammation, which may synergize with the cytokine storm associated with severe SARS-CoV-2 infection (25). Regarding other type of factors, internal and external migration have a very strong association with an increase of COVID-19 mortality, being external migration the variable with the second highest impact. This phenomenon is very particular for infectious diseases where greater movements of people move a disease from a geographical zone to another, which has occurred for instance for *Mycobacterium tuberculosis* or HIV/AIDS (26–28). The other two variables are related to urbanization, density, and economic importance of the states, specially physicians-to-population ratio. From the correlation plot, it can be seen that there is a very large positive association between physicians-to-population ratio and population density and overcrowded households, and a large negative association between this variable and both the rural and poverty proportions by state. On the other hand, the positive association between GDP and the fatality risks might also be related with states in which, due to the importance of economic activity, mobility could not be stopped after self-isolation measures, but that also do not have the health infrastructure that those states with the biggest cities have, in which, in spite of having many cases, there were less deaths.

The greatest t-CFRs, both raw and smoothed, corresponded to the states of Chihuahua, Quintana Roo, Baja California, and Tabasco (raw values of 5.11%, 4.79%, 4.34%, and 4.25%, respectively). It is noticeable how the relative risks in all these states have values between 2.5 and 3, suggesting that in these states the risks are above of what is nationally expected. We observed the presence of a spatial cluster concerning states with low risks;

however, at least till the date associated with our dataset, there was no important clustering of states with high risks using the LISA technique.

The fact that Quintana Roo is an important touristic center explains the presence of COVID-19 cases. However, according to our models, the elevated t-CFR is mostly associated with the presence of diabetes and internal migration, besides of the physicians-to-population ratio, whose possible interpretation was discussed above. In Chihuahua and Baja California, the variables with a particular positive association with t-CFR were external migration, obesity, and GDP, whereas in Tabasco these variables correspond to diabetes and internal migration.

Globally, physicians-to-population ratio, which is heavily associated with more urbanity, overcrowding households and population density, and with less poverty, has the highest positive effect on the t-CFRs, though relatively similar on all Mexico. External migration has the second highest association with the t-CFRs particularly in those states in the north in which the risks were the highest, whereas internal migration has the fourth highest association, after GDP, with particular importance in the center, south, and Yucatan peninsula. The effect of obesity and diabetes over the risks is similar; for obesity, there is relatively a similar impact on all the country, whereas for diabetes there is a little higher effect on the center, south, and Yucatan peninsula. Age between 10 and 39 years old was the only variable associated with a less risks, which agrees with previous results analyzed using Mexican data, having a relatively similar effect on all the country (4,10).

Economic effects, such as poverty, may be better studied in a disaggregated model including it as a measure at an individual level. Unfortunately, such information is unavailable in the epidemiological data set, and at most, information concerning poverty at a municipality level could be attached to each individual, being state the spatial unit containing municipalities. However; we would still be using aggregated values and in the methodology used to calculate poverty in Mexico, the state values are estimators obtained

from a representative sample, whereas the municipality values are estimated through small area estimation techniques; thus, being the former more reliable. In this sense, all analyses were performed at a state instead of at a municipal level. The reason behind this decision is that there were a lot of municipalities with zero values in the early time point we chose to focus our analysis, in both the number of COVID-19 cases and mortality and modelling such information with the probability distributions available for GGWR and other spatial linear models would not be possible. To obtain similar results we would require different tools, as zero-inflated geographical models. This could be possible future research work, including the identification of what are the characteristics of those municipalities with zero cases of the disease or no mortality. Additional future work corresponds also to analyze the data from a spatio-temporal framework, thus, the data set could be updated and the risks and associated factors followed through time for each state.

Our results are robust in terms of the model since it fits the data well and most of the statistical assumptions were satisfied; and, though there is some overdispersion, after fitting a quasi-Poisson linear model, we obtained the same results except that external migration was now significant at a 0.1 level. In terms of confounders, we fitted models with these variables as they are, as residuals, and as the first component in a Principal Component Analysis, finding in all cases that obesity, GDP, internal migration, age group of 10 to 39 years, and physicians-to-population ratio were significant.

In terms of the response, we obtained risks with the projected population in 2020 as denominator (population-fatality or mortality risks), and obtained similar results (maps) as the ones we present here, in terms of which states are on the highest quantiles, and in terms of similar clustering (29). Of notice is that using the projected values, we found a High-High cluster we did not find before in the Baja California peninsula. In fact, through the use of the Kulldorff's spatial scan statistic (30), we also identified a cluster there,

including three additional states (Sonora, Chihuahua, and Sinaloa); thus, it seems that a cluster in the north of higher risks exists, though it was not identified using the LISA methodology. Linear models using the projected values equivalent to the ones we fitted provided also similar results. For instance, in a model including the same variables as in our final model, we found the same variables significantly associated with the risks and direction of the association, except for diabetes, which was no longer significant. Analyses at an individual level were also fitted, using generalized (logistic) linear mixed models with a random intercept for state, and including mostly health related variables since socio-economic features are only at a state-level. We found that hospitalized status should be used separately from ICU and intubated, and that after taking care of multicollinearity, the results were similar as the ones at a state-level, in terms that variables significantly associated with the risks were still obesity, diabetes, and age group of 10 to 39 years, plus hypertension and cardiovascular.

It is important to notice that we are studying fatality risks associated with those individuals tested for the disease (t-CFRs); thus, care should be taken if results want to be extrapolated. As mentioned above, by using the projected population in 2020, instead of the tested individuals to model the mortality (population-fatality) risks, we obtained similar significant associations between the variables and risks, except for diabetes. However, we think this analysis is somewhat inaccurate in the sense that all health-related variables correspond to prevalence in individuals in the data set, which do not necessarily agree with those in the population, the same for age, and if we used population values by state, we would waste all the information in the epidemiological data, except for the number of deaths. Another option could have been to study case-fatality risks, thus considering the number of deaths and people infected; however, a similar problem arises, the real number of people with the disease is much larger than those being tested and results cannot be extrapolated, and this type of analysis is out of the scope of this study. Additionally, in any

analysis, the number of infected and/or deaths are even more poorly estimated when analyzing the early spread of the disease, particularly in Mexico, in which not enough population had been tested. Despite these limitations, we were able to identify some spatial predictors of fatality risks associated with COVID-19 at an early stage of the pandemic, likely reflecting factors which could have been addressed to mitigate SARS-CoV-2 spread.

In conclusion, metabolic diseases, internal and external migration, physicians-to-population ratio, GDP per capita in states without the biggest cities, and age group between 10 to 39 years old were significantly associated with early COVID-19 fatality risks in Mexico. These predictors likely influence the growth of the pandemic moving forward, but variables as prevalence of metabolic diseases cannot be easily modified in the short-term. However, the identification of important variables in Mexico associated with the risks and in specific geographical areas, could help to decide necessary public policies which could have long-term impacts on future epidemic scenarios. Even though, this is an analysis in an early stage of SARS-CoV-2 spread, it allows us to understand how the pandemic evolved within Mexico and the possible measures that should be addressed for additional waves or similar diseases in Mexico and in specific zones of the country.

ORCID

Ricardo Ramírez-Aldana: 0000-0003-4344-2928

Juan Carlos Gómez-Verján: 0000-0001-7186-8067

Omar Yaxmehen Bello-Chavolla: 0000-0003-3093-937X

Carmen García Peña: 0000-0002-9380-6964

Author contributions: Conceptualization, methodology, formal analysis, validation, and supervision: RRA. Software and data curation: OYBC and RRA. Investigation and

resources: JCGV and RRA. Analyses interpretation and writing (original draft, review, and editing): CGP, JCGV, OYBC, and RRA. Visualization: OYBC and RRA.

Conflict of Interest:

Nothing to disclose.

Funding:

This project was supported by a grant from the Secretaría de Educación, Ciencia, Tecnología e Innovación de la Ciudad de México CM-SECTEI/041/2020 “Red Colaborativa de Investigación Traslacional para el Envejecimiento Saludable de la Ciudad de México (RECITES)”

REFERENCES

1. Coronavirus Update (Live): 22,787,319 Cases and 795,367 Deaths from COVID-19 Virus Pandemic - Worldometer [Internet]. [cited 2020 Aug 20]. Available from: <https://www.worldometers.info/coronavirus/>
2. Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature*. 2020 Feb 3;579(7798):270–273.
3. Javelle E, Raoult D. COVID-19 pandemic more than a century after the Spanish flu. *Lancet Infect Dis*. 2020 Aug 11;
4. Bello-Chavolla OY, Bahena-López JP, Antonio-Villa NE, Vargas-Vázquez A, González-Díaz A, Márquez-Salinas A, et al. Predicting Mortality Due to SARS-CoV-2: A Mechanistic Score Relating Obesity and Diabetes to COVID-19 Outcomes in Mexico. *J Clin Endocrinol Metab*. 2020 Aug 1;105(8).
5. McGonagle D, Sharif K, O'Regan A, Bridgewood C. The Role of Cytokines including Interleukin-6 in COVID-19 induced Pneumonia and Macrophage Activation Syndrome-Like Disease. *Autoimmun Rev*. 2020 Jun;19(6):102537.
6. Bello-Chavolla OY, Rojas-Martinez R, Aguilar-Salinas CA, Hernández-Avila M. Epidemiology of diabetes mellitus in Mexico. *Nutr Rev*. 2017;75(suppl 1):4–12.
7. Warning For The U.S.: Diabetes And Coronavirus Cocktail Killing Young People In Mexico [Internet]. [cited 2020 Aug 20]. Available from: <https://www.forbes.com/sites/andrewwright/2020/05/24/warning-for-the-us-diabetes-and-coronavirus-cocktail-killing-young-people-in-mexico/#7f3e096748fa>
8. Zhou Y, Xu R, Hu D, Yue Y, Li Q, Xia J. Effects of human mobility restrictions on the spread of COVID-19 in Shenzhen, China: a modelling study using mobile phone data. *The Lancet Digital Health*. 2020 Aug;2(8):e417–e424.

- 526 9. Cruz-Pacheco G, Bustamante-Castañeda JF, Caputo JG, Jiménez-Corona ME,
527 Ponce-de-León-Rosales S. Dispersion of a new coronavirus sars-cov-2 by airlines in
528 2020: temporal estimates of the outbreak in mexico. *Rev Invest Clin*.
529 2020;72(3):138–143.
- 530 10. Bello-Chavolla OY, González-Díaz A, Antonio-Villa NE, Fermín-Martínez CA,
531 Márquez-Salinas A, Vargas-Vázquez A, et al. Unequal impact of structural health
532 determinants and comorbidity on COVID-19 severity and lethality in older Mexican
533 adults: Considerations beyond chronological aging. *J Gerontol A, Biol Sci Med Sci*.
534 2020 Jun 29;
- 535 11. Moroni L, Bianchi I, Lleo A. Geoepidemiology, gender and autoimmune disease.
536 *Autoimmun Rev*. 2012 May;11(6-7):A386–92.
- 537 12. Restrepo-Jiménez P, Molano-González N, Anaya J-M. Geoepidemiology of
538 Sjögren's syndrome in Latin America. *Joint Bone Spine*. 2019 Oct;86(5):620–626.
- 539 13. Chen WM, Zhou QR, Wang XM, Mao HS, Wang P, Zhou L, et al. [Integration of
540 spatial epidemiology and molecular epidemiology used for study on tuberculosis].
541 *Zhonghua Liu Xing Bing Xue Za Zhi*. 2016 Dec 10;37(12):1683–1686.
- 542 14. Datos Abiertos - Dirección General de Epidemiología | Secretaría de Salud |
543 Gobierno | gob.mx [Internet]. [cited 2020 Apr 18]. Available from:
544 <https://www.gob.mx/salud/documentos/datos-abiertos-152127?idiom=es>
- 545 15. Metodología para la medición de pobreza en México | CONEVAL [Internet]. [cited
546 2020 Aug 20]. Available from:
547 <https://www.coneval.org.mx/Medicion/MP/Paginas/Metodologia.aspx#:~:text=Metodo log%C3%ADa%20para%20la%20medici%C3%B3n%20multidimensional%20de%20la%20pobreza%20en%20M%C3%A9xico.&text=De%20esta%20forma%2C%20el%20Consejo,y%20medici%C3%B3n%20de%20la%20pobreza>.
- 551 16. Secretaria de Comunicaciones y Transportes: 5.5 Estadística Operacional de
552 Aeropuertos / Statistics by Airport [Internet]. [cited 2020 Nov 18]. Available from:
553 <http://www.sct.gob.mx/transporte-y-medicina-preventiva/aeronautica-civil/5-estadistica/55-estadistica-operacional-de-aeropuertos-statistics-by-airport/>
- 555 17. Bello-Chavolla OY, Antonio-Villa NE, Vargas-Vázquez A, Fermín-Martínez CA,
556 Márquez-Salinas A, Bahena-López JP. Profiling cases with non-respiratory
557 symptoms and asymptomatic SARS-CoV-2 infections in Mexico City. *Clin Infect Dis*.
558 2020 Aug 28;
- 559 18. Rate Transformations and Smoothing Luc Anselin | Semantic Scholar [Internet].
560 [cited 2020 Apr 17]. Available from: <https://www.semanticscholar.org/paper/Rate-Transformations-and-Smoothing-Luc-Anselin-Lozano-Koschinsky/88d8b02de84f97f556cfe0ef5a91a7df229cf363#paper-header>
- 563 19. Cressie, Noel. *Statistics For Spatial Data* (wiley Classics Library). Revised. Hoboken,
564 NJ: Wiley-interscience; 1993.
- 565 20. Moran PAP. Notes on continuous stochastic phenomena. *Biometrika*. 1950
566 Jun;37(1/2):17.

- 567 21. Anselin L. Local Indicators of Spatial Association-LISA. *Geogr Anal.* 2010 Sep
568 3;27(2):93–115.
- 569 22. Fotheringham, A. S., Brunsdon, C., and Charlton, M. E. (2002). Geographically
570 weighted regression: the analysis of spatially varying relationships. Chichester, West
571 Sussex: John Wiley and Sons.
- 572
573 23. Agresti A. Categorical Data Analysis. 3rd ed. Hoboken, NJ: Wiley; 2012.
- 574 24. Porto Tapiquén CE. Orogénesis Soluciones Geográficas. Porlamar, Venezuela 2015.
575 Estados de México". Available from <http://tapiquen-sig.jimdo.com>.
- 576 25. Bornstein SR, Dalan R, Hopkins D, Mingrone G, Boehm BO. Endocrine and
577 metabolic link to coronavirus infection. *Nat Rev Endocrinol.* 2020;16(6):297–298.
- 578 26. Pareek M, Greenaway C, Noori T, Munoz J, Zenner D. The impact of migration on
579 tuberculosis epidemiology and control in high-income countries: a review. *BMC Med.*
580 2016 Mar 23;14:48.
- 581 27. Weine SM, Kashuba AB. Labor migration and HIV risk: a systematic review of the
582 literature. *AIDS Behav.* 2012 Aug;16(6):1605–1621.
- 583 28. Gross B, Zheng Z, Liu S, Chen X, Sela A, Li J, et al. Spatio-temporal propagation of
584 COVID-19 pandemics. *medRxiv.* 2020 Mar 27;
- 585 29. Datos Abiertos de México - Proyecciones de la Población de México y de las
586 Entidades Federativas, ' ' 2016-2050 - Proyecciones de la Población de los
587 Municipios de México, 2015-2030 (base 2) [Internet]. [cited 2020 Nov 19]. Available
588 from: <https://datos.gob.mx/busca/dataset/proyecciones-de-la-poblacion-de-mexico-y-de-las-entidades-federativas-2016-2050/resource/ab8243f5-53b3-40bb-8784-2ad44231f642>
589
590
- 591 30. Kulldorff M. A spatial scan statistic. *Communications in Statistics - Theory and*
592 *Methods.* 1997 Jan;26(6):1481–1496.
- 593
594
595

A **B**

A **B** **C**

A

B

A **B** **C** **D** **E** **F** **G**