A meta-analysis on the role of pre-existing chronic disease in the cardiac complications of SARS-CoV-2 infection

Jane E. Sinclair¹, Yanshan Zhu¹, Gang Xu², Wei Ma^{3,4}, Haiyan Shi⁵, Kun-Long Ma⁶, Chun-Feng Cao⁶, Ling-Xi Kong⁶, Ke-Qiang Wan⁶, Juan Liao⁶, Hai-Qiang Wang⁷, Matt Arentz⁸, Meredith Redd¹², Linda A. Gallo¹³, & Kirsty R. Short^{1,14*}

¹ School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane, Australia

² Guangzhou First People's Hospital, Guangzhou, Guangdong, China

³ Department of Geriatric Medicine, Guangzhou First People's Hospital, Guangzhou, Guangdong, China

⁴ School of Medicine, South China University of Technology, Guangzhou, Guangdong, China

⁵ Department of Respiratory Medicine, Guangzhou Eighth People's Hospital, Guangzhou, Guangdong, China

⁶ Yongchuan Hospital of Chongqing Medical University, Yongchuan, Chongqing, China

⁷ Institute of Integrative Medicine, Shaanxi University of Chinese Medicine, Xi'an Shaanxi Province, China

⁸ Department of Global Health, University of Washington, Seattle, Washington, U.S.A.

¹² Institute for Molecular Bioscience, The University of Queensland, Brisbane, Australia

¹³ School of Biomedical Sciences, The University of Queensland, Brisbane, Australia

¹⁴ School of Public Health, The University of Queensland, Brisbane, Australia

* **Correspondence:** k.short@uq.edu.au; Tel.: +61 7 336 54226

JES: Bachelors

YZ: Masters

GX: MD

WM: MD

HS: MD

K-LM: MD

C-FC: MD

L-XK: MD

K-QW: MD

JL: MD

H-QW: MD/PhD

MA: MD

MR: PhD

LAG: PhD

KRS: PhD

30 SUMMARY

31 **Background:** Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has been associated with
 32 multiple direct and indirect cardiovascular complications. It is not yet well defined whether the patient groups at
 33 increased risk of SARS-CoV-2 severe respiratory disease experience similarly heightened incidence of
 34 cardiovascular complications. We sought to analyse the role of host co-morbidities (chronic respiratory
 35 illnesses, cardiovascular disease (CVD), hypertension or diabetes mellitus (DM)) in the acute cardiovascular
 36 complications associated with SARS-CoV-2 infection.

37 **Methods:** We retrospectively investigated published and pre-printed, publicly released, de-identified, data
 38 made available between Dec 1, 2019, and October 14, 2020. Information was accessed from PubMed, Embase,
 39 medRxiv and SSRN. 363 full-text articles were reviewed and 283 excluded for lack of original research,
 40 irrelevance to outcome, inappropriate cohort, or describing case reports of <10 patients. Data were extracted
 41 from 25 studies and the remaining 55 contacted to request appropriate data, to which four responded with data
 42 contributions. A final of 29 studies were included. This systematic review was conducted based on Preferred
 43 Reporting Items for Systematic Reviews (PRISMA) and Meta-analyses of Observational Studies in
 44 Epidemiology (MOOSE) statements. Included studies were critically appraised using Newcastle Ottawa Quality
 45 Assessment Scale (NOS). Data were extracted independently by multiple observers. When $I^2 \leq 50\%$, a
 46 fixed-effects model was selected, while a random-effects model was employed when $I^2 > 50\%$.
 47 Cardiovascular complications were measured as blood levels of cardiac biomarkers above the 99th percentile
 48 upper reference limit, new abnormalities in electrocardiography, and/or new abnormalities in echocardiography.

49 **Results:** Individual analyses of the majority of studies found that the median age was higher by approximately
 50 10 years in patients with cardiovascular complications. Pooled analyses showed the development of SARS-
 51 CoV-2 cardiovascular complications was significantly increased in patients with chronic respiratory illness
 52 (Odds Ratio (OR) 1.67[1.48,1.88]), CVD (OR 3.37[2.57,4.43]), hypertension (OR 2.68[2.11,3.41]), DM (OR
 53 1.60[1.31,1.95]) and male sex (OR 1.31[1.21,1.42]), findings that were mostly conserved during sub-analysis of
 54 studies stratified into four global geographic regions.

55 **Conclusions:** Age, chronic respiratory illness, CVD, hypertension, DM and male sex may represent prognostic
 56 factors for the development of both acute and chronic cardiovascular complications in COVID-19 disease,
 57 highlighting the need for a multidisciplinary approach to chronic disease patient management.

58

59

60

61

62 INTRODUCTION

63 Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged at the end of 2019 and has since gone
 64 on to cause the ongoing coronavirus disease 2019 (COVID-19) pandemic, infecting >50 million people
 65 worldwide. SARS-CoV-2 causes a broad range of respiratory symptoms, ranging from subclinical disease to
 66 viral pneumonia [1]. However, it is increasingly apparent that SARS-CoV-2 also causes a variety of extra-
 67 respiratory complications [1]. Specifically, SARS-CoV-2 infection has been associated with acute myocardial
 68 injury, myocarditis, arrhythmias, venous thromboembolism and other cardiac complications [2–6]. The
 69 development of these complications is likely enhanced by experimental testing of antimalarial and antiviral
 70 drugs, which have known cardiovascular side effects [4–12]. Although prevalence estimates vary, data from
 71 China suggests that approximately 7% of patients infected with SARS-CoV-2 develop cardiac injury (defined as
 72 elevated high-sensitivity cardiac troponin I or new echocardiographic; electrocardiographic (ECG)
 73 abnormalities) [13]. This prevalence increases to 25% in those hospitalised with COVID-19 [14], and 22% in
 74 patients who required intensive care unit (ICU) admission [13]. However, at present, the role of pre-existing
 75 chronic disease in the cardiac complications of COVID-19 remains to be elucidated.

76
 77 Pre-existing chronic respiratory disease, cardiovascular disease (CVD), hypertension and diabetes mellitus
 78 (DM) have each been associated with increased rates of hospitalization, ICU admission, and death from the
 79 respiratory complications of SARS-CoV-2 [13–27]. It is not yet well defined whether these patient groups are
 80 also at an increased risk of SARS-CoV-2 associated cardiac injury. Some small studies have shown that patients
 81 with pre-existing CVD are at a higher risk of myocarditis and/or myocardial infarction (MI) from SARS-CoV-2
 82 [3, 28, 29]. A retrospective, single-centre case series of 187 patients with COVID-19 also found that pre-
 83 existing CVD increased the risk of cardiac injury (as defined by troponin elevation) (54.5% vs 13.2%) and
 84 mortality compared with patients without CVD [3]. Similarly, a meta-analysis of 28 observational studies found
 85 SARS-CoV-2 associated cardiac injury biomarkers to be related to a history of hypertension but not DM [30]. In
 86 contrast, Zhu and colleagues found that individuals with type 2 DM had a greater occurrence of acute heart
 87 injury (7.3% versus 3.0%) than individuals without DM [31].

88 There is a clear need to better understand the role of patient host factors in the extra-respiratory complications of
 89 SARS-CoV-2 infection. Here, we analysed the currently available literature on the role of pre-existing chronic
 90 respiratory illnesses, CVD, hypertension and DM in the prevalence of SARS-CoV-2 associated cardiac injury.

91

92

93

94

95

96

97

98

99

METHODS

Study selection

We retrospectively investigated published and pre-printed, publicly released and de-identified data made available between Dec 1, 2019, and October 14, 2020. Information was accessed from four databases (PubMed, Embase, medRxiv and SSRN) using the following search terms: ("covid-19"[All Fields]) OR ("SARS-CoV-2"[All Fields]) AND ("cardiac"[All Fields] AND "comorbid*"[All Fields]). A total of 1076 articles were retrieved after duplicates were removed (available in either English or Chinese), following which 713 records were removed after screening by title and abstract. 363 full-text articles were reviewed. Of these, 283 articles were excluded because i) they did not include original research, ii) they were not relevant to the outcome, iii) they described an inappropriate patient population and/or iv) they described case reports of <10 patients. Of the remaining 80 studies that were deemed eligible, due to their containing data on patient pre-existing disease and cardiac complications as an outcome, 25 contained the required data format and these were extracted. The authors of the remaining 55 studies were contacted to request appropriate data (as they presented only aggregate data that did not specify the cross-over between pre-existing chronic disease and outcome), to which four groups responded with data contributions. Thus, a final of 29 studies were included in the meta-analysis [3,4,29,32-57]. The selection process is shown in Figure 1.

Data extraction and analysis

Extracted variables were aggregate data of median age, sex, underlying chronic disease (chronic respiratory illness, CVD, hypertension and DM) and the occurrence of cardiac complications as an outcome. Patients with co-morbidities were included in all applicable categories. For cases in which the data source reported more than one type of co-morbid CVD separately (such as prevalence of both chronic heart failure and coronary heart disease within their cohort), only the group with the largest number of cases was included to avoid patient double up and overrepresentation of co-morbid CVD. Analyses for each co-morbidity were performed independently, with each assuming one effect (the patient group in question) is considered as the main driver of effects. Median data for age were analysed individually using a Mann-Whitney U test. R version 4.0.0 software was used to analyse sex distribution as well as pre-existing chronic disease and cardiac outcomes through meta-analysis. Heterogeneity among studies was assessed by using the Q statistic with $*P \leq 0.05$ indicating the presence of significant heterogeneity and quantified by using the I^2 statistic. A fixed-effects model was used as when $I^2 < 50\%$ and random-effects model when $I^2 \geq 50\%$.

Definition of cardiac complications

Cardiac complications were defined as one or more of the following: blood levels of cardiac biomarkers (troponin I or creatine-kinase myocardial band) above the 99th percentile upper reference limit [58–60], and/or abnormalities observed in clinical assessments of the patients' heart function, including ECG [61–65] or echocardiography [62–66].

Quality assessment and data synthesis

The included studies were critically appraised using Newcastle Ottawa Quality Assessment Scale (NOS) [67]. High-quality studies were defined as studies fulfilling NOS score of minimum 7. Data were synthesised from 29 different and high-quality studies [3,4,29,32–57]. This systematic review was conducted based on the Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) [68] and Meta-analysis of Observational Studies in Epidemiology (MOOSE) [69] statements.

RESULTS

SARS-CoV-2 patients with pre-existing chronic illnesses are at increased risk of acute cardiac complications

We identified 1076 articles that described pre-existing chronic disease and cardiac complications of SARS-CoV-2, rejected 996 articles due to a lack of sufficient and/or appropriate data and derived a total of 80 articles. 29 of these studies were subsequently used in a meta-analysis (Figure 1), all of which were high quality (Table 1). Baseline characteristics of the patient populations as well as other relevant data are listed in Table 2. The definition of cardiac complications varied between studies, ranging from acute cardiac injury to fulminant myocarditis. 20 studies measured cardiac outcomes on the day of admission [3,29,32,34,35,37–41,43–48,50–52,57], and one at the time of patient death [33], with the remaining eight studies not specifying the time of assessment [4,36,42,49,53–55]. The studies were conducted in 10 countries in four geographical locations (U.S.A. [32,33,41,48,49], the Middle East [34,50,52], Europe [4,35–40,44,45,47], and the People’s Republic of China [3,29,42,43,46,51,53–57]) with a total of 3553 COVID-19 patients developing cardiac complications and 13,225 COVID-19 patients without cardiac complications. Patients who presented with SARS-CoV-2 associated cardiac complications were, on average, 10 years older than in those who did not (Table 2).

Figure 1. The Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) flowchart of study selection.

Table 1. Quality assessment of the included studies using the Newcastle Ottawa Scale.

Study	Selection	Comparability	Outcome	Total
Abrams et al., 2020 ³²	****	--	***	7
Arentz et al., 2020 ³³	****	--	***	7
Barman et al., 2020 ³⁴	****	--	***	7
Bertini et al., 2020 ³⁵	****	--	***	7
CAPACITY COVID, 2020 ³⁶	****	--	***	7
Duerr et al., 2020 ³⁷	****	--	***	7
Ferrante et al., 2020 ³⁸	****	--	***	7
Ghio et al., 2020 ³⁹	****	--	***	7
Guo et al., 2020 ³	****	--	***	7
Lairez et al., 2020 ⁴⁰	****	--	***	7
Lala et al., 2020 ⁴¹	****	--	***	7
Li et al., 2020 ⁴²	****	--	***	7
Liu et al., 2020 ⁴³	****	--	***	7
Lombardi et al., 2020 ⁴⁴	****	--	***	7
Lorente-Ros et al., 2020 ⁴⁵	****	--	***	7
Ma et al., 2020 ⁴⁶	****	--	***	7
Pagnesi et al., 2020 ⁴⁷	****	--	***	7
Raad et al., 2020 ⁴⁸	****	--	***	7
Sala et al., 2020 ⁴	****	--	***	7
Shah et al., 2020 ⁴⁹	****	--	***	7
Shahrokh et al., 2020 ⁵⁰	****	--	***	7
Shi et al., 2020 ²⁹	****	--	***	7
Song et al., 2020 ⁵¹	****	--	***	7
Szekely et al., 2020 ⁵²	****	--	***	7
Toraih et al., 2020 ⁵³	****	--	***	7
Wei et al., 2020 ⁵⁴	****	--	***	7
Xie et al., 2020 ⁵⁵	****	--	***	7
Xu G et al., 2020 ⁵⁶	****	--	***	7
Xu H et al., 2020 ⁵⁷	****	--	***	7

234 **Table 2.** Summary of baseline characteristics of the included studies.

Study	Population	Sample size (males)		Age (years)			CC description	Assessment	Time of assessment
		CC	No CC	CC	No CC	Findings			
Abrams et al., 2020 ³²	Mortally ill COVID-19 patients hospitalised in 3 New York City hospitals between March 1 and April 3, 2020	11 (9)	122 (74)	64 (58-72)	82 (71.8-89)	$p=0.0009$	Arrhythmic death	Last recorded cardiac rhythm	After death
Arentz et al., 2020 ³³	Critically ill COVID-19 patients admitted to Evergreen Hospital ICU, Washington State	15 (8)	6 (3)	74 (61-82)	69.5 (66-73)	$p=0.3699$	Cardio-myopathy	Radiologic and laboratory findings	Day of ICU admission and again on day 5
Barman et al., 2020 ³⁴	Consecutive COVID-19 patients hospitalised in three government hospitals in Istanbul, Turkey	150 (86)	457 (248)	68.5±13.4	56.5±15.2	$p<0.001$	Cardiac injury	Laboratory findings only	Day of hospital admission
Bertini et al., 2020 ³⁵	431 consecutive COVID-19 patients hospitalized in Italy between 10 March and 14 April 2020 who died or were treated with invasive mechanical ventilation.	24 (16)	86 (64)	71 (59-76)	68 (60-78)	$p=0.88$	Hs-TnI > 5x the upper reference limit	Laboratory findings only	Day of hospital admission
CAPACITY COVID, 2020 ³⁶	COVID-19 patients registered to the international CAPACITY-COVID collaborative consortium with inclusion current from 13 countries	752 (537)	4811 (3019)	71 (64-77)	66 (55-76)	$p<0.001$	Cardiac complication	According to criteria of the European Society of Cardiology	From hospital admission to discharge
Duerr et al., 2020 ³⁷	COVID-19 patients admitted to the University Hospital Bonn, Germany	9 (4)	10 (5)	NA	NA	NA	Pericardial effusion	Radiologic and laboratory findings	Day of hospital admission
Ferrante et al., 2020 ³⁸	Consecutive COVID-19 patients undergoing chest CT on admission to Humanitas Research Hospital in Milan, Italy	123 (95)	209 (142)	74.2 (67.8-80.1)	60.7 (51.4-70.5)	$p<0.001$	Myocardial injury	Radiologic and laboratory findings	Day of hospital admission
Ghio et al., 2020 ³⁹	Consecutive adult COVID-19 patients admitted to the Emergency Department of Fondazione IRCCS Policlinico San Matteo (Pavia, Italy) from February 21st to March 31st 2020	74 (51)	266 (188)	76.1 (69.7-82.5)	68.6 (56.4-76.4)	$p<0.001$	Myocardial injury	Laboratory findings only	Day of hospital admission
Guo et al., 2020 ³	Consecutive COVID-19 patients admitted to the Seventh Hospital of Wuhan City, China	52 (34)	135 (57)	71.4 (SD=9.43)	53.53 (SD=13.22)	$p<0.001$	Myocardial injury	Laboratory findings only	Day of hospital admission

Lairez et al., 2020 ⁴⁰	COVID-19 patients admitted to Rangueil Hospital of Toulouse University for dyspnea or chest pain from April 6, 2020, to May 1, 2020.	13 (11)	18 (16)	66±8	52±13	p=0.015	Myocardial injury	Radiologic and laboratory findings	Day of hospital admission
Lala et al., 2020 ⁴¹	COVID-19 patients admitted to five Mount Sinai Health System hospitals in New York City	530 (318)	2206 (1312)	Median=70.59%	Median=60-70%	NA	Myocardial injury	Laboratory findings only	Day of hospital admission
Li et al., 2020 ⁴²	Critical type COVID-19 patients randomly selected from Renmin Hospital of Wuhan University, China	34 (23)	48 (29)	>70%	>70%	p=0.003	Myocardial damage	Laboratory findings only	NA
Liu et al., 2020 ⁴³	COVID-19 patients from Shenzhen Third People's Hospital in Shenzhen, China	1 (1)	11 (7)	63	62 (36-65)	NA	Fulminant myocarditis, cardiac failure	Radiologic and laboratory findings	Day of hospital admission
Lombardi et al., 2020 ⁴⁴	Consecutive COVID-19 patients hospitalized in 13 Italian cardiology units from March 1 to April 9, 2020.	278 (201)	336 (234)	71.3 (12)	64.0 (13.6)	p<0.001	Myocardial injury	Laboratory findings only	Day of hospital admission
Lorente-Ros et al., 2020 ⁴⁵	Consecutive adult COVID-19 patients admitted to a large tertiary hospital in Madrid, Spain	148 (71)	559 (193)	78.7	63.4	Standardised difference = 114%	Myocardial injury	Laboratory findings only	Day of hospital admission
Ma et al., 2020 ⁴⁶	Adult patients hospitalised with COVID-19 in Yongchuan, Chongqing, China	45 (27)	39 (21)	51 (45.5-63.5)	45 (39-53)	p=0.026	NA	Radiologic and laboratory findings	Day of hospital admission
Pagnesi et al., 2020 ⁴⁷	Hospitalised non-ICU COVID-19 patients from a large tertiary centre in Milan, Italy	29 (20)	171 (111)	65 (55-76)	62 (54-74)	p=0.813	Right ventricular dysfunction	Radiologic and laboratory findings	Day of hospital admission
Raad et al., 2020 ⁴⁸	COVID-19 patients admitted to Henry Ford Health System in Southeast Michigan, USA, between March 9 and April 15, 2020.	390 (229)	630 (280)	70 (51-89)	59 (39-79)	NA	Cardiac injury	Laboratory findings only	Day of hospital admission
Sala et al., 2020 ⁴	Non-ICU COVID-19 patients admitted to seven COVID-19 units at a third-level hub centre in Milan, Italy	12 (NA)	116 (NA)	73±10	64±14	NA	Arrhythmias	Radiologic findings only	NA
Shah et al., 2020 ⁴⁹	Covid-19 patients admitted to 3 Phoebe Putney hospitals in Southwest Georgia, USA, between March 2 and June 7, 2020.	116 (62)	193 (70)	68.2±14.1	59.9±14.0	p<0.001	Cardiac injury	Laboratory findings only	NA
Shahrokh et al., 2020 ⁵⁰	COVID-19 patients presenting to the Sina Hospital emergency department in south Tehran, Iran between March and May 2020.	115 (69)	271 (167)	64.98±15.29	57.12±15.48	p<0.001	Cardiac injury	Radiologic and laboratory findings	Day of hospital admission
Shi et al.,	Consecutive severe COVID-19 inpatients of Renmin Hospital,	106	565	73	57	p<0.001	Myocardial	Laboratory	Day of hospital

2020 ²⁹	Wuhan University, China	(58)	(264)	(66-80)	(43-70)		injury	findings only	admission
Song et al., 2020 ⁵¹	Critically ill COVID-19 patients admitted to the ICU at Tongji Hospital, Wuhan, China	34 (24)	30 (18)	67.8±10.3	61.3±13.3	$p=0.033$	Myocardial injury	Laboratory findings only	Day of hospital admission
Szekely et al., 2020 ⁵²	Consecutive adult COVID-19 patients admitted to Tel Aviv Medical Center, Israel	68 (46)	32 (17)	69.8±16	65.9±20	$p=0.56$	Abnormal echocardiogram	Radiologic findings only	Day of admission
Toraih et al., 2020 ⁵³	COVID-19 patients included in studies exploring pre-existing CVD as COVID-19 risk factors, cardiac injury, ICU admission or mortality	202 (NA)	958 (NA)	NA	NA	NA	Cardiac injury	Laboratory findings only	NA
Wei et al., 2020 ⁵⁴	COVID-19 patients admitted to two treatment centres in Sichuan, China	16 (7)	85 (47)	67(61-80.5)	47(33-55)	$p<0.001$	Acute myocardial injury	Radiologic and laboratory findings	NA
Xie et al., 2020 ⁵⁵	Non-critically ill COVID-19 patients admitted to Tongji Hospital, Wuhanm China	102 (46)	517 (251)	68.9±11.7	56.4±13.6	$p<0.001$	Acute cardiac-related injury	Radiologic and laboratory findings	NA
Xu G et al., 2020 ⁵⁶	Adult patients with COVID-19 in Guangzhou, China	12 (6)	253 (116)	62.6±14.5	48.1±15.5	$p=0.002$	Acute cardiac injury	Radiologic and laboratory findings	NA
Xu H et al., 2020 ⁵⁷	COVID-19 patients consecutively hospitalized between January 2 and March 17, 2020, in the Public Health Clinical Center of Chengdu, China.	47 (23)	46 (23)	41 (31-51)	54 (40-65)	$p<0.001$	Cardiac marker abnormalities	Laboratory findings only	Day of hospital admission

CC: cardiac complications; ICU: intensive care unit; NA: data not available.

235

236

237

238

239

In a pooled analysis, chronic respiratory illness (odds ratio (OR): 1.67, 95% confidence interval (CI) [1.48,1.88]) (Fig. 2), CVD (OR: 3.37, 95% CI [2.57,4.43]) (Fig. 3), hypertension (OR: 2.68, 95% CI [2.11,3.41]) (Fig. 4), DM (OR: 1.60, 95% CI [1.31,1.95]) (Fig. 5), and male sex (OR: 1.31, 95% CI [1.21,1.42]) (Fig. 6), were each associated with a higher incidence of SARS-CoV-2 associated cardiac complications.

Figure 2: Forest plot of chronic respiratory illness as a risk factor for cardiac complications in COVID-19 patients. Fixed effect models apply when $I^2 < 50\%$.

Figure 3: Forest plot of pre-existing cardiovascular disease as a risk factor for cardiac complications in COVID-19 patients. Random effect models apply when $I^2 > 50\%$.

Figure 4: Forest plot of hypertension as a risk factor for cardiac complications in COVID-19 patients. Random effect models apply when $I^2 > 50\%$.

Figure 5: Forest plot of diabetes mellitus as a risk factor for cardiac complications in COVID-19 patients. Random effect models apply when $I^2 > 50\%$.

Figure 6: Forest plot of male sex as a risk factor for cardiac complications in COVID-19 patients. Fixed effect models apply when $I^2 < 50\%$.

As large heterogeneity was detected amongst the total combined study cohort, the cohort was subdivided based on general global region in an attempt to account for this. Cohorts were divided into studies with cohorts primarily situated in the U.S.A [32,33,41,48,49], the Middle East [34,50,52], Europe [4,35–40,44,45,47], and China [3,29,42,43,46,51,53–57].

Regionally-pooled sub-analysis of chronic respiratory illness found it to be associated with a higher incidence of SARS-CoV-2 associated cardiac complications in cohorts from the U.S.A. (OR: 1.64, 95% CI [1.28,2.09]) (Fig. 7, A), Europe (OR: 1.53, 95% CI [1.28,1.83]) (Fig. 7, C) and China (OR: 1.94, 95% CI [1.49,2.52]) (Fig. 7, D), but not the Middle East (Fig. 7, B). Regional subdivision accounted for the high heterogeneity amongst all but the Middle Eastern cohort.

Figure 7: Forest plots of chronic respiratory as a risk factor for cardiac complications in COVID-19 patients divided by geographical region. A | Studies based primarily in the United States of America. B | Studies based primarily around the Middle East. C | Studies based primarily in Europe. D | Studies based primarily in China. Fixed effect models apply when $I^2 < 50\%$ and random effects models apply when $I^2 > 50\%$.

Regionally-pooled sub-analysis of pre-existing CVD found its association with a higher incidence of SARS-CoV-2 associated cardiac complications to be conserved across all cohorts from the U.S.A. (OR: 3.55, 95% CI [2.51,5.04]) (Fig. 8, A), the Middle East (OR: 2.40, 95% CI [1.25,4.62]) (Fig. 8, B), Europe (OR: 2.99, 95% CI [1.90,4.69]) (Fig. 8, C), and China (OR: 5.52, 95% CI [3.88,7.85]) (Fig. 8, D), despite regional subdivision failing to account for high heterogeneity amongst the former three regions.

Figure 8: Forest plots of pre-existing cardiovascular disease as a risk factor for cardiac complications in COVID-19 patients divided by geographical region. A | Studies based primarily in the United States of America. B | Studies based primarily around the Middle East. C | Studies based primarily in Europe. D | Studies based primarily in China. Fixed effect models apply when $I^2 < 50\%$ and random effects models apply when $I^2 > 50\%$.

Regionally-pooled sub-analysis of hypertension found its association with a higher incidence of SARS-CoV-2 associated cardiac complications to be similarly conserved across all cohorts from the U.S.A. (OR: 2.23, 95% CI [1.22,4.08]) (Fig. 9, A), the Middle East (OR: 2.24, 95% CI [1.70,2.94]) (Fig. 9, B), Europe (OR: 2.38, 95% CI [1.46,3.88]) (Fig. 9, C), and China (OR: 3.94, 95% CI [3.10,5.01]) (Fig. 9, D), although regional subdivision could not account for high heterogeneity amongst the American and European cohorts.

Figure 9: Forest plots of hypertension as a risk factor for cardiac complications in COVID-19 patients divided by geographical region. A | Studies based primarily in the United States of America. B | Studies based primarily around the Middle East. C | Studies based primarily in Europe. D | Studies based primarily in China. Fixed effect models apply when $I^2 < 50\%$ and random effects models apply when $I^2 > 50\%$.

The association between DM and a higher incidence of SARS-CoV-2 associated cardiac complications was also conserved across all cohorts from the U.S.A. (OR: 1.54, 95% CI [1.32,1.78]) (Fig. 10, A), the Middle East (OR: 1.65, 95% CI [1.25,2.19]) (Fig. 10, B), Europe (OR: 1.50, 95% CI [1.07,2.09]) (Fig. 10, C), and China (OR: 2.04, 95% CI [1.13,3.70]) (Fig. 10, D), despite regional subdivision not accounting for high heterogeneity amongst the European and Chinese cohorts.

Figure 10: Forest plots of diabetes mellitus as a risk factor for cardiac complications in COVID-19 patients divided by geographical region. A | Studies based primarily in the United States of America. B | Studies based primarily around the Middle East. C | Studies based primarily in Europe. D | Studies based primarily in China. Fixed effect models apply when $I^2 < 50\%$ and random effects models apply when $I^2 > 50\%$.

Finally, regionally-pooled sub-analysis of male sex found it to be associated with a higher incidence of SARS-CoV-2 associated cardiac complications in cohorts from the U.S.A. (OR: 1.55, 95% CI [1.03,2.33]) (Fig. 11, A), and Europe (OR: 1.40, 95% CI [1.23,1.59]) (Fig. 11, C), but not the Middle East (Fig. 11, B) or China (Fig. 11, D). Regional subdivision accounted for the high heterogeneity amongst all but the American cohort.

Figure 11: Forest plots of male sex as a risk factor for cardiac complications in COVID-19 patients divided by geographical region. A | Studies based primarily in the United States of America. B | Studies based primarily around the Middle East. C | Studies based primarily in Europe. D | Studies based primarily in China. Fixed effect models apply when $I^2 < 50\%$ and random effects models apply when $I^2 > 50\%$.

DISCUSSION

Our meta-analysis shows that older age, male sex, and pre-existing chronic respiratory disease, CVD, hypertension and DM are all risk factors for the development of cardiac complications associated with SARS-CoV-2 infection. At present the reasons for each of these association are unclear; each is a respiratory risk factor and thus might cause an indirect cardiac effect, however it could equally be the case that they exert their own direct catalytic effect on cardiac complications.

Our analysis identified a significant association between pre-existing CVD and the risk of SARS-CoV-2 associated cardiac complications, which was conserved across global geographical regions. This is consistent with previous observations that 30% of patients with SARS-CoV-2 cardiac injury had a history of coronary heart disease [3,29]. The reason for this association remains unclear. These data may be the result of pre-existing CVD leading to an altered immune response, excessive cytokine release [5], endothelial dysfunction, increased procoagulant blood activity, rupturing of coronary atherosclerotic plaques [73], and the formation of an occlusive thrombi [74]. However, further investigations are required to validate this hypothesis.

The meta-analysis performed herein also found that chronic respiratory illness significantly increased the incidence of SARS-CoV-2 associated cardiac complications. It is possible that these data reflect the established indirect association between severe respiratory disease and subsequent acute cardiovascular event. Indeed, COVID-19 patients with pneumonia and acute respiratory distress syndrome (ARDS) display acute right ventricular hypertrophy despite an absence of pulmonary embolism or severe pulmonary hypertension [87]. However, chronic lung disease has not been consistently associated with SARS-CoV-2 ARDS and hypoxemia [88], and thus this is unlikely to be the sole explanation for the increased risk of cardiac complications in patients with chronic respiratory illness in the present analysis. SARS-CoV-2 associated cardiac complications

The association between underlying DM and cardiac complications from SARS-CoV-2 may be attributable to a multitude of other effects induced by abnormal circulating substrates, including hyperglycaemia, and/or the co-existence and complex interactions with other, even subclinical, conditions in this patient group [91]. In response to changes in substrate availability, for example, the diabetic heart becomes increasingly dependent on fatty acids as a fuel source, which decreases work efficiency, and dysregulates responses to hypoxia [92]. These metabolic changes may predispose individuals with DM to an increased risk of ischemic cardiac injury upon SARS-CoV-2 infection. The contribution of underlying DM to the risk of SARS-CoV-2 associated cardiac complications, however, depends largely on the duration of pre-existing disease [93].

There are several limitations in our review. Firstly, the limited number of studies that met the pre-defined selection criteria disallowed correction for other variables that may confound these results. For example, it is possible that our observations were confounded by the strong effect that age has on cardiovascular health. Thus, it is not possible to conclude if people with these pre-existing conditions are just more susceptible to severe COVID-19 and thus the heart complications associated with this increased severity, or if there is indeed a

cardiac-specific effect occurring. Similarly, we cannot exclude the possibility that the associations observed herein are due, at least in part, to treatments for severe COVID-19 that may damage the heart. Indeed, the experimental therapeutic hydroxychloroquine has been associated with QT interval prolongation, cardiomyopathy and heart failure [8–10]. This is likewise the case for treatments for underlying conditions, such as ACE inhibitors and ARBs which, while not known to damage the heart, may interfere with how the immune system targets the virus [78]. Information regarding which, if any, therapeutics used in these cohorts of patients was not consistently available. We anticipate that as more data on the cardiac complications become available it will be possible for future analyses to correct for these and other confounders; however, it would also be worthwhile testing the effects of each these pre-existing chronic conditions in one of the recently developed murine models of SARS-CoV-2.

Despite the above limitations, these data have important implications for patient care. Patients with one or more underlying medical conditions should be closely monitored for both severe respiratory and extra-respiratory COVID-19 complications. Furthermore, the exploratory use of medications that are independently associated with cardiac complications should be considered with caution in these patient groups and perhaps bypassed in favour of lower-risk treatment options. There is thus a clear need for a multidisciplinary, collaborative team to manage the complexity of COVID-19 illness observed in patients with one or more underlying chronic disease.

RESOURCE AVAILABILITY

Lead contact: Further information and requests for resources should be directed to and will be fulfilled by the Lead Contact, Dr. Kirsty Short (k.short@uq.edu.au).

Materials availability: This study did not generate new unique reagents.

Data and code availability: All data relevant to this study has been included in the manuscript. Data contributed by Xu et al., 2020³¹, Ma et al., 2020³², and Arentz et al., 2020³⁵, may be obtained by contacting authors of each respective study. Data retrieved from remaining studies are published and publicly available. R version 4.0.0 software was used to analyse sex distribution as well as pre-existing chronic disease and cardiac outcomes through meta-analysis. The datasets/code supporting the current study have not been deposited in a public repository but are available from the corresponding author on request.

DECLARATIONS

Acknowledgements: We want express our gratitude to the CAPACITY-COVID collaborative consortium for their data contribution:

Admiraal de Ruyter Hospital, Goes, the Netherlands: van Kesteren HAM
 Albert Schweitzer Hospital, Dordrecht, the Netherlands: van de Watering DJ
 Amphia Hospital, Breda, the Netherlands: Schaap J, Alings AMW
 Amstelland Hospital, Amstelveen, the Netherlands: Asselbergs-Koning AMH
 Amsterdam University Medical Center, Amsterdam, the Netherlands: Pinto Y, Tjong FVY
 Antonius Hospital, Sneek, the Netherlands: de Vries JK
 Antwerp University Hospital, Antwerp, Belgium: van Craenenbroeck EM, Kwakkel – van Erp HM, van Ierssel S
 AZ Maria Middelaars, Gent, Belgium: de Sutter J
 Barts Health NHS Trust, London, United Kingdom: Saxena M
 Beatrix Hospital, Gorinchem, the Netherlands: Westendorp PHM
 Bernhoven Hospital, Uden, the Netherlands: van de Wal RMA
 Bravis Hospital, Roosendaal, the Netherlands: Dorman HGR, van Boxem AJM
 Catharina Hospital, Eindhoven, the Netherlands: Tio RA
 CHU UCL Namur – Site Godinne, Yvoir, Belgium: Gabriel LG
 Deventer Hospital, Deventer, the Netherlands: Badings EA
 Diakonessenhuis, Utrecht, the Netherlands: van Ofwegen-Hanekamp CEE
 Dijklander Hospital, Hoorn, the Netherlands: Wierda E
 Dutch Network for Cardiovascular Research (WCN): Schut A
 Elizabeth-TweeSteden Hospital, Tilburg, the Netherlands: Monraats PS
 E.M.M.S Hospital, Nazareth, Israel: Hellou EH
 Erasmus University Medical Center, Rotterdam, the Netherlands: den Uil CA, Jewbali LS
 Franciscus Gasthuis, Rotterdam, the Netherlands: Nierop PR
 Franciscus Vlietland, Schiedam, the Netherlands: van der Linden MMJM
 Gelre Hospitals, Apeldoorn, the Netherlands: Groenemeijer BE
 Gelre Hospitals, Zutphen, the Netherlands: Al-Windy NYY
 Groene Hart Hospital, Gouda, the Netherlands: van Hessen MWJ
 Haaglanden Medical Center, den Haag, the Netherlands: van der Heijden DJ
 Hospital do Espírito Santo, Évora, Portugal: Ribeiro MIA
 Hospital Group Twente, Almelo, the Netherlands: Linszen GCM
 Hospital Prof. Doutor Fernando Fonesca, Amadora, Portugal: Ferreira JB
 Imperial College Healthcare NHS Trust, London, United Kingdom: Prasad S
 I.M. Sechenov First Moscow State Medical University, Moscow, Russia: Kopylov Ph. Yu., Blagova OV
 Ikazia Hospital, Rotterdam, the Netherlands: Emans ME
 Incliva Research Institute, University of Valencia, Valencia, Spain: Redón J, Forner MJ
 Isala, Zwolle, the Netherlands: Hermanides RS
 Jeroen Bosch Hospital, 's-Hertogenbosch, the Netherlands: Haerckens-Arends HE
 Jessa Hospital, Hasselt, the Netherlands: Timmermans P Jr, Messiaen P
 Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, The Netherlands: van Smeden M
 King Fahd Hospital of the University, Khobar, Saudi Arabia: Al-Ali AK, Al-Muhanna FA, Al-Rubaish AM, Almubarak YA, Alnafie AN, Alshahrani M, Alshehri A
 LangeLand Hospital, Zoetermeer, the Netherlands: van der Meer P
 Leeds Teaching Hospitals NHS Trust, Leeds, United Kingdom: Kearney M
 Leeuwarden Medical Center, Leeuwarden, the Netherlands: van den Brink FS
 Leiden University Medical Center, Leiden, the Netherlands: Siebelink HJ
 Maasstad Hospital, Rotterdam, the Netherlands: Smits PC
 Maastricht University Medical Center, Maastricht, the Netherlands: Heymans SRB
 Manchester University NHS Foundation Trust, Manchester, United Kingdom: Buch M
 Martini Hospital, Groningen, the Netherlands: Tieleman RG, Reidinga AC
 Meander Medical Center, Amersfoort, the Netherlands: Mosterd A
 Medisch Spectrum Twente, Enschede, the Netherlands: Meijs MFL, Delsing CE, van Veen HPAA
 Netherlands Heart Institute, Utrecht, the Netherlands: Hermans-van Ast W, Iperen EPA
 Northumbria Healthcare NHS Foundation Trust, Cramlington, United Kingdom: Aujayeb A
 One Day Surgery Hospital, Cairo, Egypt: Salah R
 Queen Elizabeth University Hospital, Glasgow, United Kingdom: Touyz R
 Rijnstate Hospital, Arnhem, the Netherlands: Seelig J
 Rode Kruis Hospital, Beverwijk, the Netherlands: Westendorp ICD, Veldhuis LI
 Royal Brompton and Harefield NHS Foundation Trust, London, United Kingdom: Prasad S
 Royal Devon and Exeter NHS Foundation Trust, Exeter, United Kingdom: Shore AC
 Royal Free London NHS Foundation Trust, London, United Kingdom: Captur G
 Salford Royal NHS Foundation Trust, Salford, United Kingdom: Dark P
 San Luigi Gonzaga University Hospital, Orbassano, Turin, Italy: Montagna LM, Bianco MB, Mazzilli SGM
 Saxenburgh Medical Center, Hardenberg, the Netherlands: Kaplan R, Drost JT
 Slingeland Hospital, Doetinchem, the Netherlands: Schellings DAAM

Southern Health and Social Care Trust, Portadown, United Kingdom: Moriarty A
 Spaarne Gasthuis, Haarlem, the Netherlands: Kuijper AFM
 SSR Val Rosay, Saint-Didier-au-Mont-d'Or, France: Charlotte NC
 St. Antonius Hospital, Nieuwegein, the Netherlands: ten Berg JM
 St. Jansdal, Harderwijk, the Netherlands: van der Zee PM
 Tehran Heart Center, Tehran, Iran: Shafiee A, Hedayat B, Poorhosseini H, Saneei E
 The Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle-upon-Tyne, United Kingdom: Zaman A
 Treant Zorggroep, Hogeveen, the Netherlands: Anthonio RL
 University College London Hospitals NHS Foundation Trust, London, United Kingdom: Williams B, Patel R, Bell R
 University Hospitals Bristol NHS Foundation Trust, Bristol, United Kingdom: Caputo M
 University Hospital Brussels, Brussels, Belgium: Weytjens C
 University Hospital Complex of Granada, Granada, Spain: Macías-Ruiz R
 University Hospital of Geneva, Geneva, Switzerland: Tessitore E
 University Hospitals of Leicester NHS Trust, Leicester, United Kingdom: McCann GP
 University Medical Center Groningen, Groningen, the Netherlands: van Gilst WH
 University Medical Center Utrecht, Utrecht, the Netherlands: Asselbergs FW, van der Harst P, Linschoten M
 Van Weel-Bethesda Hospital, Dirksland, the Netherlands: Wu KW
 Zaans Medical Center, Zaandam, the Netherlands: Verschure DO
 Zuyderland Medical Center, Heerlen, the Netherlands: Kietse-laer BLJH

Author contributions: Idea and study design: KRS, LAG; Data collection and analysis: JES, YZ, GX, WM, HS, K-LM, C-FC, L-XK, K-QW, JL, MA, KRS; Writing the article: JES, KRS; Draft revision: JES, GX, WM, HS, K-LM, C-FC, L-XK, K-QW, JL, MA, MR, LAG, KRS; All authors have read the manuscript and have approved this submission.

Declaration of interests: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Funding: KRS was supported by the Australian Research Council [DE180100512]. Collection of data contributed by Xu et al. was supported by the Guangzhou Health Research Project, the Guangzhou Health Committee (NO.20191A010014).

Ethics approval and consent to participate: The meta-analysis portion of this project was reviewed by the Human Ethics Research Office, The University of Queensland (CRICOS provider number: 00025B; application number: 2020001352) and deemed to be exempt from ethics review under the National Statement on Ethical Conduct in Human Research (National Statement §5.1.22).

REFERENCES

1. Clerkin KJ, Fried JA, Raikhelkar J et al. COVID-19 and cardiovascular disease. *Circulation*. 2020;141:1648-1655. doi:10.1161/CIRCULATIONAHA.120.046941.
2. Huang C, Wang Y, Li X et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497-506. doi:10.1016/S0140-6736(20)30183-5.
3. Guo T, Fan Y, Chen M et al. Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA Cardiol*. 2020; doi:10.1001/jamacardio.2020.1017.
4. Sala S, Peretto G, Gramegna M et al. Acute myocarditis presenting as a reverse Tako-Tsubo syndrome in a patient with SARS-CoV-2 respiratory infection. *Eur Heart J*. 2020;41(19):1861-1862. doi:10.1093/eurheartj/ehaa286.
5. Zheng Y, Ma Y, Zhang J, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol*. 2020;17:259-260. doi:10.1038/s41569-020-0360-5.
6. Inciardi RM, Lip L, Zacccone G et al. Cardiac involvement in a patient with coronavirus disease 2019 (COVID-19). *JAMA Cardiol*. 2020; doi:10.1001/jamacardio.2020.1096.

7. Edelson DP, Sasson C, Chan PS, et al. Interim guidance for basic and advanced life support in adults, children, and neonates with suspected or confirmed COVID-19: from the Emergency Cardiovascular Care Committee and Get With the Guidelines® -Resuscitation Adult and Pediatric Task Forces of the American Heart Association in collaboration with the American Academy of Pediatrics, American Association for Respiratory Care, American College of Emergency Physicians, the Society of Critical Care Anesthesiologists: supporting organisations: American Association of Critical Care Nurses and National EMS Physicians. *Circulation*. 2020; doi:10.1161/CIRCULATIONAHA.120.047463.
8. Page RL, 2nd, O'Bryant CL, Cheng D et al. Drugs that may cause or exacerbate heart failure: a scientific statement from the American Heart Association. *Circulation*. 2016;134(6):e32-69. doi:10.1161/cir.0000000000000426.
9. Guzik TJ, Mohiddin SA, Dimarco A et al. COVID-19 and the cardiovascular system: implications for risk assessment, diagnosis, and treatment options. *Cardiovasc Res*. 2020; doi:10.1093/cvr/cvaa106.
10. Chatre C, Roubille F, Vernhet H, Jorgensen C, Pers Y-M. Cardiac complications attributed to chloroquine and hydroxychloroquine: a systematic review of the literature. *Drug Saf*. 2018;41(10):919-931. doi:10.1007/s40264-018-0689-4.
11. Castagné B, Viprey M, Martin J, Schott A-M, Cucherat M, Soubrier M. Cardiovascular safety of tocilizumab: a systematic review and network meta-analysis. *PLoS ONE*. 2019; doi:10.1371/journal.pone.0220178.
12. Corrales-Medina VF, Alvarez KN, Weissfeld LA et al. Association between hospitalization for pneumonia and subsequent risk of cardiovascular disease. *JAMA*. 2015;313(3):264-274. doi:10.1001/jama.2014.18229.
13. Wang D, Hu B, Hu C et al. Clinical characteristics of 138 hospitalised patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020;323(11):1061-1069. doi:10.1001/jama.2020.1585.
14. Giustino G., Croft L.B., Stefanini G.G. Characterization of myocardial injury in patients with COVID-19. *J Am Coll Cardiol*. 2020;76:2043–2055.
15. Lippi G, Henry BM. Chronic obstructive pulmonary disease is associated with severe coronavirus disease 2019 (COVID-19). *Respir Med*. 2020; doi:10.1016/j.rmed.2020.105941.
16. Li B, Yang J, Zhao F et al. Prevalence and impact of cardiovascular metabolic diseases on COVID-19 in China. *Clin Res Cardiol*. 2020;109:521-538. doi:10.1007/s00392-020-01626-9.
17. Zhou F, Yu T, Du R et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054-1062. doi:10.1016/S0140-6736(20)30566-3.
18. Yang J, Zheng Y, Gou X et al. Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis. *Int J Infect Dis*. 2020;94:91-95. doi:10.1016/j.ijid.2020.03.017.
19. Chen N, Zhou M, Dong X et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507-513. doi:10.1016/S0140-6736(20)30211-7.
20. Epidemiology Working Group for NCIP Epidemic Response, Chinese Center for Disease Control and Prevention. [The epidemiological characteristics of an outbreak of 2019 novel coronavirus disease (COVID-19) in China]. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020;41(2):145-151. doi:10.3760/cma.j.issn.0254-6450.2020.02.003.
21. Yang X, Yu Y, Xu J et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centred, retrospective, observational study. *Lancet Respir Med*. 2020;8(5):475-481. doi:10.1016/S2213-2600(20)30079-5.
22. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;323(13):1239-1242. doi:10.1001/jama.2020.2648.
23. Targher G, Mantovani A, Wang X-B et al. Patients with diabetes are at higher risk for severe illness from COVID-19. *Diabetes Metab*. 2020; doi:10.1016/j.diabet.2020.05.001
24. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med*. 2020;8(4):e21. doi:10.1016/S2213-2600(20)30116-8.
25. Wu C, Chen X, Cai Y et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med*. 2020; doi:10.1001/jamainternmed.2020.0994.
26. Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med*. 2020;46:846-848.

- doi:10.1007/s00134-020-05991-x.
27. Huang I, Lim MA, Pranata R. Diabetes mellitus is associated with increased mortality and severity of disease in COVID-19 pneumonia – as systematic review, meta-analysis, and meta-regression. *Diabetes Metab Syndr*. 2020;14(4):395-403. doi:10.1016/j.dsx.2020.04.018.
28. Hui H, Zhang Y, Yang X et al. Clinical and radiographic features of cardiac injury in patients with 2019 novel coronavirus pneumonia. *medRxiv*. 2020; doi:10.1101/2020.02.24.20027052.
29. Shi S, Qin M, Cai Y, et al. Characteristics and clinical significance of myocardial injury in patients with severe coronavirus disease 2019. *Eur Heart J*. 2020; doi:10.1093/eurheartj/ehaa408.
30. Li J-W, Han T-W, Woodward M et al. The impact of 2019 novel coronavirus on heart injury: a systematic review and meta-analysis. *Prog Cardiovasc Dis*. 2020; doi:10.1016/j.pcad.2020.04.008.
31. Xhu L, She Z-G, Cheng X et al. Association of blood glucose control and outcomes in patients with COVID-19 and pre-existing type 2 diabetes. *Cell Metab*. 2020; doi:10.1016/j.cmet.2020.04.021.
32. Abrams M, Wan E.Y., Waase M.P., et al. Clinical and cardiac characteristics of COVID-19 mortalities in a diverse New York City cohort. *J Cardiovasc Electrophysiol*. 2020. doi: 10.1111/jce.14772.
33. Arentz M, Yim E, Klaff L et al. Characteristics and outcomes of 21 critically ill patients with COVID-19 in Washington State. *JAMA*. 2020;323(16):1612-1614. doi:10.1001/jama.2020.4326.
34. Barman HA, Atici A, Sahin I, et al. Prognostic significance of cardiac injury in COVID-19 patients with and without coronary artery disease. *Coronary Artery Dis*. 2020; doi: 10.1097/MCA.0000000000000914.
35. Bertini M., Ferrari R., Guardigli G., Malagu M., Vitali F., Zucchetti O., et al. (2020). Electrocardiographic features of 431 consecutive, critically ill COVID-19 patients: an insight into the mechanisms of cardiac involvement. *ESC*, doi: 10.1093/europace/ehaa258.
36. Linschoten M., Asselbergs F.W. (2020). CAPACITY-COVID: a European Registry to determine the role of cardiovascular disease in the COVID-19 pandemic. doi:10.1093/eurheartj/ehaa280. The Netherlands. Accessed 13 October 2020. <https://capacity-covid.eu/>
37. Duerr G.D., Heine A., Hamiko M., Zimmer S., Luetkens J.A., Nattermann J., et al. (2020). Parameters predicting COVID-19-induced myocardial injury and mortality. *Life Sci*, 260, 118400.
38. Ferrante G., Fazzari F., Cozzi O., Maurina M., Bragato R., D'Orazio F., et al. (2020). Risk factors for myocardial injury and death in patients with COVID-19: insights from a cohort study with chest computed tomography. *ESC*, doi: 10.1093/cvr/cvaa193.
39. Ghio S., Baldi E., Vicentini A., Lenti M.V., Sabatino A.D., Matteo A.D., et al. (2020). Cardiac involvement at presentation in patients hospitalized with COVID-19 and their outcome in a tertiary referral hospital in Northern Italy. *Intern Emerg Med*, doi: 10.1007/s11739-020-02493-y.
40. Lairez O., Blanchard V., Houard V., Vardon-Boune F., Lemasle M., Cariou E., et al. (2020). Cardiac imaging phenotype in patients with coronavirus disease 2019 (COVID-19): results of the cocarde study. *Int J Cardiovasc Imag*, doi: 10.1007/s10554-020-02010-4.
41. Lala A., Johnson K.W., Januzzi J.L., Russak A.J., Paranjpe I., Richter F., et al. (2020). Prevalence and impact of myocardial injury in patients hospitalized with COVID-19 infection. *J Am College Cardiol*, doi: 10.1016/j.jacc.2020.06.007.
42. Li L., Zhang S., He B., Chen X., Zhao Q. (2020). Retrospective study of risk factors for myocardial damage in patients with critical coronavirus disease 2019 in Wuhan. *J Am Heart Assoc*, 10.1161/JAHA.120.016706.
43. Liu Y, Yang Y, Zhang C et al. Clinical and biochemical indexes from 2019-nCov infected patients linked to viral loads and lung injury. *Sci China Life Sci*. 2020;63:364-374. doi:10.1007/s11427-020-1643-8.
44. Lombardi C.M., Carubelli V., Iorio A., Inciardi R.M., Bellasi A., Canale C., et al. (2020). Association of troponin levels with mortality in Italian patients hospitalized with coronavirus disease 2019: Results of a multicenter study. *JAMA Cardiol*, doi: 10.1001/jamacardio.2020.3538.
45. Lorente-Ros A., Ruiz J.M.M., Rincon L.M., Perez R.O., Rivas S., Martinez-Moya R., et al. (2020). Myocardial injury determination improves risk stratification and predicts mortality in COVID-19 patients. *Cardiol J*, doi: 10.5603/CJ.a2020.0089.
46. Ma K-L, Liu Z-H, Cao C-F et al. COVID-19 myocarditis and severity factors: an adult cohort study. *medRxiv*. 2020; doi:10.1101/2020.03.19.20034124.
47. Pagnesi M., Baldetti L., Beneduce A., Calvo F., Gramegna M., Pazzanese V., et al. (2020). Pulmonary hypertension and right ventricular involvement in hospitalised patients with COVID-19. *Heart*, doi: 10.1136/heartjnl-2020-317355.
48. Raad M., Dabbagh M., Gorgis S., Yan J., Chehab O., Dagher C., et al. (2020). Cardiac injury patterns and inpatients outcomes among patients admitted with COVID-19. *Elsevier*, doi: 10.1016/j.amjcard.2020.07.040.
49. Shah P., Doshi R., Chenna A., Owens R., Cobb A., Ivey H., et al. (2020). Prognostic value of elevated cardiac troponin I in hospitalized COVID-19 patients. *Elsevier*, 10.1016/j.amjcard.2020.08.041.

50. Shahrokh K.S., Oraii A., Soleimani A., Hadadi A., Shajari Z., Montazeri M., *et al.* (2020). The association between cardiac injury and outcomes in hospitalised patients with COVID-19. *Intern Emerg Med*, doi: 10.1007/s11739-020-02466-1.
51. Song Y., Gao P., Ran T., Qian H., Guo F., Chang L., *et al.* (2020). High inflammatory burden: a potential cause of myocardial injury in critically ill patients with COVID-19. *Front Cardiovasc Med*, doi: 10.3389/fcvm.2020.00128.
52. Szekely Y., Lichter Y., Taieb P., Banai A., Hochstadt A., Merdler I., *et al.* (2020). Spectrum of cardiac manifestatos in COVID-19. *Circulation*, 142, 342–353.
53. Toraih E.A., Elshazli R.M., Hussein M.H., Elgaml A., Amin M., El-Mowafy M., *et al.* (2020). Association of cardiac biomarkers and comorbidities with increased mortality, severity, and cardiac injury in COVID-19 patients. *J Med Virol*, doi: 10.1002/jmv.26166.
54. Wei J-F., Huang F-Y., Xiong T-Y., Liu Q., Chen H., Wang H., *et al.* (2020). Acute myocardial injury is common in patients with COVID-19 and impairs their prognosis. *Heart*, 106, 1154–1159.
55. Xie Y., Chen S., Wang X., Li B., Zhang T., He X., *et al.* (2020). Early diagnosis and clinical significance of acute cardiac injury – Under the iceberg: A retrospective cohort study of 619 non-critically ill hospitalized COVID-19 pneumonia patients. *medRxiv*, doi: 10.1101/2020.07.06.20147256.
56. Xu G, Zhao J, Zhang F et al. Favorable outcomes of elderly COVID-19 patients in Guangzhou, China: a retrospective, observational study. Research Square. 2020; doi:10.21203/rs.3.rs-26511/v1.
57. Xu H., Hou K., Xu R., Li Z., Fu H., Wen L., *et al.* (2020). Clinical characteristics and risk factors of cardiac involvement in COVID-19. *J Am Heart Assoc*, doi: 10.1161/JAHA.120.016807.
58. Wang X, Jiang H, Wu P et al. Epidemiology of avian influenza A H7N9 virus in human beings across five epidemics in mainland China, 2013-17: an epidemiological study of laboratory-confirmed case series. *Lancet Infect Dis*. 2017;17(8):822-832. doi:10.1016/S1473-3099(17)30323-7.
59. Yu D, Xiang G, Zhu W et al. The re-emergence of highly pathogenic avian influenza H7N9 viruses in humans in mainland China, 2019. *Euro Surveill*. 2019;24(21):1900273. doi:10.2807/1560-7917.ES.2019.24.21.1900273.
60. Thygesen K, Alpert JS, Jaffe AS et al. Fourth universal definition of myocardial infarction (2018). *Circulation*. 2018;138(20):e618-651. doi:10.1161/CIR.0000000000000617.
61. Gao H-N, Ly H-Z, Cao B et al. Clinical findings in 111 cases of influenza A (H7N9) virus infection. *N Engl J Med*. 2013;368:2277-2285. doi:10.1056/nejmoa1305584.
62. Jeyanathan T, Overgaard C, MGeer A. Cardiac complications of influenza infection in 3 adults. *CMAJ*. 2013;185(7):581-584. doi:10.1503/cmaj.110807.
63. Wang J, Xu H, Yang X et al. Cardiac complications associated with the influenza viruses A subtype H7N9 or pandemic H1N1 in critically ill patients under intensive care. *Braz J Infect Dis*. 2017;21(1):12-18. doi:10.1016/j.bjid.2016.10.005.
64. Estabragh ZR, Mamas A. The cardiovascular manifestations of influenza: a systematic review. *Int J Cardiol*. 2013;167(6):2397-2403. doi:10.1016/j.ijcard.2013.01.274.
65. Ito T, Akamatsu K, Ukimura A et al. The prevalence and findings of subclinical influenza-associated cardiac abnormalities among Japanese patients. *Intern Med*. 2018;57(13):1819-1826. doi:10.2169/internalmedicine.0316-17.
66. Chacko B, Peter JV, Pichamutha K et al. Cardiac manifestations in patients with pandemic (H1N1) 2009 virus infection needing intensive care. *J Crit Care*. 2012;27(1):e1-106. doi:10.1016/j.jccr.2011.05.016.
67. Wells GA, Shea B, O'Connell D et al. 2019. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp. Accessed 27 July 2020.
68. Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred reporting items for systematic reviews and meta-analysis: the PRISMA statement. *PLoS Med*. 2009;6(7):e1000097. doi:10.1371/journal.pmed.1000097.
69. Stroup AF, Berlin JA, Morton SC et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA*. 2000;283(15):2008-2012. doi:10.1001/jama.283.15.2008.
70. Nehring SM., Goyal A., Bansal P., Patel B.C. (2020). C Reactive Protein (CRP) [Updated 2020 Jun 5]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK441843/>.
71. Ekelund S, Manfred K. Choosing cut-offs for NT-proBNP. 2011. <https://acutecaretesting.org/-/media/acutecaretesting/files/pdf/choosing-cut-offs-for-ntprobnp.pdf>
72. Lindsey J.B., Kennedy K.F., Stolker J.M., Gilchrist I.C, Mukherjee D., Marso S.P., *et al.* (2011). Prognostic implications of Creatine Kinase-MB elevation after percutaneous coronary intervention. *Circ Cardiovasc Interv*, 4, 474-480.

73. Madjid M, Vela D, Khalili-Tabrizi H, Casscells SW, Litovsky S. Systemic infections cause exaggerated local inflammation in atherosclerotic coronary arteries: clues to the triggering effect of acute infections on acute coronary syndromes. *Tex Heart Inst J*. 2007;34(1):11-18. PMID:17420787.
74. Corrales-Medina VF, Musher DM, Shachkina S, Chirinos JA. Acute pneumonia and the cardiovascular system. *Lancet*. 2013;381(9865):496-505. doi:10.1016/S0140-6736(12)61266-5.
75. Ishiyama Y, Gallagher PE, Averill DB, Tallant EA, Brosnihan KB, Ferrario CM. Upregulation of angiotensin-converting enzyme 2 after myocardial infarction by blockade of angiotensin II receptors. *Hypertension*. 2004;43(5):970-976. doi:10.1161/01.HYP.0000124667.34652.1a.
76. Wang J, He W, Guo L et al. The ACE2-Ang (1-7)-Mas receptor axis attenuates cardiac remodeling and fibrosis in post-myocardial infarction. *Molec Med Rep*. 2017;16(2):1973-1981. doi:10.3892/mmr.2017.6848.
77. Ferrario CM, Jessup J, Chappell MC et al. Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation*. 2005;111(20):2605-2610. doi:10.1161/circulationaha.104.510461.
78. Gheblawia M, Wang K, Viveirosa A et al. Angiotensin converting enzyme 2: SARSCoV-2 receptor and regulator of the renin-angiotensin system: celebrating the 20th anniversary of the discovery of ACE2. *Circ Res*. 2020; doi:10.1161/CIRCRESAHA.120.317015.
79. Patel VB, Zhong JC, Grant MB, Oudit GY. Role of the ACE2/Angiotensin 1-7 axis of the renin-angiotensin system in heart failure. *Circ Res*. 2016;118(8):1313-1326. doi:10.1161/CIRCRESAHA.116.307708.
80. Marchiano S, Hsiang T-Y, Higashi T, et al. SARS-CoV-2 infects human pluripotent stem cell-derived cardiomyocytes, impairing electrical and mechanical function. *bioRxiv*. 2020; doi: 10.1101/2020.08.30.274464.
81. Lindner D, Fitzek A, Brauninger H, et al. Association of cardiac infection with SARS-CoV-2 in confirmed COVID-19 autopsy cases. *JAMA Cardiol*. 2020; doi: 10.1001/jamacardio.2020.3551.
82. Han MLK, McLaughlin VV, Criner GJ, Martinez FJ. Pulmonary diseases and the heart. *Circulation*. 2007;116(25):2992-3005. doi:10.1161/CIRCULATIONAHA.106.685206.
83. Forfia PR, Vaidya A, Wiegers SE. Pulmonary heart disease: the heart-lung interaction and its impact on patient phenotypes. *Pulm Circ*. 2013;3(1):5-19. doi:10.4103/2045-8932.109910.
84. Barnes M, Heywood AE, Mahimbo A, Rahman B, Newall AT, Macintyre CR. Acute myocardial infarction and influenza: a meta-analysis of case-control studies. *Heart*. 2015;101(21):1738-1747. doi:10.1136/heartjnl-2015-307691.
85. Carter P, Lagan J, Fortune C et al. Association of cardiovascular disease with respiratory disease. *J Am Coll Cardiol*. 2019;73(17):2166-2177. doi:10.1016/j.jacc.2018.11.063.
86. Ingebrigsten TS, Marott JL, Vestbo J, Nordestgaard BG, Lange P. Coronary heart disease and heart failure in asthma, COPD and asthma-COPD overlap. *BMJ Open Res*. 2020;7:e000470. doi:10.1136/bmjresp-2019-000470.
87. Martinez-Mateo V, Fernandez-Anguita MJ, Paule A. Electrocardiographic signs of acute right ventricular hypertrophy in patients with COVID-19 pneumonia: A clinical case series. *Electrocardiology*. 2020;62:100-102.
88. Halpin DMG, Faner R, Sibila O, Badia JR, Agusti A. Do chronic respiratory diseases or their treatment affect the risk of SARS-CoV-2 infection? *Lancet*. 2020; doi:10.1016/S2213-2600(20)30167-3.
89. Bermejo-Martin JF, Martin-Fernandez M, Lopez-Mestanza C, Duque P, Almansa R. Shared features of endothelial dysfunction between sepsis and its preceding risk factors (aging and chronic disease). *J Clin Med*. 2018;7(11):400. doi:10.3390/jcm7110400.
90. Chen L, Li X, Chen M, Feng Y, Xiong C. The ACE2 expression in human heart indicated new potential mechanism of heart injury among patients infected with SARS-CoV-2. *Cardiovasc Res*. 2020;116(6):1097-1100. doi:10.1093/cvr/cvaa078.
91. Heather LC, Clarke K. Metabolism, hypoxia, and the diabetic heart. *J Mol Cell Cardiol*. 2011;50(4):598-605. doi:10.1016/j.yjmcc.2011.01.007.
92. Iqbal T, Welsby PJ, Howarth FC, Bidasee K, Adeghate E, Singh J. Effects of diabetes-induced hyperglycaemia in the heart: biochemical and structural alterations. In: Turan B, Dhalla N, editors. *Diabetic Cardiomyopathy: Advances in Biochemistry in Health and Disease*, vol 9. New York: Springer; 2014. ISBN:9781461493174.
93. Rask-Madsen C, King GL. Vascular complications of diabetes: mechanisms of injury and protective factors. *Cell Metab*. 2013;17(1):20-33. doi:10.1016/j.cmet.2012.11.012.