Burden of intestinal parasitic infections in children and its association with hand washing practice in Ethiopia: a systematic review and meta-analysis

Fasil Wagnew^{1*}, Aster Tadesse¹, Amanuel Abajobir²

¹College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia

²Maternal and Child Wellbeing, African Population and Health Research Center, Nairobi, Kenya

*corresponding author: Fasil Wagnew

Email addresses

FW: fasilw.n@gmail.com

AT: astutadesse84@gmail.com

AA: amanuel.abajobir@uq.net.au

27 Abstract

28 **Background:** Intestinal parasitic infections (IPIs) are a major public health challenges in
29 developing countries including Ethiopia, although few studies previously estimated the
30 magnitude of IPIs and associated factors in the country. Reports from these scarce studies
31 were also widely varied and remained inconsistent. This study thus aimed to synthesize the
32 pooled magnitude of IPIs and factors affecting it.

33 **Methods:** Internationally broad based medical database including MEDLINE/PubMed,
34 EMBASE, PsychINFO and Web of Science, and Google Scholar for grey literature were
35 exhaustively searched using *a priori* set criteria to identify studies estimating the prevalence
36 of IPIs among children from 2000-2018. PRISMA guideline was used to systematically
37 review and meta-analyze these studies. Details of study characteristics including sample size,
38 magnitude of effect sizes (including odds ratios (ORs)) and standard errors were extracted.
39 Random-effects model was used to calculate pooled estimates in Stata/se version-14. I^2 and
40 meta-bias statistics assessed heterogeneity and Egger's test for publication bias. Sub-group
41 analyses were also carried out based on age of children and regions.

42 **Results:** Forty-three studies were included in the final analysis (N = 20,008 children). The
43 overall prevalence of IPIs, with one or more species, was 48.2% (95% CI: 40.1, 56.3) in
44 Ethiopian children. Based on sub-group analyses, the highest prevalence of IPIs was observed
45 among school-age children (52.4% (95% CI, 41.3, 63.5)) and in Amhara regional state
46 (52.1% (95% CI, 37.3-66.8)). The odds of having IPIs was nearly six times higher in children
47 who were not practicing hand washing as compared to their counterparts (pooled OR = 5.6
48 (95% CI: 3.4,9.3). Funnel plot analysis and Egger's test detected no publication bias.

49 **Conclusion:** On aggregate, the pooled prevalence of IPIs among Ethiopian children is
50 significantly high. Not hand washing before eating was a risk factor for IPIs. The
51 establishment of applicable sanitation services and health education will help reduce the
52 magnitude of IPIs and promote a healthier childhood.

53

54 **Keywords:** Intestinal parasitic infections, systematic review and meta-analysis, Ethiopia

55

56

57 Background

58 Intestinal parasitic infections (IPIs) are serious public health challenges globally [1] whereby
 59 one-third of the total population is infected with intestinal parasites (IP), and an estimated 4.5
 60 billion people are at risk of infection with one of the three common IPIs – ascaris
 61 lumbricoides (800–1000 million), *hookworm* (700- 900 million) and *trichuris trichiura* (500
 62 million) [2-4] contributing for significant illnesses and disability-adjusted life year lost [5].
 63 Sadly, around 12% of global burden of disease caused by IPIs occur among school-age
 64 children in developing countries [7] especially in sub-Saharan African countries ranging from
 65 27.7% to 95% [8-12], and attributable to major causes of morbidity and mortality among
 66 these children [1].

67 Multifactorial causes contribute to the escalation of IPIs in children including overcrowding,
 68 lack of safe water, poor personal hygiene, poor hand washing and nutritional disorders [20,
 69 21]. Indeed, IPIs can be consequences of poor socio-economic status and also contribute to a
 70 substantial economic burden [6]. That is, IPIs are closely interrelated with poor
 71 socioeconomic status of the family, lack of sanitation, inadequate access to pure water and
 72 waste disposal especially in low- and middle-income countries [13-15]. In particular, under-
 73 five and school-aged children in these settings are categorized as high-risk population for IPIs
 74 [16]. Furthermore, these infections are causes and consequences for anemia and other
 75 comorbidities as well as physical and cognitive developmental delays [17, 18] resulting in
 76 suboptimal child survival, poor growth and development as well as meager educational
 77 performance [8, 10, 18]. In addition, chronic forms of IPIs are associated with the
 78 transmission and severity of other infections and/or diseases such as leprosy and human
 79 immune-deficiency virus (HIV) [19].

80 Interestingly, these infections have recently been receiving a high priority from global
 81 community. Cognizant to this, the World Health Organization (WHO) has permitted
 82 preventive chemoprophylaxis as worldwide mainstay strategy to control the communicability
 83 of these infections [22]. In this strategy, scheduled administration of safe and efficacious
 84 deworming drugs for risk groups is recommended, setting a minimum target at 75%, and
 85 maximum 100% for success rate [22-24]. In fact, general cleanliness or sanitation are
 86 imperative indicators to achieve these standards at a community level [25].

87 Numerous studies have studied the magnitude of IPIs in preschool and school-age children in
 88 Ethiopia. However, there is a significant variability in the reports and an aggregate figure is
 89 implicated to know the ‘real’ burden of IPIs. We thus carried out a systematic review and

meta-analysis on the prevalence of IPIs and its association with common indicators including hand washing practice in children. These analyses were carried out in the context of in-depth methodological application and rigorous statistical analyses that could be helpful for estimating national level burden of IPIs as previous studies are characterized by high spatial variation.

Methods

Study design and search strategy

Using broad-based computerized international databases, we employed systematic searches to find all published studies on the prevalence of IPIs in Ethiopian children from 2000-2020. Databases included MEDLINE/PubMed, EMBASE, Psych INFO and Web of Science, and Google Scholar for grey literature. Reference lists of eligible articles, hand searches for grey literature and relevant scientific conference websites were also appraised to enhance electronic searches. Observational studies conducted on the prevalence of IPIs among children in Ethiopia were selected for this systematic review and meta-analysis. Search protocol was developed by using such keywords as “prevalence” OR “Epidemiology” AND “intestinal parasite,” “intestinal helminthes”, “intestinal protozoa”, “soil-transmitted helminthes” AND “associated factors” OR “predictors” AND “Ethiopia”. To make a certain scientific rigor, we strictly adhered to the Preferred Reporting of Systematic Reviews and Meta-Analysis (PRISMA) guideline [26].

Inclusion criteria

All studies written in English, estimating the prevalence of IPIs in Ethiopian children and of high-quality scores were entered into the final meta-analysis.

Exclusion Criteria

Studies that did not estimate the prevalence of IPIs, articles without definite sample size or full-text, those that did not achieve the minimum quality scores, case reports, case series, and experimental studies were excluded from the meta-analysis.

Data Extraction

Two reviewers (FW and AT) screened the downloaded titles and abstracts using the eligibility criteria. Discussions and mutual consensus were in place when possible arguments raised between the reviewers. Two reviewers (FW and AT) then assessed full-texts of potentially eligible papers. We made some efforts to communicate the authors whenever further information was needed. Numerator and denominator data and beta coefficients and

their standard errors (if given) were used to calculate ORs, where ORs with 95% CI were not available. The data extraction format included first author, study year, region of study, study design and sample size, and prevalence of IPIs.

Quality Appraisal

Articles were assessed for quality, with only high-quality studies included in the analysis. Two authors (FW, AT) independently evaluated the quality of each relevant article. The reviewer's cross-checked their quality appraisal scores and resolved any disparity before calculating the final appraisal score. Moreover, Kappa statistic was used to measure agreement between reviewers to determine uniformity of amongst eligible full-text articles using guidelines proposed by Landis and Koch [31]: < 0.20 = slight agreement; $0.21-0.40$ = fair agreement; $0.41-0.60$ = moderate agreement; $0.61-0.80$ = substantial agreement; and > 0.80 = almost perfect agreement. Finally, AA reviewed overall quality of extracted data and edited the manuscript. Newcastle-Ottawa Scale adapted for prevalence studies methodological quality assessment tool was also used [27]. The tool has three components in general—the first component graded as five stars gives a due emphasis on the methodological quality of each original study. The second component of the tool explores the comparability of the study. The third component focuses on the outcome and statistical analysis of studies. Articles with a score of ≥ 6 out of 10 were considered as high-quality. Accordingly, all eligible studies had high quality scores.

Data analysis

Information about study designs, study sample and country were summarized using Microsoft Excel and then exported to STATA/se version-14 for analysis. Meta-analysis of the magnitude of IPIs was carried out using a random-effects model, generating a pooled prevalence with 95% CIs. Heterogeneity among included studies was assessed using Cochran's Q and I^2 statistic [28]. I^2 statistic estimates the percentage of heterogeneity across studies. Forest plots were also used to estimate the pool effect sizes and weight of overall and each included study with respective CI to offer a graphic summary of the data. Publication bias was determined based on the symmetry of funnel plots [29], and Egger's test [30]. Sub-group analysis was carried out, based on age of participants and regional states, because of the presence of significant heterogeneity across the included studies.

Results

Our search yielded 2,896 articles; of these, 442 duplications and 2,411 unrelated articles were excluded. Kappa coefficient revealed that agreement rate between the two reviewers for the included articles was 0.81. Fifteen full-text articles [32-46] were excluded due to unmet outcome of interest or location. Finally, 43 eligible studies representing 20,008 participants met the inclusion criteria for this review that determine the burden of IPIs (Figure 1).

Descriptive characteristics of the included studies

Descriptive characteristics for the included epidemiological studies is presented in Table 1. Of these, 42 studies were cross-sectional in design and only one study was case-control [47]. Children were classified based on presence versus absence of an IPI for any species. The largest sample size was reported by the study conducted by Grimes J et al [50] and smallest for the study done by Mulatu G [16]. Similarly, the prevalence of IPI was highest for the study conducted by Worku L et al [48] and lowest for the study conducted by Zemene T [49] (Table 1).

Prevalence of IPIs in children

The overall prevalence of IPIs with one or more species was 48.2% (95% CI: 40.1, 56.3%) in Ethiopian children (Figure 2). We performed subgroup analyses based on regional states and age of participants as there was significant heterogeneity across included studies. For instance, the prevalence of IPIs was 52.4% (95% CI: 41.3, 63.5) in school-age children and 52.1% (95% CI: 37.3, 66.8) of children with IPIs lived in Amhara regional state (Table 2). Overall Egger's test revealed no statistically significant publication bias, p-value = 0.62 (Additional file 1).

Association of residence and hand washing practice with IPIs

Five studies were included to determine the association of hand washing before eating on IPIs (Table 3). Children without hand washing before eating experienced six times more likely the occurrence of IPIs (OR: 5.6 (95% CI: 3.4, 9.3)) as compared to hand washing before eating (Figure 3). However, the association between residence (urban versus rural) showed a non-significant association with IPIs (OR: 1.1 (95% CI: 0.5, 2.1) (Figure 4).

Discussions

This study estimated the pooled prevalence of IPIs with one or more species in Ethiopian children and its association with residence and hand washing practice, using the documented data from systematic literature review, which was assembled from different regions of the country. After diligent search through scientific databases, scientific reports and grey literature, 43 studies were included in the final analysis. Accordingly, the overall pooled prevalence of IPIs among Ethiopian children was 48.2%. This finding was in line with previous studies that revealed 42.5% IPIs in Syria [51], 47.6% in Afghanistan [52] and 38% in Iran [53]. In contrast, the prevalence of IPIs observed in current study is higher compared to studies in Thailand that reported 38.8% [54], in Shush country (5.14%) [55], in Turkey (17%) [56] and in Korea (10.5%) [57]. This observed variation could be due to socio-economic status, poor hygienic practice and sanitary facilities, inadequate safe drinking water and environmental factors as well as different methods used for the diagnosis of IPIs.

We performed sub-group analysis based on the age of participants and regional states of the country. Accordingly, the evidence revealed that the prevalence of IPIs among school-age children was two-fold higher than that in under five-children. The difference observed between under-five and school-age children might be because of the following reasons—first, close contact between students and unhygienic conditions in the school compound; second, variation in sanitation and human development indices; and finally, due to implementation of continuing deworming programme for under-five children by ministry of health. Likewise, the highest prevalence of IPIs were observed in Amhara regional states and the lowest prevalence was in Tigray state. This regional discrepancy might be because of cultural habits (sanitary status) among the regions, variations in methodology and sample size. The advancement of public health measures and policy implementation could also vary across regions that might attribute for the difference.

In an effort to examine the association between residence and hand washing before eating and IPIs, hand washing before eating was associated with a significant reduction in the occurrence of IPIs. Indeed, IPIs is frequently caused by *hookworms*, *ascaris lumbricoides* and *trichuris trichiura* and affected over one billion individuals globally [58]. This could be due to probable to be hand contaminated with cysts of *giardia*, *cryptosporidium* and other species. The meager association between residence and IPIs in this meta-analysis is supported

by previous studies [60, 61]. This might be because of variations among studies, although Egger's test for publication bias showed no statistically significant evidence, $p\text{-value} = 0.06$.

The findings of this study may have both clinical and public health implications towards scaling up effective and efficient management of IPIs by informing decision makers and practitioners. In other words, understanding the magnitude and distribution of IPIs is an important precondition for planning and evaluating intervention programs. For instance, safe water and personal hygiene as a basic human right will indisputably result in major health improvements and prevention of different diseases including parasitic infections [59]. That is, since almost all of IPIs are feco-oral communicable infections, delivery of tap water and latrines, health education on personal hygiene and environmental sanitation are crucial to control and reduce IPIs in the country.

However, these findings need to be considered in the context of some important limitations; these include: (i) significant heterogeneity across studies might have contributed for meager association between residence and IPIs, although presumably children in urban areas have better access to personal and environmental hygiene; (ii) this study did not determine other possible risk factors contributing to the occurrence of IPIs; (iii) the gold standard diagnosis of Pinworm infection is Gram staining, though several studies had employed only wet smear; and (iv) stool samples were taken at once but standard diagnosis usually take at least 3 samples.

Conclusion

This meta-analysis summarized a high pooled prevalence of IPIs in Ethiopian children especially among school-age children. Establishment of applicable sanitation services and health education for hand washing will help make a healthier childhood.

List of abbreviations

AA	Amanuel Abajobir
AT	Aster Tadesse
FW	Fasil Wagnew
IPIs	Intestinal Parasitic Infections
OR	Odds Ratio

Declarations

Ethics approval and consent to participate

Not applicable

Consent to publication

Not applicable

Availability of data and material

-The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

-PRISMA checklist

Competing interests

The authors declare that they have no competing interests.

Funding

Not applicable

Authors' contribution

FW involved in the conception of research idea; (FW, AT, AA) undertook data extraction, analysis, interpretation, and manuscript write-up. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to acknowledge the Debre Markos University library for providing us with a wide range of available online databases.

References

1. Mehraj V, Hatcher J, Akhtar S, Rafique G, Beg MA: **Prevalence and factors associated with intestinal parasitic infection among children in an urban slum of Karachi.** *PloS one* 2008, **3**(11):e3680.
2. Horton J: **Human gastrointestinal helminth infections: are they now neglected diseases?** *Trends in parasitology* 2003, **19**(11):527-531.
3. Utzinger J, Keiser J: **Schistosomiasis and soil-transmitted helminthiasis: common drugs for treatment and control.** *Expert opinion on pharmacotherapy* 2004, **5**(2):263-285.
4. Organization WH: **Prevention and control of intestinal parasitic infections: report of a WHO Expert Committee [meeting held in Geneva from 3 to 7 March 1986].** 1987.
5. Okyay P, Ertug S, Gultekin B, Onen O, Beser E: **Intestinal parasites prevalence and related factors in school children, a western city sample-Turkey.** *BMC public health* 2004, **4**(1):64.
6. Chan MS: **The global burden of intestinal nematode infections—fifty years on.** *Parasitology today* 1997, **13**(11):438-443.

- 272 7. Hailegebriel T: **Prevalence of intestinal parasitic infections and associated risk factors**
273 **among students at Dona Berber primary school, Bahir Dar, Ethiopia.** *BMC infectious*
274 *diseases* 2017, **17**(1):362.
- 275 8. Abdel-Aziz MA, Afifi AA, Malik EM, Adam I: **Intestinal protozoa and intestinal**
276 **helminthic infections among schoolchildren in Central Sudan.** *Asian Pacific Journal of*
277 *Tropical Medicine* 2010, **3**(4):292-293.
- 278 9. Ayalew A, Debebe T, Worku A: **Prevalence and risk factors of intestinal parasites among**
279 **Delgi school children, North Gondar, Ethiopia.** *Journal of Parasitology and Vector*
280 *Biology* 2011, **3**(5):75-81.
- 281 10. Emile N, Bosco NJ, Karine B: **Prevalence of intestinal parasitic infections and associated**
282 **risk factors among Kigali Institute of Education students in Kigali, Rwanda.** *Tropical*
283 *biomedicine* 2013, **30**(4):718-726.
- 284 11. Erismann S, Diagbouga S, Odermatt P, Knoblauch AM, Gerold J, Shrestha A, Grissoum T,
285 Kaboré A, Schindler C, Utzinger J: **Prevalence of intestinal parasitic infections and**
286 **associated risk factors among schoolchildren in the Plateau Central and Centre-Ouest**
287 **regions of Burkina Faso.** *Parasites & vectors* 2016, **9**(1):554.
- 288 12. Speich B, Marti H, Ame SM, Ali SM, Bogoch II, Utzinger J, Albonico M, Keiser J:
289 **Prevalence of intestinal protozoa infection among school-aged children on Pemba**
290 **Island, Tanzania, and effect of single-dose albendazole, nitazoxanide and albendazole-**
291 **nitazoxanide.** *Parasites & vectors* 2013, **6**(1):3.
- 292 13. Brooker S: **Estimating the global distribution and disease burden of intestinal nematode**
293 **infections: adding up the numbers—a review.** *International journal for parasitology* 2010,
294 **40**(10):1137-1144.
- 295 14. Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J: **Helminth infections:**
296 **the great neglected tropical diseases.** *The Journal of clinical investigation* 2008,
297 **118**(4):1311-1321.
- 298 15. Utzinger J, Bergquist R, Olveda R, Zhou X-N: **Important helminth infections in Southeast**
299 **Asia: diversity, potential for control and prospects for elimination.** In: *Advances in*
300 *parasitology. Volume 72*, edn.: Elsevier; 2010: 1-30.
- 301 16. Mulatu G, Zeynudin A, Zemene E, Debalke S, Beyene G: **Intestinal parasitic infections**
302 **among children under five years of age presenting with diarrhoeal diseases to two public**
303 **health facilities in Hawassa, South Ethiopia.** *Infectious diseases of poverty* 2015, **4**(1):49.
- 304 17. Brooker S, Clements AC, Bundy DA: **Global epidemiology, ecology and control of soil-**
305 **transmitted helminth infections.** *Advances in parasitology* 2006, **62**:221-261.
- 306 18. King CH, Bertino A-M: **Asymmetries of poverty: why global burden of disease valuations**
307 **underestimate the burden of neglected tropical diseases.** *PLoS Neglected Tropical*
308 *Diseases* 2008, **2**(3):e209.
- 309 19. Gelaw A, Anagaw B, Nigussie B, Silesh B, Yirga A, Alem M, Endris M, Gelaw B:
310 **Prevalence of intestinal parasitic infections and risk factors among schoolchildren at the**
311 **University of Gondar Community School, Northwest Ethiopia: a cross-sectional study.**
312 *BMC public health* 2013, **13**(1):304.
- 313 20. Alum A, Rubino JR, Ijaz MK: **The global war against intestinal parasites—should we use**
314 **a holistic approach?** *International Journal of Infectious Diseases* 2010, **14**(9):e732-e738.
- 315 21. Tiwari BR, Chaudhary R, Adhikari N, Jayaswal S, Poudel T, Rijal K: **Prevalence of**
316 **intestinal parasitic infections among school children of Dadeldhura District.** *Nepal JHAS*
317 2013, **3**(1):14-16.
- 318 22. Organization WH: **Prevention and control of schistosomiasis and soil-transmitted**
319 **helminthiasis: report of a WHO expert committee.** 2002.
- 320 23. Hotez PJ, Molyneux DH, Fenwick A, Kumaresan J, Sachs SE, Sachs JD, Savioli L: **Control**
321 **of neglected tropical diseases.** *New England Journal of Medicine* 2007, **357**(10):1018-1027.
- 322 24. Ismail M: **Preventive chemotherapy in human helminthiasis, Coordinating use of**
323 **anthelmintic drugs in control interventions: A manual for health professionals and**
324 **programme managers.** *Indian Journal of Medical Research* 2007, **126**(3):235.

25. Phiri K, Whitty C, Graham S, Ssembatya-Lule G: **Urban/rural differences in prevalence and risk factors for intestinal helminth infection in southern Malawi.** *Annals of Tropical Medicine & Parasitology* 2000, **94**(4):381-387.
26. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gotzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D: **The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration.** *J Clin Epidemiol* 2009, **62**(10):e1-34.
27. **Newcastle-Ottawa Scale customized for cross-sectional studies.** In. available from https://static-content.springer.com/esm/.../12889_2012_5111.
28. Higgins JP, Thompson SG, Deeks JJ, Altman DG: **Measuring inconsistency in meta-analyses.** *BMJ* 2003, **327**(7414):557-560.
29. Borenstein M, Hedges LV, Higgins J, Rothstein HR: **A basic introduction to fixed-effect and random-effects models for meta-analysis.** *Research Synthesis Methods* 2010, **1**(2):97-111.
30. Begg CB, Mazumdar M: **Operating characteristics of a rank correlation test for publication bias.** *Biometrics* 1994:1088-1101.
31. Landis JR, Koch GG: **The measurement of observer agreement for categorical data.** *biometrics* 1977:159-174.
32. Akinbo FO, Okaka C, Omoregie R: **Prevalence of intestinal parasitic infections among HIV patients in Benin City, Nigeria.** *Libyan Journal of Medicine* 2010, **5**(1):5506.
33. Awole M, Gebre-Selassie S, Kassa T, Kibru G: **Prevalence of intestinal parasites in HIV-infected adult patients in Southwestern Ethiopia.** *Ethiopian Journal of Health Development* 2003, **17**(1):71-78.
34. Chan M, Medley G, Jamison D, Bundy D: **The evaluation of potential global morbidity attributable to intestinal nematode infections.** *Parasitology* 1994, **109**(3):373-387.
35. Ericsson CD, Steffen R, Siddiqui AA, Berk SL: **Diagnosis of Strongyloides stercoralis infection.** *Clinical Infectious Diseases* 2001, **33**(7):1040-1047.
36. Harhay MO, Horton J, Olharto PL: **Epidemiology and control of human gastrointestinal parasites in children.** *Expert review of anti-infective therapy* 2010, **8**(2):219-234.
37. Kassu A, Mengistu G, Ayele B, Diro E, Mekonnen F, Ketema D, Moges F, Mesfin T, Getachew A, Ergicho B: **HIV and intestinal parasites in adult TB patients in a teaching hospital in Northwest Ethiopia.** *Tropical doctor* 2007, **37**(4):222-224.
38. Kassu A, Tsegaye A, Wolday D, Petros B, Aklilu M, Sanders E, Fontanet A, Van Baarle D, Hamann D, De Wit TR: **Role of incidental and/or cured intestinal parasitic infections on profile of CD4+ and CD8+ T cell subsets and activation status in HIV-1 infected and uninfected adult Ethiopians.** *Clinical & Experimental Immunology* 2003, **132**(1):113-119.
39. Legesse M, Erko B: **Zoonotic intestinal parasites in Papio anubis (baboon) and Cercopithecus aethiops (vervet) from four localities in Ethiopia.** *Acta Tropica* 2004, **90**(3):231-236.
40. Mengistu A, Gebre-Selassie S, Kassa T: **Prevalence of intestinal parasitic infections among urban dwellers in southwest Ethiopia.** *Ethiopian Journal of Health Development* 2007, **21**(1):12-17.
41. Scrivener S, Yemaneberhan H, Zebeignus M, Tilahun D, Girma S, Ali S, McElroy P, Custovic A, Woodcock A, Pritchard D: **Independent effects of intestinal parasite infection and domestic allergen exposure on risk of wheeze in Ethiopia: a nested case-control study.** *The Lancet* 2001, **358**(9292):1493-1499.
42. Sissay MM, Ugglä A, Waller PJ: **Epidemiology and seasonal dynamics of gastrointestinal nematode infections of sheep in a semi-arid region of eastern Ethiopia.** *Veterinary Parasitology* 2007, **143**(3-4):311-321.
43. Sissay MM, Ugglä A, Waller PJ: **Prevalence and seasonal incidence of larval and adult cestode infections of sheep and goats in eastern Ethiopia.** *Tropical Animal Health and Production* 2008, **40**(6):387-394.
44. Aiemojoy K, Gebresillasie S, Stoller NE, Shiferaw A, Tadesse Z, Chanyalew M, Aragie S, Callahan K, Keenan JD: **Epidemiology of soil-transmitted helminth and intestinal**

379 **Protozoan infections in preschool-aged children in the amhara region of Ethiopia.** *The*
380 *American journal of tropical medicine and hygiene* 2017, **96**(4):866-872.

381 45. Maru DS: **Prevalence of intestinal parasitic infections and associated risk factors among**
382 **school children in Adigrat town, northern Ethiopia.** *Int J Emerg Trends in Sci Technol*
383 2015, **4**(1):4943-4948.

384 46. Sayyari A, Imanzadeh F, Bagheri Yazdi S, Karami H, Yaghoobi M: **Prevalence of intestinal**
385 **parasitic infections in the Islamic Republic of Iran.** 2005.

386 47. Firdu T, Abunna F, Girma M: **Intestinal protozoal parasites in diarrheal children and**
387 **associated risk factors at Yirgalem Hospital, Ethiopia: A case-control study.**
388 *International scholarly research notices* 2014, **2014**.

389 48. Worku L, Damte D, Endris M, Tesfa H, Aemero M: **Schistosoma mansoni infection and**
390 **associated determinant factors among school children in Sanja Town, Northwest**
391 **Ethiopia.** *Journal of parasitology research* 2014, **2014**.

392 49. Zemene T, Shiferaw MB: **Prevalence of intestinal parasitic infections in children under**
393 **the age of 5 years attending the Debre Birhan referral hospital, North Shoa, Ethiopia.**
394 *BMC research notes* 2018, **11**(1):58.

395 50. Grimes JE, Tadesse G, Gardiner IA, Yard E, Wuletaw Y, Templeton MR, Harrison WE,
396 Drake LJ: **Sanitation, hookworm, anemia, stunting, and wasting in primary school**
397 **children in southern Ethiopia: Baseline results from a study in 30 schools.** *PLoS*
398 *neglected tropical diseases* 2017, **11**(10):e0005948.

399 51. Al-kafri A, Harba A: **Intestinal parasites in basic education pupils in urban and rural**
400 **Idlib.** *Syrian Clinical Laboratory Revues* 2009, **5**(2):2-5.

401 52. Gabrielli A, Ramsan M, Naumann C, Tsogzolmaa D, Bojang B, Khoshal M, Connolly M,
402 Stothard J, Montresor A, Savioli L: **Soil-transmitted helminths and haemoglobin status**
403 **among Afghan children in World Food Programme assisted schools.** *Journal of*
404 *helminthology* 2005, **79**(4):381-384.

405 53. Daryani A, Hosseini-Teshnizi S, Hosseini S-A, Ahmadpour E, Sarvi S, Amouei A, Mizani A,
406 Gholami S, Sharif M: **Intestinal parasitic infections in Iranian preschool and school**
407 **children: A systematic review and meta-analysis.** *Acta tropica* 2017, **169**:69-83.

408 54. Nithikathkul C, Changsap B, Wannapinyosheep S, Poister C, Boontan P: **The prevalence of**
409 **Enterobius vermicularis among primary school students in Samut Prakan Province,**
410 **Thailand.** *Southeast Asian journal of tropical medicine and public health* 2001, **32**:133-137.

411 55. Fallahizadeh S, Feiz-Haddad MH, Kazemi F, Afrisham R: **Prevalence of Intestinal Parasitic**
412 **Infections in Shush County, Southwest of Iran during 2014-2016.** *International Journal of*
413 *Infection* 2017, **4**(3).

414 56. Çeliksöz A, Güler N, Güler G, Öztöpe AY, Degerli S: **Prevalence of intestinal parasites in**
415 **three socioeconomically-different regions of Sivas, Turkey.** *Journal of Health, Population*
416 *and Nutrition* 2005:184-191.

417 57. Lee S-E, Lee J-H, Ju J-W, Lee W-J, Cho S-H: **Prevalence of Enterobius vermicularis**
418 **among preschool children in Gimhae-si, Gyeongsangnam-do, Korea.** *The Korean journal*
419 *of parasitology* 2011, **49**(2):183.

420 58. Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, Hotez PJ: **Soil-**
421 **transmitted helminth infections: ascariasis, trichuriasis, and hookworm.** *The Lancet*
422 2006, **367**(9521):1521-1532.

423 59. Bartram J, Cairncross S: **Hygiene, sanitation, and water: forgotten foundations of health.**
424 *PLoS medicine* 2010, **7**(11):e1000367.

425 60. Strunz EC, Addiss DG, Stocks ME, Ogden S, Utzinger J, Freeman MC: **Water, sanitation,**
426 **hygiene, and soil-transmitted helminth infection: a systematic review and meta-analysis.**
427 *PLoS medicine* 2014, **11**(3):e1001620.

428 61. Ziegelbauer K, Speich B, Mäusezahl D, Bos R, Keiser J, Utzinger J: **Effect of sanitation on**
429 **soil-transmitted helminth infection: systematic review and meta-analysis.** *PLoS medicine*
430 2012, **9**(1):e1001162.

431 62. Ameya G, Tsalla T, Getu F, Getu E: **Antimicrobial susceptibility pattern, and associated**
432 **factors of Salmonella and Shigella infections among under five children in Arba Minch,**
433 **South Ethiopia.** *Annals of clinical microbiology and antimicrobials* 2018, **17**(1):1.

- 434 63. Tadesse G: **The prevalence of intestinal helminthic infections and associated risk factors**
435 **among school children in Babile town, eastern Ethiopia.** *Ethiopian Journal of Health*
436 *Development* 2005, **19**(2):140-147.
- 437 64. de Lucio A, Amor-Aramendía A, Bailo B, Saugar JM, Anegagrie M, Arroyo A, López-
438 Quintana B, Zewdie D, Ayehubizu Z, Yizengaw E: **Prevalence and genetic diversity of**
439 **Giardia duodenalis and Cryptosporidium spp. among school children in a rural area of**
440 **the Amhara Region, North-West Ethiopia.** *PloS one* 2016, **11**(7):e0159992.
- 441 65. Abossie A, Seid M: **Assessment of the prevalence of intestinal parasitosis and associated**
442 **risk factors among primary school children in Chenchu town, Southern Ethiopia.** *BMC*
443 *Public Health* 2014, **14**(1):166.
- 444 66. Alemu A, Tegegne Y, Damte D, Melku M: **Schistosoma mansoni and soil-transmitted**
445 **helminths among preschool-aged children in Chuahit, Dembia district, Northwest**
446 **Ethiopia: prevalence, intensity of infection and associated risk factors.** *BMC Public*
447 *Health* 2016, **16**(1):422.
- 448 67. Mathewos B, Alemu A, Woldeyohannes D, Alemu A, Addis Z, Tiruneh M, Aimero M, Kassu
449 A: **Current status of soil transmitted helminths and Schistosoma mansoni infection**
450 **among children in two primary schools in North Gondar, Northwest Ethiopia: a cross**
451 **sectional study.** *BMC Research Notes* 2014, **7**(1):88.
- 452 68. Tadege B, Shimelis T: **Infections with Schistosoma mansoni and geohelminths among**
453 **school children dwelling along the shore of the Lake Hawassa, southern Ethiopia.** *PloS*
454 *one* 2017, **12**(7):e0181547.
- 455 69. Beyene G, Tasew H: **Prevalence of intestinal parasite, Shigella and Salmonella species**
456 **among diarrheal children in Jimma health center, Jimma southwest Ethiopia: a cross**
457 **sectional study.** *Annals of clinical microbiology and antimicrobials* 2014, **13**(1):10.
- 458 70. Hailu T, Alemu M, Abera B, Mulu W, Yizengaw E, Genanew A, Beredie F: **Multivariate**
459 **analysis of factors associated with Schistosoma mansoni and hookworm infection among**
460 **primary school children in rural Bahir Dar, Northwest Ethiopia.** *Tropical diseases, travel*
461 *medicine and vaccines* 2018, **4**(1):4.
- 462 71. Reji P, Belay G, Erko B, Legesse M, Belay M: **Intestinal parasitic infections and**
463 **malnutrition amongst first-cycle primary schoolchildren in Adama, Ethiopia.** *African*
464 *journal of primary health care & family medicine* 2011, **3**(1).
- 465 72. Tefera E, Belay T, Mekonnen SK, Zeynudin A, Belachew T: **Prevalence and intensity of**
466 **soil transmitted helminths among school children of Mendera Elementary School,**
467 **Jimma, Southwest Ethiopia.** *The Pan African medical journal* 2017, **27**.
- 468 73. Alemayehu B, Tomass Z: **Prevalence of intestinal Helminthiasis and associated risk**
469 **factors among school children in Dawro zone, southern Ethiopia.** *J Biol Agr Heal Care*
470 2015, **5**:76-82.
- 471 74. Degarege A, Erko B: **Prevalence of intestinal parasitic infections among children under**
472 **five years of age with emphasis on Schistosoma mansoni in Wonji Shoa Sugar Estate,**
473 **Ethiopia.** *PloS one* 2014, **9**(10):e109793.
- 474 75. Begna T, Solomon T, Yohannes ZENEBA EA: **Intestinal parasitic infections and**
475 **nutritional status among primary school children in Delo-mena district, South Eastern**
476 **Ethiopia.** *Iranian journal of parasitology* 2016, **11**(4):549.
- 477 76. Feleke DG, Arega S, Teklewini M, Kindie K, Gedefie A: **Schistosoma mansoni and other**
478 **helminthes infections at Haike primary school children, North-East, Ethiopia: a cross-**
479 **sectional study.** *BMC research notes* 2017, **10**(1):609.
- 480 77. Anuar TS, Salleh FM, Moktar N: **Soil-transmitted helminth infections and associated risk**
481 **factors in three Orang Asli tribes in Peninsular Malaysia.** *Scientific reports* 2014, **4**:4101.
- 482 78. Yami A, Mamo Y, Kebede S: **Prevalence and predictors of intestinal helminthiasis among**
483 **school children in Jimma zone; a cross-sectional study.** *Ethiopian journal of health*
484 *sciences* 2011, **21**(3).
- 485 79. Alemu A, Atnafu A, Addis Z, Shiferaw Y, Teklu T, Mathewos B, Birhan W, Gebretsadik S,
486 Gelaw B: **Soil transmitted helminths and Schistosoma mansoni infections among school**
487 **children in Zarima town, northwest Ethiopia.** *BMC infectious diseases* 2011, **11**(1):189.

80. Amare B, Ali J, Moges B, Yismaw G, Belyhun Y, Gebretsadik S, Woldeyohannes D, Tafess K, Abate E, Endris M: **Nutritional status, intestinal parasite infection and allergy among school children in Northwest Ethiopia.** *BMC pediatrics* 2013, **13**(1):7.
81. Dejene T: **Impact of irrigation on the prevalence of intestinal parasite infections with emphasis on schistosomiasis in Hintallo-Wejerat, North Ethiopia.** *Ethiopian Journal of Health Sciences* 2008, **18**(2).
82. Derso A, Nibret E, Munshea A: **Prevalence of intestinal parasitic infections and associated risk factors among pregnant women attending antenatal care center at Felege Hiwot Referral Hospital, northwest Ethiopia.** *BMC infectious diseases* 2016, **16**(1):530.
83. Legesse M, Erko B: **Prevalence of intestinal parasites among schoolchildren in a rural area close to the southeast of Lake Langano, Ethiopia.** *The Ethiopian Journal of Health Development (EJHD)* 2017, **18**(2).
84. Abera A, Nibret E: **Prevalence of gastrointestinal helminthic infections and associated risk factors among schoolchildren in Tilili town, northwest Ethiopia.** *Asian Pacific journal of tropical medicine* 2014, **7**(7):525-530.
85. Tulu B, Taye S, Amsalu E: **Prevalence and its associated risk factors of intestinal parasitic infections among Yadot primary school children of South Eastern Ethiopia: a cross-sectional study.** *BMC research notes* 2014, **7**(1):848.
86. Feleke H, Medhin G, Abebe A, Beyene B, Kloos H, Asrat D: **Enteric pathogens and associated risk factors among under-five children with and without diarrhea in Wegera District, Northwestern Ethiopia.** *The Pan African Medical Journal* 2018, **29**.
87. Wegayehu T, Adamu H, Petros B: **Prevalence of Giardia duodenalis and Cryptosporidium species infections among children and cattle in North Shewa Zone, Ethiopia.** *BMC infectious diseases* 2013, **13**(1):419.
88. Amor A, Rodriguez E, Saugar JM, Arroyo A, López-Quintana B, Abera B, Yimer M, Yizengaw E, Zewdie D, Ayehubizu Z: **High prevalence of Strongyloides stercoralis in school-aged children in a rural highland of north-western Ethiopia: the role of intensive diagnostic work-up.** *Parasites & vectors* 2016, **9**(1):617.
89. Bitew AA, Abera B, Seyoum W, Endale B, Kiber T, Goshu G, Admass A: **Soil-transmitted helminths and Schistosoma mansoni infections in Ethiopian Orthodox Church students around lake tana, northwest Ethiopia.** *PloS one* 2016, **11**(5):e0155915.
90. Abate A, Kibret B, Bekalu E, Abera S, Teklu T, Yalew A, Endris M, Worku L, Tekeste Z: **Cross-sectional study on the prevalence of intestinal parasites and associated risk factors in Teda Health Centre, Northwest Ethiopia.** *ISRN parasitology* 2013, **2013**.
91. Shumbej T, Belay T, Mekonnen Z, Tefera T, Zemene E: **Soil-transmitted helminths and associated factors among pre-school children in Butajira Town, South-Central Ethiopia: A community-based cross-sectional study.** *PloS one* 2015, **10**(8):e0136342.
92. Ayalew D, Boelee E, Endeshaw T, Petros B: **Cryptosporidium and Giardia infection and drinking water sources among children in Lege Dini, Ethiopia.** *Tropical Medicine & International Health* 2008, **13**(4):472-475.
93. Nyantekyi LA, Legesse M, Belay M, Tadesse K, Manaye K, Macias C, Erko B: **Intestinal parasitic infections among under-five children and maternal awareness about the infections in Shesha Kekele, Wondo Genet, Southern Ethiopia.** *Ethiopian Journal of Health Development* 2010, **24**(3).
94. Gizaw Z, Adane T, Azanaw J, Addisu A, Haile D: **Childhood intestinal parasitic infection and sanitation predictors in rural Dembiya, northwest Ethiopia.** *Environmental Health and Preventive Medicine* 2018, **23**(1):26.
95. Eshetu T, Sibhatu G, Megiso M, Abere A, Baynes HW, Biadgo B, Zeleke AJ: **Intestinal Parasitosis and Their Associated Factors among People Living with HIV at University of Gondar Hospital, Northwest-Ethiopia.** *Ethiopian journal of health sciences* 2017, **27**(4):411-420.

Figure legends

Figure 1:- Flowchart diagram describing selection of studies for the burden of IPIs and association with residence and hand washing practice in Ethiopian children, 2018

Figure 2:- Forest Plot of the 31 studies that quantitatively assessed the pooled prevalence of IPIs, 2018.

Figure 3: Forest Plot of the 6 studies that quantitatively assessed the effect of hand washing before eating on the occurrences of IPIs, 2018

Figure 4:- Forest Plot of the 6 studies that quantitatively assessed the effect of residence on the occurrences of IPIs, 2018.

Additional Files

Additional file 1:- Funnel plot of the included studies for checking publication bias.

Table 1: Descriptive summary of 31 included studies on the prevalence of intestinal parasitic infections and association with residence and hand washing in Ethiopian children.

Authors name	year	Region	Design	Age of subjects	Sample size	No of people with outcome	Prevalence (%)	Quality score
Gemechu et al[62]	2017	SNNP	CS	<5yr	167	29	17.45	7
Girum Tadesse [63]	2001	SNNP	CS	school age	415	113	27.2	9
AidadeLucio et al[64]	2016	Amhara	CS	school age	393	234	59.6	7
Tamirat Hailegebriel[7]	2016	Amhara	CS	school age	359	235	65.5	7
Abossie A et al[65]	2012	SNNP	CS	school age	400	324	81	8
Zemene T et al[49]	2014	Amhara	CS	<5yr	217	38	17.4	7
Agezew A et al[66]	2015	Amhara	CS	<5yr	401	141	35.2	8
Mathewos B et al[67]	2012	Amhara	CS	school age	261	174	66.7	8
Gelaw A et al[19]	2012	Amhara	CS	school age	304	104	34.2	8
Tadege B et al[68]	2015	SNNP	CS	school age	374	254	67.9	8

Mulatu G [16]	2011	SNNP	CS	<5yr	158	42	26.6	7
Beyene G et al [69]	2012	Oromia	CS	school age	260	129	49.6	6
Hailu T et al [70]	2017	Amhara	CS	school age	409	193	47.2	7
Reji P et al [71]	2008	Oromia	CS	school age	358	127	35.5	8
Teferae E et al [72]	2010	Oromia	CS	school age	715	346	48.4	7
Alemayehu B et al [73]	2015	SNNP	CS	school age	503	363	72.2	6
Yirgalem G et al [74]	2013	Oromia	CS	<5yr	374	91	24.3	7
Firdu T et al [47]	2011	SNNP	Case-control	school age	230	84	36.52	8
Worku L et al [48]	2013	Amhara	CS	school age	385	372	96.6	9
Begna T et al [75]	2013	SNNP	CS	school age	492	131	26.6	9
Grimes J et al [50]	2017	SNNP	CS	school age	3729	874	23	7
Daniel G et al [76]	2017	Oromia	CS	school age	279	85	30.5	7
Tilahun A et al [77]	2010	Amhara	CS	school age	384	211	54.9	7
Yami A et al [78]	2008	Oromia	CS	school age	855	403	47.1	9
Abebe Alemu et al [79]	2009	Amhara	CS	school age	319	263	82.4	7
Amare B [80]	2008	Amhara	CS	school age	405	92	22.7	8
Tadesse D [81]	2000	Tigray	CS	school age	800	285	35.6	8
Derso A [82]	2012	Amhara	CS	school age	385	170	44.2	8
Ayalew A et al [9]	2010	Amhara	CS	school age	704	562	79.8	6
Legesse M et al [83]	2003	Oromia	CS	school age	259	217	83.8	6
Alamneh A et al [84]	2014	Amhara	CS	school age	380	170	44.2	7
Tulu et al. [85]	2014	Amhara	CS	School age	340	89	26.2	8
Feleke et al. [86]	2018	Amhara	CS	<5	225	49	21.8	6

Wegayehu et al. [87]	2011	Amhara	CS	school age	284	81	28.5	7
Aranzazu A et al [88]	2016	Amhara	CS	School age	396	306	77.5	6
Bitew et al. [89]	2016	Amhara	CS	School age	384	251	65.6	7
Mulusew A	2014	Amhara	CS	School age	357	245	68.4	7
Abate et al [90]	2013	Amhara	CS	School age	410	255	62.3	7
Aida de Lucio [64]	2016	Amhara	CS	School age	393	216	55	8
Teha Shumbej [91]	2015	SNNP	CS	<5	377	88	23.3	6
Ayalew et al. [92]	2008	Dire-Dawa	CS	School age	655	311	47.5	7
Liza A. [93]	2010	SNNP	CS	<5	288	245	85	8
Gizaw et al. [94]	2018	Amhara	CS	<5	225	58	25.8	6

555 CS=Cross-Sectional, SNNP= Southern Nations, Nationalities, and Peoples

556 **Table 2: Subgroup analysis based on regions and age of participants among diabetes**
557 **patients**

558

Variables	Characteristics	Estimates with 95% CI	P-value
Age of participant	Under-five children	24.23(17.44-31.02)	<0.001
	School-age children	52.43(41.37-63.5)	
Region	Amhara	52.1(37.3-66.8)	<0.001
	Oromia	45.1(30.9-59.3)	
	SNNP	43.9(24.9-62.9)	
	Tigray	35.6(32.2-38.9)	

559 SNNP= Southern Nations, Nationalities, and Peoples

560 **Table 3 the effect of hand washing before eating on intestinal parasitic infections in**
561 **Ethiopian children.**

Authors and year	Region	Hand washing	<u>Intestinal parasites</u>		OR(with 95% CI)
		Before eating	Positive	negative	
Gemechu A et al,2017[62]	Southern	Yes	7	83	1
		NO	22	25	4.70(1.88-11.75)
Ayalew A et al,2010[9]	Amhara	Yes	261	110	1
		No	301	32	3.96(2.58-6)
Abera A et al,2014[84]	Amhara	Yes	135	191	1
		No	35	24	2.06(1.17-3.62)
Tadege B et al,2015[68]	SNNP	Yes	57	156	1
		No	140	21	14.02(8-24.3)
Alemu A et al,2015[66]	Amhara	Yes	6	96	1
		No	86	213	6.46(2.73-15.3)
Hailu et al, 2017[70]	Amhara	Yes	128	227	1
		No	40	14	2.90(1.52-5.53)
Teha et al. 2015[91]	SNNP	Yes	67	186	1
		No	21	191	4.4(2.6-7.6)
Gizew et al. 2018[94]	Amhara	Yes	10	92	1
		No	48	75	5.8(2.7-12.4)
Tegegne E. et al. 2017[95]	Amhara	Yes	22	148	1
		No	43	10	28.9(12.7-65.7)

562 `SNNP=Southern Nations, Nationalities, and Peoples

563

564

565

566

NOTE: Weights are from random effects analysis

Study

%

ID

ES (95% CI)

Weight

Gemechu A et al,2017

4.70 (1.80, 12.28)

9.29

Ayalew A et al,2010

3.90 (2.54, 5.98)

12.65

Abera A et al,2014

2.00 (1.14, 3.52)

11.86

Tadege B et al,2015

14.00 (8.08, 24.26)

11.95

Alemu A et al,2015

6.40 (2.70, 15.15)

9.93

Hailu et al, 2017

2.90 (1.52, 5.53)

11.35

Teha et al. 2015

4.40 (2.59, 7.48)

12.07

Gizew et al 2018

5.80 (2.75, 12.23)

10.69

Tegegne E. et al 2017

28.90 (12.72, 65.68)

10.20

Overall (I-squared = 82.5%, p = 0.000)

5.63 (3.39, 9.37)

100.00

NOTE: Weights are from random effects analysis

.1

1

10