

Supplemental Table 3. Sensitivity analysis applying ranges of thresholds to filter in SNPs in the genetic instrument.

Outcome	P value threshold to filter SNPs	N of SNPs remained	beta (95% CI)	P by inverse variance weighted	MR-Egger intercept P
White matter volume (normalized)	$< 1 \times 10^{-9}$	57	-0.064 (-0.119, -0.01)	0.02	0.095
	$< 1 \times 10^{-10}$	47	-0.092 (-0.149, -0.035)	0.002	0.992
	$< 1 \times 10^{-11}$	40	-0.087 (-0.147, -0.028)	0.004	0.907
	$< 1 \times 10^{-12}$	38	-0.088 (-0.148, -0.028)	0.004	0.922
	$< 1 \times 10^{-13}$	32	-0.091 (-0.154, -0.028)	0.005	0.668
	$< 1 \times 10^{-14}$	29	-0.091 (-0.156, -0.026)	0.006	0.631
	$< 1 \times 10^{-15}$	28	-0.09 (-0.156, -0.024)	0.007	0.615
	$< 1 \times 10^{-16}$	24	-0.074 (-0.143, -0.005)	0.034	0.444
	$< 1 \times 10^{-17}$	19	-0.075 (-0.149, -0.002)	0.045	0.382
	$< 1 \times 10^{-18}$	18	-0.085 (-0.16, -0.011)	0.025	0.560
White matter volume (unnormalized)	$< 1 \times 10^{-9}$	57	-0.033 (-0.061, -0.005)	0.022	0.091
	$< 1 \times 10^{-10}$	47	-0.047 (-0.076, -0.017)	0.002	0.990
	$< 1 \times 10^{-11}$	40	-0.045 (-0.075, -0.014)	0.004	0.922
	$< 1 \times 10^{-12}$	38	-0.044 (-0.076, -0.013)	0.005	0.920
	$< 1 \times 10^{-13}$	32	-0.046 (-0.078, -0.013)	0.006	0.684
	$< 1 \times 10^{-14}$	29	-0.046 (-0.079, -0.012)	0.008	0.632
	$< 1 \times 10^{-15}$	28	-0.045 (-0.079, -0.011)	0.009	0.616
	$< 1 \times 10^{-16}$	24	-0.037 (-0.072, -0.001)	0.042	0.431
	$< 1 \times 10^{-17}$	19	-0.038 (-0.076, 0.000)	0.053	0.337
	$< 1 \times 10^{-18}$	18	-0.043 (-0.082, -0.004)	0.030	0.507
Grey matter volume (normalized)	$< 1 \times 10^{-9}$	57	-0.016 (-0.056, 0.024)	0.438	0.151
	$< 1 \times 10^{-10}$	47	-0.030 (-0.072, 0.012)	0.164	0.945
	$< 1 \times 10^{-11}$	40	-0.032 (-0.076, 0.012)	0.150	0.939
	$< 1 \times 10^{-12}$	38	-0.029 (-0.074, 0.015)	0.196	0.873
	$< 1 \times 10^{-13}$	32	-0.045 (-0.092, 0.002)	0.058	0.847
	$< 1 \times 10^{-14}$	29	-0.048 (-0.096, 0.000)	0.049	0.990
	$< 1 \times 10^{-15}$	28	-0.046 (-0.095, 0.002)	0.062	0.942
	$< 1 \times 10^{-16}$	24	-0.044 (-0.094, 0.007)	0.091	0.903
	$< 1 \times 10^{-17}$	19	-0.04 (-0.094, 0.014)	0.149	0.428
	$< 1 \times 10^{-18}$	18	-0.037 (-0.093, 0.018)	0.184	0.491
Grey matter volume (unnormalized)	$< 1 \times 10^{-9}$	57	-0.009 (-0.037, 0.018)	0.504	0.107
	$< 1 \times 10^{-10}$	47	-0.019 (-0.048, 0.010)	0.202	0.969
	$< 1 \times 10^{-11}$	40	-0.020 (-0.050, 0.010)	0.188	0.955
	$< 1 \times 10^{-12}$	38	-0.018 (-0.048, 0.013)	0.256	0.81
	$< 1 \times 10^{-13}$	32	-0.028 (-0.060, 0.004)	0.085	0.793
	$< 1 \times 10^{-14}$	29	-0.031 (-0.064, 0.002)	0.065	0.989
	$< 1 \times 10^{-15}$	28	-0.029 (-0.062, 0.004)	0.085	0.924
	$< 1 \times 10^{-16}$	24	-0.028 (-0.062, 0.007)	0.119	0.915
	$< 1 \times 10^{-17}$	19	-0.026 (-0.063, 0.011)	0.168	0.496
	$< 1 \times 10^{-18}$	18	-0.025 (-0.062, 0.013)	0.202	0.558

CI = confidence interval

The SNPs reaching the designated P value threshold remained in the genetic instrument for atrial fibrillation in each analysis. The causal estimates were derived by the inverse variance weighted method, and the MR-Egger intercept P values were calculated to detect the presence of significant pleiotropy.

Supplemental Table 6. Causal estimates from stroke on brain white or grey matter volume.

Outcome	Genetic instrument	MR method	MR-Egger intercept P value	beta (95% CI)	P value
Genetic instrument for stroke including all genome-wide significant SNPs reported in the previous GWAS (N = 27)	White matter volume (normalized)	Inverse variance weighted	0.931	-0.003 (-0.059, 0.054)	0.925
		MR-Egger		-0.006 (-0.103, 0.091)	0.903
		Weighted median		0.007 (-0.075, 0.089)	0.864
	White matter volume (unnormalized)	Inverse variance weighted	0.845	-0.002 (-0.031, 0.027)	0.877
		MR-Egger		-0.006 (-0.055, 0.043)	0.807
		Weighted median		0.004 (-0.038, 0.045)	0.865
	Grey matter volume (normalized)	Inverse variance weighted	0.114	0.019 (-0.022, 0.061)	0.36
		MR-Egger		-0.024 (-0.091, 0.043)	0.482
		Weighted median		0.006 (-0.054, 0.066)	0.841
	Grey matter volume (unnormalized)	Inverse variance weighted	0.076	0.013 (-0.016, 0.041)	0.377
		MR-Egger		-0.021 (-0.067, 0.025)	0.375
		Weighted median		-0.007 (-0.049, 0.035)	0.742
Genetic instrument for stroke after excluding confounder-associated SNPs (N = 18)	White matter volume (normalized)	Inverse variance weighted	0.930	0.011 (-0.050, 0.073)	0.719
		MR-Egger		0.015 (-0.085, 0.115)	0.773
		Weighted median		0.014 (-0.066, 0.095)	0.729
	White matter volume (unnormalized)	Inverse variance weighted	0.963	0.005 (-0.027, 0.037)	0.749
		MR-Egger		0.004 (-0.047, 0.056)	0.874
		Weighted median		0.007 (-0.039, 0.052)	0.774
	Grey matter volume (normalized)	Inverse variance weighted	0.259	0.014 (-0.032, 0.06)	0.547
		MR-Egger		-0.02 (-0.094, 0.053)	0.594
		Weighted median		0.006 (-0.059, 0.071)	0.848
	Grey matter volume (unnormalized)	Inverse variance weighted	0.176	0.011 (-0.020, 0.042)	0.492
		MR-Egger		-0.018 (-0.068, 0.033)	0.503
		Weighted median		0.000 (-0.043, 0.042)	0.984

CI = confidence interval