1 Investigating the origin of the Belgian second SARS-CoV-2 wave by using
2 (pre)admission screening samples.

3 Naesens Reinout^{1,2}, Heireman Laura¹, Vandamme Sarah³, Willems Philippe^{4,5}, Van Herendael
 4 Bruno^{4,5}, Verstrepen Walter¹, De Schouwer Pieter¹, Bruynseels Peggy^{1,2}

5 Affiliations:

6 ^{1.} Department of Medical Microbiology, ZiekenhuisNetwerk Antwerpen, B-2020
 7 Antwerp, Belgium

8 ^{2.} Department of Infection Prevention and Control, ZiekenhuisNetwerk Antwerpen, B-
 9 2020 Antwerp, Belgium

10 ^{3.} Department of Medical Microbiology, University Hospital Antwerp, B-2650 Edegem,
 11 Belgium

12 ^{4.} Department of Medical Microbiology, GasthuisZusters Antwerpen, B-2060 Antwerp,
 13 Belgium

14 ^{5.} Department of Infection Prevention and Control, GasthuisZusters Antwerpen, B-2060
 15 Antwerp, Belgium

16

17 **Corresponding Author:** Reinout Naesens **Corresponding Author Email:**
 18 reinout.naesens@zna.be

19   **Abstract**

20   The goal of this study was to estimate rates of SARS-CoV-2 carriership and viral loads in the
21   general Antwerp population and to compare the estimated prevalences and incidences with
22   governmental data (numbers of detected positive cases, stringency measure index) in order to
23   evaluate the dynamics leading to the second wave. We used (pre)admission screening results
24   from the major Antwerp hospitals for estimating community prevalences and incidences.
25   43.545 samples were included (April – November 2020). High SARS-CoV-2 carriership rates
26   (mean week prevalence of 1.3%) were found in the general Antwerp population. 35.4% of
27   positive cases carried high viral loads. Only a small proportion (15.3%) of the viral circulation
28   was detected by the nationally implemented testing policy. In the weeks before the second
29   Belgian wave, increasing prevalences and incidences were found, together with country-wide
30   easing of restriction measures. In our opinion these findings have led to origin of the second
31   viral wave.

32   **Keywords:** SARS-CoV-2; Belgium; second wave; driver; contact-tracing; prevalence;
33   incidence

34   **Conflict of interest:** None

35   **Funding statement:** No funding

36 INTRODUCTION

37 Since March 2020, the world has been in the grips of a pandemic caused by the Severe Acute
38 Respiratory Syndrome virus type 2 (SARS-CoV-2) leading to excess mortality and morbidity.
39 The virus has a detrimental impact on both economic and psycho-social well-being¹.

40 Belgium was highly impacted, having had to deal with two waves up to November 2020: the
41 first from the beginning of March to the end of May, the second during the fall (showing a
42 decline in November). The first wave was contained by implementing a full lockdown starting
43 on 14 March 2020. The first measures to ease restrictions were introduced on 4 June 2020.

44 Despite warnings from various Belgian key expert virologists and epidemiologists in
45 September 2020, the government decided to ease restrictions further, having been encouraged
46 by various pressure groups and public opinion. Belgium faced its second wave as from
47 October 2020 (Figure 1).

48 Until now, little is known about the dynamics and prevalence of asymptomatic carriership and
49 viral loads in the general Belgian population². Furthermore, the role of the different age
50 categories in the viral spread has not been fully elucidated³.

51 Since the major Antwerp hospitals implemented extensive (pre-)admission screening of
52 asymptomatic cases in April 2020 (eg. prior to elective surgery or endoscopic procedures),
53 and have continued to do so, they possess valuable information regarding the actual
54 prevalence of the virus in the general population throughout the pandemic period.

55 The goal of this study was to determine rates of carriership and viral loads in the general
56 population and to evaluate the dynamics leading to the onset of the second Belgian wave.

57

58 METHODS

59 DATA COLLECTION

60 During the period 27 April to 15 November 2020, we collected SARS-CoV-2 positivity data
 61 from (pre)admission screenings from the following hospitals: the Antwerp University
 62 Hospital (UZA), the multiple site public hospital ZiekenhuisNetwerk Antwerpen (ZNA), and
 63 the major multiple site private Antwerp hospital GemeenschapsZusters Antwerpen (GZA).
 64 Together they account for approximately 90% of hospital bed capacity (3697 hospital beds) in
 65 the city of Antwerp.

66 (Pre-)admission screenings were defined as follows: pre-hospital screenings prior to elective
 67 admissions (GZA, UZA, ZNA), and systematic admission screenings in asymptomatic
 68 patients admitted for other reasons than COVID-19 (GZA, UZA). Screenings of high/low risk
 69 contacts or individuals with symptoms were excluded.

70 A distinction in test indications was obtained by implementing different sample flows per
 71 indication (pre-admission screenings were performed by a nurse at patients' homes for the
 72 inclusions from UZA and ZNA, and screenings performed in a drive-through swab laboratory,
 73 or during a pre-visit at the swab laboratory, from UZA and GZA respectively). Unique test
 74 codes were also used by GZA and UZA for registering the swab samples in the Laboratory
 75 Information System (LIS).

76 Samples were analysed in the laboratory attached to the hospital of the prescribing physician.
 77 All laboratories are certified by the Belgian government (BELAC). The following molecular
 78 techniques and platforms were used:

79 In the clinical laboratory of GZA, nucleic acids were extracted on STARlet IVD® (Seegene
 80 INC., Seoul, Korea) using the Viral DNA/RNA C-kit. Subsequently SARS-CoV-2 qPCR
 81 (CDC N1 target) was performed on a Quantstudio 7 flex qPCR cycler® of Thermo Fisher

82 Scientific (Waltham, MA, USA) according to the protocol published by the Centers of
83 Disease Control and Prevention (CDC).

84 The laboratory of UZA used the Xpert Xpress SARS-CoV-2 test detecting E and N2 target on
85 the GeneXpert® Platform (Cepheid, USA) as per manufacturer's instructions. Alternatively,
86 an in-house PCR was performed detecting E target according to Corman *et al.* 2020 with
87 extraction on NucliSens EasyMag® (Biomérieux, France) or QiaSymphony® (Qiagen, The
88 Netherlands) and amplification with Cobas LightCycler 480 II® or Cobas z480 (Roche
89 Molecular Diagnostics, Switzerland). Alternatively, a BD MAX SARS-CoV-2 kit was used
90 detecting N1 and N2 target with an all-in one extraction amplification on the BD MAX®
91 platform (Becton Dickinson, USA).

92 In the laboratory of ZNA, nucleic acids were extracted on MagNA Pure 96 using the MagNA
93 Pure 96 DNA and Viral NA Small Volume Kit (Roche Molecular Diagnostics, Switzerland).
94 Subsequently SARS-CoV-2 qPCR (N1 target) was performed on a LightCycler 480 qPCR
95 cyclor according to the protocol published by the CDC. Alternatively, nucleic acids were
96 extracted on KingFisher Flex using the MagMAX Viral/Pathogen II Nucleic Acid Isolation
97 Kit. These extracts were analysed on a Quantstudio 5 qPCR cyclor using the TaqPath™
98 COVID-19 CE-IVD RT-PCR Kit (ThermoFisher Scientific).

99 The parameters postal code, age, PCR result (associated with PCR platform used) and cycle
100 threshold (Ct) value (or alternatively Crossing point (Cp)) if available (GZA, ZNA) were
101 retrieved from the LIS for each patient, and transferred to a central ZNA-based and secured
102 database.

103 The following data were collected from the Belgian governmental scientific institution,
104 Sciensano⁴: absolute numbers (weekly) of positively detected cases for Antwerp province,

admitted patients at the hospital for Antwerp province, and returning travelers in August 2020 with positivity rate (only available on national level).

The following data were collected from the Flemish Agency for Care and Health (governmental agency responsible for the contact-tracing policy) for Antwerp province by using the information from the web-based and secured database (Controle-toren)⁵: percentages (weekly) of successfully (by phone) contacted positive cases in the Antwerp province. Age distributions for the Antwerp province were obtained from Statistiek Vlaanderen⁶.

We used the Oxford COVID-19 Government Response Stringency index to estimate governmental measures. The index goes from zero (no restrictions) to hundred (strictest)⁷.

DATA ANALYSES AND DEFINITIONS

Cases with domicile outside the Antwerp province were excluded for further analysis. SARS-CoV-2 carriership was estimated by calculation of the weekly asymptomatic positivity rate in the study population. The incidence was calculated by using an estimated duration of PCR positiveness of seventeen days as was found by Cevik and colleagues⁸. The percentage of the cohort infected by SARS-CoV-2 (for the whole investigated period) was calculated by adding up all estimated week incidences. By dividing the absolute weekly numbers of positively tested cases in the province by the number of province inhabitants (1.858 million) in the province, the percentage of positively tested patients in the Antwerp province was calculated. The percentage of weekly new COVID-19 hospital admitted patients in the Antwerp province was calculated by dividing the absolute weekly number of admitted patients by the number of inhabitants in the province.

Viral loads (only available for GZA and ZNA) were calculated by extrapolation of the Ct/Cp values corresponding to 5000 genome equivalents per mL of the AccuPlex™ SARS-CoV-2 full genome reference material (Seracare). A high viral load was defined as $\geq 256\,000$ viral copies/mL⁹. Ratios of positive cases (on a weekly basis; all inclusions) with high viral loads were plotted on the same timeline as the plotted estimated incidences.

Mean positivity ratios per age decade (0-10 y, 11-20 y, 21-30 y, 31-40 y, 41-50 y, 51-60 y, 61-70 y, 71-80 y, 81-90 y, 90+ y) were investigated for the investigated cohort. Positivity ratios of carriership were calculated for preschools (0-4 y), primary schools (5-11 y) and secondary schools (12-18 y). The mean positivity ratio was also assessed for (pre)school children (0-18 y) and the elderly (+80 y). Box-whisker plots represent the viral load among different age categories.

Ethical approvals were obtained by the hospital Institutional Boards (GZA: Approval N° 200906RETRO; UZA: Approval N°001355; ZNA: Approval N° 5416).

RESULTS

Overall, 43,545 cases were identified of which 38,763 cases (89.0%) were included. The results of 4782 cases (11.0%) were excluded since domicile (based on postal code) was outside Antwerp province. GZA, UZA, and ZNA accounted for respectively 53.2%, 34.6% and 10.7% of included cases. The overall SARS-CoV-2 positivity rate was 1.3% (n=520). Younger age groups (<21 years old) were underrepresented in the investigated cohort as compared to the Antwerp province (9.0 % versus 22.3%; $p < 0.01$). Age groups of > 70 years old were overrepresented in the cohort (25.0 % versus 14.1 %, $p < 0.01$). The distribution of cases was not equally dispersed throughout Antwerp province with an underweight of cases from the city of Antwerp versus the Antwerp province (20.8% of cases had domicile in the city of Antwerp, versus 26.8 % of inhabitants for the whole Antwerp province $p < 0.01$). Cases with domicile in the city of Antwerp (n = 8067) had an overall positivity rate of 1.8%, whereas cases with domicile outside the city of Antwerp (n = 30696) had an overall positivity rate of 1.2% ($p < 0.01$).

Estimated carriership (with high viral loads), estimated weekly new cases and number of admissions are shown in Figure 1. Calculated weekly cohort carriership varied between 0.1 % (week of 29 June) and 8.6 % (week of 26 October). The mean week percentage was 1.5%. The calculated ratio of carriership with high viral loads varied between 0% (week of 11 May, 18 May, 1 June, 22 June, 29 June, 6 July, 13 July and 24 August) and 6.7 % (week of 26 October). The mean week ratio was 0.8%. Overall, the percentage of positive carriers with high viral loads was 35.4 %. The calculated weekly incidence varied between 0.0% (see Figure 2) and 6.5 % (week of 19 October). The mean week incidence was 0.7%. Overall, we calculated that 20.3 % of the investigated cohort became infected with SARS-CoV-2 in the

period from 27 April to 15 November. 3.1 % of Antwerp province inhabitants tested positive in the investigated period (Sciensano data). In August 2020, 1.4 million Belgians (12.2% of Belgian population) filled in a Passenger Locator Form. 2% of travelers tested positive for SARS-CoV-2. On population level, this equals a potential increase of 0.2% in SARS-CoV-2 carrier-rate.

The percentage of weekly new hospital admissions varied between 0.00% (see Figure 1) and 0.02% (week of 26 October, 2 November, and 9 November) of the Antwerp province population. The mean ratio of weekly detected cases versus the estimated weekly incidence was 29.1 % (range: 2.9 % - 100 %). The percentage of successfully contacted positive cases varied between 54.0% (week of 22 June) and 94.0 % (week of 26 October) for the Antwerp province with a mean percentage of successfully contacted positive cases of 83.0% (data only available from 1 June to 15 November since registration only started on 1 June).

The SARS-CoV-2 positivity rate according to age is 0.9% (95 % Confidence Interval (CI): 0.4 – 1.5 %) (n= 1292) for daycare children (0-4 years), 1.0% (95 % CI: 0.5 – 1.9 %) (n= 877) for primary school children (5-12 years), 0.8% (95 % CI: 0.3 – 1.6 %) (n = 878) for secondary school children (13-18 years), 0.9% (95 % CI: 0.6 – 1.3 %) (n= 3047) for overall school-aged civilians (0-18 years), 1.4% (95 % CI: 1.3 – 1.5 %) (n = 35637) for adults (>18 years) and 2.6% (95 % CI: 2.1 – 3.2 %) (n = 4033) for the elderly (>80 years). The distribution of viral loads and ratios of carriers with high viral loads among different age categories are presented in Figure 2 and 3 respectively.

The results of the Oxford Stringency Index are shown in Figure 1.

DISCUSSION

Numerous reports provide data indicating that asymptomatic (or pre-symptomatic) subjects can transmit COVID-19 with high efficiency^{10,11}. In the living review paper of Buitrago-Garcia, the secondary attack rate is only slightly lower in contacts of people with asymptomatic infection than in those with symptomatic infection (relative risk 0.35, 95% CI 0.10-1.27)¹². Mathematical modelling studies (not peer reviewed) have suggested that asymptomatic individuals might be major drivers for the growth of the COVID-19 pandemic¹³.

We found high positivity ratios of asymptomatic carriership in our investigated cohort of Antwerp inhabitants throughout the investigated pandemic period (27 April to 14 November 2020). Lowest positivity ratios were seen at the end of June (week of 29 June): after a period of a full lockdown and only minor easing of containment measures, SARS-CoV-2 positivity ratios were found to be at levels of 1 out of 1000 samples. Similar or even far lower prevalences led to massive screening programs in China in order to stop further viral spread¹⁴.

Taking into account the SARS-CoV-2 key transmission epidemiological parameters (estimated R_0 factor of 2.0 and a dispersion factor of 0.10), a strategy accepting prevalences above 1% (the overall average in our cohort being 1.3 %) together with the simultaneously reopening of a locked community were, in our opinion, the perfect ingredients for rapid escalation towards a subsequent second national wave¹⁵. This finding is important as Belgium has chosen a strategy of accepting significant circulation of the virus within the population without having an accurate insight of past and actual viral prevalences (in contrast to eg. Luxemburg and China, where thorough search and/or isolation programs have been set up)^{14,16}. Also, pressure is already increasing to further ease the contingency measures taken in order to control the second wave.

The finding that, due to the implemented testing and tracing strategy, only marginal numbers of positive cases were detected (we estimated that, by extrapolation of cohort data, 20.3 % of the Antwerp population contracted the virus in the study period, but only 3.1 % (15.3% of estimated positive cases) were effectively tested positively) and subsequently traced (83, range: 54 % - 94 %), even in the periods of upward movement of the curves, was probably another important contributing factor to the uncontrollable rapid further viral spread seen in October 2020. The hypothesis of underestimated viral circulation in Belgium has also been put forward by Herzog and colleagues, by analyzing epi-serological data¹⁷. Since we only measured the base-line (population wide) prevalence, our estimation may even be an underestimation of the real viral circulation, since outbreaks and clusters are not represented in our analysis, but may have been picked up in documented testing numbers (Sciensano). On the other hand, the underrepresentation of the younger age groups versus the overrepresentation of the older age groups in the investigated cohort leads to an upward bias.

Furthermore, backward tracing has not been implemented optimally to date by the governmental task force¹⁸. We argue that the Belgian testing and tracing strategy was not able to contain and monitor the pandemic in Belgium adequately, this being one of the major ingredients of a failing containment policy leading to the second national wave.

The province of Antwerp was one of the three Belgian provinces (out of 10) with highest number of tests per inhabitants^{4,5}. There is no reason for postulating an exceptional role of Antwerp province in Belgium as being a highly affected province. Inversely, the southern part of the country was affected more severely in the second wave, as shown in the governmental Sciensano data.

Interestingly, we also found higher proportions of carriers in the city of Antwerp (1.8%) than in the province of Antwerp (1.2%; $p < 0.01$). This reflects a national trend: the second wave was more prevalent in crowded bigger capital cities⁴.

With regard to positivity ratios according to age category, we found high numbers (around 1 % throughout the investigated period) in school children and young adults (0-30 years).

Positivity ratios of young adults (21-40 years-old) were significantly higher than the adult population aged 51-70 years ($p < 0.01$).

Although the role of children as main drivers of transmission is still debated, our finding may be another argument for active transmission in these age groups^{20,21,22}. A recent seroprevalence survey in two Belgian municipalities (one highly affected, the other hardly affected) also showed equivalent proportions of infections in primary school children versus adolescents in the first three years of secondary education²³.

On the other hand, if children were equally susceptible, one might expect even higher rates of carriership, since children were less restricted in the number of contacts (summercamps, schools were open from 1 September 2020). Since we were not able to correct for number of contacts, we cannot further determine the value of this hypothesis.

With regard to age distribution, another interesting finding is the high overall percentage of positivity ratios in the elderly (+70 y). In our hypothesis, this is the reflection of the augmented transmission risks in residential care homes.

We also found significant proportions of carriers with high viral loads (on average 35.4 % of all inclusions). In the week of 26 October, we estimate that 7.0% of Antwerp inhabitants were carrying high viral loads. Evaluating the whole investigated period, on average 0.7% of the population was carrying high viral loads. Although the link of viral load with contagiousness has not been established, this might have been an important factor in the spread of the virus¹⁹.

Viral loads in younger age groups were not lower than viral loads in the working age population. (Figure 2) In another non-peer-reviewed publication, it was also shown that there is no significant difference between viral loads in 1-20 year-olds as compared to 21-100 year-

old adults²⁴. Furthermore, another study suggests that the viral load in children below 5 years of age with mild to moderate COVID-19 symptoms is higher than in older children and adults²⁵.

In our hypothesis, positive children may have further introduced the virus into households, which suggests that this is a potentially underestimated factor in the derailing of the situation in September 2020.

Interestingly, we found significantly higher proportions of carriers with high viral loads in the elder age categories. The combination of higher prevalences and higher ratios of carriers with high viral loads in this age group might be an important driver of the abundance of outbreaks and associated excess morbidity and mortality seen in retirement homes.

Overall, not recognizing the presence and potential important role of the asymptomatic carriers with higher viral loads may also have been another ingredient which led to the subsequent second wave.

In our opinion, the role of the returning travelers was very limited. In August 12.2 % of the Belgian population filled in a Passenger Locator form. About 2% of returning travelers tested positive. On a population level, this may have led to an increase of 0.2% in carriership rates.

On the positive side, we found that by implementing strict contingency measures (i.e. following the tightening and easing advice of experts) even substantial viral circulation of around 1% can be controlled: the ratio of positive carriers was contained in the period between 27 April and 14 September (19 weeks) without overstressing health care capacities. However, the efforts required to maintain controllable containment were enormous, thereby putting high pressure on social (limited number of contacts) and economic well-being (Oxford Stringency Index not below 50.0 between 27 April and 14 September).

Finally, we come to our last driver of the second wave, probably the most important one. Mid to end September, key Belgian epidemiologists, virologists and biostatisticians warned the temporary government of the possible rapid escalation in numbers of new infections and admissions, based on available data. The government decided not to follow their advice, and implemented a strategy that eased restriction measures (Oxford Stringency Index on 14 September: 52.8; 5 October: 47.2) in a time when there was a steadily increasing number of new infections (incidence increased from 0.9% the week of 14 September to 2.3% the week of 5 October). They were being encouraged to carry out this strategy by various pressure groups. In October, Belgium was facing a second wave with a rapid increase in numbers of COVID admissions, thereby highly impacting the health care system and hospital facilities.

The limitations of this study include that (weak positive) asymptomatic cases could have corresponded to previous infections with persistent existence of detectable RNA. This may have biased the results. However, the rapid decrease of positivity ratios after the implementation of extra measures (second pandemic wave) suggests we predominantly included active infections. Although the phenomenon of prolonged shedders does certainly exist, our data suggest that it is relatively insignificant and does not affect the major findings in this paper. Also, the median duration of positive results has been shown to be two to three weeks⁴. Another limitation is that we did not follow up positive/negative cases. We did not link the data to secondary COVID-19 cases, thus not establishing the contagiousness of positive cases. The screenings were linked to hospital procedures/admissions. Although we can accept these screenings as random samples from the community, the cohort is not a representative sample from the community: data points were not equally distributed in the Antwerp province, and age distributions were not representative for the Antwerp province (underestimation of younger age groups, and overestimation of older age groups).

In conclusion, we found that SARS-CoV-2 was significantly present in the Antwerp province population in the period after the first wave. The virus circulated in all age groups with a predominance in the elderly. 1.3% of the investigated cohort tested positive for SARS-CoV-2 with 35% carrying high viral loads. The testing and contact-tracing policy only detected a small proportion of the real viral circulation. The restriction measures were eased in a period of increasing incidences. The combination of these factors were, in our opinion, the ingredients for the onset of the second wave. If Belgium chooses to accept a certain viral circulation, a very strict barometer has to be put into place, in order to prevent further worsening of the pandemic situation. The positivity rates in (pre)admission screening results may be an important tool;

Acknowledgements: None

Author Bio:

Dr. Reinout Naesens is Head of the Infection Prevention and Control Department and member of the Medical Microbiology Department of the ZNA group, the largest public hospital in Belgium.

REFERENCES

1. World Health Organisation, World Health Organisation situation reports, <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>. Accessed 2 december 2020.
2. Byambasuren O, Cardona M, Bell K, Clark J, McLaws M-L, Glasziou P. Estimating the extent of true asymptomatic COVID-19 and its potential for community transmission: systematic review and meta-analysis. medRxiv. 2020; (published online September 13.) (preprint) doi: <https://doi.org/10.1101/2020.05.10.20097543>
3. Castagnoli R, Votto M, Licari A, Brambilla I, Bruno R, Perlini S, et al. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection in Children and Adolescents: A Systematic Review. JAMA Pediatr. 2020;174:882-889. doi:10.1001/jamapediatrics.2020.1467.
4. Sciensano, <https://epistat.wiv-isp.be/covid/>. Accessed 2 december 2020.
5. Toezicht Volksgezondheid, <https://zorgatlas.vlaanderen.be>. Accessed 2 december 2020.
6. Statistiek Vlaanderen, <https://www.statistiekvlaanderen.be/nl/bevolking-naar-leeftijd-en-geslacht>. Accessed 2 december 2020.
7. Oxford COVID-19 Government Response Tracker, <https://covidtracker.bsg.ox.ac.uk/>. Accessed 2 december 2020.
8. Cevik M, Tate M, Lloyd O, Maraolo AE. SARS-CoV-2, SARS-CoV, and MERS-CoV viral load dynamics, duration of viral shedding, and infectiousness: a sytematic review and meta-analysis. Lancet Microb. 2020. Online first. doi: [https://doi.org/10.1016/S2666-5247\(20\)30172-5](https://doi.org/10.1016/S2666-5247(20)30172-5).

9. Bullard J, Dust K, Funk D, Strong JE, Alexander D, Garnett L, et al. Predicting infectious SARS-CoV-2 from diagnostic samples. *Clin Infect Dis*. 2020;22:ciaa638. doi: 10.1093/cid/ciaa638.
10. Koh WC, Naing L, Chaw L, Rosledzana MA, Alikhan MF, Jamaludin SA, et al. What do we know about SARS-CoV-2 transmission? A systematic review and meta-analysis of the secondary attack rate and associated risk factors. *PLoS One*. 2020;15(10):e0240205. doi: 10.1371/journal.pone.0240205.
11. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Euro Surveill*. 2020;25(10):2000180. doi: 10.2807/1560-7917.ES.2020.25.10.2000180.
12. Buitrago-Garcia DC, Egli-Gany D, Counotte MJ, Hossmann S., Imeri H, Ipekci AM, et al. Asymptomatic SARS-CoV-2 infections: a living systematic review and meta-analysis. *medRxiv*. 2020; (published online July 28.) (preprint) doi: <https://doi.org/10.1101/2020.04.25.20079103>.
13. Dobrovolny HM. Modeling the role of asymptomatics in infection spread with application to SARS-CoV-2. *PLoS One* 2020; 15:e0236976. doi: 10.1371/journal.pone.0236976.
14. Health Commission of Hubei Province. Daily report on epidemic situation of COVID-19 in Hubei province. <http://wjw.hubei.gov.cn/bmdt/ztl/fkxxgzbdgrfyyq/xxfb/index.shtml>. Accessed 2 december 2020.
15. Kucharski AJ, Russell TW, Diamond C, Liu Y, Edmunds J, Funk S, Eggo RM; Centre for Mathematical Modelling of Infectious Diseases COVID-19 working group. Early

- 373 dynamics of transmission and control of COVID-19: a mathematical modelling study.
374 Lancet Infect Dis. 2020; pii:S1473-3099(20)30144-4.
- 375 16. <https://covid19.public.lu/en/testing.html> Accessed 2 december 2020.
- 376 17. Herzog S, Abrams S, Wouters I, Ekinici E, Pateet L, Coppens A, et al. Seroprevalence
377 of IgG antibodies against SARS coronavirus 2 in Belgium – a serial prospective cross-
378 sectional nationwide study of residual samples. medRxiv. 2020; (published online
379 October 1.) (preprint) doi: <https://doi.org/10.1101/2020.06.08.20125179>.
- 380 18. Endo A; Leclerc QJ, Knight GM, Medley GF, Atkins KE, Funk S, et al. Implication of
381 backward contact tracing in the presence of overdispersed transmission in COVID-19
382 outbreaks. Wellcome Open Res. 2020;5:239. doi: 10.12688/wellcomeopenres.16344.1.
- 383 19. Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, et al. SARSCoV-2 Viral Load
384 in Upper Respiratory Specimens of Infected Patients. N Engl J Med. 2020;382:1177-
385 1179. doi: <https://www.nejm.org/doi/full/10.1056/NEJMc2001737>.
- 386 20. Jones TC, Mühlemann B, Veith T, Biele G, Zuchowski M, Hoffmann J, et al. An
387 analysis of SARS-CoV-2 viral load by patient age. medRxiv. 2020; (preprint) doi:
388 <https://doi.org/10.1101/2020.06.08.20125484>.
- 389 21. Viner RM, Russell SJ, Croker H, Packer J, Ward J, Stansfield C, et al. School closure
390 and management practices during coronavirus outbreaks including COVID-19: a rapid
391 systematic review. Lancet Child & Adolescent Health. 2020;4(5):397-404. doi:
392 10.1016/S2352-4642(20)30095-X.
- 393 22. Heavey L, Casey G, Kelly C, Kelly D, McDarby G. No evidence of secondary
394 transmission of COVID-19 from children attending school in Ireland, 2020. Euro
395 Surveillance: bulletin Europeen sur les maladies transmissibles = European
396 communicable disease bulletin. 2020;25(21). doi: 10.2807/1560-
397 7917.ES.2020.25.21.2000903.

- 398 23. Sciensano, [https://www.sciensano.be/sites/default/files/limburg-validation-sars-](https://www.sciensano.be/sites/default/files/limburg-validation-sars-cov2_report_20201112_final.pdf)
399 [cov2_report_20201112_final.pdf](https://www.sciensano.be/sites/default/files/limburg-validation-sars-cov2_report_20201112_final.pdf). Accessed 2 December 2020.
- 400 24. Jones TC, Mühlemann B, Veith T, Zuchowski M, Hofmann J, Stein A, et al. An
401 analysis of SARS-CoV-2 viral load by patient age. Available from:
402 [https://zoonosen.charite.de/fileadmin/user_upload/microsites/m_cc05/virologie-](https://zoonosen.charite.de/fileadmin/user_upload/microsites/m_cc05/virologie-ccm/dateien_upload/Weitere_Dateien/analysis-of-SARS-CoV-2-viral-load-by-patient-age.pdf)
403 [ccm/dateien_upload/Weitere_Dateien/analysis-of-SARS-CoV-2-viral-load-by-patient-](https://zoonosen.charite.de/fileadmin/user_upload/microsites/m_cc05/virologie-ccm/dateien_upload/Weitere_Dateien/analysis-of-SARS-CoV-2-viral-load-by-patient-age.pdf)
404 [age.pdf](https://zoonosen.charite.de/fileadmin/user_upload/microsites/m_cc05/virologie-ccm/dateien_upload/Weitere_Dateien/analysis-of-SARS-CoV-2-viral-load-by-patient-age.pdf). Accessed 2 December 2020.
- 405 25. Heald-Sargent T, Muller WJ, Zheng X, Rippe J, Patel AB, Kociolek LK. Age-Related
406 Differences in Nasopharyngeal Severe Acute Respiratory Syndrome Coronavirus 2
407 (SARS-CoV-2) Levels in Patients With Mild to Moderate Coronavirus Disease 2019
408 (COVID-19). JAMA Pediatr. 2020;174(9):902-903. doi:
409 [10.1001/jamapediatrics.2020.3651](https://doi.org/10.1001/jamapediatrics.2020.3651).

410 Table 1 Age distribution of included cases and age distribution of the province of Antwerp.

Age category	Number of inclusions	Percentage of inclusions	Percentage of inhabitants in the Antwerp province	<i>p</i>-value (difference in rates of inclusions versus the Antwerp province)
0-10 y	1971	5.1	11.3	<i>p</i> <0.01
11-20 y	1493	3.9	11.0	<i>p</i> <0.01
21-30 y	3942	10.2	12.1	<i>p</i> <0.01
31-40 y	5110	13.2	13.1	<i>p</i> =0.83
41-50 y	4376	11.3	12.7	<i>p</i> <0.01
51-60 y	5993	15.5	14.0	<i>p</i> <0.01
61-70 y	6194	16.0	11.8	<i>p</i> <0.01
71-80 y	5651	14.6	8.2	<i>p</i> <0.01
81-90 y	3450	8.9	4.9	<i>p</i> <0.01
+90 y	583	1.5	1.0	<i>p</i> =0.23

411

412 Table 2. Overall positivity rate of SARS-CoV-2 in asymptomatic carriers according to age
413 groups.

414

Age category	Overall percentage of positive tests	95% CI
0 - 10 y	1.0	0.6 to 1.5
11 - 20 y	1.3	0.8 to 2.0
21 - 30 y	1.6	1.3 to 2.1
31 - 40 y	1.6	1.3 to 2.0
41 - 50 y	1.2	0.9 to 1.6
51 - 60 y	1.0	0.7 to 1.2
61 - 70 y	0.9	0.7 to 1.1
71 - 80 y	1.2	0.9 to 1.5
81 - 90 y	2.7	2.2 to 3.3
> 90 y	2.6	1.4 to 4.2

415

---

416 Figure 1. Estimated prevalence and incidence of SARS-CoV-2 carriers, estimated prevalence
417 of carriers with high viral loads, percentages of cases with confirmed a positive SARS-CoV-2
418 test and percentages of confirmed hospital admitted COVID cases in the Antwerp province.

419 Figure 2. Viral loads according to age groups in asymptomatic carriers (viral loads only
420 available for GZA and ZNA).

421 Figure 3. Proportion of asymptomatic carriers with high viral loads according to age category.

medRxiv preprint doi: <https://doi.org/10.1101/2020.12.09.20246462>; this version posted December 24, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Distribution of log copies/mL SARS-CoV-2 in asymptomatic patients among different age categories

Age category	Proportion of carriers with high viral loads (%)	95 % CI
0-10y	26.3	9.1-51.2
11-20y	21.1	6.1-45.6
21-30y	26.2	16.0-38.5
31-40y	21.0	12.7-31.5
41-50y	23.1	12.5-36.8
51-60y	19.0	9.9-31.4
61-70y	61.1	46.9-74.1
71-80y	43.1	30.8-56.0
81-90y	64.1	53.5-73.9
+90y	60.0	32.2-83.7