Use of Diagnosis Codes to Find Blood Transfusion Adverse Events in Electronic Health Records

Roselie A. Bright¹; Susan J. Bright-Ponte²; Lee Anne Palmer²; Summer K. Rankin³; Sergey Blok³

¹ Office of the Commissioner, Food and Drug Administration (FDA), Silver Spring, MD, USA.

Corresponding author: Roselie.Bright@fda.hhs.gov, 240-429-3972.

² Center for Veterinary Medicine, FDA, Rockville, MD, USA

³ Booz Allen Hamilton, McLean, VA, USA

ABSTRACT

Background: Electronic health records (EHRs) and big data tools offer the opportunity for surveillance of adverse events (patient harm associated with medical care). We chose the case of transfusion adverse events (TAEs) and potential TAEs (PTAEs) because 1.) real dates were obscured in the study data, and 2.) there was emerging recognition of new types during the study data period.

Objective: We aimed to use the structured data in electronic health records (EHRs) to find TAEs and PTAEs among adults.

Methods: We used 49,331 adult admissions involving critical care at a major teaching hospital, 2001-2012, in the MIMIC-III EHRs database. We formed a T (defined as packed red blood cells, platelets, or plasma) group of 21,443 admissions vs. 25,468 comparison (C) admissions. The ICD-9-CM diagnosis codes were compared for T vs. C, described, and tested with statistical tools.

Results: TAEs such as transfusion associated circulatory overload (TACO; 12 T cases; rate ratio (RR) 15.61; 95% CI 2.49 to 98) were found. There were also PTAEs similar to TAEs, such as fluid overload disorder (361 T admissions; RR 2.24; 95% CI 1.88 to 2.65), similar to TACO. Some diagnoses could have been sequelae of TAEs, including nontraumatic compartment syndrome of abdomen (52 T cases; RR 6.76; 95% CI 3.40 to 14.9) possibly being a consequence of TACO.

Conclusions: Surveillance for diagnosis codes that could be TAE sequelae or unrecognized TAE might be useful supplements to existing medical product adverse event programs.

KEYWORDS

Electronic health records; big data; patient harm; patient safety; public health; product surveillance, postmarketing; diagnosis codes; proof of concept study; critical care; blood transfusion; humans

ACKNOWLEDGEMENTS

We thank enthusiastic support by our FDA and Booz Allen Hamilton supervisors, Department of Health and Human Services innovation programs (Ignite Accelerator and Data Science CoLab), and Alistair Johnson, DPhil, of the MIMIC-III program, Massachusetts Institute of Technology. George Plopper, PhD, Booz Allen Hamilton, provided project and consultation support. Many FDA colleagues offered ideas and feedback regarding the selection of the case of blood transfusion and the final paper. All authors had access to the data. Drs. Bright, Bright-Ponte, and Palmer are responsible for the study topic, design, and

interpretation. Drs. Rankin and Bright formed the study groups. Drs. Bright and Blok are responsible for data analysis.

Financial support: The research was done with FDA support and under contract HHSF223201510027B between FDA and Booz Allen Hamilton Inc.

Conflicts of interest: None of the authors have other relevant financial interests.

Disclaimer: The opinions are those of the authors and do not represent official policy of either the FDA or Booz Allen Hamilton.

ABBREVIATIONS USED MORE THAN ONCE

95% LCL:	normal approximate 95% lower confidence limits
95% UCL:	normal approximate 95% upper confidence limits
AE:	adverse events
C:	comparison group of admissions
EHRs:	electronic health records
FDA:	Food and Drug Administration
FET:	Fisher's exact test
GI:	gastrointestinal
ICD9:	International Statistical Classification of Diseases and Related Health Problems 9-CM
MIMIC-III:	Medical Information Mart for Intensive Care III
NOS:	Not otherwise specified
PTAE:	potential transfusion adverse event
RD:	rate difference
RR:	rate ratio
T:	group of transfusion admissions
TACO:	transfusion associated circulatory overload
TAE:	transfusion adverse event
TRALI:	transfusion related acute lung injury

INTRODUCTION

FDA relies on spontaneous reports of patient harm (known as adverse events (AE)) from manufacturers, healthcare providers, and the general public to learn of AEs that may need regulatory action [1]. Published deficiencies of these reports [2-10] include varying representativeness of harm and problems. Now that electronic healthcare records (EHRs) are very common [11], we have an opportunity to leverage them as sources of AE information [4, 12-14].

Challenges to the automated use of diagnosis codes for surveillance include:

- ICD [International Classification of Diseases] codes may be “invalid, insensitive or non-specific” [15].
- EHRs are “...subject to access restrictions...” [2].
- “...[N]ot all events and outcomes are consistently captured ...” [13].

The most directly useful program would analyze EHRs [1, 6] in real time. The data available to us (critical care admissions) did not allow calendar time analyses, so we compared blood transfusion admissions (T) vs. comparison admissions (C) because:

- An earlier version of the dataset showed a higher risk of near-term mortality for red blood cell transfusion patients, compared to non-transfused patients [16].
- During the time of the dataset, new transfusion AE (TAE) types, transfusion related acute lung injury (TRALI) and transfusion associated cardiac overload (TACO), were being recognized among transfusion experts and prompted new guidelines to reduce the use of blood transfusion [17].
- During the dataset time, far fewer reports were coming to FDA than would have been expected, considering the level of professional concern [18-20].

The focus of this study is on diagnosis codes in electronic health records (EHR) because it is a relatively available source of information in EHRs and are readily computable. Many studies have looked for predetermined unwelcome conditions; for example, emergency department visits due to opioid overdose [21], motor vehicle crash injury [22], and serious infections [23]. We studied the utility of this method in this setting so that we can compare the results to using another surveillance method with the same data.

The objective of this study was to observe the utility of diagnosis codes to find TAEs that could be further evaluated (outside this study’s scope) with a formal analysis to demonstrate causality.

METHODS

We used EHRs for critical care admissions within an adult hospital, Beth Israel Deaconess Medical Center, Boston, MA. The Massachusetts Institute of Technology (MIT) worked with the hospital to process EHRs, including unstructured notes, from 2001 to 2012 into a privacy-protected dataset, Medical Information Mart for Intensive Care III (MIMIC-III) [24, 25], that is publicly available to those meeting human subjects research requirements. The research was designated not human subjects research by the FDA Institutional Review Board under Code of Federal Regulations Title 45 Part 46 [26].

We removed admissions with patient age < 16 years on the date of admission, as well as admissions without any notes, from the total of 58,976 admissions, resulting in 49,284 admissions.

The transfused group (T) of 21,443 admissions had at least one transfusion (red blood cells, platelets, or plasma) “input code” in the “Inputs” table or International Statistical Classification of Diseases and Related Health Problems (ICD) 9-CM Volume 3 procedure code in the “Procedures” table, listed in Appendix Table A1 (Codes for transfusion and exclusion groups). Among the remaining patients, we identified 2,373 admissions whose only indication of possible transfusion are the codes listed in Appendix Table A1 and excluded them from further study. The other 25,468 admissions formed the comparison (C) group.

We counted the numbers of admissions in T and C for all ICD9 diagnoses [27] that occurred at least as often in T as in C, using Python packages SciPy Stats [28] and Pandas [29]. Most of the codes were present in the MIMIC-III dictionary of ICD9 diagnosis codes. For those that were not, names were found on a public website and enclosed in brackets [30]. When the choice of appropriate code name was not clear, we examined the procedure codes for the relevant admissions, and if necessary, the discharge notes to adjudicate the correct name.

We manually reviewed the information that was available for admissions that received an explicit TAE code despite being assigned to C.

For each diagnosis code we calculated the rate difference (RD), rate ratio (RR), their normal approximate 95% lower and upper confidence limits (95% LCL and 95% UCL), and p values [31]. When any of the rate numerators had a value of 5 or less, we calculated the Fishers Exact Test (FET) p value with Python and then derived the confidence limits [32].

Highly related codes were presented adjacent to each other in Tables 2 through 9 and subtotaled. All 524 diagnoses in the Tables met one of the following criteria [33-86]:

- Known TAE
- Hypercoagulation, embolism, or anemia diagnosis
- $RR > 2$
- $RR > 1.5$ and actual (T minus C) counts over 100
- The FET p value is < 0.05 (calculated when either of the T or C counts < 6)

TAE codes in our study were compared to TAE codes in published literature with a Fisher’s Exact Test calculator [87].

Our presentation of exact p values as low as “<0.0001” allows the reader to calculate their own adjustment of alpha based on their clinical interpretation of the interdependence of the diagnosis codes. If all 524 presented diagnoses are completely dependent, the alpha level to use is 0.0000979 [88]. Specific adjusted alpha levels assuming complete dependence are given with the results in tables organized by TAE-oriented system: iatrogenic (Table 2), circulation (3), bleeding and anemia (4), immune, neoplasm, and infection (5), lungs and oral (6), fluids regulation (7), digestion and hormones (8), and connective, nervous and reproductive (9).

We used published studies of TAEs in Medicare data [19, 20, 89, 90] to calculate expected TAE rates in MIMIC-III.

RESULTS

Table 1 shows the explanations for all admissions that were assigned to the C group and had an explicit TAE diagnosis code. For six of the twelve admissions, the notes say transfusion occurred. For four, the notes say there was transfusion just before admission to the hospital or CCU. For the remaining two, there was no mention of transfusion in the notes, so the TAE diagnosis codes were not explained.

The remaining tables have 524 diagnosis codes. The following TAE codes were found in the T group:

- ICD9 code 27661, Transfusion associated circulatory overload, in Table 7
- ICD9 code 27951, Acute graft-versus-host disease, in Table 5, which could apply to transplants or transfusion, in Table 5
- ICD9 code 78066, Febrile nonhemolytic transfusion reaction, in Table 5
- ICD9 code 9998, Other and unspecified transfusion reaction not elsewhere classified, in Table 2
- ICD9 code 99989, Other transfusion reaction, in Table 2

Most of them were at least one of: predated admission (53), known reasons or associated with reasons to transfuse (315), known risk factors for TAE (12), sequelae of TAE (120), clinically very similar to known TAE (120), or known TAE (13).

For two explicit TAE codes, codes for similar conditions were also statistically significantly more common in T.

- For febrile nonhemolytic transfusion reaction in Table 5, 78061 (Fever presenting with conditions classified elsewhere) was also much more common in T than C and was assigned much more often (196 admissions vs 3).
- For transfusion associated circulatory overload in Table 7, 2766 (fluid overload disorder) was also much more common in T than C and was assigned much more often (571 admissions vs 13).

There were codes for conditions similar to TAE that were statistically significantly more common in T:

- Some of the hypercoagulation and emboli disorders in Table 3 and bleeding in Table 4 may have been alloimmune thrombocytopenic purpura [35].
- Some of the anemia diagnoses in Table 4 may have been hemolytic transfusion reactions [37, 40].
- Some of the infections in Table 5 may have been transmitted via transfusion 43, 75-79].
- Codes in Table 6 that might have signified TRALI [65] were:
 - 5119 (Unspecified pleural effusion) [80]
 - 5118 (Other specified forms of pleural effusion except tuberculous) [80]
 - 9973 (Respiratory complications not elsewhere classified) [49, 72]
 - If transfusion was associated with the trauma or surgery [72]
 - 5185 (Pulmonary insufficiency following trauma and surgery)
 - 51851 (Acute respiratory failure following trauma and surgery)
 - 51852 (Other pulmonary insufficiency, not elsewhere classified, following trauma and surgery)

Many other codes may have been consequences of TAE, for example:

- Kidney injury [44, 70] in Table 7, such as:
 - 5845 (Acute kidney failure with lesion of tubular necrosis)
 - 5848 (Acute kidney failure with other specified pathological lesion in kidney)
- Esophagus injury or gastritis in Table 8 possibly due to intubation, such as:
 - 53010 (Esophagitis, unspecified) [43, 49, 86]
 - 53540 (Other specified gastritis, without mention of hemorrhage) [43]

- Pressure ulcer [81] due to extended bed rest in Table 9:
 - 70703 (Pressure ulcer, lower back)
 - 70711 (Ulcer of thigh)

As one would expect, many of the diagnosis codes that were statistically significantly more common in T than C were, or could have been indicators of, reasons to transfuse (including preexisting conditions:

- Loss of blood due to spontaneous reasons, trauma, or surgery [33-35, 37, 47, 48], in Table 4
- Malignant cancers [37, 43, 47, 48, 70] in Table 5

Some codes might have been preexisting conditions that raise the risk or severity of TR, for example:

- 28262 (Hb-SS disease with crisis) [65] in Table 4
- 2690 (Deficiency of vitamin K) [37] in Table 8

Some codes may represent both reasons to transfuse and consequences of transfusion, for example:

- Emboli and thrombi in Table 3 [76]
- Bleeding in Table 4 [35, 37, 47, 48]
- Hemolytic anemia in Table 4 [37, 76]

There are many codes that we were not able to explain. Most were vague, while some were specific, for example:

- Benign neoplasms in Table 5
- 5933 (Stricture or kinking of ureter) in Table 7
- 73342 (Aseptic necrosis of head and neck of femur) in Table 9.

Many medical care AEs occur at higher frequency in hospital critical care [91, 92].

As would be expected, many diagnosis codes that were associated with transfusion were reasons to transfuse or predated the admission.

A relatively small number were explicit TAE codes. Some codes were for conditions that are the same as explicit TAE codes, but without the attribution and many of them may have been TAE. The strategy of looking for codes for conditions similar to each other has been used to analyze adverse event reports [93]. One interpretation of this phenomenon is knowledge that it is TAE but reluctance to code as such; another could be lack of awareness that the condition was a TAE, and a remaining possibility is that the potential TAE occurred before transfusion.

Our findings on two TAE codes were statistically significantly less than would have been expected from Medicare studies of the frequencies of TRALI and febrile non-haemolytic transfusion reaction (ICD9 code 78066) [20, 90] (see Table 10). Our results were not statistically significantly different for TACO and posttransfusion purpura (ICD9 code 28741) [19, 89].

Our results showed that allowing additional diagnoses that could indicate TAE could be useful for widening the net for detection of signals of TAEs.

Some diagnoses may have been consequent to TAEs.

A number of diagnoses were not explained.

Our investigation of TAE codes in C demonstrated that codes alone may not provide the full clinical picture that is in the notes. Other studies have reported that while codes can provide more information than AE reports [4], the codes themselves provide an incomplete picture [12]. Our main measures of RD

and RR could be underestimates of the risks of T if the loss of information due to reliance on codes is evenly distributed across the data or otherwise biased.

The analysis of codes is merely to generate signals of AE. All signals require further investigation to assess causality because diagnosis codes generally do not provide the sequence of events before, and never during, hospitalization or assess competing explanations of the associations.

CONCLUSIONS

This study successfully demonstrated simple surveillance of patient harm by using diagnosis codes in a large healthcare database, as a supplement to current post-marketing surveillance methods. In routine practice, automation of the interpretation phase could point human monitors to the codes that warrant further investigation by clinical experts.

The limitations of full reliance on diagnosis codes motivate exploring automated methods of using the notes for surveillance of AEs.

We look forward to accessible methods for public health analysts to observe patterns in patient care and experience in real time, by detecting both anticipated and unanticipated signals in electronic health records that indicate potential safety concerns with medical care. Such a method could supplement existing safety surveillance methods.

Table 1. Explanations for all admissions that were assigned to the comparison group and had an explicit transfusion adverse event (TAE) diagnosis code (International Statistical Classification of Diseases and Related Health Problems 9-CM). Bracketed diagnosis code names were not in the MIMIC-III database, therefore were looked up [30] and confirmed in the notes.

ICD9 diagnosis code and name	# admissions	Explanation
27661 <i>Transfusion associated circulatory overload</i>	1	Notes indicate transfusion occurred.
[999.8 <i>Other and unspecified transfusion reaction not elsewhere classified. 999.80 is Transfusion reaction, unspecified.</i>]	7	<ul style="list-style-type: none"> • For 4 admissions, notes say transfusion occurred. • For 1 admission, notes do not mention transfusion or TAE. • For 1 admission, notes mention transfusion just before admission to this hospital. • For 1 admission, notes say patient was transfused and had transfusion related acute lung injury (TRALI); also had diagnosis codes 51881 (Acute respiratory failure) and 5184 (Acute lung edema NOS).
99989 <i>Other transfusion reaction</i>	4	<ul style="list-style-type: none"> • For 2 admissions, notes say patients were transfused in the hospital but not critical care unit. • For 1 admission, notes say patient was transfused before hospitalization. • For 1 admission, there was no mention of transfusion in the notes.

Table 2. Analysis of T (transfusion) and C (comparison) groups for iatrogenic diagnosis codes. If all diagnoses in the table are dependent, alpha is 0.0057. Bracketed diagnosis code names were not in the MIMIC-III database, therefore were looked up [30] and confirmed in some notes.

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
Unspecified or other transfusion reaction (alpha = 0.025)																		
9998 [Other and unspecified transfusion reaction not elsewhere classified; or Transfusion reaction, unspecified.]						y	73	7	100	11	427	343	511	11.82	6.33	24.45	< 0.0001	
99989 Other transfusion reaction						y	27	4			315	244	387	13.56	6.25	29.45	< 0.0001	
Fistula (alpha = 0.010)																		
5374 Fistula of stomach or duodenum		y					21	4	266	124	796	638	954	2.79	2.25	3.48	< 0.0001	
56981 Fistula of intestine, excluding rectum and anus		y					119	52			84	1	166	6.83	2.20	21.14	0.0009	[33, 34]
5961 Intestinovesical fistula		y					17	9			368	264	473	2.98	2.13	4.21	< 0.0001	[33, 34]
6191 Digestive-genital tract fistula, female		y					21	13			47	6	88	2.46	1.04	6.25	0.0242	[33, 34]
9986 Persistent postoperative fistula		y					88	46			51	5	98	2.10	1.00	4.57	0.0314	[33, 34]
9999 Other and unspecified complications of medical care, not elsewhere classified			y	y	y	y	133	51			437	329	546	3.39	2.44	4.78	< 0.0001	
V4282 Peripheral stem cells replaced by transplant			y				46	25			125	57	192	2.39	1.44	4.06	0.0003	[35]

Table 3. Analysis of T (transfusion) and C (comparison) groups for circulation diagnosis codes. If all diagnoses in the table are dependent, alpha is 0.00069. Bracketed diagnosis code names were not in the MIMIC-III database, therefore were looked up [30] and confirmed in some notes.

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference	
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Known TAE	Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit			Upper 95% confidence limit
Heart valve disorders (alpha = 0.0039)									4637	2999	10871	10170	11570	2.01	1.92	2.11	< 0.0001		
3940 Mitral stenosis			y					19	9		56	14	99	2.75	1.18	6.89	0.0092	[33, 34, 36]	
3941 Rheumatic mitral insufficiency			y					37	14		122	65	180	3.44	1.82	6.88	< 0.0001	[33, 34, 36]	
4240 Mitral valve disorders			y					1505	1308		2328	1903	2754	1.50	1.39	1.61	< 0.0001	[33, 34, 36]	
3960 Mitral valve stenosis and aortic valve stenosis			y					39	13		135	78	193	3.90	2.04	7.97	< 0.0001	[33, 34, 36]	
3962 Mitral valve insufficiency and aortic valve stenosis			y					259	118		785	629	940	2.85	2.29	3.58	< 0.0001	[33, 34, 36]	
3963 Mitral valve insufficiency and aortic valve insufficiency			y					215	118		580	433	726	2.37	1.88	2.99	< 0.0001	[33, 34, 36]	
3968 Multiple involvement of mitral and aortic valves			y					75	37		217	132	302	2.64	1.76	4.02	< 0.0001	[33, 34, 36]	
4241 Aortic valve disorders			y					1520	879		3937	3544	4329	2.25	2.07	2.45	< 0.0001	[33, 34, 36]	
7464 Congenital insufficiency of aortic valve	y		y					164	98		413	284	543	2.18	1.68	2.82	< 0.0001	[33, 34, 36]	
39891 Rheumatic heart failure (congestive)			y					309	123		1000	833	1167	3.27	2.64	4.06	< 0.0001	[33, 34, 36]	
99602 Mechanical complication due to heart valve prosthesis			y					49	6		207	148	266	10.62	4.55	30.34	< 0.0001	[33, 34, 36]	
99671 Other complications due to heart valve prosthesis			y					35	12		120	65	175	3.79	1.92	8.03	< 0.0001	[36, 37]	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
V433 Heart valve replaced by other means			y				411	264			970	762	1178	2.02	1.73	2.37	< 0.0001	[36, 37]
Non-valvular disorders of the structure of the heart (alpha = 0.0039)																		
41042 Acute myocardial infarction of other inferior wall, subsequent episode of care			y				17	9			47	6	88	2.46	2.29	2.92	0.024	[36, 38]
41071 Subendocardial infarction, initial episode of care			y				1767	1221			3862	3424	4300	1.88	1.75	2.03	< 0.0001	[36, 38]
41072 Subendocardial infarction, subsequent episode of care			y				172	103			433	300	566	2.17	1.69	2.80	< 0.0001	[36, 38]
4111 Intermediate coronary syndrome			y				917	591			2157	1846	2469	2.02	1.82	2.24	< 0.0001	[36, 38]
99603 Mechanical complication due to coronary bypass graft			y				9	0			42	-7	91				0.0039	[36, 37]
99672 Other complications due to other cardiac device, implant, and graft			y	y			282	176			684	513	856	2.08	1.72	2.53	< 0.0001	[36, 37]
V174 [Family history of other cardiovascular diseases.]	y						15	7			45	7	82	2.79	1.07	8.08	0.019	
41419 Other aneurysm of heart			y				7	0			33	-13	79				0.016	[36]
4200 Acute pericarditis in diseases classified elsewhere			y				11	2			44	-22	110	7.15	1.32	38.83	0.022	[36]
4219 Acute endocarditis, unspecified			y				8	1			34	-23	90	10.40	1.12	96.60	0.039	[36]
4230 Hemopericardium			y				62	25			199	125	274	3.23	2.00	5.36	< 0.0001	[36]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
4231 Adhesive pericarditis			y				22	10			67	21	112	2.86	1.30	6.77	0.0039	[36]
42971 Acquired cardiac septal defect			y				37	8			144	90	198	6.02	2.76	14.95	< 0.0001	[36]
Disorders of heart rhythm (alpha = 0.017)																		
									7529	6444	12005	11060	12950	1.52	1.47	1.57	< 0.0001	
42653 Other bilateral bundle branch block			y				16	7			50	11	88	2.97	1.16	8.54	0.012	[36, 38]
42731 Atrial fibrillation			y				6711	5882			10205	9310	11100	1.48	1.43	1.54	< 0.0001	[36, 38]
4275 Cardiac arrest			y				802	555			1750	1455	2045	1.88	1.68	2.10	< 0.0001	[36, 38]
Myocardial infarction (alpha = 0.010)																		
									3675	2479	8249	7621	8878	1.93	1.83	2.03	< 0.0001	
41042 Acute myocardial infarction of other inferior wall, subsequent episode of care			y			y	17	9			47	6	88	2.46	1.04	6.25	0.024	[36, 38]
41071 Subendocardial infarction, initial episode of care			y			y	1767	1221			3862	3424	4300	1.88	1.75	2.03	< 0.0001	[36, 38]
41072 Subendocardial infarction, subsequent episode of care			y			y	172	103			433	300	566	2.17	1.69	2.80	< 0.0001	[36, 38]
4111 Intermediate coronary syndrome			y			y	917	591			2157	1846	2469	2.02	1.82	2.24	< 0.0001	[36, 38]
4275 Cardiac arrest			y			y	802	555			1750	1455	2045	1.88	1.68	2.10	< 0.0001	[36, 38]
Other heart disorders (alpha = 0.025)																		
									1502	773	4233	3851	4615	2.53	2.32	2.76	< 0.0001	
4294 Functional disturbances following cardiac surgery			y	y			13	8			32	-5	69	2.11	0.81	5.88	0.088	[36, 37, 39]
9971 Cardiac complications, not elsewhere classified			y	y	y		1489	765			4201	3820	4581	2.53	2.32	2.77	< 0.0001	[36, 39]
Atherosclerosis (alpha = 0.013)																		
									481	231	1415	1201	1629	2.71	2.31	3.18	< 0.0001	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
4400 Atherosclerosis of aorta	y						180	96			495	362	628	2.44	1.89	3.16	< 0.0001	[36]
44024 Atherosclerosis of native arteries of the extremities with gangrene	y						220	106			646	501	791	2.70	2.13	3.44	< 0.0001	[36]
44031 Atherosclerosis of autologous vein bypass graft of the extremities	y						34	15			105	49	161	2.95	1.56	5.83	0.0003	[36]
4408 Atherosclerosis of other specified arteries	y						47	14			169	106	232	4.37	2.36	8.59	< 0.0001	[36]
Weakened vascular wall (alpha = 0032)																		
									1805	848	5377	4964	5790	2.77	2.55	3.01	< 0.0001	
4412 Thoracic aneurysm without mention of rupture	y		y				302	164			820	647	993	2.39	1.97	2.91	< 0.0001	[36]
4414 Abdominal aneurysm without mention of rupture	y		y				362	283			673	470	877	1.66	1.42	1.95	< 0.0001	[36]
4417 Thoracoabdominal aneurysm, without mention of rupture	y		y				45	20			138	74	203	2.93	1.69	5.23	< 0.0001	[36]
4420 Aneurysm of artery of upper extremity	y		y				8	1			34	-23	90	10.40	1.12	96.60	0.039	[36]
4422 Aneurysm of iliac artery	y		y				51	28			137	66	209	2.37	1.47	3.90	0.0002	[36]
4423 Aneurysm of artery of lower extremity	y		y				102	37			343	249	438	3.59	2.44	5.37	< 0.0001	[36]
44283 Aneurysm of splenic artery	y		y				12	3			45	-29	119	5.20	1.12	24.19	0.035	[36]
44284 Aneurysm of other visceral artery	y		y				38	7			152	98	206	7.06	3.11	18.74	< 0.0001	[36]
4549 Asymptomatic varicose veins			y				25	14			66	16	116	2.32	1.16	4.83	0.0093	[40, 41]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference	
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Known TAE	Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit			Upper 95% confidence limit
4550 Internal hemorrhoids without mention of complication		y						156	55		530	414	647	3.69	2.70	5.11	< 0.0001	[42]	
4552 Internal hemorrhoids with other complication		y						74	27		248	168	329	3.56	2.27	5.76	< 0.0001	[42]	
4553 External hemorrhoids without mention of complication		y						50	18		169	103	235	3.61	2.07	6.58	< 0.0001	[42]	
4555 External hemorrhoids with other complication		y						29	14		85	32	138	2.69	1.38	5.52	0.0015	[42]	
4561 Esophageal varices without mention of bleeding	y							91	31		313	225	402	3.82	2.51	5.94	< 0.0001	[36]	
45621 Esophageal varices in diseases classified elsewhere, without mention of bleeding	y							305	113		1017	853	1181	3.51	2.82	4.40	< 0.0001	[36]	
4568 Varices of other sites		y						155	33		605	495	714	6.11	4.17	9.19	< 0.0001	[36, 42]	
Other Diseases of Arteries, Arterioles, And Capillaries (alpha = 0.025)										97	56	252	152	351	2.25	1.60	3.19	< 0.0001	
44629 Other specified hypersensitivity angiitis			y					16	7		50	11	88	2.97	1.16	8.54	0.0116	[43]	
4471 Stricture of artery			y					81	49		202	111	293	2.15	1.49	3.13	< 0.0001	[36]	
Other vascular complication (alpha = 0.013)										676	364	1847	1589	2106	2.42	2.12	2.75	< 0.0001	
9961 Mechanical complication of other vascular device, implant, and graft			y					98	57		253	153	352	2.24	1.60	3.16	< 0.0001	[37]	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
99674 Other complications due to other vascular device, implant, and graft		y					363	211			936	744	1128	2.24	1.88	2.66	< 0.0001	[38, 39]
9972 Peripheral vascular complications, not elsewhere classified					y		179	86			526	396	657	2.71	2.08	3.54	< 0.0001	[34, 36, 44]
99779 Vascular complications of other vessels					y		36	10			132	78	186	4.68	2.28	10.58	< 0.0001	[34, 36, 44]
Thrombosis and embolism (alpha = 0.0037)																		
									1100	496	3351	3031	3672	2.88	2.59	3.21	< 0.0001	
V5863 Long-term (current) use of antiplatelet/antithrombotic	y		y				16	6			53	16	91	3.47	1.29	10.82	0.0056	[45]
4440 [Embolism and thrombosis of abdominal aorta.]			y		y		33	20			82	24	141	2.15	1.20	3.95	0.0058	[40]
44421 Arterial embolism and thrombosis of upper extremity			y		y		42	20			124	61	187	2.73	1.57	4.91	0.0001	[40]
44422 Arterial embolism and thrombosis of lower extremity			y		y		146	55			484	370	597	3.45	2.52	4.80	< 0.0001	[40]
44481 Embolism and thrombosis of iliac artery			y		y		36	17			107	49	165	2.75	1.51	5.23	0.0003	[40]
44489 Embolism and thrombosis of other specified artery			y		y		70	30			219	139	299	3.03	1.95	4.82	< 0.0001	[40]
4449 Embolism and thrombosis of unspecified artery			y		y		6	2			21	-51	93	3.90	0.32	47.40	0.2890	[40]
4466 Thrombotic microangiopathy			y		y		31	8			116	66	166	5.04	2.27	12.69	< 0.0001	[40]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
452 Portal vein thrombosis			y	y			194	62			682	554	811	4.07	3.04	5.51	< 0.0001	[40]
45340 Acute venous embolism and thrombosis of unspecified deep vessels of lower extremity			y	y			58	37			138	60	216	2.04	1.33	3.17	0.0005	[40]
4538 [Acute venous embolism and thrombosis of other specified veins.]			y	y			404	202			1160	962	1357	2.60	2.19	3.10	< 0.0001	[40]
45385 Acute venous embolism and thrombosis of subclavian veins			y	y			19	11			49	5	93	2.25	1.02	5.23	0.0282	[40]
45386 Acute venous embolism and thrombosis of internal jugular veins			y	y			24	14			62	12	111	2.23	1.11	4.66	0.0143	[40]
449 Septic arterial embolism			y	y			21	12			55	9	101	2.28	1.07	5.07	0.0194	[40]
Could raise risk of or due to thrombosis or embolism (alpha = 0.013)																		
									519	200	1703	1488	1918	3.37	2.86	3.99	< 0.0001	
E9348 Other agents affecting blood constituents causing adverse effects in therapeutic use			y	y			31	15			91	36	145	2.69	1.41	5.36	0.0011	
4549 Asymptomatic varicose veins	y			y			25	14			66	16	116	2.32	1.16	4.83	0.0093	[40, 41]
5570 Acute vascular insufficiency of intestine	y				y		371	123			1289	1111	1467	3.92	3.19	4.85	< 0.0001	[37, 40, 46]
5579 Unspecified vascular insufficiency of intestine	y				y		92	48			257	162	352	2.49	1.74	3.61	< 0.0001	[37, 40, 46]

Table 4. Analysis of T (transfusion) and C (comparison) groups for bleeding and anemia diagnosis

codes. If all diagnoses in the table are dependent, alpha is 0.00026. Bracketed diagnosis code names were not in the MIMIC-III database, therefore were looked up [30] and confirmed in some notes.

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference	
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit			
Bleeding or hemorrhage not for traumatic or injury or surgery reasons (alpha = 0.00092)									6962	1941	25507	24750	26260	4.66	4.44	4.91	<0.0001		
7827 Spontaneous ecchymoses			y					15	8			41	3	80	2.44	0.97	6.64	0.035	[37]
36281 Retinal hemorrhage			y					19	3			78	1	155	8.24	2.40	28.30	0.00	[37]
44101 Dissection of aorta, thoracic			y					151	77			428	307	549	2.55	1.93	3.40	<0.0001	[33, 34, 37]
4411 Thoracic aneurysm, ruptured			y					20	6			72	31	113	4.34	1.68	13.19	0.0006	[33, 34, 37]
4413 Abdominal aneurysm, ruptured			y					67	9			280	210	350	9.68	4.81	22.09	<0.0001	[33, 34, 37]
44322 Dissection of iliac artery			y					23	7			82	38	126	4.27	1.78	11.79	0.0002	[33, 34, 37]
44329 Dissection of other artery			y					31	15			91	36	145	2.69	1.41	5.36	0.0011	[33, 34, 37]
4472 Rupture of artery			y					23	5			89	1	178	5.98	2.09	17.14	0.00	[33, 34, 37]
4550 Internal hemorrhoids without mention of complication	y		y					156	55			530	414	647	3.69	2.70	5.11	<0.0001	[42]
4552 Internal hemorrhoids with other complication	y		y					74	27			248	168	329	3.56	2.27	5.76	<0.0001	[42]
4553 External hemorrhoids without mention of complication	y		y					50	18			169	103	235	3.61	2.07	6.58	<0.0001	[42]
4555 External hemorrhoids with other complication	y		y					29	14			85	32	138	2.69	1.38	5.52	0.0015	[42]
4568 Varices of other sites	y		y					155	33			605	495	714	6.11	4.17	9.19	<0.0001	[36, 42]
4560 Esophageal varices with bleeding			y		y			30	3			129	44	214	13.01	4.66	36.31	0.00	[35, 37, 47, 48]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
45620 Esophageal varices in diseases classified elsewhere, with bleeding		y		y			300	52			1213	1062	1363	7.50	5.57	10.28	<0.0001	[35, 37, 47, 48]
4590 Hemorrhage, unspecified		y		y			93	22			355	269	441	5.50	3.43	9.19	<0.0001	[35, 37, 47, 48]
53021 Ulcer of esophagus with bleeding		y		y	y		45	11			170	110	230	5.32	2.71	11.41	<0.0001	[35, 37, 47, 48, 49, 50]
5307 Gastroesophageal laceration-hemorrhage syndrome		y		y			121	70			313	203	424	2.25	1.66	3.06	<0.0001	[35, 37, 47, 48, 49]
53082 Esophageal hemorrhage		y		y			67	18			248	174	322	4.84	2.84	8.66	<0.0001	[35, 37, 47, 48, 49]
53100 Acute gastric ulcer with hemorrhage, without mention of obstruction		y		y			55	6			235	172	298	11.92	5.14	33.89	<0.0001	[35, 37, 47, 48]
53110 Acute gastric ulcer with perforation, without mention of obstruction		y		y			8	1			34	-23	90	10.40	1.12	96.60	0.04	[35, 37, 47, 48]
53140 Chronic or unspecified gastric ulcer with hemorrhage, without mention of obstruction	y	y		y			231	38			941	810	1073	7.91	5.59	11.46	<0.0001	[35, 37, 47, 48]
53150 Chronic or unspecified gastric ulcer with perforation, without mention of obstruction	y	y		y			12	3			45	-29	119	5.20	1.12	24.19	0.04	[35, 37, 47, 48]
53200 Acute duodenal ulcer with hemorrhage, without mention of obstruction		y		y			73	1			337	229	444	94.94	37.93	237.63	0.00	[35, 37, 47, 48]
53240 Chronic or unspecified duodenal ulcer with hemorrhage, without mention of obstruction	y	y		y			229	42			917	785	1049	7.09	5.09	10.10	<0.0001	[35, 37, 47, 48]
53250 Chronic or unspecified duodenal ulcer with perforation, without mention of obstruction	y	y		y			17	4			65	-16	146	5.53	1.59	19.22	0.01	[35, 37, 47, 48]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
53260 Chronic or unspecified duodenal ulcer with hemorrhage and perforation, without mention of obstruction	y	y	y				11	0			51	0	103				0.00	[35, 37, 47, 48]
53340 Chronic or unspecified peptic ulcer of unspecified site with hemorrhage, without mention of obstruction	y	y	y				9	1			38	-19	95	11.71	1.43	95.53	0.02	[35, 37, 47, 48]
53400 Acute gastrojejunal ulcer with hemorrhage, without mention of obstruction		y	y				7	0			33	-13	79				0.02	[35, 37, 47, 48]
53440 Chronic or unspecified gastrojejunal ulcer with hemorrhage, without mention of obstruction	y	y	y				29	7			110	62	158	5.39	2.31	14.57	<0.0001	[35, 37, 47, 48]
53501 Acute gastritis, with hemorrhage		y	y				34	5			141	46	235	8.84	3.62	21.61	0.00	[35, 37, 47, 48]
53511 Atrophic gastritis, with hemorrhage		y	y				8	1			34	-23	90	10.40	1.12	96.60	0.04	[35, 37, 47, 48]
53541 Other specified gastritis, with hemorrhage		y	y				85	22			318	235	400	5.02	3.11	8.44	<0.0001	[35, 37, 47, 48]
53551 Unspecified gastritis and gastroduodenitis, with hemorrhage		y	y				107	42			348	251	446	3.31	2.30	4.85	<0.0001	[35, 37, 47, 48]
53561 Duodenitis, with hemorrhage		y	y				45	4			196	97	294	14.63	6.34	33.78	0.00	[35, 37, 47, 48]
5371 Gastric diverticulum	y	y					7	0			33	-13	79				0.02	[51]
53782 Angiodysplasia of stomach and duodenum without mention of hemorrhage		y					42	11			156	98	215	4.97	2.51	10.70	<0.0001	[52]
53783 Angiodysplasia of stomach and duodenum with hemorrhage		y	y				113	14			477	387	567	10.50	6.00	19.82	<0.0001	[35, 37, 47, 48]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
53784 Dieulafoy lesion (hemorrhagic) of stomach and duodenum		y		y			46	4			200	101	299	14.96	6.52	34.30	0.00	[35, 37, 47, 48]
56212 Diverticulosis of colon with hemorrhage		y		y			293	41			1219	1073	1366	9.29	6.69	13.22	<0.0001	[35, 37, 47, 48]
56213 Diverticulitis of colon with hemorrhage		y		y			23	7			82	38	126	4.27	1.78	11.79	0.0002	[35, 37, 47, 48]
56881 Hemoperitoneum (nontraumatic)		y					174	29			707	593	822	7.80	5.25	12.00	<0.0001	[53]
5690 Anal and rectal polyp		y					13	2			53	-14	121	8.45	1.77	40.28	0.01	[33, 34, 40, 54]
5693 Hemorrhage of rectum and anus		y		y			66	37			175	94	257	2.32	1.53	3.57	<0.0001	[35, 37, 47, 48]
56983 Perforation of intestine		y					158	72			479	357	600	2.85	2.15	3.83	<0.0001	[33, 34, 37, 52]
56985 Angiodysplasia of intestine with hemorrhage		y		y			91	7			399	320	479	16.91	7.88	43.24	<0.0001	[35, 37, 47, 48]
56986 Dieulafoy lesion (hemorrhagic) of intestine		y		y			7	0			33	-13	79				0.02	[35, 37, 47, 48]
5723 Portal hypertension		y					757	249			2637	2383	2892	3.95	3.42	4.58	<0.0001	[33, 34, 37, 55]
5724 Hepatorenal syndrome		y					301	45			1242	1093	1391	8.70	6.34	12.18	<0.0001	[56]
5780 Hematemesis		y		y			288	138			848	683	1014	2.71	2.21	3.35	<0.0001	[35, 37, 47, 48]
5781 Blood in stool		y		y			521	163			1845	1636	2055	4.16	3.48	4.99	<0.0001	[35, 37, 47, 48]
5789 Hemorrhage of gastrointestinal tract, unspecified		y		y			1158	237			4551	4251	4850	6.35	5.52	7.34	<0.0001	[35, 37, 47, 48]
66612 Other immediate postpartum hemorrhage, delivered, with mention of postpartum complication		y		y			21	0			98	35	161				0.00	[35, 37, 47, 48]
66624 Delayed and secondary postpartum hemorrhage, postpartum condition or complication		y		y			8	0			37	-10	85				0.01	[35, 37, 47, 48]
7847 Epistaxis			y		y		202	99			587	448	726	2.65	2.08	3.41	<0.0001	[57]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	<i>Known TAE</i>					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
7863 [Hemoptysis.]					y		244	156			579	418	739	2.03	1.66	2.50	<0.0001	[35, 37, 47, 48, 58]
Surgery that typically results in bleeding or hemorrhage (alpha = 0.0039)																		
99652 Mechanical complication due to graft of other tissue, not elsewhere classified		y					15	5			52	-36	140	3.90	1.05	14.46	0.04	[33, 34]
99799 Complications affecting other specified body systems, not elsewhere classified				y			14	7			40	3	77	2.60	0.98	7.61	0.032	[36]
E8798 Other specified procedures as the cause of abnormal reaction of patient, or of later complication, without mention of misadventure at time of procedure	y						796	690			1238	929	1547	1.50	1.35	1.66	<0.0001	[33, 34]
9982 Accidental puncture or laceration during a procedure, not elsewhere classified		y					629	294			1879	1636	2123	2.78	2.42	3.21	<0.0001	[33, 34, 37]
9983 [Disruption of operation wound.]		y	y				24	6			90	47	134	5.20	2.07	15.56	0.0001	[33, 34, 37, 40, 47]
99831 Disruption of internal operation (surgical) wound		y	y				124	35			453	352	554	4.61	3.14	6.91	<0.0001	[33, 34, 37, 40, 47]
99832 Disruption of external operation (surgical) wound		y	y				172	59			591	469	712	3.79	2.81	5.19	<0.0001	[33, 34, 37, 47]
99889 Other specified complications of procedures not elsewhere classified		y	y				127	80			305	190	421	2.06	1.55	1.55	<0.0001	[33, 34, 37, 47]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference	
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Known TAE	Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit			Upper 95% confidence limit
E8781 Surgical operation with implant of artificial internal device causing abnormal patient reaction, or later complication, without mention of misadventure at time of operation			y		y			411	278			920	710	1130	1.92	1.65	2.25	<0.0001	[33, 34, 37, 47]
E8782 Surgical operation with anastomosis, bypass, or graft, with natural or artificial tissues used as implant causing abnormal patient reaction, or later complication, without mention of misadventure at time of operation			y		y			826	302			2769	2500	3038	3.56	3.11	4.07	<0.0001	[33, 34, 37, 47]
E8788 Other specified surgical operations and procedures causing abnormal patient reaction, or later complication, without mention of misadventure at time of operation			y		y			764	567			1530	1237	1822	1.75	1.57	1.96	<0.0001	[33, 34, 37, 47]
V5331 Fitting and adjustment of cardiac pacemaker			y		y			95	42			292	199	386	2.94	2.03	4.34	<0.0001	[33, 34, 37, 47]
V6441 Laparoscopic surgical procedure converted to open procedure			y		y			89	56			214	118	311	2.07	1.46	2.94	<0.0001	[33, 34, 37, 47]
Injury or trauma that typically results in bleeding or hemorrhage (alpha = 0.00065)										5340	2045	17570	16880	18260	3.40	3.23	3.58	<0.0001	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	<i>Known TAE</i>					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
80425 Closed fractures involving skull or face with other bones with subarachnoid, subdural, and extradural hemorrhage, with prolonged (more than 24 hours) loss of consciousness, without return to pre-existing conscious level		y					7	0			33	-13	79				0.02	[33, 34, 37]
8603 Traumatic hemothorax with open wound into thorax		y					15	5			52	-36	140	3.90	1.05	14.46	0.04	[33, 34, 37]
8605 Traumatic pneumothorax with open wound into thorax		y					16	6			53	16	91	3.47	1.29	10.82	0.0056	[33, 34, 37]
86122 Laceration of lung without mention of open wound into thorax		y					13	6			39	4	74	2.82	1.00	9.04	0.0282	[33, 34, 37]
86132 Laceration of lung with open wound into thorax		y					14	2			58	-10	126	9.10	2.01	41.18	0.00	[33, 34, 37]
8621 Injury to diaphragm, with open wound into cavity		y					14	6			44	8	80	3.03	1.10	9.63	0.0167	[33, 34, 37]
8900 Open wound of hip and thigh, without mention of complication		y					15	7			45	7	82	2.79	1.07	8.08	0.0194	[33, 34, 37]
8901 Open wound of hip and thigh, complicated		y					8	0			37	-10	85				0.01	[33, 34, 37]
90089 Injury to other specified blood vessels of head and neck		y					14	7			40	3	77	2.60	0.98	7.61	0.032	[33, 34, 37]
9010 Injury to thoracic aorta		y					23	5			89	1	178	5.98	2.09	17.14	0.00	[33, 34, 37]
90253 Injury to iliac artery		y					11	0			51	0	103				0.00	[33, 34, 37]
90289 Injury to other specified blood vessels of abdomen and pelvis		y					14	1			62	0	123	18.21	3.27	101.48	0.00	[33, 34, 37]
90441 Injury to popliteal artery		y					8	1			34	-23	90	10.40	1.12	96.60	0.04	[33, 34, 37]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
9047 Injury to other specified blood vessels of lower extremity		y					10	1			43	-15	101	13.01	1.77	95.69	0.01	[33, 34, 37]
9582 Secondary and recurrent hemorrhage		y					15	0			70	14	126				0.00	[33, 34, 37]
95892 Traumatic compartment syndrome of lower extremity					y		26	9			89	42	136	3.76	1.71	9.11	0.0002	[59]
95893 Traumatic compartment syndrome of abdomen					y		10	0			47	-3	97				0.00	[60]
99702 Iatrogenic cerebrovascular infarction or hemorrhage		y					279	134			821	658	984	2.71	2.20	3.35	<0.0001	[33, 34, 37]
99811 Hemorrhage complicating a procedure		y					1320	204			5424	5111	5737	8.42	7.26	9.80	<0.0001	[33, 34, 37]
99812 Hematoma complicating a procedure		y					885	329			2948	2668	3227	3.50	3.08	3.98	<0.0001	[33, 34, 37]
72992 Nontraumatic hematoma of soft tissue		y					64	21			223	149	297	3.96	2.39	6.83	<0.0001	[33, 34, 37]
E8700 Accidental cut, puncture, perforation or hemorrhage during surgical operation		y					119	63			329	221	437	2.46	1.80	3.39	<0.0001	[33, 34, 37]
E8708 Accidental cut, puncture, perforation or hemorrhage during other specified medical care		y					52	30			135	62	208	2.25	1.41	3.66	0.0003	[33, 34, 37]
E8152 Other motor vehicle traffic accident involving collision on the highway injuring motorcyclist		y					14	6			44	8	80	3.03	1.10	9.63	0.0167	[33, 34, 37]
E8230 Other motor vehicle nontraffic accident involving collision with stationary object injuring driver of motor vehicle other than motorcycle		y					40	21			111	49	174	2.48	1.43	4.42	0.0005	[33, 34, 37]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
E9353 Salicylates causing adverse effects in therapeutic use		y					23	12			64	17	112	2.49	1.19	5.50	0.0079	[33, 34, 37]
8602 Traumatic hemothorax without mention of open wound into thorax		y			y		72	26			243	163	322	3.60	2.27	5.88	<0.0001	[33, 34, 37, 61]
8604 Traumatic pneumothorax without mention of open wound into thorax		y			y		87	51			223	129	317	2.22	1.55	3.20	<0.0001	[33, 34, 37, 61]
8620 Injury to diaphragm, without mention of open wound into cavity		y					22	7			77	34	121	4.09	1.69	11.33	0.0004	[33, 34, 37]
6271 Postmenopausal bleeding		y					15	8			41	3	80	2.44	0.97	6.64	0.0354	[33, 34, 37]
64421 Early onset of delivery, delivered, with or without mention of antepartum condition		y					19	6			67	27	107	4.12	1.58	12.60	0.001	[33, 34, 37]
86389 Injury to other gastrointestinal sites, without mention of open wound into cavity		y					25	16			59	8	111	2.03	1.04	4.07	0.0237	[33, 34, 37]
86803 Injury to other intra-abdominal organs without mention of open wound into cavity, peritoneum		y					87	43			252	160	343	2.63	1.81	3.89	<0.0001	[33, 34, 37]
86804 Injury to other intra-abdominal organs without mention of open wound into cavity, retroperitoneum		y					61	23			202	129	276	3.45	2.10	5.84	<0.0001	[33, 34, 37]
92232 Contusion of buttock		y					28	4			116	29	203	9.10	3.34	24.80	0.00	[33, 34, 37]
80620 Closed fracture of T1-T6 level with unspecified spinal cord injury		y					24	4			98	13	182	7.80	2.68	22.69	0.00	[33, 34, 37]
80621 Closed fracture of T1-T6 level with complete lesion of cord		y					9	0			42	-7	91				0.00	[33, 34, 37]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
80625 Closed fracture of T7-T12 level with unspecified spinal cord injury		y					17	6			58	19	96	3.68	1.39	11.41	0.0032	[33, 34, 37]
80629 Closed fracture of T7-T12 level with other specified spinal cord injury		y					14	2			58	-10	126	9.10	2.01	41.18	0.00	[33, 34, 37]
8064 Closed fracture of lumbar spine with spinal cord injury		y					18	10			48	6	91	2.34	1.02	5.68	0.0263	[33, 34, 37]
8056 Closed fracture of sacrum and coccyx without mention of spinal cord injury		y					107	54			305	204	407	2.58	1.84	3.64	<0.0001	[33, 34, 37]
81201 Closed fracture of surgical neck of humerus		y					26	13			75	25	125	2.60	1.29	5.51	0.0035	[33, 34, 37]
81209 Other closed fracture of upper end of humerus		y					22	13			56	9	103	2.20	1.06	4.76	0.0207	[33, 34, 37]
81221 Closed fracture of shaft of humerus		y					21	13			51	5	98	2.10	1.00	4.57	0.0314	[33, 34, 37]
81250 Open fracture of unspecified part of lower end of humerus		y					7	0			33	-13	79				0.02	[33, 34, 37]
81311 Open fracture of olecranon process of ulna		y					11	1			48	-11	106	14.31	2.12	96.57	0.01	[33, 34, 37]
8082 Closed fracture of pubis		y					180	78			560	431	689	3.00	2.29	3.97	<0.0001	[33, 34, 37]
8083 Open fracture of pubis		y					12	1			52	-7	112	15.61	2.49	97.93	0.00	[33, 34, 37]
80841 Closed fracture of ilium		y					42	17			135	73	197	3.21	1.79	6.02	<0.0001	[33, 34, 37]
80843 Multiple closed pelvic fractures with disruption of pelvic circle		y					23	8			79	34	123	3.74	1.61	9.67	0.0006	[33, 34, 37]
80849 Closed fracture of other specified part of pelvis		y					8	0			37	-10	85				0.01	[33, 34, 37]
80853 Multiple open pelvic fractures with disruption of pelvic circle		y					9	1			38	-19	95	11.71	1.43	95.53	0.02	[33, 34, 37]
8088 Closed unspecified fracture of pelvis		y					13	6			39	4	74	2.82	1.00	9.04	0.0282	[33, 34, 37]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	<i>Known TAE</i>					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
8080 Closed fracture of acetabulum		y					139	52			462	351	573	3.48	2.51	4.88	<0.0001	[33, 34, 37]
82021 Closed fracture of intertrochanteric section of neck of femur		y					98	43			303	208	398	2.96	2.05	4.35	<0.0001	[33, 34, 37]
82022 Closed fracture of subtrochanteric section of neck of femur		y					49	11			189	127	251	5.79	2.97	12.36	<0.0001	[33, 34, 37]
8208 Closed fracture of unspecified part of neck of femur		y					45	27			113	45	181	2.17	1.32	3.63	0.0011	[33, 34, 37]
82101 Closed fracture of shaft of femur		y					63	17			233	161	305	4.82	2.79	8.79	<0.0001	[33, 34, 37]
82111 Open fracture of shaft of femur		y					22	2			95	21	170	14.31	4.10	49.91	0.00	[33, 34, 37]
82121 Closed fracture of condyle, femoral		y					13	6			39	4	74	2.82	1.00	9.04	0.0282	[33, 34, 37]
82123 Closed supracondylar fracture of femur		y					24	10			76	29	123	3.12	1.44	7.31	0.0014	[33, 34, 37]
82131 Open fracture of condyle, femoral		y					6	0			28	-17	73				0.03	[33, 34, 37]
82133 Open supracondylar fracture of femur		y					8	1			34	-23	90	10.40	1.12	96.60	0.04	[33, 34, 37]
92400 Contusion of thigh		y					25	15			63	12	113	2.17	1.10	4.42	0.0152	[33, 34, 37]
8220 Closed fracture of patella		y					22	12			60	13	106	2.38	1.13	5.286	0.0125	[33, 34, 37]
8221 Open fracture of patella		y					6	0			28	-17	73				0.03	[33, 34, 37]
82300 Closed fracture of upper end of tibia alone		y					44	19			137	73	201	3.01	1.72	5.46	<0.0001	[33, 34, 37]
82302 Closed fracture of upper end of fibula with tibia		y					28	11			91	41	141	3.31	1.60	7.37	0.0004	[33, 34, 37]
82312 Open fracture of upper end of fibula with tibia		y					9	1			38	-19	95	11.71	1.43	95.53	0.02	[33, 34, 37]
82330 Open fracture of shaft of tibia alone		y					10	2			39	-26	105	6.50	1.10	38.43	0.04	[33, 34, 37]
82332 Open fracture of shaft of fibula with tibia		y					30	12			97	45	149	3.25	1.62	6.97	0.0003	[33, 34, 37]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference	
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit			
82382 Closed fracture of unspecified part of fibula with tibia		y					24	4			98	13	182	7.80	2.68	22.69	0.00	[33, 34, 37]	
82392 Open fracture of unspecified part of fibula with tibia		y					15	8			41	3	80	2.44	0.97	6.64	0.035	[33, 34, 37]	
8240 Fracture of medial malleolus, closed		y					27	11			86	37	136	3.19	1.53	7.13	0.0006	[33, 34, 37]	
8242 Fracture of lateral malleolus, closed		y					15	8			41	3	80	2.44	0.97	6.64	0.035	[33, 34, 37]	
8245 Bimalleolar fracture, open		y					14	1			62	0	123	18.21	3.27	101.48	0.00	[33, 34, 37]	
82521 Closed fracture of astragalus		y					13	6			39	4	74	2.82	1.00	9.04	0.028	[33, 34, 37]	
9588 Other early complications of trauma		y					32	5			131	38	225	8.32	3.33	20.79	0.00	[33, 34, 37]	
E8497 Accidents occurring in residential institution		y					551	474			870	613	1127	1.51	1.33	1.71	<0.0001	[33, 34, 37]	
Conditions that typically involve surgery and bleeding (alpha = 0.0037)										3936	2605	9015	8366	9663	1.97	1.87	2.07	<0.0001	
3361 Vascular myelopathies		y					39	22			103	40	166	2.31	1.33	4.08	0.0013	[33, 34, 62]	
3363 Myelopathy in other diseases classified elsewhere		y					70	43			172	87	258	2.12	1.43	3.17	0.0001	[33, 34, 62]	
3940 Mitral stenosis		y					19	9			56	14	99	2.75	1.18	6.89	0.0092	[33, 34, 36]	
3941 Rheumatic mitral insufficiency		y					37	14			122	65	180	3.44	1.82	6.88	<0.0001	[33, 34, 36]	
3960 Mitral valve stenosis and aortic valve stenosis		y					39	13			135	78	193	3.90	2.04	7.97	<0.0001	[33, 34, 36]	
3962 Mitral valve insufficiency and aortic valve stenosis		y					259	118			785	629	940	2.85	2.29	3.58	<0.0001	[33, 34, 36]	
3963 Mitral valve insufficiency and aortic valve insufficiency		y					215	118			580	433	726	2.37	1.88	2.99	<0.0001	[33, 34, 36]	
3968 Multiple involvement of mitral and aortic valves		y					75	37			217	132	302	2.64	1.76	4.02	<0.0001	[33, 34, 36]	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	<i>Known TAE</i>					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
41419 Other aneurysm of heart		y					7	0			33	-13	79				0.02	[33, 34, 36]
4240 Mitral valve disorders		y					1505	1308			2328	1903	2754	1.50	1.39	1.61	<0.0001	[33, 34, 36]
4241 Aortic valve disorders		y					1520	879			3937	3544	4329	2.25	2.07	2.45	<0.0001	[33, 34, 36]
4295 Rupture of chordae tendineae		y					48	13			177	115	240	4.80	2.56	9.66	<0.0001	[33, 34, 37]
42971 Acquired cardiac septal defect		y					37	8			144	90	198	6.02	2.76	14.95	<0.0001	[33, 34, 63]
5933 Stricture or kinking of ureter		y					23	5			89	1	178	5.98	2.09	17.14	0.00	[33, 34]
Iron disorders (alpha = 0.025)																		
									869	202	3328	3066	3591	5.60	4.80	6.55	<0.0001	
2750 [Disorders of iron metabolism.]		y	y				40	24			100	36	165	2.17	1.28	3.76	0.0021	[64]
2800 Iron deficiency anemia secondary to blood loss (chronic)		y	y				829	178			3228	2973	3482	6.06	5.15	7.16	<0.0001	[37]
Sickle cell disease (alpha = 0.05)																		
									10	2	39	-26	105	6.50	1.10	38.43	0.04	
28262 Hb-SS disease with crisis			y				10	2			39	-26	105	6.50	1.10	38.43	0.04	[65]
Acquired hemolytic anemia (alpha = 0.013)																		
									136	37	502	396	607	4.78	3.30	7.08	<0.0001	
2830 Autoimmune hemolytic anemias		y	y				45	13			163	102	224	4.50	2.39	9.10	<0.0001	[37, 40, 66]
28311 Hemolytic-uremic syndrome		y	y				17	0			79	21	138				0.00	[37, 40, 66]
28319 Other non-autoimmune hemolytic anemias		y	y				20	9			61	18	104	2.89	1.26	7.21	0.0056	[37, 40, 66]
2839 Acquired hemolytic anemia, unspecified		y	y				54	15			198	132	265	4.68	2.60	8.93	<0.0001	[37, 40, 66]
Aplastic anemia and other bone marrow failure syndromes (alpha = 0.013)																		
									692	301	2148	1895	2400	2.99	2.61	3.43	<0.0001	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference	
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Known TAE	Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit			Upper 95% confidence limit
2841 [Pancytopenia.]			y		y			288	182		690	517	864	2.06	1.70	2.49	<0.0001	[37]	
2848 [Other specified aplastic anemias.]			y					270	51		1076	933	1220	6.89	5.09	9.48	<0.0001	[37]	
28489 Other specified aplastic anemias			y					40	20		115	53	177	2.60	1.49	4.70	0.0003	[37]	
99685 Complications of transplanted bone marrow			y					94	48		266	171	362	2.55	1.78	3.69	<0.0001	[37, 48]	
Other and unspecified anemias (alpha = 0.013)										4426	845	17611	17030	18190	6.81	6.33	7.34	<0.0001	
64821 Anemia of mother, delivered, with or without mention of antepartum condition			y		y			12	3		45	-29	119	5.20	1.12	24.19	0.035	[37, 44, 67]	
64822 Anemia of mother, delivered, with mention of postpartum complication			y		y			19	1		85	18	152	24.71	5.43	112.50	<0.0001	[37, 44, 67]	
2851 Acute posthemorrhagic anemia			y		y			3870	637		15764	15230	16300	7.90	7.26	8.61	<0.0001	[37, 44, 67]	
79001 Precipitous drop in hematocrit			y		y			216	81		717	579	855	3.47	2.67	4.53	<0.0001	[37, 44, 67]	
Neutropenia (alpha = 0.013)										369	154	1169	985	1352	3.12	2.58	3.79	<0.0001	
2880 [Neutropenia.]			y					110	26		420	326	513	5.50	3.56	8.79	<0.0001	[37]	
28800 Neutropenia, unspecified			y					163	78		480	356	605	2.72	2.06	3.61	<0.0001	[37]	
28803 Drug induced neutropenia			y					81	44		220	130	310	2.39	1.64	3.54	<0.0001	[37]	
28804 Neutropenia due to infection			y					15	6		48	12	85	3.25	1.19	10.23	0.0097	[37, 68]	
Spleen problems (alpha = 0.0073)										247	98	800	652	949	3.28	2.58	4.19	<0.0001	
2894 Hypersplenism			y					18	3		73	-3	149	7.80	2.21	27.60	0.00	[37, 47]	
28951 Chronic congestive splenomegaly			y					15	4		56	-25	137	4.88	1.29	18.40	0.020	[37, 47]	
28959 Other diseases of spleen			y					83	34		265	179	352	3.17	2.11	4.88	<0.0001	[37, 47]	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	<i>Known TAE</i>					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
86502 Injury to spleen without mention of open wound into cavity, capsular tears, without major disruption of parenchyma		y					28	13			84	33	135	2.80	1.40	5.89	0.0013	[37, 47]
86504 Injury to spleen without mention of open wound into cavity, massive parenchymal disruption		y					27	3			115	33	198	11.71	4.02	34.10	0.00	[37, 47]
86509 Other injury into spleen without mention of open wound into cavity		y					25	15			63	12	113	2.17	1.10	4.42	0.0152	[37, 47]
7892 Splenomegaly		y					51	26			145	74	215	2.55	1.56	4.26	0.0001	[37, 47]
Coagulation problems that promote bleeding (alpha = 0.0047)																		
							5096	2395	15177	14480	15870	2.77	2.64	2.91	<0.0001			
2866 Defibrination syndrome		y	y				387	65			1572	1401	1742	7.74	5.94	10.23	<0.0001	[69]
2867 Acquired coagulation factor deficiency		y	y				373	93			1406	1233	1579	5.22	4.15	6.62	<0.0001	[37, 40, 47, 69]
2869 Other and unspecified coagulation defects		y	y				751	256			2584	2330	2839	3.82	3.31	4.41	<0.0001	[37, 40, 47, 69]
2873 [Primary thrombocytopenia.]	y	y					36	17			107	49	165	2.75	1.51	5.23	0.0003	[45]
28731 Immune thrombocytopenic purpura		y	y				66	32			193	114	272	2.68	1.73	4.23	<0.0001	[35, 37, 40, 47]
2874 [Secondary thrombocytopenia.]		y	y				398	181			1207	1014	1400	2.86	2.39	3.43	<0.0001	[35, 37, 40, 47]
28749 Other secondary thrombocytopenia		y	y				79	51			186	94	277	2.01	1.40	2.92	0.0001	[35, 37, 40, 47]
2875 Thrombocytopenia, unspecified		y	y				1928	1103			5036	4595	5477	2.27	2.11	2.45	<0.0001	[35, 37, 40, 47]
66632 Postpartum coagulation defects, delivered, with mention of postpartum complication		y	y				10	0			47	-3	97				0.00	[35, 37, 40, 47]
79092 Abnormal coagulation profile		y	y				591	394			1343	1092	1595	1.95	1.71	2.22	<0.0001	[35, 37, 40, 47]
E9342 Anticoagulants causing adverse effects in therapeutic use		y	y				477	203			1497	1288	1706	3.06	2.59	3.62	<0.0001	[35, 37, 40, 47]

Table 5. Analysis of T (transfusion) and C (comparison) groups for immune, neoplasm, and infection diagnosis codes. If all diagnoses in the table are dependent, alpha is 0.00049. Bracketed diagnosis code names were not in the MIMIC-III database, therefore were looked up [30] and confirmed in some notes.

ICD9 diagnosis code and name	Predates current admission	Interpretation					# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference											
		?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Known TAE	Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit			Upper 95% confidence limit										
27951 Acute graft-versus-host disease			y		y		y	9	1			38	-19	95	11.71	1.43	95.53	0.021	[40, 47]										
Inflammation (alpha = 0.025)																				106	36	365	270	461	3.83	2.60	5.76	< 0.0001	
99594 Systemic inflammatory response syndrome due to noninfectious process with acute organ dysfunction		y						90	27			323	236	410	4.34	2.79	6.94	< 0.0001	[43]										
37200 Acute conjunctivitis, unspecified						y		16	9			42	2	82	2.31	0.96	5.94	0.038	[43]										
Fever (alpha = 0.025)																				129	70	351	238	464	2.40	1.78	3.25	< 0.0001	
78061 Fever presenting with conditions classified elsewhere					y			126	70			337	224	449	2.34	1.73	3.18	< 0.0001	[47, 70-72]										
78066 Febrile nonhemolytic transfusion reaction						y		3	0			14	-30	58				0.25	[70, 71]										
44629 Other specified hypersensitivity angitis					y			16	7			50	11	88	2.97	1.16	8.54	0.012	[43]										
Malignant neoplasm (alpha = 0.0030)																				891	370	2828	2544	3113	3.13	2.77	3.55	< 0.0001	
1533 Malignant neoplasm of sigmoid colon			y			y		23	14			57	8	106	2.14	1.05	4.49	0.022	[37, 43, 48, 70]										
1534 Malignant neoplasm of cecum			y			y		20	10			57	14	101	2.60	1.16	6.22	0.010	[37, 43, 48, 70]										
1540 Malignant neoplasm of rectosigmoid junction			y			y		16	4			60	-21	141	5.20	1.44	18.79	0.012	[37, 43, 48, 70]										
1974 Secondary malignant neoplasm of small intestine including duodenum			y			y		65	36			174	94	255	2.35	1.54	3.63	< 0.0001	[37, 43, 48, 70]										

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
1975 Secondary malignant neoplasm of large intestine and rectum			y			y	50	25			144	74	213	2.60	1.58	4.39	0.0001	[37, 43, 48, 70]
1830 Malignant neoplasm of ovary			y			y	50	31			122	50	194	2.10	1.31	3.40	0.0009	[37, 43, 48, 70]
1986 Secondary malignant neoplasm of ovary			y			y	14	6			44	8	80	3.03	1.10	9.63	0.017	[37, 43, 48, 70]
1888 Malignant neoplasm of other specified sites of bladder			y			y	43	26			107	41	174	2.15	1.29	3.65	0.0016	[37, 43, 48, 70]
1981 Secondary malignant neoplasm of other urinary organs			y			y	29	13			89	37	141	2.90	1.46	6.08	0.0008	[37, 43, 48, 70]
19882 Secondary malignant neoplasm of genital organs			y			y	32	13			103	49	156	3.20	1.63	6.65	0.0002	[37, 43, 48, 70]
20400 Acute lymphoid leukemia, without mention of having achieved remission			y		y		27	12			83	33	133	2.93	1.43	6.34	0.0012	[37, 67, 73]
20500 Acute myeloid leukemia, without mention of having achieved remission			y		y		163	32			645	534	757	6.62	4.51	10.01	< 0.0001	[37, 47]
20501 Acute myeloid leukemia, in remission	y						20	10			57	14	101	2.60	1.16	6.22	0.010	
20510 Chronic myeloid leukemia, without mention of having achieved remission			y		y		42	23			113	49	178	2.37	1.40	4.14	0.0006	[37]
27788 Tumor lysis syndrome		y					20	10			57	14	101	2.60	1.16	6.22	0.010	[37, 43, 48, 70, 74]
1550 Malignant neoplasm of liver, primary			y			y	237	100			747	600	894	3.08	2.43	3.93	< 0.0001	[37, 43, 48, 70]
1552 Malignant neoplasm of liver, not specified as primary or secondary			y			y	40	5			169	115	222	10.40	4.11	33.78	< 0.0001	[37, 43, 48, 70]
Benign neoplasms (alpha = 0.0043)																		
									663	211	2335	2098	2572	4.09	3.49	4.79	< 0.0001	
2111 Benign neoplasm of stomach		y					58	22			192	120	263	3.43	2.07	5.88	< 0.0001	
2112 Benign neoplasm of duodenum, jejunum, and ileum		y					37	9			140	86	195	5.35	2.53	12.60	< 0.0001	

ICD9 diagnosis code and name	Interpretation							# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference	
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit			
2113 Benign neoplasm of colon		y						245	74			877	734	1020	4.31	3.31	5.66	< 0.0001		
2114 Benign neoplasm of rectum and anal canal		y						24	10			76	29	123	3.12	1.44	7.31	0.0014		
2115 Benign neoplasm of liver and biliary passages		y						11	6			30	-3	63	2.38	0.81	7.85	0.077		
2180 Submucous leiomyoma of uterus			y	y				6	0			28	-17	73					0.031	
2181 Intramural leiomyoma of uterus		y						14	6			44	8	80	3.03	1.10	9.63	0.017		
2230 Benign neoplasm of kidney, except pelvis		y						8	1			34	-23	90	10.40	1.12	96.60	0.039		
2352 Neoplasm of uncertain behavior of stomach, intestines, and rectum		y						10	0			47	-3	97					0.0020	
2381 Neoplasm of uncertain behavior of connective and other soft tissue		y						13	3			50	-24	124	5.64	1.29	24.59	0.021		
2387 [Neoplasm of uncertain behavior of other lymphatic and hematopoietic tissues.]		y						96	24			362	274	449	5.20	3.30	8.51	< 0.0001		
23875 Myelodysplastic syndrome, unspecified		y						141	56	457	344	569	3.27	2.39	4.55	< 0.0001				
V1000 Personal history of malignant neoplasm of gastrointestinal tract, unspecified	y							20	11			54	9	98	2.36	1.08	5.46	0.018		
Septicemia (alpha = 0.0085)										2656	2062	4992	4442	5543	1.68	1.58	1.78	< 0.0001		
3811 Methicillin susceptible Staphylococcus aureus septicemia					y			332	180			903	722	1084	2.40	1.99	2.89	< 0.0001	[75]	
3819 Other staphylococcal septicemia					y			89	58			207	110	304	2.00	1.42	2.83	< 0.0001	[75]	
3843 Septicemia due to pseudomonas					y			84	41			245	155	334	2.66	1.81	3.97	< 0.0001	[75]	
3844 Septicemia due to serratia					y			18	9			52	10	93	2.60	1.11	6.57	0.015	[75]	
388 Other specified septicemias						y		143	61			448	334	563	3.05	2.24	4.19	< 0.0001	[43]	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
389 Unspecified septicemia						y	1990	1713			3138	2650	3626	1.51	1.42	1.61	< 0.0001	[43]
Bacterial pneumonia (alpha = 0.0085)																		
48241 Methicillin susceptible pneumonia due to Staphylococcus aureus						y	521	261	1268	554	3927	3585	4269	2.98	2.69	3.30	< 0.0001	
48283 Pneumonia due to other gram-negative bacteria						y	189	75			1494	1270	1718	2.60	2.23	3.02	< 0.0001	[43]
48282 Pneumonia due to escherichia coli [E. coli]						y y	77	25			612	482	743	3.28	2.50	4.34	< 0.0001	[43]
4820 Pneumonia due to Klebsiella pneumoniae						y	168	52			269	189	350	4.01	2.52	6.57	< 0.0001	[43, 76]
4821 Pneumonia due to Pseudomonas						y y	296	131			597	478	716	4.20	3.06	5.85	< 0.0001	[43]
5111 Pleurisy with effusion, with mention of a bacterial cause other than tuberculosis						y	17	10			911	745	1076	2.94	2.38	3.64	< 0.0001	[43]
											43	2	85	2.21	0.96	5.40	0.041	[43]
Other bacterial diseases (alpha = 0.0073)																		
4110 Staphylococcus infection in conditions classified elsewhere and of unspecified site, staphylococcus, unspecified						y	36	21	2470	1660	5567	5051	6082	1.94	1.82	2.06	< 0.0001	
4186 Helicobacter pylori [H. pylori]						y	159	45			93	32	153	2.23	1.27	4.02	0.0027	[43]
4210 Acute and subacute bacterial endocarditis						y	241	135			580	466	695	4.60	3.28	6.55	< 0.0001	[43]
56723 Spontaneous bacterial peritonitis						y	137	49			640	484	795	2.32	1.87	2.89	< 0.0001	[43]
7907 Bacteremia						y	835	630			463	354	573	3.64	2.61	5.15	< 0.0001	[43]
417 Pseudomonas infection in conditions classified elsewhere and of unspecified site						y y	246	155			1635	1328	1942	1.72	1.55	1.91	< 0.0001	[43]
845 Intestinal infection due to Clostridium difficile						y	816	625			591	431	752	2.06	1.68	2.54	< 0.0001	[43]
											1564	1260	1869	1.70	1.53	1.89	< 0.0001	[43]
Viral hepatitis (alpha = 0.0085)																		
									402	165	1283	1092	1474	3.17	2.64	3.82	< 0.0001	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference	
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Known TAE	Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit			Upper 95% confidence limit
7022 Chronic viral hepatitis B with hepatic coma without hepatitis delta	y							15	4		56	-25	137	4.88	1.29	18.40	0.019		
7044 Chronic hepatitis C with hepatic coma	y							129	45		440	335	546	3.73	2.64	5.36	< 0.0001		
7020 Viral hepatitis B with hepatic coma, acute or unspecified, without mention of hepatitis delta					y			18	3		73	-3	149	7.80	2.21	27.60	0.0015	[77]	
7041 Acute hepatitis C with hepatic coma					y			12	2		49	-18	115	7.80	1.54	39.47	0.013	[78, 79]	
7070 [Unspecified viral hepatitis C without hepatic coma.]					y			173	97		459	327	591	2.32	1.80	3.01	< 0.0001	[77]	
7071 Unspecified viral hepatitis C with hepatic coma					y			55	14		206	140	273	5.11	2.80	9.95	< 0.0001	[77]	
Cytomegalovirus or herpes virus (alpha = 0.017)										99	35	336	243	429	3.68	2.48	5.57	< 0.0001	
785 Cytomegaloviral disease					y			62	28		189	113	265	2.88	1.82	4.67	< 0.0001	[78, 79]	
4841 Pneumonia in cytomegalic inclusion disease						y		14	3		55	-20	129	6.07	1.47	25.09	0.013	[43]	
5479 Herpes simplex with other specified complications						y		23	4		93	9	177	7.48	2.52	22.17	0.00031	[43]	
Candidiasis (alpha = 0.013)										254	85	880	732	1027	3.89	3.03	5.03	< 0.0001	
1124 Candidiasis of lung						y		27	9		94	46	142	3.90	1.78	9.43	0.0001	[43]	
1125 Disseminated candidiasis						y		135	36		500	396	605	4.88	3.36	7.25	< 0.0001	[43]	
11289 Other candidiasis of other specified sites						y		76	35		229	145	313	2.82	1.87	4.34	< 0.0001	[43]	
1129 Candidiasis of unspecified site						y		16	5		57	20	93	4.16	1.46	14.53	0.0025	[43]	
Other fungal diseases (alpha = 0.017)										199	68	684	553	815	3.81	2.88	5.09	< 0.0001	
1173 Aspergillosis						y		76	39		215	129	301	2.53	1.70	3.83	< 0.0001	[43]	
1179 Other and unspecified mycoses						y		107	27		402	309	495	5.15	3.35	8.18	< 0.0001	[43]	
4847 Pneumonia in other systemic mycoses						y		16	2		67	-2	137	10.40	2.51	43.20	0.0013	[43]	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference		
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Known TAE	Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit			Upper 95% confidence limit	
Infection unspecified organism (alpha = 0.0015)											7215	4544	17354	16480	18220	2.07	1.99	2.14	< 0.0001	
7854 Gangrene						y		73	47		172	84	260	2.02	1.38	2.98	0.0001	[43]		
71105 Pyogenic arthritis, pelvic region and thigh						y		19	8		60	18	102	3.09	1.29	8.16	0.0048	[43]		
71102 Pyogenic arthritis, upper arm						y		4	2		11	-99	122	2.60	0.03	225	0.69	[43]		
71103 Pyogenic arthritis, forearm						y		10	6		25	-7	57	2.17	0.71	7.26	0.12	[43]		
71109 Pyogenic arthritis, multiple sites						y		7	0		33	-13	79				0.016	[43]		
73015 Chronic osteomyelitis, pelvic region and thigh	y					y		13	6		39	4	74	2.82	1.00	9.04	0.028	[43]		
73007 Acute osteomyelitis, ankle and foot						y		65	41		156	74	239	2.06	1.37	3.13	0.0002	[43]		
73008 Acute osteomyelitis, other specified sites						y		68	38		181	98	263	2.33	1.54	3.56	< 0.0001	[43]		
73028 Unspecified osteomyelitis, other specified sites						y		71	44		173	87	259	2.10	1.42	3.13	0.0001	[43]		
5100 Empyema with fistula						y		37	24		86	24	149	2.01	1.17	3.50	0.0068	[80]		
5109 Empyema without mention of fistula						y		125	64		353	243	464	2.54	1.86	3.49	< 0.0001	[80]		
28804 Neutropenia due to infection						y		15	6		48	12	85	3.25	1.19	10.23	0.0097	[37, 68]		
3241 Intraspinal abscess						y		88	44		253	161	345	2.60	1.79	3.83	< 0.0001	[43]		
99731 Ventilator associated pneumonia						y		347	214		851	661	1041	2.11	1.77	2.51	< 0.0001	[72, 81]		
99762 Infection (chronic) of amputation stump						y		46	25		125	57	192	2.39	1.44	4.06	0.0003	[43]		
99851 Infected postoperative seroma						y		16	8	46	7	85	2.60	1.05	7.02	0.022	[43]			
99859 Other postoperative infection						y		780	394	2225	1950	2499	2.57	2.28	2.91	< 0.0001	[43]			
9993 [Other infection due to medical care not elsewhere classified.]						y		15	8	41	3	80	2.44	0.97	6.64	0.035	[43]			

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
V090 Infection with microorganisms resistant to penicillins						y	416	215			1169	968	1370	2.52	2.13	2.98	< 0.0001	[43]
V0980 Infection with microorganisms without mention of resistance to multiple drugs						y	126	59			376	267	485	2.78	2.02	3.85	< 0.0001	[43]
V0981 Infection with microorganisms with resistance to multiple drugs						y	18	9			52	10	93	2.60	1.11	6.57	0.015	[43]
53641 Infection of gastrostomy					y		17	9			47	6	88	2.46	1.04	6.25	0.024	[43]
5401 Acute appendicitis with peritoneal abscess					y		14	7			40	3	77	2.60	0.98	7.61	0.032	[43]
56722 Peritoneal abscess					y		147	90			363	239	486	2.12	1.62	2.79	< 0.0001	[43]
56731 Psoas muscle abscess					y		24	11			72	25	120	2.84	1.34	6.42	0.0027	[43]
56738 Other retroperitoneal abscess					y		24	5			94	5	183	6.24	2.22	17.54	0.00054	[43]
64781 Other specified infectious and parasitic diseases of mother, delivered, with or without mention of antepartum condition					y		6	0			28	-17	73				0.031	[43]
99661 Infection and inflammatory reaction due to cardiac device, implant, and graft					y		94	50			259	163	355	2.45	1.72	3.52	< 0.0001	[43]
99662 Infection and inflammatory reaction due to other vascular device, implant, and graft					y		623	319			1762	1516	2008	2.54	2.22	2.92	< 0.0001	[43]
99666 Infection and inflammatory reaction due to internal joint prosthesis					y		38	14			127	69	185	3.53	1.87	7.05	< 0.0001	[43]
99667 Infection and inflammatory reaction due to other internal orthopedic device, implant, and graft					y		45	26			117	49	184	2.25	1.36	3.80	0.0007	[43]
99592 Severe sepsis					y		2277	1630			4774	4273	5275	1.82	1.70	1.94	< 0.0001	[43]
78552 Septic shock					y		1492	1090			3049	2642	3457	1.78	1.65	1.93	< 0.0001	[76]

ICD9 diagnosis code and name	Interpretation						Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
72886 Necrotizing fasciitis						Y	145	71	220	2.31	1.46	3.71	0.0001	[82]

Table 6. Analysis of T (transfusion) and C (comparison) groups for lungs and oral diagnosis codes. If all diagnoses in the table are dependent, alpha is 0.0034. Bracketed diagnosis code names were not in the MIMIC-III database, therefore were looked up [30] and confirmed in some notes.

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
Pleural effusion (alpha = 0.025)																		
5118 [Other specified forms of pleural effusion except tuberculous.]					y		135	31	1682	1177	3624	3195	4052	1.86	1.72	2.00	< 0.0001	[80]
5119 Unspecified pleural effusion					y		1547	1146			3105	2689	3521	1.76	1.63	1.90	< 0.0001	[80]
Other diseases of respiratory system (alpha = 0.0073)																		
5121 Iatrogenic pneumothorax					y		464	297	2289	960	7232	6776	7689	3.10	2.87	3.35	< 0.0001	[80, 81]
5185 [Pulmonary insufficiency following trauma and surgery.]				y			1359	436			4774	4435	5114	4.05	3.64	4.53	< 0.0001	[72]
51851 Acute respiratory failure following trauma and surgery				y			116	55			344	239	449	2.74	1.97	3.85	< 0.0001	[72]
51852 Other pulmonary insufficiency, not elsewhere classified, following trauma and surgery				y			48	27			127	58	196	2.31	1.41	3.85	< 0.0001	[72]
5192 Mediastinitis					y		30	18			75	20	131	2.17	1.17	4.13	0.0078	[83]
7864 Abnormal sputum		y					10	1			43	-15	101	13.01	1.77	95.69	0.012	
9973 [Respiratory complications not elsewhere classified.]				y	y		262	126			770	612	928	2.70	2.18	3.37	< 0.0001	[49, 72]
Diseases of oral cavity, salivary glands, and jaws (alpha = 0.0085)																		
52100 Dental caries, unspecified	y						65	28	165	76	497	373	621	2.82	2.14	3.76	< 0.0001	
52109 Other dental caries	y						13	6			39	4	74	2.82	1.00	9.04	0.028	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	<i>Known TAE</i>					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
5224 Acute apical periodontitis of pulpal origin					y		22	14			52	4	100	2.04	1.00	4.32	0.033	[43]
52510 Acquired absence of teeth, unspecified	y						9	1			38	-19	95	11.71	1.43	95.53	0.021	
52800 Stomatitis and mucositis, unspecified					y		24	10			76	29	123	3.12	1.44	7.31	0.0014	[43]
52801 Mucositis (ulcerative) due to antineoplastic therapy					y		32	17			88	32	144	2.45	1.32	4.70	0.002	[43]

Table 7. Analysis of T (transfusion) and C (comparison) groups for diagnosis codes that relate to fluids regulation. If all diagnoses in the table are dependent, alpha is 0.0032. Bracketed diagnosis code names were not in the MIMIC-III database, therefore were looked up [30] and confirmed in some notes.

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with cod	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
<i>27661 Transfusion associated circulatory overload</i>						y	12	1			52	-7	112	15.61	2.49	97.93	0.0034	[40, 71]
Fluid overload and consequence (alpha = 0.0064)										610	294	1791	1550	2032	2.70	2.34	3.11	<0.0001
2766 [Fluid overload disorder.]						y	361	210			931	739	1122	2.24	1.88	2.66	<0.0001	[33, 37, 40, 71]
60886 Edema of male genital organs		y					43	18			136	73	199	3.11	1.75	5.72	<0.0001	[33, 34, 84]
4578 Other noninfectious disorders of lymphatic channels				y			30	10			104	53	155	3.90	1.86	8.95	0.0001	
72972 Nontraumatic compartment syndrome of lower extremity					y		33	6			132	82	182	7.15	2.96	20.88	<0.0001	[85]
72973 Nontraumatic compartment syndrome of abdomen					y		52	10			207	144	270	6.76	3.40	14.93	<0.0001	[60]
95892 Traumatic compartment syndrome of lower extremity					y		26	9			89	42	136	3.76	1.71	9.11	0.0002	[59]
95893 Traumatic compartment syndrome of abdomen					y		10	0			47	-3	97				0.0020	[60]
72886 Necrotizing fasciitis					y		55	31			145	71	220	2.31	1.46	3.71	0.0001	[82]
Kidney problem (alpha = 0.013)										1717	725	5408	5012	5804	3.08	2.82	3.36	<0.0001
5845 Acute kidney failure with lesion of tubular necrosis				y	y		1613	667			5131	4748	5513	3.15	2.87	3.45	<0.0001	[44, 70]
5848 Acute kidney failure with other specified pathological lesion in kidney				y	y		36	19			100	40	159	2.46	1.38	4.55	0.001	[44, 70]
59381 Vascular disorders of kidney				y	y		53	30			140	67	213	2.30	1.44	3.72	0.0002	[44, 70]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with cod	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
86601 Injury to kidney without mention of open wound into cavity, hematoma without rupture of capsule					y	y	15	9			38	-2	77	2.17	0.89	5.60	0.06	[44, 70]
Disorders of urinary system other than kidneys (alpha = 0.017)																		
5933 Stricture or kinking of ureter		y					23	5			89	1	178	5.98	2.09	17.13	0.0009	[33, 34]
59582 Irradiation cystitis						y	15	8			41	3	80	2.44	0.97	6.64	0.035	[43]
9975 Urinary complications, not elsewhere classified		y					308	136			949	780	1118	2.95	2.40	3.63	<0.0001	

Table 8. Analysis of T (transfusion) and C (comparison) groups for digestion and hormones diagnosis

codes. If all diagnoses in the table are dependent, alpha is 0.00071. Bracketed diagnosis code names were not in the MIMIC-III database, therefore were looked up [30] and confirmed in some notes.

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference										
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit												
Diseases of Esophagus, Stomach, and Duodenum (alpha = 0.0032)																			942	413	2912	2617	3207	2.97	2.64	3.34	<0.0001	
53010 Esophagitis, unspecified						y		120	78			280	167	393	2.00	1.49	2.70	<0.0001	[43, 49, 86]									
53011 Reflux esophagitis						y		20	9			61	18	104	2.89	1.26	7.21	0.0056	[43, 86]									
53012 Acute esophagitis						y		23	7			82	38	126	4.27	1.78	11.79	0.0002	[43, 49, 86]									
5302 [Ulcer of esophagus.]						y		22	14			52	4	100	2.04	1.00	4.32	0.033	[43, 49, 86]									
53020 Ulcer of esophagus without bleeding						y		39	21			107	45	169	2.42	1.39	4.32	0.0008	[43, 49, 86]									
86222 Injury to esophagus without mention of open wound into cavity						y		7	0			33	-13	79				0.016	[43, 49, 86]									
53170 Chronic gastric ulcer without mention of hemorrhage or perforation, without mention of obstruction	y							18	5			66	-21	153	4.68	1.43	225	0.012										
53190 Gastric ulcer, unspecified as acute or chronic, without mention of hemorrhage or perforation, without mention of obstruction						y		113	46			362	261	463	3.19	2.25	4.60	<0.0001	[43]									
53290 Duodenal ulcer, unspecified as acute or chronic, without hemorrhage or perforation, without mention of obstruction						y		58	32			156	80	232	2.36	1.51	3.75	0.0001	[43]									
53540 Other specified gastritis, without mention of hemorrhage						y		71	44			173	87	259	2.10	1.42	3.13	0.0001	[43]									
53560 Duodenitis, without mention of hemorrhage						y		97	55		255	156	354	2.29	1.63	3.25	<0.0001	[43]										
5371 Gastric diverticulum								7	0		33	-13	79				0.016	[51]										

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
5374 Fistula of stomach or duodenum			y				21	4			84	1	166	6.83	2.20	225	0.0009	[33, 34]
53782 Angiodysplasia of stomach and duodenum without mention of hemorrhage			y				42	11			156	98	215	4.97	2.51	10.70	<0.0001	[52]
53789 Other specified disorders of stomach and duodenum					y		259	78			928	781	1075	4.32	3.34	5.64	<0.0001	[86]
5379 Unspecified disorder of stomach and duodenum					y		25	9			84	38	131	3.61	1.63	8.80	0.0004	[86]
Small and large intestine, rectum, and anus disorders (alpha = 0.0030)																		
5552 Regional enteritis of small intestine with large intestine					y		21	11	2304	1228	6341	5865	6818	2.44	2.28	2.62	<0.0001	
56210 Diverticulosis of colon (without mention of hemorrhage)	y						471	365			888	656	1120	1.68	1.46	1.93	<0.0001	
5570 Acute vascular insufficiency of intestine			y				371	123			1289	1111	1467	3.92	3.19	4.85	<0.0001	[37, 40, 46]
5571 Chronic vascular insufficiency of intestine	y						96	29			344	254	433	4.31	2.82	6.77	<0.0001	[37, 40, 46]
5579 Unspecified vascular insufficiency of intestine			y				92	48			257	162	352	2.49	1.74	3.61	<0.0001	[37, 40, 46]
5601 Paralytic ileus					y		662	390			1689	1429	1949	2.21	1.95	2.51	<0.0001	[37, 86]
5650 Anal fissure			y				15	8			41	3	80	2.44	0.97	6.64	0.035	[52]
5690 Anal and rectal polyp			y				13	2			53	-14	121	8.45	1.77	225	0.0074	[33, 34, 40, 54]
56941 Ulcer of anus and rectum			y				39	11			142	86	199	4.61	2.32	9.98	<0.0001	[52]
56949 Other specified disorders of rectum and anus			y				39	17			121	61	181	2.98	1.65	5.62	0.0001	
56981 Fistula of intestine, excluding rectum and anus			y				119	52			368	264	473	2.98	2.13	4.21	<0.0001	[33, 34]
56982 Ulceration of intestine			y				41	10			155	98	213	5.33	2.63	11.94	<0.0001	[33, 34, 37, 52]
56983 Perforation of intestine			y				158	72			479	357	600	2.85	2.15	3.83	<0.0001	[33, 34, 37, 52]
56984 Angiodysplasia of intestine (without mention of hemorrhage)			y				23	6			86	43	129	4.99	1.97	14.97	<0.0001	[52]

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
56989 Other specified disorders of intestine			y				48	23			141	74	209	2.71	1.62	4.68	<0.0001	[52]
5699 Unspecified disorder of intestine			y				12	7			31	-4	66	2.23	0.81	6.68	0.0834	[52]
V453 Intestinal bypass or anastomosis status	y						84	54			198	104	292	2.02	1.42	2.90	<0.0001	[33, 34]
Liver disorders (alpha = 0.0034)																		
									4614	1786	15113	14470	15750	3.36	3.18	3.55	<0.0001	
570 Acute and subacute necrosis of liver			y				758	305			2441	2180	2703	3.23	2.83	3.70	<0.0001	[37, 45]
5712 Alcoholic cirrhosis of liver			y				831	369			2552	2275	2830	2.93	2.59	3.32	<0.0001	[37, 45]
57149 Other chronic hepatitis			y				24	8			83	38	129	3.90	1.70	10.05	0.0003	[37, 45]
5715 Cirrhosis of liver without mention of alcohol			y				867	420			2537	2250	2825	2.68	2.39	3.02	<0.0001	[37, 45]
5722 Hepatic encephalopathy			y				501	163			1752	1545	1958	4.00	3.34	4.80	<0.0001	
5723 Portal hypertension			y				757	249			2637	2383	2892	3.95	3.42	4.58	<0.0001	[33, 34, 37, 55]
5724 Hepatorenal syndrome			y				301	45			1242	1093	1391	8.70	6.34	12.18	<0.0001	[56]
5728 Other sequelae of chronic liver disease			y				156	43			573	460	686	4.72	3.35	6.78	<0.0001	[37, 45]
5730 Chronic passive congestion of liver			y				48	25			134	66	203	2.50	1.51	4.23	0.0001	[37, 45]
5734 Hepatic infarction			y				21	1			94	26	163	27.31	6.36	225	<0.0001	[37, 45]
86401 Injury to liver without mention of open wound into cavity, hematoma and contusion			y				18	9			52	10	93	2.60	1.11	6.57	0.0150	[37, 45]
86404 Injury to liver without mention of open wound into cavity, laceration, major			y				17	8			51	11	91	2.76	1.13	7.40	0.0133	[37, 45]
86405 Injury to liver without mention of open wound into cavity laceration, unspecified			y				73	47			172	11	91	2.02	1.13	7.40	<0.0001	[37, 45]
99682 Complications of transplanted liver			y				142	35			537	430	643	5.28	3.62	7.87	<0.0001	[37, 45]
7824 Jaundice, unspecified, not of newborn			y	y			100	59			255	154	356	2.20	1.58	3.10	<0.0001	[37, 45]
Gallbladder disorders (alpha = 0.0073)																		
									543	285	1510	1280	1741	2.48	2.14	2.87	<0.0001	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
57410 Calculus of gallbladder with other cholecystitis, without mention of obstruction	y						107	60			284	180	387	2.32	1.68	3.24	<0.0001	
5750 Acute cholecystitis						y	115	61			318	211	424	2.45	1.78	3.40	<0.0001	[43, 86]
57510 Cholecystitis, unspecified						y	24	13			65	17	114	2.40	1.18	5.14	0.0086	[43, 86]
57511 Chronic cholecystitis	y						80	36			244	158	330	2.89	1.93	4.41	<0.0001	[43, 86]
57512 Acute and chronic cholecystitis	y					y	22	13			56	9	103	2.20	1.06	4.76	0.021	[43, 86]
5768 Other specified disorders of biliary tract			y				181	95			503	370	637	2.48	1.92	3.21	<0.0001	
57490 Calculus of gallbladder and bile duct without cholecystitis, without mention of obstruction	y						14	7			40	3	77	2.60	0.98	7.62	0.032	
Nonspecific digestive disorders (alpha = 0.025)										587	272	1762	1527	1997	2.81	2.43	3.25	<0.0001
9974 [Digestive system complications not elsewhere classified.]		y					513	235			1550	1331	1769	2.84	1.78	2.37	<0.0001	
99749 Other digestive system complications		y					74	37			212	128	297	2.60	1.73	3.97	<0.0001	
Abdominal non-organ disorders (alpha = 0.0051)										1814	713	5903	5500	6306	3.31	3.03	3.61	<0.0001
5672 [Other suppurative peritonitis.]						y	181	36			715	597	833	6.54	4.55	9.63	<0.0001	[43]
56721 Peritonitis (acute) generalized						y	60	27			183	108	258	2.89	1.81	4.74	<0.0001	[43]
56729 Other suppurative peritonitis						y	90	38			283	193	374	3.08	2.09	4.63	<0.0001	[43]
5678 [Other specified peritonitis.]						y	27	5			108	17	199	7.02	2.63	225	0.0001	[43]
56781 Choleperitonitis						y	16	4			60	-21	141	5.20	1.44	225	0.0118	[43]
56789 Other specified peritonitis						y	25	9			84	38	131	3.61	1.63	8.80	<0.0001	[43]
5679 Unspecified peritonitis						y	62	31			178	101	255	2.60	1.66	4.14	<0.0001	[43]
5680 Peritoneal adhesions (postoperative) (postinfection)	y						275	129			820	659	981	2.77	2.24	3.44	<0.0001	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference	
	Predates current admission	?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit			
7895 [Ascites.]			y				430	104			1632	1447	1818	5.38	4.33	6.73	<0.0001	[33, 70]	
78959 Other ascites			y				648	330			1839	1588	2089	2.55	2.23	2.93	<0.0001	[33, 70]	
Malnutrition (alpha = 0.017)										846	490	2188	1895	2481	2.25	2.01	2.52	<0.0001	
262 Other severe protein-calorie malnutrition				?			38	22			98	36	160	2.25	1.30	3.99	0.0019		
2639 Unspecified protein-calorie malnutrition				?			792	462			2037	1753	2321	2.23	1.99	2.51	<0.0001		
2690 Deficiency of vitamin K			y				16	6			53	16	91	3.47	1.29	10.82	0.0056	[37]	
2738 Other disorders of plasma protein metabolism		y					86	53			211	117	306	2.11	1.48	3.03	<0.0001		
2554 [Corticoadrenal insufficiency.]							179	110			440	304	577	2.12	1.66	2.71	<0.0001		

Table 9. Analysis of T (transfusion) and C (comparison) groups for connective, nervous, and reproductive diagnosis codes. If all diagnoses in the table are dependent, alpha is 0.0023. TAE: transfusion adverse event. Bracketed diagnosis code names were not in the MIMIC-III database, therefore were looked up [30] and confirmed in some notes.

ICD9 diagnosis code and name	Interpretation							# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE	Rate difference per 100,000					Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit			
Bone or joint disorder (alpha = 0.0073)										168	74	518	393	643	2.95	2.23	3.94	< 0.0001	
7200 Ankylosing spondylitis	y						30	17			79	24	134	2.30	1.23	4.44	0.0049		
7209 Unspecified inflammatory spondylopathy					y		8	0			37	-10	85				0.0078	[43]	
73342 Aseptic necrosis of head and neck of femur		y					34	22			80	20	140	2.01	1.14	3.61	0.0092		
9964 [Mechanical complication of internal orthopedic device implant and graft.]			y				19	8			60	18	101	3.09	1.29	8.16	0.0048		
99644 Peri-prosthetic fracture around prosthetic joint			y				18	2			77	6	148	11.71	3.02	45.37	0.0004		
7313 Major osseous defects			y				23	7			82	38	126	4.27	1.78	11.79	0.0002		
73382 Nonunion of fracture	y						36	18			103	44	162	2.60	1.44	4.87	0.0006		
Skin disorder (alpha = 0.017)										786	593	1539	1242	1837	1.72	1.55	1.92	< 0.0001	
70703 Pressure ulcer, lower back						y	728	559			1391	1103	1678	1.69	1.51	1.89	< 0.0001	[81]	
70711 Ulcer of thigh						y	18	9			52	10	93	2.60	1.11	6.57	0.015	[81]	
7098 Other specified disorders of skin					y		40	25			97	32	162	2.08	1.23	3.58	0.0033	[37, 81]	
Diseases of The Nervous System And Sense Organs (alpha = 0.010)										336	154	1015	837	1192	2.84	2.34	3.46	< 0.0001	
2939 Unspecified transient mental disorder in conditions classified elsewhere		y					83	45			226	135	316	2.40	1.65	3.53	< 0.0001		
3361 Vascular myelopathies							39	22			103	40	166	2.31	1.33	4.08	0.0013	[33, 34, 62]	
3363 Myelopathy in other diseases classified elsewhere							70	43			172	87	258	2.12	1.43	3.17	0.0001	[33, 34, 62]	

ICD9 diagnosis code and name	Interpretation						# T with code	# C with code	Subtotal # T with code	Subtotal # C with code	Rate difference statistics			Rate ratio statistics			p value	Reference
	Predates current admission?	Could be reason to T	Raises risk or severity of TAE	Could be TAE	Could be caused by TAE	Known TAE					Rate difference per 100,000	Lower 95% confidence limit	Upper 95% confidence limit	Rate ratio	Lower 95% confidence limit	Upper 95% confidence limit		
E9359 Unspecified analgesic and antipyretic causing adverse effects in therapeutic use							108	33			385	290	481	4.26	2.86	6.49	< 0.0001	
35981 Critical illness myopathy				y	y		36	11			128	74	183	4.26	2.12	9.27	< 0.0001	[36, 43]
Diseases of The Genitourinary System (alpha = 0.017)																		
									68	36	188	106	270	2.46	1.62	3.79	< 0.0001	
605 Redundant prepuce and phimosis							17	11			40	-3	82	2.01	0.89	4.75	0.0656	
6146 Pelvic peritoneal adhesions, female (postoperative) (postinfection)					y		22	10			67	21	112	2.86	1.30	6.77	0.0039	[43]
6238 Other specified noninflammatory disorders of vagina	y						29	15			81	28	135	2.51	1.30	5.05	0.0027	
Complications of Pregnancy, Childbirth, and the Puerperium (alpha = 0.013)																		
									62	15	235	165	306	5.38	3.02	10.18	< 0.0001	
64891 Other current conditions classifiable elsewhere of mother, delivered, with or without mention of antepartum condition							17	5			61	-26	149	4.42	1.30	14.99	0.0169	
65421 Previous cesarean delivery, delivered, with or without mention of antepartum condition	y						10	2			39	-26	105	6.50	1.10	38.43	0.0385	
67432 Other complications of obstetrical surgical wounds, delivered, with mention of postpartum complication				y			6	0			28	-17	73				0.0312	[48]
V270 Outcome of delivery, single liveborn		y					29	8			107	58	155	4.71	2.10	11.94	< 0.0001	

Table 10. Comparison of MIMIC-III diagnoses of TAEs (transfusion adverse events) with published Medicare data from varying time periods 2007-2012. The expected number applies the rate found in Medicare data to the study T (transfusion) group.

Transfusion adverse event (TAE)	# TAE in T	Medicare data			Expected # in T	Fisher's exact test (FET) p value
		# TAE/ # transfused	Calendar years	Reference		
Transfusion related acute lung injury (TRALI)	0	2556/ 11,378,264	2007-2011	[20]	4.8	0.019
Transfusion associated circulatory overload (TACO)	12	1340/ 2,147,038	2011	[19]	13.4	0.089
Posttransfusion purpura	0	78/ 4,336,338	2011-2012	[89]	0.39	1
Febrile non-haemolytic transfusion reaction	3	2517/ 4,336,338	2011-2012	[90]	12.5	0.004

REFERENCES

1. Brewer T, Colditz GA. Postmarketing surveillance and adverse drug reactions: current perspectives and future needs. *JAMA*. 1999 Mar 3;281(9):824-9. PMID:10071004. DOI: 10.1001/jama.281.9.824.
2. Scott HD, Thacher-Renshaw A, Rosenbaum SE, et al. Physician reporting of adverse drug reactions: Results of the Rhode Island Adverse Drug Reaction Reporting Project. *JAMA* 1990;263:1785-1788. PMID:2313850.
3. Hoang T, Liu J, Pratt N, et al. Authenticity and credibility aware detection of adverse drug events from social media. *Int J Med Inform*. 2018Dec;120:101-115. PMID:30409335. DOI:10.1016/j.ijmedinf.2018.09.002.
4. Bright RA, Nelson RC. Automated support for pharmacovigilance: a proposed system. *Pharmacoepidemiol and Drug Safety*. 2002;11(2):121-125. PMID: 11998536. DOI:10.1002/pds.684.
5. Samore MH, Evans RS, Lassen A, et al. Surveillance of medical device-related hazards and adverse events in hospitalized patients. *JAMA* 2004;291:325-34. PMID: 14734595. DOI:10.1001/jama.291.3.325.
6. Bright RA. Strategy for surveillance of adverse drug events. *Food Drug Law J* 2007;62(3):605-615. PMID:17915403.
7. Classen D, Li M, Miller S, Ladner D. An electronic health record-based real-time analytics program for patient safety surveillance and improvement. *Health Aff (Millwood)*. 2018Nov;37(11):1805-1812. PMID: 30395491. DOI:10.1377/hlthaff.2018.0728.
8. Wang L, Rastegar-Mojarad M, Ji Z, et al. Detecting pharmacovigilance signals combining electronic medical records with spontaneous reports: a case study of conventional disease-modifying antirheumatic drugs for rheumatoid arthritis. *Front Pharmacol*. 2018Aug7;9:875. PMID:30131701. DOI:10.3389/fphar.2018.00875.
9. Alghamdi AA, Keers RN, Sutherland A, Ashcroft DM. Prevalence and nature of medication errors and preventable adverse drug events in paediatric and neonatal intensive care settings: a systematic review. *Drug Saf*. 2019Dec;42(12):1423-1436. PMID:31410745. DOI:10.1007/s40264-019-00856-9.
10. Molina FJ, Rivera PT, Cardona A, et al. Adverse events in critical care: Search and active detection through the trigger tool. *World J Crit Care Med*. 2018Feb4;7(1):9-15. PMID:29430403. DOI:10.5492/wjccm.v7.i1.9.

11. Report to Congress: Update on the adoption of health information technology and related efforts to facilitate the electronic use and exchange of health information. Office of the National Coordinator for Health Information Technology, US Department of Health and Human Services. 2016 Feb. https://www.healthit.gov/sites/default/files/Attachment_1_-_2-26-16_RTC_Health_IT_Progress.pdf.
12. Melton GB, Hripcsak G. Automated detection of adverse events using natural language processing of discharge summaries. *J Am Med Inform Assoc*. 2005;12:448-457. PMID:15802475. DOI:10.1197/jamia.M1794.
13. Classen D, Li M, Miller S, Ladner D. An electronic health record-based real-time analytics program for patient safety surveillance and improvement. *Health Aff (Millwood)*. 2018Nov;37(11):1805-1812. PMID:30395491. DOI:10.1377/hlthaff.2018.0728.
14. Patadia VK, Schuemie MJ, Coloma PM, et al. Can electronic health records databases complement spontaneous reporting system databases? A historical-reconstruction of the association of rofecoxib and acute myocardial infarction. *Front Pharmacol*. 2018Jun 6;9:594. PMID:29928230. DOI:10.3389/fphar.2018.00594.
15. Zhou L, Siddiqui T, Seliger SL, et al. Text preprocessing for improving hypoglycemia detection from clinical notes - A case study of patients with diabetes. *Int J Med Inform*. 2019Sep;129:374-380. PMID:31445280. DOI:10.1016/j.ijmedinf.2019.06.020.
16. Dejam A, Malley BE, Feng M, et al. The effect of age and clinical circumstances on the outcome of red blood cell transfusion in critically ill patients. *Crit Care* 2014;18(4):487. PMID:25175389. DOI:[10.1186/s13054-014-0487-z](https://doi.org/10.1186/s13054-014-0487-z).
17. Carson JL, Grossman BJ, Kleinman S, et al. Red blood cell transfusion: a clinical practice guideline from the AABB. *Ann Intern Med*. 2012; 57:49-58. PMID:22751760. DOI:10.7326/0003-4819-157-1-201206190-00429.
18. Holness L, Knippen MA, Simmons L, Lachenbruch PA. Fatalities caused by TRALI. *Transfus Med Rev*. 2004;18(3):184-188. PMID:15248168. DOI:10.1016/j.tmr.2004.03.004.
19. Menis M, Anderson SA, Forshee RA, et al. Transfusion-associated circulatory overload (TACO) and potential risk factors among the inpatient US elderly as recorded in Medicare administrative databases during 2011. *Vox Sang*. 2014Feb;106(2):144-52. PMID:23848234. DOI:10.1111/vox.12070.

20. Menis M, Anderson SA, Forshee RA, et al. Transfusion-related acute lung injury and potential risk factors among the inpatient US elderly as recorded in Medicare claims data, during 2007 through 2011. *Transfusion*. 2014;54:2182-2193. PMID:24673344 DOI:10.1111/trf.12626.
21. Reardon JM, Harmon KJ, Schult GC, Staton CA, Waller AE. Use of diagnosis codes for detection of clinically significant opioid poisoning in the emergency department: A retrospective analysis of a surveillance case definition. *BMC Emerg Med*. 2016Feb8;16:11. PMID:26856978. DOI:10.1186/s12873-016-0075-4.
22. Hargrove J, Waller A. Motor vehicle crash case definitions and how they impact injury surveillance. *N C Med J*. 2018Nov-Dec;79(6):351-357. PMID: 30397080. DOI:10.18043/ncm.79.6.351.
23. Wiese AD, Griffin MR, Stein CM, et al. Validation of discharge diagnosis codes to identify serious infections among middle age and older adults. *BMJ Open*. 2018Jun19;8(6):e020857. PMID:29921683. DOI:10.1136/bmjopen-2017-020857.
24. Johnson AEW, Pollard TJ, Shen L, et al. MIMIC-III, a freely accessible critical care database. *Sci Data*. 2016;3:160035. PMID: 27219127. DOI:[10.1038/sdata.2016.35](https://doi.org/10.1038/sdata.2016.35).
25. MIMIC-III Critical Care Database. <https://mimic.physionet.org/about/mimic/>.
26. Code of Federal Regulations Title 45 Part 46 Protection of Human Subjects, Subpart A—Basic HHS Policy for Protection of Human Research Subjects, §46.101 (b) (4). 2000 Oct 1. <https://www.govinfo.gov/content/pkg/CFR-2000-title45-vol1/pdf/CFR-2000-title45-vol1-part46.pdf>.
27. Slee VN. The International Classification of Diseases: Ninth Revision (ICD-9). *Ann Intern Med*. 1978;88(3):424-426. PMID:629506. DOI:10.7326/0003-4819-88-3-424.
28. Statistical functions (scipy.stats). The SciPy Community. 2019. <https://docs.scipy.org/doc/scipy/reference/stats.html>.
29. Pandas. Python Software Foundation. <https://pypi.org/project/pandas/>.
30. The Web's Free ICD-9-CM Medical Coding Reference. ICD9Data.com. <http://www.icd9data.com/>.
31. Comparison of two rates. MedCalc Software Ltd. 2020. https://www.scistat.com/statisticaltests/rate_comparison.php.
32. Altman DG, Bland JM. How to obtain the confidence interval from a P value. *BMJ* 2011;343-344. PMID:21824904. DOI:10.1136/bmj.d2090.

33. Stanworth SJ, Tinmouth AT. Plasma transfusion and use of albumin. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;287-297. ISBN:978-1-444-35886-5.
34. Omi EC, Gamelli RL. Transfusion therapy in the care of trauma and burn patients. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;589-603. ISBN:978-1-444-35886-5.
35. Arnold DM, Smith JW, Warkentin TE. Management of immune thrombocytopenia. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;344-375. ISBN:978-1-444-35886-5.
36. Juffermans NP, Walsh TS. Introduction. IN: Juffermans NP, Walsh TS, eds. *Transfusion in the Intensive Care Unit*. Springer, NY, 2015;1. DOI:10.1007/978-3-319-08735-1_1.
37. Kennedy MS, Wu HM. Transfusion therapy and transfusion in transplantation. IN: Harmening D, ed. *Modern Blood Banking and Transfusion Practices*, 5th ed. F.A. Davis Company, Philadelphia, 2005;303-321. ISBN:9780803612488.
38. Carvalho B, Quiney NF. 'Near-miss' hyperkalaemic cardiac arrest associated with rapid blood transfusion. *Anaesthesia*. 1999Nov;54(11):1094-6. PMID:10540099. DOI:10.1046/j.1365-2044.1999.01109.x.
39. Heafner T, Bews K, Kalra M, et al. Transfusion timing and postoperative myocardial infarction and death in patients undergoing common vascular procedures. *Ann Vasc Surg*. 2019. PMID:31626929. DOI:10.1016/j.avsg.2019.08.090.
40. Schmidt AE, Refaai MA, Blumberg N. Platelet transfusion and thrombosis: More questions than answers. *Semin Thromb Hemost*. 2016 Mar;42(2):118-24. PMID:26716501. DOI:10.1055/s-0035-1564833.
41. Chang S-L, Huang Y-L, Lee M-C, et al. Association of varicose veins with incident venous thromboembolism and peripheral artery disease. *JAMA*. 2018; 319(8):807-817. PMID:29486040. DOI:10.1001/jama.2018.0246.
42. Rossetti A, Buchs NC, Breguet R, et al. Transarterial embolization in acute colonic bleeding: review of 11 years of experience and long-term results. *Int J Colorectal Dis*. 2013Jun;28(6):777-82. PMID:23208010. DOI:10.1007/s00384-012-1621-5.

43. Vamvakas EC, Bordin JO, Blajchman MA. Immunomodularity and proinflammatory effects of allogeneic blood transfusion. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;699-717. ISBN:978-1-444-35886-5.
44. Davenport RD. Hemolytic transfusion reactions. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;811- 825. ISBN:978-1-444-35886-5.
45. Raife TJ, Rose JS, Lentz SR. Bleeding from acquired coagulation defects and antithrombotic therapy. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;376-390. ISBN:978-1-444-35886-5.
46. Debus ES, Mueller-Huelsbeck S, Koelbel T, Larena-Avellaneda A. Intestinal ischemia. *Int J Colorectal Dis*. 2011 Sep;26(9):1087-97. PMID: 21541663 DOI:10.1007/s00384-011-1196-6.
47. Marks PW. Thrombocytopenia and platelet transfusion. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;199-210. ISBN:978-1-444-35886-5.
48. Carson JL, Hébert P. Anemia and red blood cell transfusion. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;131-148. ISBN:978-1-444-35886-5.
49. Halaweh O, Usmani SZ, Abouasaleh R. Barrett's esophagitis related bronchoesophageal fistula - the diagnostic value of persistent air leak in the ventilated setting. *McGill J Med* 2007; 10(2):96-98. PMID: 18523545. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2323478/>.
50. Shears M, Alhazzani W, Marshall JC, et al. Stress ulcer prophylaxis in critical illness: a Canadian survey. *Can J Anaesth*. 2016Jun; 63(6):718-724. PMID:26911559. DOI:10.1007/s12630-016-0612-3.
51. Rashid F, Aber A, Iftikhar SY. A review on gastric diverticulum. *World J Emerg Surg*. 2012;7:1. PMID: 22257431. DOI:10.1186/1749-7922-7-1.
52. Warkentin TE, Moore JC, Anand SS, et al. Gastrointestinal bleeding, angiodysplasia, cardiovascular disease, and acquired von Willebrand syndrome. *Transfus Med Rev*. 2003;17(4):272-286. PMID:14571395. DOI: 10.1016/S0887-7963(03)00037-3.
53. Ivascu FA, Janczyk RJ, Bair HA, Bendick PJ, Howells GA. Spontaneous retroperitoneal hemorrhage. *Am J Surg*. 2005;189(3):345-347. PMID:15792766. DOI:10.1016/j.amjsurg.2004.11.020.

54. Wang CY, Won CW, Shieh MJ. Aggressive colonoscopic approaches to lower intestinal bleeding. *Gastroenterol Jpn*. 1991;26:125–128. PMID:1884944. DOI:10.1007/BF02779280.
55. Bacon BR. Cirrhosis and its complications. In: Jameson JL, Fauci AS, Kasper DL, et al, eds. *Harrison's Principles of Internal Medicine*, 20th ed. McGraw-Hill Medical. ISBN:978-1-259-64404-7. Accessed 2019. <https://accessmedicine.mhmedical.com/content.aspx?bookid=2129§ionid=192283819>.
56. Briglia AE, Anania FA. Hepatorenal syndrome: Definition, pathophysiology, and intervention. *Crit Care Clin*. 2002 Apr;18(2):345-73. PMID:12053838. DOI:10.1016/s0749-0704(01)00003-3.
57. Morgan DJ, Kellerman R. Epistaxis: evaluation and treatment. *Prim Care*. 2014 Mar;41(1):63-73. PMID:24439881. DOI: 10.1016/j.pop.2013.10.007.
58. Ittrich H, Bockhorn M, Klose H, Simon M. The diagnosis and treatment of hemoptysis. *Dtsch Arztebl Int*. 2017 Jun; 114(21):371-381. PMID:28625277. DOI: 10.3238/arztebl.2017.0371.
59. Gourgiotis S, Villias C, Germanos S, et al. Acute limb compartment syndrome: A review. *J Surg Education*. 2007; 64(3):178-186. PMID:17574182. DOI:10.1016/j.jsurg.2007.03.006.
60. Sosa G, Gandham N, Landeras V, et al. Abdominal compartment syndrome. *Dis Mon*. 2019 Jan;65(1):5-19. PMID:30454823. DOI:10.1016/j.disamonth.2018.04.003.
61. Selvan K, Edriss H, Sigler M, Nugent KM. Complications and resource utilization associated with mechanical ventilation in a medical intensive care unit in 2013. *J Intensive Care Med*. 2017 Feb; 32(2):146-150. PMID:26474803. DOI: 10.1177/0885066615612413.
62. Rabinstein AA. Vascular myelopathies. *Continuum (Minneap Minn)*. 2015 Feb;21(1 Spinal Cord Disorders):67-83. PMID:25651218. DOI:10.1212/01.CON.0000461085.79241.e0.
63. Radford MJ, Johnson RA, Daggett WM Jr, et al. Ventricular septal rupture: a review of clinical and physiologic features and an analysis of survival. *Circulation*. 1981;64(3):545-53. PMID: 7020978. DOI: 10.1161/01.cir.64.3.545.
64. Sheth S. Transfusional iron overload. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;858-869. ISBN:978-1-444-35886-5.
65. Sharma S, Sharma P, Tyler LN. Transfusion of blood and blood products: indications and complications. *Am Fam Physician*. 2011 Mar 15;83(6):719-24. PMID: 21404983. <https://www.aafp.org/afp/2011/0315/p719.html>.

66. Harmening DM, Steffey NB, Prihoda LA, Green REB. Autoimmune hemolytic anemias. IN: Harmening D, ed. *Modern Blood Banking and Transfusion Practices*, 5th ed. F.A. Davis Company, Philadelphia, 2005;396-430. ISBN:9780803612488.
67. Duffy TP. Autoimmune hemolytic anemia and paroxysmal nocturnal hemoglobinuria. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;321-343. ISBN:978-1-444-35886-5.
68. Gibson C, Berliner N. How we evaluate and treat neutropenia in adults. *Blood*. 2014Aug;124(8):1251-8. PMID:24869938. DOI:[10.1182/blood-2014-02-482612](https://doi.org/10.1182/blood-2014-02-482612).
69. Hellstern P. Composition and mechanisms of blood coagulation and fibrinolysis. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;247-259. ISBN:978-1-444-35886-5.
70. Tormey CA, Edward L. Snyder EL. Transfusion support for the oncology patient. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;482-497. ISBN:978-1-444-35886-5.
71. Pomper GJ. Febrile, allergic, and nonimmune transfusion reactions. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;826-846. ISBN:978-1-444-35886-5.
72. Wallis JP, Sachs UJH. Transfusion-related acute lung injury. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;870-884. ISBN:978-1-444-35886-5.
73. Vianna NJ. *Lymphoreticular Malignancies: Epidemiologic and Related Aspects*. Medical & Technical Publishing Co. Ltd, Lancaster, UK, 1975; ISBN:839108133.
74. Muslimani A, Chisti MM, Nadeau L, et al. How we treat tumor lysis syndrome. *Oncology J* 2011;25(4):369-75. PMID:21618960. <https://www.cancernetwork.com/view/how-we-treat-tumor-lysis-syndrome>.
75. Park YA, Brecher ME. Bacterial contamination of blood products. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;771-790. ISBN:978-1-444-35886-5.

76. Cook LO, Wise SC, Larison PJ. Adverse effects of blood transfusion. IN: Harmening D, ed. *Modern Blood Banking and Transfusion Practices*, 5th ed. F.A. Davis Company, Philadelphia, 2005;336-358. ISBN:9780803612488.
77. Alter HJ, Esteban-Mur JI. Transfusion-transmitted hepatitis. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;718-745. ISBN:978-1-444-35886-5.
78. Barbara JAJ, Dow BC. Retroviruses and other viruses. IN: Simon TL, Snyder EL, Solheim BG, et al, eds. *Rossi's Principles of Transfusion Medicine*, 4th ed. Blackwell Publishing Ltd. 2009;746-759. ISBN:978-1-444-35886-5.
79. Williams EF, Jarreau PC, Zitzmann MP. Transfusion-transmitted disease. IN: Harmening D, ed. *Modern Blood Banking and Transfusion Practices*, 5th ed. F.A. Davis Company, Philadelphia, 2005;359-382. ISBN:9780803612488.
80. Finley C, Clifton J, FitzGerald JM, Yee J. Empyema: An increasing concern in Canada. *Can Respir J*. 2008 Mar; 15(2): 85–89. PMID:18354748. DOI:10.1155/2008/975312.
81. Kalanuria AA, Ziai W, Mirski M. Ventilator-associated pneumonia in the ICU. *Crit Care*. 2014; 18(2): 208. PMID:25029020. DOI:10.1186/cc13775.
82. Wang J-M, Lim H_K. Necrotizing fasciitis: eight-year experience and literature review. *Braz J Infect Dis*. 2014;18 (2):137-143. PMID:24275377. DOI:10.1016/j.bjid.2013.08.003.
83. Cross MR, Greenwald MF, Dahhan A. Esophageal perforation and acute bacterial mediastinitis: Other causes of chest pain that can be easily missed. *Medicine (Baltimore)*. 2015 Aug; 94(32): e1232. PMID: 26266352. DOI:10.1097/MD.0000000000001232.
84. Cheville AL, McGarvey CL, et al. Lymphedema management. *Semin Radiat Oncol*. 2003 Jul;13(3):290-301. PMID:12903017. DOI:10.1016/S1053-4296(03)00035-3.
85. Karaoren G, Bakan N, Tomruk SG, et al. Bilaterally symmetrical lower extremity compartment syndrome following massive transfusion. *Case Rep Orthop*. 2016;2016:2718421. PMID:26885421. DOI: 10.1155/2016/2718421.
86. Mutlu GM, Mutlu EA, Factor P. GI complications in patients receiving mechanical ventilation. *Chest*. 2001 Apr;119(4):1222-41. PMID:11296191. DOI:10.1378/chest.119.4.1222.

87. Stangroom J. Easy Fisher exact test calculator. Social Science Statistics. 2020.
<https://www.socscistatistics.com/tests/fisher/default2.aspx>.
88. Dunn OJ. Confidence intervals for the means of dependent, normally distributed variables. *J Am Stat Assoc*. 1959;54:613-21. <https://www.jstor.org/stable/pdf/2282541.pdf>.
89. Menis M, Forshee RA, Anderson SA, et al. Posttransfusion purpura occurrence and potential risk factors among the inpatient US elderly, as recorded in large Medicare databases during 2011 through 2012. *Transfusion* 2015;55:284-295. PMID:25065878. DOI:10.1111/trf.12782.
90. Menis M, Forshee RA, Anderson SA, et al. Febrile non-haemolytic transfusion reaction occurrence and potential risk factors among the U.S. elderly transfused in the inpatient setting, as recorded in Medicare databases during 2011-2012. *Vox Sang*. 2015;108:251-261. PMID: 25470076. DOI:10.1111/vox.12215.
91. Bates DW, Cullen DJ, Laird N. et al. Incidence of adverse drug events and potential adverse drug events: implications for prevention. *JAMA*.1995;274:29-34. PMID:7791255. DOI:10.1001/jama.1995.03530010043033.
92. Kane-Gill SL, Kirisci L, Verrico MM, Rothschild JM. Analysis of risk factors for adverse drug events in critically ill patients. *Crit Care Med*. 2012;40(3):823–828. PMID:22036859. DOI:10.1097/CCM.0b013e318236f473.
93. Rivkees SA, Szarfman A. Dissimilar hepatotoxicity profiles of propylthiouracil and methimazole in children. *J Clin Endocrinol Metab*. 2010Jul;95(7):3260-3267. PMID:20427502. DOI:10.1210/jc.2009-2546.

Appendix to Bright et al, “Use of Diagnosis Codes for Blood Transfusion Adverse Events in Electronic Health Records”

Table A1. Codes for definitions of T (transfused) and Excluded groups. See main paper for methods.

Group	ICD9 code	Item code	Long name
T	9901		Exchange transfusion
T	9903		Other transfusion of whole blood
T	9904		Transfusion of packed cells
T	9905		Transfusion of platelets
T	9907		Transfusion of other serum
T		783	Blood Products
T		5649	PRBCs
T		5661	PLTS
T		5751	PRBCs
T		6174	PLT transfusion
T		7597	PRBC
T		30001	Packed 'BC's
T		30002	Whole Blood
T		30004	Washed P'BC's
T		30005	Fresh Frozen Plasma
T		30006	Platelets
T		30103	OR FFP
T		30104	OR Packed 'BC's
T		30105	OR Platelets
T		30179	P'BC's
T		30180	Fresh Froz Plasma
T		42239	CC7 PRBC
T		42323	er ffp
T		42324	er prbc
T		42588	VICU PRBC
T		42740	prbc in er
T		43009	ffp pacu
T		43010	Packed cells pacu
T		44044	FFP Drip

Group	ICD9 code	Item code	Long name
T		44172	FFP GTT
T		44236	E.R. FFP
T		44560	blood
T		44819	FFP ON FARR 2
T		44905	ed blood products
T		45669	ED FFP
T		46122	ER in FFP
T		46124	er in prbc
T		46407	ED PRBC
T		46410	angio FFP
T		46418	ER ffp
T		46530	Floor FFP
T		46612	E.R. prbc
T		46684	ER FFP
T		220969	Filterrythrocytestes
T		220970	Fresh Frozen Plasma
T		220971	ESDEP (Solvent / Detergent Virus-Inactivated Plasma)
T		220972	Octopals
T		220996	Packed Red Cells
T		221004	Thrombocyte suspension
T		221013	Whole Blood
T		225168	Packed Red Blood Cells
T		225170	Platelets
T		226367	OR FFP Intake
T		226368	OR Packed RBC Intake
T		226369	OR Platelet Intake
T		227070	PACU Packed RBC Intake
T		227071	PACU Platelet Intake
T		227072	PACU FFP Intake
Excluded	9900		Perioperative autologous transfusion of whole blood or blood components
Excluded	9902		Transfusion of previously collected autologous blood
Excluded		30053	Cell Saver
Excluded		30092	Autotransfusion
Excluded		30094	Other Blood Products
Excluded		30106	OR Autologous Blood
Excluded		40801	plasma pheresis

Group	ICD9 code	Item code	Long name
Excluded		41381	PLASMAPHERESIS
Excluded		41383	PLASMAPHORESIS
Excluded		41810	plasmaphoresis
Excluded		42538	cell saver/or
Excluded		42629	CELL SAVER IN OR
Excluded		42784	cell saver OR
Excluded		42801	cell saver in or
Excluded		42830	OR CELL SAVER
Excluded		42866	cell saver in OR
Excluded		44064	plasma phoresis
Excluded		44511	Plasmapheresis
Excluded		44515	PLASMAPHERESIS IN
Excluded		44517	pheresis ivf
Excluded		44528	plasmapheresis
Excluded		44856	plasmaphersis
Excluded		44861	Pheresis
Excluded		44873	plasmapharesis
Excluded		44878	Pheresis Intake
Excluded		44912	pheresis.
Excluded		44919	Plasmaphoresis
Excluded		45214	OR cellsaver
Excluded		45379	OR CELL Saver
Excluded		45839	partial exchange
Excluded		45878	cellsaver
Excluded		46078	OR cell saver
Excluded		46222	Plasma pheresis
Excluded		46384	plasma pheresis in
Excluded		46599	OR Cell Saver
Excluded		221193	Auto transfusion
Excluded		225173	Cell Saver
Excluded		226370	OR Autologous Blood Intake
Excluded		226372	OR Cell Saver Intake
Excluded		227532	Plasma Pheresis
Excluded		227551	Plasma Pheresis.