Efficacy of B₁₂ fortified nutrient bar and yogurt in improving plasma B₁₂ concentrations – results from two double blind randomised placebo controlled trials

Chittaranjan Yajnik^{1*}, Sonal Kasture¹, Vaishali Kantikar¹, Himangi Lubree¹, Dattatray Bhat¹,
Deepa Raut¹, Nilam Memane¹, Aboli Bhalerao¹, Rasika Ladkat¹, Pallavi Yajnik¹, Sudhir
Tomar², Tejas Limaye¹, Sanat Phatak¹

Affiliations:

1. King Edward Memorial Hospital Research Center, Pune, Maharashtra, India.
2. ICAR-National Dairy Research Institute, Karnal, Haryana, India.

***Corresponding Author:**

Prof. Chittaranjan Yajnik,

Diabetes Unit, King Edward Memorial Hospital Research Center,

Sardar Moodiar Road, Rasta Peth, Pune – 411011.

Maharashtra, India.

csyajnik@gmail.com

Abstract

Background: Dietary vitamin B₁₂ (B₁₂) deficiency is common in Indians. It may affect hematologic and neurocognitive systems and maternal deficiency predisposes offspring to neural tube defects and non-communicable disease. Long-term tablet supplementation is not sustainable.

Objective: To study efficacy of B₁₂ fortified nutrient bar and yogurt in improving plasma B₁₂ concentrations in children and adults.

Methods: Two double-blind, placebo-controlled randomised directly observed therapy (DOT) trials were conducted: 1. Healthy children (10–13Y) were fed nutrient bar fortified with B₁₂ (2 mcg), multiple micronutrients (B₁₂ 1.8 mcg) or placebo for 120 days. 2. Healthy adults (18–50Y) were fed yogurt fortified with B₁₂ (2 mcg) or *Propionibacterium* (1×10^8 cfu/g) or placebo for 120 days. B₁₂, folate, homocysteine and hemoglobin concentrations and anthropometry were measured before and post intervention.

Results: We randomised 164 children and 118 adults; adherence was 96% and 82% respectively. In children, B₁₂ fortified bars increased B₁₂ concentrations significantly above baseline (B₁₂ alone: median +91 pmol/l, B₁₂+ multiple micronutrients: +82 pmol/l) compared to placebo. In adults, B₁₂ fortified yogurt increased B₁₂ significantly (median +38 pmol/l) compared to placebo, but *Propionibacterium* did not. In both trials, homocysteine fell significantly with B₁₂ supplementation. There was no significant difference in different groups in anthropometry and hemoglobin.

Conclusions: B₁₂ fortified foods are effective in improving B₁₂ status in Indian children and adults. They could be used to improve vitamin B₁₂ status in the national programs for children, adolescents and women of reproductive age. They could also be used as over the counter products.

Keywords: Vitamin B₁₂ deficiency, B₁₂ fortified foods, nutrient bar and yogurt, randomized controlled trial, India

Brief Highlights:

Vitamin B12 deficiency is common in India. Culturally acceptable fortified foods will help reduce it. We performed two RCTs (children and adults) with B12 fortified nutrient bar and yogurt at near RDA doses and found significant improvement in B12 status. This has important implications for nutritional policy.

Introduction

Low Vitamin B₁₂ (B₁₂) status is widely prevalent in vegetarian populations¹. Though asymptomatic in many individuals, deficiency is associated with haematologic, neurocognitive and cardiovascular manifestations in some². Low maternal B₁₂ status during pregnancy is linked to increased risk of neural tube defects (NTDs), preeclampsia, gestational diabetes, preterm delivery, fetal growth restriction, as well as to increased risk of future adiposity and insulin insensitivity in the child³.

Only bacteria synthesize B₁₂ naturally⁴. Animals get their B₁₂ from food and water colonised with B₁₂ producing microorganisms. Foods rich in B₁₂ include meat, liver, fish, eggs, and milk. Plant foods do not contain B₁₂. Low B₁₂ status in developing countries is largely attributed to low intake of animal-origin foods, either because of cultural and religious ethical practices (Hindu, Jain and Buddhist), or due to poverty, which precludes expensive non-vegetarian foods from diet⁵. Low B₁₂ status is common in Indians, both in those living in India as well as those migrated abroad⁶. This is largely attributable to vegetarianism but not to pernicious anemia (malabsorption due to lack of intrinsic factor). Our research showed that Vit B₁₂ deficiency is common in and around Pune, India⁶, despite normal B₁₂ absorption in > 85% of the population⁷. B₁₂ status improved in community based trials by supplementation with capsules containing 2 mcg B₁₂ per day⁸ and drinking 400 ml of milk daily⁹.

Increasing intake of animal origin foods to improve vit B₁₂ status has obvious practical limitations in Indians. Though low dose supplements are successful in a trial setting, they are unlikely to have a big impact in public health owing to poor long-term adherence to medications in asymptomatic individuals. Food fortification may be an effective way to improve B₁₂ status in Indians and other vegetarian populations. Fortification of breakfast cereals and milk with B₁₂ is common in many countries but not in India. In recent years,

probiotics including *Propionibacterium* have been claimed to improve B₁₂ status¹⁰. This could also be of help in Indians and warrants further research.

As part of our research to improve B₁₂ status of the population, we tested two B₁₂ fortified food items - i) A nutrient bar in school children, and ii) Yogurt in adults, for their efficacy to improve B₁₂ status. We also tested a *Propionibacterium* species claimed to produce substantial quantities of B₁₂ as a separate fortificant for yogurt.

Methods:

Study Design and Intervention: We conducted two double blind, placebo-controlled, randomized, directly observed therapy (DOT) trials. We excluded those with chronic medical illness, those on vitamin supplementation and those with very low B₁₂ concentration (<100 pmol/l) or anemia (hemoglobin <10 g/dl) for ethical reasons and were advised appropriate treatment. Randomization was computer generated, and stratification was based on baseline B₁₂ and hemoglobin concentrations (below and above median) to ensure comparable distribution. All participants were instructed not to take any vitamin supplements during the trial. Adherence was calculated for each participant by calculating the number of days of attendance during the trial.

Nutrient bar trial: This was conducted in a village school (Pabal, Pune District, approximately 70 km from Pune) in children aged 10–13 years. School authorities approved participation in the trial. We arranged a meeting with the school children and their parents to explain the trial and invited them to sign an assent and consent. Included children were randomized to 3 groups to receive one bar per day, fortified with a) 2mcg B₁₂ or b) multiple micronutrients (MMN, including 1.82 mcg B₁₂ (UNICEF guidelines)¹¹, or c) no added micronutrients (placebo). All bars were similar in appearance and taste. The children ate the

bars in the school in the morning recess, directly observed by research staff. The duration of supplementation was 120 consecutive school days, from September 2011 to March 2012.

Yogurt trial: The yogurt study was conducted in volunteers aged 18–50 years at the KEM Hospital, Pune. Hospital staff members and their friends were invited to enroll through notice board advertisement. Eligible participants were randomised into 3 groups to receive 100 gm yogurt fortified with a) 2µg B₁₂, b) 1×10⁸cfu/g *Propionibacterium*, and c) without any additions (placebo). The strain used for fortification (*Propionibacterium freudenreichi* subsp. *freudenreichii* ATCC 6207, GRAS certified) was provided by the National Dairy Research Institute (NDRI, Karnal, Haryana, India) and was recommended because it was known to produce B₁₂ in vitro¹². The three yogurt preparations were similar in appearance, taste, flavor and smell. The yogurt was eaten under supervision for 120 consecutive working days between December 2013 and May 2014.

In nutrient bar trial, informed written assent was obtained from the children, and consent was signed by the parents. In yogurt trial, participants provided written consent. Both trials were approved by KEM Hospital Research Centre Ethics Committee. For the nutrient bar trial, permissions were obtained from the school authorities and the District Health Officer. The trials were registered with the Clinical Trials Registry of India (CTRI/2012/07/002799 - nutrient bar, CTRI/2015/04/005703 -yogurt). Adverse events were recorded.

Measurements: Height was measured to the nearest 0.1 cm using a stadiometer (CMS Instruments, London, UK) and body weight to the nearest 0.01 kg using an electronic weighing scale (Model no. HD-358, Tanita Corporation, Japan). At baseline, non-fasting venous blood sample was collected in EDTA vacutainers for measurements of hemogram, B₁₂, folate and homocysteine. Hemogram was measured on a Beckman Coulter Analyzer (AC.T diffTM, Miami, Florida, USA) on the same day. Plasma aliquots were stored (-70° C) until further analysis. B₁₂ was measured by a microbiological assay using a colistin sulfate-

resistant strain of *Lactobacillus leichmannii* (CV <8%)^{13, 14}. Plasma total homocysteine (homocysteine) was measured by fluorescence derivative of monobromobimane using HPLC (inter and intra batch CV <4%). Plasma folate was measured by a microbiological assay using a chloramphenicol-resistant strain of *Lactobacillus Casei* (inter and intra CV <8%)^{15, 16}. The blood and clinical measurements were repeated at the end of the trial.

Statistical analysis: Data are presented as median (25th – 75th centile) for continuous variables and as percentages for categorical variables. Skewed variables were log normalized before analysis. Significance of the differences between baseline and end of trial levels of clinical and biochemical measurements were tested by paired t test for continuous variables and by chi-square test for categorical variables. Significance of difference from placebo group was tested by t-test. Analysis was by intention to treat (ITT) method. The participants who were lost to follow-up were analyzed by ‘last observation carried forward’ (LOCF) method.

For children, we calculated sample size required in each group to demonstrate a 50% rise in plasma B₁₂ concentrations at 5% significance level and found that 55 individuals allowing a dropout rate of 10% will provide a power of more than 90%. For adults, we calculated sample size required in each group to demonstrate a 30% rise in plasma B₁₂ concentrations at 5% significance level and found that 45 individuals allowing a dropout rate of 10% will provide a power of more than 80%.

Results:

Nutrient bar trial: We approached 180 school children and their parents, of whom 178 (99%) agreed to participate. Fourteen children were excluded: 13 with B₁₂ <100 pmol/l and 1 anemic; they were prescribed appropriate treatment (figure -1a). The 164 randomized children (57% girls) were 11.3 (10.9 – 12.0) years old, 139.4 (134.5 – 144.5) cm tall and with

a weight of 28.8 (25.2 – 33.4) kg; there was no significant difference between groups. Plasma vitamin B₁₂, folate and homocysteine concentrations were comparable. Table-1

After intervention, B₁₂ rose significantly by median 91 pmol/l and 82 pmol/l above the baseline in B₁₂ alone and B₁₂ + MMN group respectively (p<0.001) while there was a no change in the placebo group (figure -2). There was a small rise in folate concentrations in the placebo and B₁₂ groups, and substantial rise in the MMN group. Homocysteine reduced by median 1.4 and 3.8 µmol/l in the B₁₂ alone and B₁₂+ MMN groups respectively (p<0.001), and increased by median 1.9 µmol/l (p<0.001) in the placebo group (figure -2). Hemoglobin concentrations did not change with the intervention. There was an average gain of 4.3 (3.6 – 5.2) cm in height and 2.9 (2.0 – 4.4) kg in weight post intervention but no significant difference between the groups.

Comparison with placebo: The post intervention concentrations of B₁₂ and rise above baseline were higher in both B₁₂ supplementation groups compared to those in the placebo group. Rise and post intervention concentrations of folate were higher in the MMN group compared to the placebo and B₁₂ group. Fall in homocysteine concentrations was significantly greater in both the B₁₂ and MMN groups compared to the placebo group (figure -2).

Mean adherence for daily nutrient bar consumption was high and similar in three groups (≥95%). One child dropped out during the trial because of change of school (figure -1a). There were a total of four serious adverse events (hospital admissions for incidental illnesses), none related to the intervention product.

Yogurt trial: Of 178 adult volunteers, 148 consented to participate. Thirty were excluded: 24 with B₁₂ <100 pmol/l, 2 with anaemia and 4 due to medical disorders (figure -1b). The 118 randomized adults (81% women) were 27.0 (27.8 – 35.0) years old, 157.5 (152.8 – 162.5) cm tall, with a weight of 56.5 (49.4 – 64.6) kg and BMI of 22.9 (19.7 – 26.0) kg/m², there was no

significant difference between groups. Plasma vitamin B₁₂, folate and homocysteine concentrations were comparable. Table-2

After intervention, B₁₂ rose significantly by median 38 pmol/l above the baseline in the B₁₂ group (p<0.001) and homocysteine decreased by median 2.7 µmol/l (p<0.001) (figure -2).

There were no significant changes in B₁₂ or homocysteine in the *Propionibacterium* and the placebo groups. There was a small rise in folate concentrations in all groups, but no significant changes in hemoglobin concentrations and weight in any group.

Comparison with placebo: The rise in B₁₂ concentration and fall in homocysteine concentration from baseline was higher in B₁₂ fortified yogurt group compared to the placebo and *Propionibacterium* fortified groups. The fall in the homocysteine concentration was relatively smaller and not significantly different between groups (figure -2).

During the intervention 6 participants discontinued from the B₁₂, 9 from *Propionibacterium* and 9 from plain yogurt group; all for reasons not related to intervention. Ninety four (82%) participants completed the trial (figure -1b). Mean adherence for daily yogurt consumption was moderate and similar in three groups (≥82%). There was one serious adverse event (hospital admission for incidental illness) not related to the intervention product.

Discussion:

Our results show that regular consumption of B₁₂ fortified nutrient bar or yogurt for 4 months significantly improved B₁₂ status. Our population has high prevalence of B₁₂ deficiency, and we used near RDA doses to make it public health relevant. DOT design ensured high adherence and allowed us to explore the full potential of fortification with near RDA doses. The response was proportional to the dose of B₁₂ per kg body weight, and therefore, higher in children compared to adults. Placebo comparison helped highlight the specific effects of vitamin B₁₂ interventions. Nutrient bar and yogurt are two commonly eaten food items in

India, and were well accepted by the participants. Our results make it an attractive public health proposition. The *Propionibacterium* probiotic at the recommended dose did not influence B₁₂ status.

Improvement in B₁₂ status reflected in reduction of circulating homocysteine concentrations, indicating an improvement in the methylation status of the body in both the children and the adults. Addition of other micronutrients (folic acid, B6 and B2) to B₁₂ had an additional effect on lowering of homocysteine in children. There was no effect on hemoglobin concentration in these trials, which might reflect a requirement for a relatively higher dose of these vitamins for hematological effects. In another study of more severely B₁₂ deficient adolescent girls we were able to show improvement in hemoglobin concentration (and peripheral nerve function) with similar daily dose of vit B₁₂ but over a longer period of 11 months.¹⁷ Our results support fortification of commonly eaten food items as vehicles for improving B₁₂ status in populations with low B₁₂ status who do not have the problem of defective absorption.

There are very few studies of exclusive B₁₂ fortification to improve the vitamin status. They have been done in pre-school children (MMN)^{18,19} and in elderly (B₁₂)²⁰, the two groups commonly thought to be at risk of low B₁₂ status. The vehicles for B₁₂ were rice, milk, wheat flour (consumed as bread), common salt and breakfast cereals.^{18, 19, 21-23} In a population with high burden of low B₁₂ status, a multipronged food fortification program may be more successful than isolated food product fortification to improve the B₁₂ status. In our own and others' experience, supplementation with tablets in asymptomatic individuals suffers rapid reduction in adherence. Such supplementation also has complex logistic requirements in field operations in large populations.

We performed our investigations in a population with high prevalence of low B₁₂ status, making it relevant to many such populations in India and other developing countries.²⁴

Participation rates were high. The placebo-controlled, randomized design ensured a causal association for both the trials. Use of DOT approach ensured near complete compliance ($\geq 95\%$) in children, thus allowing full potential of the intervention to be exploited. Our findings will be directly applicable to vulnerable populations (children, adolescents, young adults and pregnant women) where low B₁₂ status may have major implications for growth, development and reproductive health. Possible weaknesses include small sample size and a relatively short period of intervention. Relatively lower adherence in the adults (average 82%) is a reflection of the real-life situation in working middle class and not a reflection on the investigational product. Despite these, there was a rewarding improvement in B₁₂ status.

In summary, we used fortified versions of two commonly consumed food items to improve B₁₂ status in deficient populations. This approach appears superior to use of vitamin tablets, which has proved difficult to sustain over longer periods of time in asymptomatic population. Both food items additionally provide calories, proteins and other nutrients and therefore could be easily adopted in national programs of feeding malnourished children and adults, as well as in the Mid-day Meal programs in schools. There is a scope to perform a longer trial to investigate benefits to physical, cognitive and reproductive outcomes. Both items could be made at low cost by small-scale home industry promoting local economy. Future trials in community will help estimate effectiveness of these food fortifications in deficient populations.

Conflicts of interests: All authors declare no competing interests.

Acknowledgements: The study was funded by Department of Bioechnology (DBT), New Delhi, India. DSM India kindly provided nutrient bars for the study. Hi Tech Biosciences

India Limited, Pune produced yogurt for the trial. We thank the team at Diabetes Unit, KEM Hospital Research Centre, Pune for practical assistance during the trial (Charudatta Joglekar, Sonali Wagle, Vidya Mudliar, Komal Adawani, Rucha Wagh). We thank Dr. Arun Nanivadekar and Dr Mohan Gupte for their guidance. We are grateful to the participants of both the trials.

Authorship:

CY, ST, PY conceptualized and planned the study. SK, VK, HL, RL, PY and CY were involved in the trial conduct. DB, DR, NM performed all the laboratory measurements. Statistical analysis was done by TL and AB. CY, TL, SP and DB wrote the manuscript.

References

1. Green R, Allen LH, Bjørke-Monsen AL, et al. Vitamin B₁₂ deficiency. *Nat Rev Dis Primers*. 2017;3:17040.
2. Stabler SP. Clinical practice. Vitamin B₁₂ deficiency. *N Engl J Med*. 2013;368(2):149–160.
3. Yajnik C. Vitamin B₁₂: An Intergenerational Story. *Nestle Nutr Inst Workshop Ser*. 2020;93:91–102.
4. Smith AD, Warren MJ, Refsum H. Vitamin B₁₂. *Adv Food Nutr Res*. 2018;83:215–279.
5. Allen LH. Causes of vitamin B₁₂ and folate deficiency. *Food Nutr Bull*. 2008;29:(2 Suppl):S20-S37.
6. Yajnik CS, Deshpande SS, Lubree HG, et al. Vitamin B12 deficiency and hyperhomocysteinemia in rural and urban Indians. *J Assoc Physicians India*. 2006;54:775-782.
7. Bhat DS, Thuse N, Joglekar C, et al. 6E-3 Absorption of oral vitamin B12 in Indians; the Pune Maternal Nutrition Study. *Early Hum Dev*. 2007;83.10.1016/S0378-3782(07)70167-6.
8. Deshmukh US, Joglekar CV, Lubree HG, et al. Effect of physiological doses of oral vitamin B12 on plasma homocysteine: a randomized, placebo-controlled, double-blind trial in India. *Eur J Clin Nutr*. 2010;64(5):495-502.
9. Naik S, Bhide V, Babhulkar A, et al. Daily milk intake improves vitamin B-12 status in young vegetarian Indians: an intervention trial. *Nutr J*. 2013;12:136.
10. Poonam, Pophaly SD, Tomar SK, De S, Singh R. Multifaceted attributes of dairy propionibacteria: a review. *World J Microbiol Biotechnol*. 2012;28(11):3081-3095.

11. World Health Organization. Preventing and controlling micronutrient deficiencies in populations affected by an emergency. Joint statement by the World Health Organization, the World Food Programme and the United Nations Children's Fund. <https://tinyurl.com/y9xfhgpg> . Published: 2007. Accessed March, 2020.
12. Fang H, Kang J, Zhang D. Microbial production of vitamin B12: a review and future perspectives. *Microb Cell Fact*. 2017;16(1):15.
13. Kelleher BP, Walshe KG, Scott JM, O'Brian SD. Microbiological assay for vitamin B12 with use of a colistin-sulfate-resistant organism. *Clin Chem*. 1987;33(1):52-54.
14. Kelleher BP, Broin SD. Microbiological assay for vitamin B12 performed in 96-well microtitre plates. *J Clin Pathol*. 1991;44(7):592-595.
15. Horne DW, Patterson D. Lactobacillus casei microbiological assay of folic acid derivatives in 96-well microtiter plates. *Clin Chem*. 1988;34(11):2357-2359.
16. Tamura T, Freeberg LE, Cornwell PE. Inhibition of EDTA of growth of Lactobacillus casei in the folate microbiological assay and its reversal by added manganese or iron. *Clin Chem*. 1990;36(11):1993.
17. Yajnik CS, Behere RV, Bhat DS, et al. A physiological dose of oral vitamin B-12 improves hematological, biochemical-metabolic indices and peripheral nerve function in B-12 deficient Indian adolescent women. *PLoS One*. 2019;14(10):e0223000.
18. Thankachan P, Rah JH, Thomas T, et al. Multiple micronutrient-fortified rice affects physical performance and plasma vitamin B-12 and homocysteine concentrations of Indian school children. *J Nutr*. 2012;142(5):846-852.
19. Kuriyan R, Thankachan P, Selvam S, et al. The effects of regular consumption of a multiple micronutrient fortified milk beverage on the micronutrient status of school children and on their mental and physical performance. *Clin Nutr*. 2016;35(1):190-198.

20. Brito A, Verdugo R, Hertrampf E, et al. Vitamin B-12 treatment of asymptomatic, deficient, elderly Chileans improves conductivity in myelinated peripheral nerves, but high serum folate impairs vitamin B-12 status response assessed by the combined indicator of vitamin B-12 status. *Am J Clin Nutr*. 2016;103(1):250-257.
21. Garrod MG, Buchholz BA, Miller JW, Haack KW, Green R, Allen LH. Vitamin B12 Added as a Fortificant to Flour Retains High Bioavailability when Baked in Bread. *Nucl Instrum Methods Phys Res B*. 2019;438:136-140.
22. Vinodkumar M, Rajagopalan S. Multiple micronutrient fortification of salt. *Eur J Clin Nutr*. 2009;63(3):437-445.
23. Powers HJ, Stephens M, Russell J, Hill MH. Fortified breakfast cereal consumed daily for 12 wk leads to a significant improvement in micronutrient intake and micronutrient status in adolescent girls: a randomised controlled trial. *Nutr J*. 2016;15(1):69.
24. McLean E, de Benoist B, Allen LH. Review of the magnitude of folate and vitamin B12 deficiencies worldwide. *Food Nutr Bull*. 2008;29(2 Suppl):S38-S51.

Table 1: Nutrient bar trial - comparison within and between groups

		Placebo (n=55)			MMN (n=55)			B-12 (n=54)			Significance of difference between post intervention concentrations [#]	
		Baseline	120 days	P*	Baseline	120 days	P*	Baseline	120 days	P*	Placebo & MMN	Placebo & B12
Biochemistry	Vitamin B12 (pM)	183 (141, 230)	192 (135, 277)	0.562	173 (132, 224)	251 (210, 310)	<0.001	188 (156, 227)	282 (208, 373)	<0.001	<0.001	<0.001
	Folate (nm/L)	20.8 (15.6, 27.3)	21.4 (17.0, 28.2)	0.049	19.0 (15.7, 24.8)	45.8 (35.0, 58.3)	<0.001	19.6 (16.1, 22.8)	20.4 (16.4, 31.8)	0.012	<0.001	0.967
	tHcy (μmol/L)	16.4 (14.3, 21.3)	18.9 (15.6, 24.1)	<0.001	17 (13.1, 20.6)	12.7 (10.6, 14.4)	<0.001	15.7 (12.6, 20.7)	14.2 (12.5, 16.3)	<0.001	<0.001	<0.001
	Hemoglobin (g/dl)	13.7 (13.4, 14.2)	13.9 (13.2, 14.3)	0.791	13.7 (13.0, 14.1)	13.5 (12.9, 14.2)	0.980	13.8 (13.1, 14.3)	13.9 (13.4, 14.4)	0.251	0.154	0.946
Compliance			95%			96%			96%			

Numbers are median (25th-75th centile). Differences within the groups are tested by *paired T-test (parametric) and between the groups by

[#]Mann-Whitney test (non-parametric). tHcy: Total Homocysteine.

(The analysis was by intention to treat and those lost to follow-up were ascribed final value by last observation carried forward method

Table 2: Yoghurt trial: comparison within and between groups

		Placebo (n=40)			Propionibacterium (n=39)			B12 (n=39)			Significance of difference between post intervention concentrations [#]	
		Baseline	120 days	P*	Baseline	120 days	P*	Baseline	120 days	P*	Placebo & Propionibacterium	Placebo & B12
Biochemistry	B12 (pM)	151 (123, 239)	167 (120, 234)	0.670	167 (125, 214)	168 (142, 216)	0.460	142 (118, 185)	207 (154, 278)	<0.001	0.212	0.057
	Folate (nm/L)	14 (11.0, 22.3)	18.8 (14.5, 24.4)	0.062	18.1 (13.7, 23.6)	20.5 (17.3, 28.4)	0.009	16 (12.1, 22.6)	19 (17.2, 27.8)	0.001	0.104	0.994
	tHcy (μmol/L)	19.0 (11.9, 34.9)	19.3 (12.0, 32.6)	0.675	16.9 (12.0, 25.0)	17.3 (12.1, 23.4)	0.815	22.7 (14.3, 43.5)	17.3 (12.9, 29.8)	0.001	0.426	0.720
	Hemoglobin (g/dl)	11.5 (10.0, 12.4)	11.9 (10.5, 13.0)	0.015	11.2 (9.5, 12.0)	11.3 (10.1, 12.3)	0.018	11.4 (10.4, 13.4)	11.6 (10.2, 14.0)	0.539	0.511	0.935
Compliance			77%			84%			85%			

Numbers are median (25th – 75th centile). Differences within the groups are tested by *paired T-test (parametric) and between the groups by

[#]Mann-Whitney test (non-parametric). tHcy: Total Homocysteine.

(The analysis was by intention to treat and those lost to follow-up were ascribed final value by last observation carried forward method)

Figure 1

Figure 1 shows flow of participants in two trials. A. Nutrient bar trial was conducted in village school children. B. Yogurt trial was conducted in healthy adult volunteers in an urban setup

Figure 2

Figure 2 (box plot) shows change in plasma vitamin B12 and homocysteine concentrations after supplementation with fortified food products for 120 days. Differences between the groups are tested by Mann-Whitney test. P-values are given as * <0.05, **<0.01, ***<0.001, ns: non significant. MMN: Multi Micro Nutrient (1.8 mcg vitamin B12), B-12: Vitamin B12 (2.0 mcg), Propionibacterium: *Propionibacterium freudenreichi* (1×10^8 cfu/g)