Visual Impairment and Risk of Dementia: the UK Biobank Study

Running title: VI and dementia

Authors

Zhuoting Zhu, MD PhD¹

Danli Shi, MD²

Huan Liao, MD³

Jason Ha, MBBS(Hons), BMedSc(Hons)⁴

Xianwen Shang, PhD^{1,4}

Yu Huang, MD PhD¹

Xueli Zhang, PhD¹

Yu Jiang, MD²

Longyue Li, MBBS⁵

Honghua Yu, MD PhD¹

Wenyi Hu, MBBS¹

Wei Wang, MD PhD²

Xiaohong Yang, MD PhD¹

Mingguang He, MD PhD^{1,2,4}

Affiliations

1. Department of Ophthalmology, Guangdong Academy of Medical Sciences, Guangdong Provincial People's Hospital, Guangzhou, China.
2. State Key Laboratory of Ophthalmology, Zhongshan Ophthalmic Center, Sun Yat-sen University, Guangzhou, China.
3. Neural Regeneration Group, Institute of Reconstructive Neurobiology, University of Bonn, Bonn, Germany.
4. Centre for Eye Research, Melbourne University, East Melbourne, Victoria, Australia.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

5. Sun Yat-sen University, Guangzhou, China.

Word count: Abstract: 149; Whole paper: 2485.

Tables: 3; Supplement Table: 3.

Corresponding author

Mingguang He, MD PhD

Email: mingguang.he@unimelb.edu.au

Xiaohong Yang, MD PhD

Email: syyangxh@scut.edu.cn

Wei Wang, MD PhD

Email: wangw289@mail.sysu.edu.cn

ABSTRACT

INTRODUCTION: The association between visual impairment (VI) and the risk of dementia has been poorly understood. We sought to investigate the VI-dementia relationship in the UK Biobank Study.

METHODS: A total of 117,187 volunteers (aged 40-69 years) deemed free of dementia at baseline were included. Habitual distance visual acuity worse than 0.3 logMAR units in the better-seeing eye was used to define VI. The incident dementia was based on electronically linked hospital inpatient and death records.

RESULTS: During a median follow up of 5.96 years, the presence of VI was significantly associated with incident dementia (HR=1.78, 95% CI: 1.18-2.68, P=0.006). There was a clear trend between the severity of VI and the risk of dementia (P for trend=0.002).

DISCUSSION: Visually impaired individuals were more likely to develop incident dementia, with a progressively greater risk among those with worse visual acuity. Our findings highlight the value of regular vision screening and elimination of VI.

Key words: visual impairment, dementia, ageing

HIGHLIGHTS

1. The association between VI and dementia has been poorly understood;
2. VI is associated with incident dementia in non-demented adults;
3. There is a clear trend between the severity of VI and the risk of dementia;
4. VI may be a marker of increased dementia risk.

RESEARCH IN CONTEXT

SYSTEMATIC REVIEW: We searched and reviewed the literature using traditional sources (e.g., PubMed and GoogleScholar). While the association between VI and cognitive function/decline are increasingly studies, investigation of the association between VI and the risk of dementia has been largely overlooked.

INTERPRETATION: We found that visually impaired individuals were more likely to develop incident dementia, with a progressively greater risk among those with worse visual acuity. Our findings imply that VI may be an important marker of dementia.

FUTURE DIRECTIONS: These findings call for more studies to investigate (a) the role of visual acuity changes on the risk of dementia; (b) the relationship between other components of visual function and incident dementia; (c) the relationship between eye diseases and incident dementia; and (d) the potential benefits of vision rehabilitation on dementia prevention.

Introduction

The number of individuals with visual impairment (VI) and blindness is projected to more than double by 2050 as a result of shifting demographics and ageing population.^{1,2} In addition to vision loss, visually impaired individuals are more likely to suffer from mental disorders (e.g., depression),³ social isolation,⁴ unintentional injuries,⁵ functional disabilities,⁶ and poor survival.^{7, 8}

Dementia is an increasing threat to global health, affecting nearly 46.8 million people worldwide, with this number expected triple by 2050.⁹ Despite this heavy burden, effective treatments for dementia are yet to be developed. Therefore, prevention, early detection and management are essential to address the enormous burden of dementia.

There is mounting evidence that suggests that VI and cognitive decline may be closely related.¹⁰⁻²² Nevertheless, the association between VI and risk of incident dementia has been poorly understood.²³⁻³¹ Besides, the relatively small sample size, focus of late-onset dementia, self-reported visual function and bias of identifying dementia cases in previous studies may mask the true association between VI and incident dementia.

Therefore, we aimed to investigate the association of objectively determined VI and VI severity with incident dementia in a large-scale sample aged 40-69 years from the UK Biobank Study.

Methods

Study Sample

The UK Biobank is a large-scale prospective cohort study, enrolling more than 500,000 individuals (40-69 years) from across the UK, with baseline recruitment taking place during 2006-2010. Details of the rationale, design, and assessments used in the UK Biobank Study have been previously described elsewhere.³² Briefly,

approximately 9.2 million people aged 40-69 years in the UK's National Health Service (NHS) residing near one of 22 assessment centers were invited. A total of 502,505 individuals (response rate of 5.5%) agreed to participate and visited the assessment centers. During the baseline assessment, participants answered comprehensive questionnaires, underwent physical measures, and provided biological samples. They also gave consent to link to their health-related records. Ophthalmic assessments, including logMAR visual acuity (VA), autorefraction, intraocular pressure (IOP), keratometry, corneal biomechanics, and spectral-domain Optical Coherence Tomography (OCT) imaging, were introduced to the baseline assessment in 2009 for 6 assessment centers.

The UK Biobank Study's ethical approval had been granted by the National Information Governance Board for Health and Social Care and the NHS North West Multicenter Research Ethics Committee. Written informed consent was obtained from all participants.

Visual Acuity Testing

The detailed procedure of VA testing for the UK Biobank Study has been described elsewhere.³³ Presenting distance VA was measured with the habitual correction (if any) at 4 m using the logarithm of the minimum angle of resolution (logMAR) chart (Precision Vision, LaSalle, Illinois, USA) on a computer screen. Presenting VA was scored as the total number of correctly read lines, converted to logMAR units. In the present analysis, VI was defined as the presenting VA worse than 0.3 logMAR units (Snellen 20/40) in the better-seeing eye. Based on the presenting VA in the better-seeing eye, the severity of VI was classified as mild ($0.3 < \log\text{MAR} \leq 0.6$; $20/40 < \text{Snellen} \leq 20/80$), moderate ($0.6 < \log\text{MAR} \leq 0.7$; $20/80 < \text{Snellen} \leq 20/100$) and severe ($0.7 < \log\text{MAR}$; $\text{Snellen} < 20/100$).

Ascertainment of Incident Dementia

Dementia cases in the UK Biobank Study were ascertained by combining data from

participants' medical history and record linkage to hospital admissions data and the national death register. Detailed information regarding the algorithms used to combine data from different sources to identify dementia can be found at the following link: https://biobank.ndph.ox.ac.uk/showcase/showcase/docs/alg_outcome_dementia.pdf.

In the present analysis of incident dementia, we excluded all participants who already had a diagnosis of dementia in the hospital admissions data or self-reported dementia at the point of study entry. Follow-up time was calculated as the duration between the date of the first assessment and censored at the date of incident dementia, date of death, date of loss to follow-up, or 1st March 2016, whichever occurred earliest.

Covariates

Factors known to be associated with dementia were included as potential confounders in the present analysis. These variables included age at baseline assessment, sex, ethnicity (recorded as white and non-white), Townsend deprivation indices (an area-based proxy measure for socioeconomic status), education attainment (recorded as college or university degree, and others), family history of dementia (a marker of biological vulnerability), smoking status (recorded as current/previous and never), physical activity level (recorded as above moderate/vigorous/walking recommendation or not), and comorbidities (depression, diabetes, hypertension, and hyperlipidemia).

Self-reporting and/or the score on the Patient Health Questionnaire (PHQ, the first two items) of at least 3 were used to identify participants with depression.³⁴ Hypertension was defined to include those participants who had self-reported, or doctor-diagnosed hypertension, were taking antihypertensive drugs, or had a systolic blood pressure of at least 130 mmHg or a diastolic blood pressure of at least 80 mmHg averaged over two measurements. Diabetes was defined to include those participants who had self-reported or doctor-diagnosed diabetes mellitus, were taking anti-hyperglycaemic medications or using insulin, or had a glycosylated hemoglobin level of $\geq 6.5\%$. Hyperlipidemia was defined to include participants with

doctor-diagnosed hyperlipidemia, were taking lipid-lowering drugs, or had a total cholesterol level ≥ 6.21 mmol/L.

Statistical Analysis

Descriptive statistics, including means and standard deviations (SDs), numbers and percentages, were used to report baseline characteristics of study participants. The unpaired t tests were used to compare means between two groups on continuous variables, and Pearson's χ^2 tests to compare distributions between two groups on categorical variables. The log-rank test was used to compare distributions of incident dementia between VI and non-VI groups. Hazard ratios (HRs) and 95% confidence intervals (CIs) were estimated based on the Cox proportional hazards regression models. Age- and sex-adjusted Cox proportional hazards regression models were used to identify covariates strongly associated with incident dementia. The covariates that were found to be associated with incident dementia in the age- and sex-adjusted models were then adjusted in the multivariable models.

We performed sensitivity analysis classifying VI using two different cutoffs: a presenting VA worse than 0.6 logMAR units (Snellen 20/80) and worse than 0.7 logMAR units (Snellen 20/100) in the better-seeing eye. To reduce the likelihood of accidental inclusion of prevalent cases, we also conducted sensitivity analysis after excluding dementia cases whose date of first recorded diagnosis occurred within one years from the date of baseline assessment.

The proportional hazards assumption was satisfied graphically. All tests were two-sided, and statistical significance was set at a P value of less than 0.05. All analyses were performed using Stata version 13 (StataCorp).

Data availability

The UK Biobank is an open-access resource to researchers through registration of

proposed research. (see ukbiobank.ac.uk/register-apply). Access to the UK Biobank data was granted after registration. The application ID was 62443.

Results

Study Sample

Of the 502,505 participants enrolled in the baseline UK Biobank Study between 2006 and 2010, VA was measured in 117,252 (23.3%) participants. Supplementary Table 1 demonstrates the baseline characteristics of participants with and without VA data. In summary, participants with VA data tended to be slightly older and living in a socioeconomically deprived area compared with those without VA data. There was a similar proportion of men and women, but compared to those without VA data, there was a slightly lower proportion of individuals who were current/former smokers, a slightly higher representation of non-white ethnicity, college or university degree, family history of dementia, above physical activity recommendation, and comorbidities than those without VA data. In the analysis of incident dementia, we excluded participants diagnosed with dementia prior to baseline assessment (n=52) and those who self-reported at the baseline nurse interview (n=13). A total of 117,187 participants were included in the final analysis.

Among the 117,187 participants who were not diagnosed with dementia at the baseline assessment, the mean (SD) age was 56.8 (8.11) years with 54.4% being female. Other baseline characteristics for the study population are shown in Table 1. A total of 4,018 (3.43%) participants had VI and 113,169 (96.6%) did not have VI. Table 1 provides the baseline characteristics for the VI and comparator groups. Participants with VI were more likely to be older, of non-white ethnicity, living in a socioeconomically deprived area and less educated than those without VI. Relative to the comparison group, a slightly lower proportion of visually impaired participants had a family history of dementia, but a higher proportion had depression, diabetes mellitus, and hypertension.

Incident Dementia

The median (interquartile range, IQR) follow-up duration post baseline enrollment was 5.96 (5.77-6.23) years. A total of 438 participants (0.37%) were ultimately classified as having incident dementia. The baseline characteristics stratified by dementia status at follow-up are shown in the Table 2. Age- and gender-adjusted Cox proportional hazards regression models showed that ethnicity, education attainment, Townsend deprivation index, physical activity level, family history of dementia, comorbidities (depression and diabetes mellitus) at baseline were significant risk factors for the incident dementia.

Participants with VI at baseline were more likely to develop dementia at follow-up compared with those without VI (0.87% vs 0.36%, log-rank test, $z = 28.5$, $P < 0.001$). The age- and gender-adjusted Cox proportional hazard regression model found that VI was associated with a 2.01-fold higher risk of subsequent dementia (95% CI: 1.43-2.85, $P < 0.001$, Table 3). After further adjusting for potential confounders, the presence of VI was independently associated with a 78% higher risk of incident dementia (95% CI: 1.18-2.68, $P = 0.006$). There was a significant trend toward higher dementia risk across the VI severity groups (P for trend = 0.002), where the greatest risk for dementia was among those with severe VI (HR=3.53, 95% CI: 1.31-9.49, $P = 0.013$).

Sensitivity Analysis

We observed similar findings of sensitivity analyses using different thresholds for classifying VI, where the risk of dementia was greatest among participants with a presenting VA worse than 0.7 logMAR units for VI (HR=3.42, 95% CI: 1.27-9.19, $P=0.015$), followed by a cutoff of 0.6 logMAR units for VI (HR=2.39, 95% CI: 0.99-5.80, $P=0.052$, Supplementary Table 2). Sensitivity analysis excluding participants with dementia cases firstly diagnosed within one year after the baseline assessment showed comparable results to the main analysis, again verifying the

robustness of our findings (Supplementary Table 3).

Discussion

In this large community-based population of 117,187 initially dementia-free individuals, we found that visually impaired participants were more likely to develop incident dementia, even after adjusting for potential confounders. We also found that the increased risk of dementia occurred in a graded fashion by VI severity. Our findings highlighted the value of regular vision screening and elimination of VI among middle-aged and elderly individuals.

Our findings have been corroborated by previous studies. The English Longitudinal Study of Ageing,²³ the Ginkgo Evaluation of Memory Study,²⁴ Health and Retirement Study cohort,²⁵ and Women's Health Initiative²⁶ reported that the presence of VI was independently associated with a higher incidence of dementia during follow-up periods of 7 to 18 years. Interestingly, the Three-City study suggested that poor vision was an indicator of dementia risk at short (the first 2 years after inclusion) and middle-term (from 2 to 4 years after inclusion), but not at long-term (beyond 4 years after inclusion).²⁷ Data from the Korean National Health Insurance Service²⁸ and the cohort of the Hong Kong Elderly Health Centres²⁹ showed that individuals with severe VI were at the greatest risk of incident dementia, thereby further supporting the association between VI and incident dementia. Nevertheless, other studies presented data refuting these conclusions: a case-control study in Germany, Michalowsky et al. found that VI was not significantly associated with the risk of dementia.³⁰ Likewise, the association between poor visual acuity and the risk of developing dementia did not reach significance in the Health, Aging, and Body Composition Study.³¹ Discrepancy in study design, statistical methodology, demographic characteristics, definition of VI or poor vision, as well as the detection of dementia may partly explain the inconsistency among studies' results.

Several mechanisms have been postulated to explain the association between VI and

the risk of dementia, either in favor of VI being the manifestation of neurodegeneration, greater cognitive load and lower cognitive efficiency, functional disabilities, or shared pathogenesis between VI and dementia. Firstly, it has been suggested that VI may be one of the first manifestations of dementia.³⁵ Secondly, visually impaired individuals may place more stress on neural resources to optimally perform visual tasks, thus increasing the cognitive loads.³⁶ Alternatively, based on the visual deprivation hypothesis, reduced visual inputs and stimulation may compromise the cognitive efficiency.³⁷ It remains possible that the association between VI and dementia may be due to the intermediate factors, such as depression, functional disabilities, social isolation, and decreased engagement in physical activity and cognitively stimulating activities.³⁸⁻⁴⁰ Last but not the least, the shared pathogenesis between the causes of VI and dementia may also explain this association. Beta-amyloid deposits and genetic risk factors for dementia have been implicated in patients with age-related macular degeneration, one of the leading causes of VI.⁴¹

From a health policy perspective, our findings highlight the value of regular vision screening and elimination of VI. Firstly, vision assessment is a widely available and low-cost test. In addition, with the advent of the smartphone-based eye care services,⁴² the accessibility and uptake of visual acuity testing have greatly improved, particularly in the disadvantaged areas. Given the strong evidence of the association between VI and incident dementia, visual acuity testing may be a valuable tool to identify individuals at higher risk of cognitive decline and dementia. Secondly, our findings imply the potential benefits of the vision training and VI elimination in dementia prevention. It has been indicated that vision-related training (e.g., the field of view training) and low vision rehabilitation (e.g., cataract surgery) may improve cognitive function.^{43, 44} Further studies are needed to investigate the benefits of vision training and vision rehabilitation in dementia prevention.

Despite the advantages of large sample size, long-term follow-up, complete adjustment for confounders, the age range in the analytical sample involving cases of

early-onset dementia, and the use of routinely updated health-related records identifying incident dementia in the present analysis, several limitations should be borne in mind. First, we only explored the association between baseline visual acuity and the risk of dementia. Further studies are needed to investigate the effects of visual acuity changes and other components of visual function (e.g., contrast sensitivity) on the risk of dementia. Second, causes of VI were not available in the present analysis, thus preventing us from investigating associations between specific causes of VI and the risks of dementia. Third, due to the observational design, we could not draw the causal inferences. Nevertheless, we observed similar results in the sensitivity analysis, where dementia cases identified within one year after baseline assessment were excluded. Chances of reverse causality might be trivial. Last but not the least, although a broad array of potential confounders were accounted for in the statistical analysis, we could not completely exclude residual confounding.

In summary, we found that visually impaired individuals were more likely to develop incident dementia, with a progressively greater risk among those with worse visual acuity. Our findings highlight the value of regular vision screening and vision rehabilitation. Further research is warranted to confirm our findings and further investigate the effects of vision rehabilitation on dementia prevention.

Funding: The present work was supported by the Fundamental Research Funds of the State Key Laboratory of Ophthalmology, Project of Investigation on Health Status of Employees in Financial Industry in Guangzhou, China (Z012014075), Science and Technology Program of Guangzhou, China (202002020049). Prof. Mingguang He receives support from the University of Melbourne at Research Accelerator Program and the CERA Foundation. The Centre for Eye Research Australia receives Operational Infrastructure Support from the Victorian State Government. The sponsor or funding organization had no role in the design or conduct of this research.

Competing interests: The author(s) have no proprietary or commercial interest in any

materials discussed in this article.

Abbreviations and Acronyms:

VI = visual impairment; **NHS** = National Health Service; **VA** = visual acuity; **IOP** = intraocular pressure; **OCT** = Optical Coherence Tomography; **PHQ** = Patient Health Questionnaire; **SD** = standard deviations; **HR** = hazard ratios; **CI** = confidence intervals; **IQR** = interquartile range.

Author Contributions:

Study concept and design: Zhu ZT, Shi DL, Liao H, He MG, Yang XH.

Acquisition, analysis, or interpretation: All authors.

Drafting of the manuscript: Zhu ZT, Shi DL, Liao H, Shang XW.

Critical revision of the manuscript for important intellectual content: Jason Ha, Wang W, He MG, Yang XH.

Statistical analysis: Zhu ZT, Shi DL, Shang XW.

Obtained funding: He MG, Yang XH.

Administrative, technical, or material support: Zhu ZT, Shang XW, Wang W, He MG, Yang XH.

Study supervision: He MG, Yang XH.

References:

1. Bourne RRA, Flaxman SR, Braithwaite T, et al. Magnitude, temporal trends, and projections of the global prevalence of blindness and distance and near vision impairment: a systematic review and meta-analysis. *Lancet Glob Health* 2017;5(9):e888-e97.
2. Congdon NG, Friedman DS, Lietman T. Important causes of visual impairment in the world today. *JAMA* 2003;290(15):2057-60.
3. Zhang X, Bullard KM, Cotch MF, et al. Association between depression and functional vision loss in persons 20 years of age or older in the United States, NHANES 2005-2008. *JAMA Ophthalmol* 2013;131(5):573-81.
4. Jones GC, Rovner BW, Crews JE, Danielson ML. Effects of depressive symptoms on health behavior practices among older adults with vision loss. *Rehabil*

Psychol 2009;54(2):164-72.

5. Ivers RQ, Norton R, Cumming RG, et al. Visual impairment and risk of hip fracture. *Am J Epidemiol* 2000;152(7):633-9.
6. Crews JE, Campbell VA. Vision impairment and hearing loss among community-dwelling older Americans: implications for health and functioning. *Am J Public Health* 2004;94(5):823-9.
7. McCarty CA, Nanjan MB, Taylor HR. Vision impairment predicts 5 year mortality. *Br J Ophthalmol* 2001;85(3):322-6.
8. Lee DJ, Gomez-Marin O, Lam BL, Zheng DD. Visual acuity impairment and mortality in US adults. *Arch Ophthalmol* 2002;120(11):1544-50.
9. MJ P. World Alzheimer Report 2015: The Global Impact Of Dementia: An Analysis Of Prevalence, Incidence, Cost And Trends: Alzheimer's Disease Int. 2015.
10. Lin MY, Gutierrez PR, Stone KL, et al. Vision impairment and combined vision and hearing impairment predict cognitive and functional decline in older women. *J Am Geriatr Soc* 2004;52(12):1996-2002.
11. Clemons TE, Rankin MW, McBee WL, Age-Related Eye Disease Study Research G. Cognitive impairment in the Age-Related Eye Disease Study: AREDS report no. 16. *Arch Ophthalmol* 2006;124(4):537-43.
12. Pham TQ, Kifley A, Mitchell P, Wang JJ. Relation of age-related macular degeneration and cognitive impairment in an older population. *Gerontology* 2006;52(6):353-8.
13. Whitson HE, Cousins SW, Burchett BM, et al. The combined effect of visual impairment and cognitive impairment on disability in older people. *J Am Geriatr Soc* 2007;55(6):885-91.
14. Ong SY, Cheung CY, Li X, et al. Visual impairment, age-related eye diseases, and cognitive function: the Singapore Malay Eye study. *Arch Ophthalmol* 2012;130(7):895-900.
15. Woo SJ, Park KH, Ahn J, et al. Cognitive impairment in age-related macular degeneration and geographic atrophy. *Ophthalmology* 2012;119(10):2094-101.
16. Dupuis K, Pichora-Fuller MK, Chasteen AL, et al. Effects of hearing and vision impairments on the Montreal Cognitive Assessment. *Neuropsychol Dev Cogn B Aging Neuropsychol Cogn* 2015;22(4):413-37.
17. Court H, McLean G, Guthrie B, et al. Visual impairment is associated with physical and mental comorbidities in older adults: a cross-sectional study. *BMC Med* 2014;12:181.
18. Yamada Y, Denking MD, Onder G, et al. Dual Sensory Impairment and Cognitive Decline: The Results From the Shelter Study. *J Gerontol A Biol Sci Med Sci* 2016;71(1):117-23.
19. Chen SP, Bhattacharya J, Pershing S. Association of Vision Loss With Cognition in Older Adults. *JAMA Ophthalmol* 2017;135(9):963-70.
20. Ward ME, Gelfand JM, Lui LY, et al. Reduced contrast sensitivity among older women is associated with increased risk of cognitive impairment. *Ann Neurol* 2018;83(4):730-8.
21. Zhu Z, Liao H, Wang W, et al. Association between age-related macular

- degeneration and subjective cognitive complaints. *Br J Ophthalmol* 2019.
22. Lim ZW, Chee ML, Da Soh Z, et al. Association Between Visual Impairment and Decline in Cognitive Function in a Multiethnic Asian Population. *JAMA Netw Open* 2020;3(4):e203560.
 23. Davies-Kershaw HR, Hackett RA, Cadar D, et al. Vision Impairment and Risk of Dementia: Findings from the English Longitudinal Study of Ageing. *J Am Geriatr Soc* 2018;66(9):1823-9.
 24. Hwang PH, Longstreth WT, Jr., Brenowitz WD, et al. Dual sensory impairment in older adults and risk of dementia from the GEM Study. *Alzheimers Dement (Amst)* 2020;12(1):e12054.
 25. Maharani A, Dawes P, Nazroo J, et al. Associations Between Self-Reported Sensory Impairment and Risk of Cognitive Decline and Impairment in the Health and Retirement Study Cohort. *J Gerontol B Psychol Sci Soc Sci* 2020;75(6):1230-42.
 26. Tran EM, Stefanick ML, Henderson VW, et al. Association of Visual Impairment With Risk of Incident Dementia in a Women's Health Initiative Population. *JAMA Ophthalmol* 2020.
 27. Nael V, Peres K, Dartigues JF, et al. Vision loss and 12-year risk of dementia in older adults: the 3C cohort study. *Eur J Epidemiol* 2019;34(2):141-52.
 28. Paik JS, Ha M, Jung YH, et al. Low vision and the risk of dementia: a nationwide population-based cohort study. *Sci Rep* 2020;10(1):9109.
 29. Lee ATC, Richards M, Chan WC, et al. Higher dementia incidence in older adults with poor visual acuity. *J Gerontol A Biol Sci Med Sci* 2020.
 30. Michalowsky B, Hoffmann W, Kostev K. Association Between Hearing and Vision Impairment and Risk of Dementia: Results of a Case-Control Study Based on Secondary Data. *Front Aging Neurosci* 2019;11:363.
 31. Brenowitz WD, Kaup AR, Lin FR, Yaffe K. Multiple Sensory Impairment Is Associated With Increased Risk of Dementia Among Black and White Older Adults. *J Gerontol A Biol Sci Med Sci* 2019;74(6):890-6.
 32. Sudlow C, Gallacher J, Allen N, et al. UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. *PLoS Med* 2015;12(3):e1001779.
 33. Chua SYL, Thomas D, Allen N, et al. Cohort profile: design and methods in the eye and vision consortium of UK Biobank. *BMJ Open* 2019;9(2):e025077.
 34. Lowe B, Kroenke K, Grafe K. Detecting and monitoring depression with a two-item questionnaire (PHQ-2). *J Psychosom Res* 2005;58(2):163-71.
 35. Armstrong RA. Alzheimer's Disease and the Eye. *Journal of Optometry* 2009;2(3):103-11.
 36. Roberts KL, Allen HA. Perception and Cognition in the Ageing Brain: A Brief Review of the Short- and Long-Term Links between Perceptual and Cognitive Decline. *Front Aging Neurosci* 2016;8:39.
 37. Swaab DF, Dubelaar EJ, Hofman MA, et al. Brain aging and Alzheimer's disease; use it or lose it. *Prog Brain Res* 2002;138:343-73.
 38. Fischer ME, Cruickshanks KJ, Klein BE, et al. Multiple sensory impairment and quality of life. *Ophthalmic Epidemiol* 2009;16(6):346-53.

39. Laitinen A, Koskinen S, Harkanen T, et al. A nationwide population-based survey on visual acuity, near vision, and self-reported visual function in the adult population in Finland. *Ophthalmology* 2005;112(12):2227-37.
40. Kiely KM, Anstey KJ, Luszcz MA. Dual sensory loss and depressive symptoms: the importance of hearing, daily functioning, and activity engagement. *Front Hum Neurosci* 2013;7:837.
41. Ohno-Matsui K. Parallel findings in age-related macular degeneration and Alzheimer's disease. *Prog Retin Eye Res* 2011;30(4):217-38.
42. Li Z, Liu R, Xiao O, et al. Progression of Myopic Maculopathy in Highly Myopic Chinese Eyes. *Invest Ophthalmol Vis Sci* 2019;60(4):1096-104.
43. Edwards JD, Fausto BA, Tetlow AM, et al. Systematic review and meta-analyses of useful field of view cognitive training. *Neurosci Biobehav Rev* 2018;84:72-91.
44. Ishii K, Kabata T, Oshika T. The impact of cataract surgery on cognitive impairment and depressive mental status in elderly patients. *Am J Ophthalmol* 2008;146(3):404-9.

Table 1. Baseline Characteristics of the Study Participants Stratified by Visual Impairment Status.

Baseline Characteristics	Total	Non-VI Group	VI Group	OR (95% CI) ^a
N	117,187	113,169	4,018	-
Age, mean (SD), yrs	56.8 (8.11)	56.8 (8.12)	58.8 (7.57)	1.03 (1.03-1.04)
Gender, No. (%)				
Female	63,745 (54.4)	61,533 (54.4)	2,212 (55.0)	1 [Reference]
Male	53,442 (45.6)	51,636 (45.6)	1,806 (45.0)	0.96 (0.90-1.02)
Ethnicity, No. (%)				
White	104,249 (89.0)	100,945 (89.2)	3,304 (82.2)	1 [Reference]
Others	12,938 (11.0)	12,224 (10.8)	714 (17.8)	2.08 (1.91-2.26)
Townsend index, mean (SD)	-0.93 (3.01)	-0.96 (3.00)	-0.12 (3.34)	1.10 (1.09-1.11)
Education level, No. (%)				
College or university degree	40,557 (34.6)	39,504 (34.9)	1,053 (26.2)	1 [Reference]
Others	76,630 (65.4)	73,665 (65.1)	2,965 (73.8)	1.42 (1.32-1.53)
Smoking status, No. (%)				
Never	64,601 (55.5)	62,410 (55.5)	2,191 (55.4)	1 [Reference]
Former/current	51,795 (44.5)	50,035 (44.5)	1,760 (44.6)	0.96 (0.90-1.02)
Physical activity, No. (%)				

Not meeting recommendation	16,867 (17.7)	16,292 (17.7)	575 (18.4)	1 [Reference]
Meeting recommendation	78,268 (82.3)	75,711 (82.3)	2,557 (81.6)	0.93 (0.85-1.02)
Family history of dementia, No.				
(%)				
No	99,228 (84.7)	95,760 (84.6)	3,468 (86.3)	1 [Reference]
Yes	17,959 (15.3)	17,409 (15.4)	550 (13.7)	0.79 (0.72-0.87)
History of depression, No. (%)				
No	105,197 (89.8)	101,692 (89.9)	3,505 (87.2)	1 [Reference]
Yes	11,990 (10.2)	11,477 (10.1)	513 (12.8)	1.38 (1.26-1.52)
History of diabetes, No. (%)				
No	109,346 (93.3)	105,706 (93.4)	3,640 (90.6)	1 [Reference]
Yes	7,841 (6.69)	7,463 (6.59)	378 (9.41)	1.37 (1.23-1.53)
History of hypertension, No.				
(%)				
No	29,972 (25.6)	29,141 (25.8)	831 (20.7)	1 [Reference]
Yes	87,215 (74.4)	84,028 (74.2)	3,187 (79.3)	1.17 (1.08-1.27)
History of hyperlipidemia, No.				
(%)				
No	62,918 (53.7)	60,857 (53.8)	2,061 (51.3)	1 [Reference]
Yes	54,269 (46.3)	52,312 (46.2)	1,957 (48.7)	0.96 (0.90-1.03)

SD = standard deviation; VI = visual impairment; OR = odds ratio; CI = confidence interval.

^a Logistic regression models adjusted for age and gender.

Table 2. Baseline Characteristics Stratified by Dementia Status at Follow-up.

Baseline Characteristics	Non-Dementia Group	Dementia Group	HR (95% CI) ^a
N	116,749	438	-
Age, mean (SD), yrs	56.8 (8.10)	64.0 (5.13)	1.18 (1.16-1.21)
Gender, No. (%)			
Female	63,550 (54.4)	195 (44.5)	1 [Reference]
Male	53,199 (45.6)	243 (55.5)	1.39 (1.15-1.68)
Ethnicity, No. (%)			
White	103,857 (89.0)	392 (89.5)	1 [Reference]
Others	12,892 (11.0)	46 (10.5)	1.49 (1.10-2.02)
Townsend index, mean (SD)	-0.93 (3.01)	-0.76 (3.17)	1.06 (1.03-1.09)
Education level, No. (%)			
College or university degree	40,471 (34.7)	86 (19.6)	1 [Reference]
Others	76,278 (65.3)	352 (80.4)	1.73 (1.36-2.19)
Smoking status, No. (%)			
Never	64,394 (55.5)	207 (47.7)	1 [Reference]
Former/current	51,568 (44.5)	227 (52.3)	1.12 (0.93-1.36)
Physical activity, No. (%)			
Not meeting recommendation	16,788 (17.7)	79 (24.7)	1 [Reference]
Meeting recommendation	78,027 (82.3)	241 (75.3)	0.58 (0.45-0.75)
Family history of dementia, No. (%)			
No	98,897 (84.7)	331 (75.6)	1 [Reference]
Yes	17,852 (15.3)	107 (24.4)	1.45 (1.16-1.80)
History of depression, No. (%)			
No	104,841 (89.8)	356 (81.3)	1 [Reference]
Yes	11,908 (10.2)	82 (18.7)	2.75 (2.16-3.50)
History of diabetes, No. (%)			

No	108,988 (93.4)	358 (81.7)	1 [Reference]
Yes	7,761 (6.65)	80 (18.3)	2.30 (1.80-2.94)
History of hypertension, No. (%)			
No	29,909 (25.6)	63 (14.4)	1 [Reference]
Yes	86,840 (74.4)	375 (85.6)	1.14 (0.87-1.50)
History of hyperlipidemia, No. (%)			
No	62,738 (53.7)	180 (41.1)	1 [Reference]
Yes	54,011 (46.3)	258 (58.9)	1.08 (0.89-1.31)

SD = standard deviation; HR = hazard ratio; CI = confidence interval.

^a Cox proportional hazards regression models adjusted for age and gender.

Table 3. Cox Proportional Hazards Models for Incident Dementia by Visual Impairment Status.

VI Status	Age- and Gender-adjusted Model		Multivariable Model ^a	
	HR (95% CI)	P value	HR (95% CI)	P value
VI status				
None	1 [Reference]	-	1 [Reference]	-
VI	2.01 (1.43-2.85)	<0.001	1.78 (1.18-2.68)	0.006
VI severity				
None	1 [Reference]	-	1 [Reference]	-
Mild VI	2.00 (1.38-2.91)	<0.001	1.66 (1.05-2.62)	0.002
Moderate VI	0.91 (0.13-6.48)		1.12 (0.16-8.01)	
Severe VI	3.05 (1.14-8.18)		3.53 (1.31-9.49)	

VI = visual impairment; HR = hazard ratio; CI = confidence interval.

^a Cox proportional hazards regression models adjusted for age, gender, race/ethnicity, education level, Townsend index, physical activity level, family history of dementia, history of diabetes mellitus, hypertension, and depression.