Changes in the Health of Adolescent Athletes: A Comparison of Health Measures Collected Before and During the CoVID-19 Pandemic

Timothy A. McGuine PhD, ATC, Kevin Biese MA, ATC, Scott J. Hetzel MS, Labina Petrovska,
Stephanie Kliethermes PhD, Claudia L. Reardon MD David R. Bell PhD, ATC, M. Alison Brooks
MD, MPH, Andrew M. Watson MD, MS

Affiliations: This work was performed at the University of Wisconsin-Madison

Corresponding author: Tim McGuine, University of Wisconsin School of Medicine and Public
Health, 1685 Highland Avenue, Madison, WI 53705-2281, mcguine@ortho.wisc.edu

Conflict of Interest Disclosures (includes financial disclosures): The authors have no
conflicts of interest to disclose.

Funding: No additional funding was secured for this study.

ABSTRACT

Context: In the spring of 2020, schools closed to in-person teaching and sports were cancelled to control the transmission of CoVID-19. The changes that took place to the physical and mental health among young athletes during this time remain unknown, however.

Objective: Identify changes in the health (mental health, physical activity and quality of life) of athletes that occurred during the CoVID-19 pandemic.

Design: Cross sectional study.

Setting: Sample recruited via social media.

Patients or Other Participants: 3243 Wisconsin adolescent athletes (age=16.2±1.2 yrs., female=58% female) completed an online survey in May 2020 (DuringCoVID-19). Health measures for this cohort were compared with previously reported data for Wisconsin adolescent athletes (n=5231, age=15.7±1.2, 47% female) collected in 2016–2018 (PreCoVID-19).

Main Outcome Measure(s): Demographic information included: sex, grade and sports played. Health assessments included the Patient Health Questionnaire-9 Item (PHQ-9) to identify depression symptoms, the Pediatric Functional Activity Brief Scale (PFABS) for physical activity, and the Pediatric Quality of Life Inventory 4.0 (PedsQL) for health related quality of life (HRQoL). Univariable comparisons of these variables between groups were made via t-tests or chi-square tests. Means and 95% confidence intervals (CI) for each group were estimated by survey weighted ANOVA models.

RESULTS: Compared to PreCoVID-19, a larger proportion of the During-CoVID-19 participants reported rates of moderate to severe levels of depression (9.7% vs 32.9%, $p<0.001$). During-CoVID-19 participants reported 50% lower (worse) PFABS scores (mean:12.2 [95%CI: 11.9, 12.5] vs 24.7 [24.5, 24.9] $p<0.001$) and lower (worse) PedsQL total scores compared to the PreCoVID-19 participants (78.4 [78.0, 78.8] vs. 90.9 [90.5, 91.3] $p<0.001$).

CONCLUSIONS: During the CoVID-19 pandemic, adolescent athletes reported increased symptoms of depression, decreased physical activity and decreased quality of life compared to adolescent athletes in previous years.

Key words: Public health, COVID-19, High School, Depression,

Key points:

- 1) Adolescent athletes during CoVID-19 were three times more likely to report moderate to severe symptoms of depression compared to data collected prior to CoVID-19.
- 2) Adolescent athletes during CoVID-19 reported significantly lower physical activity and quality of life scores compared to high school athletes prior to the CoVID-19 pandemic
- 3) Post CoVID-19 policies should be implemented to improve the health of adolescent athletes in the US.

INTRODUCTION

An estimated 8.4 million US high school students participate in interscholastic athletics.¹ Adolescent sport participation is recognized to have profound positive influences on the health and well-being of adolescent students as evidenced by higher academic achievement, greater levels of healthy physical activity and decreased levels of anxiety and depression compared to students who do not participate in athletics.²⁻⁵ Additionally, high school sport participation is one of the most important factors for life-long physical activity and health.⁶⁻¹²

During the winter and spring of 2020, CoVID-19, the disease caused by the novel SARS-CoV-2 coronavirus, reached pandemic proportions throughout the world. Across the US, schools were closed and athletic seasons were cancelled to slow the spread of the disease. Experts have suggested that efforts to limit the spread of CoVID-19 may have profound societal, economic, and psychosocial consequences for students and need to be further studied.¹³⁻¹⁷

A recent study reported that female athletes, athletes in grade 12, team sport participants and athletes from areas with higher % of the population under age 18 in poverty reported lower levels of physical activity, greater symptoms of anxiety and depression, and lower HRQoL in May of 2020 during the nationwide shutdown.¹⁸ Describing the prevalence of mental health disorders during the CoVID-19-related restrictions is useful, but does not fully describe how specific populations of athletes have been impacted. We are not aware of research to date that has specifically documented mental and physical health *changes* in adolescent athletes during the pandemic. Quantifying these changes will allow health care providers to implement strategies to improve the health of adolescent athletes moving forward.²⁵

An estimated 93,000 (33%) of the 284,000 Wisconsin high school students compete in interscholastic sports each year.¹ At the start of the 2019/2020 school year, 478 Wisconsin schools offered interscholastic athletics and fielded teams competing in fifteen to twenty sports.¹⁹

These student athletes typically competed on one to three athletic teams throughout the school year (fall, winter and spring sport seasons), but many also continued to train with their teammates when their sport is not in season.^{20,21} Over half of these student athletes also competed on club sport teams not affiliated with their school during the school year.^{22,23}

In March 2020, Wisconsin government and health officials mandated a two-week state-wide school closure to slow the spread of CoVID-19. The closure order was eventually extended twice to run through the end of the school year.²⁴ While high school students' academic work moved online, all in-person teaching and interscholastic athletics for students in grades 9–12 were cancelled. Additionally, adolescent club sport teams were restricted from participating during this time. While necessary to slow the community spread of the virus, the CoVID-19 mitigation strategies may nonetheless have negative effects on the physical and mental health of adolescents.²⁶

Quantifying the changes to the health of adolescent athletes that occurred during the CoVID-19 pandemic is critical to understand how these issues impacted athletes in order to to improve management in the future.²⁷⁻²⁹ This information will allow sports medicine providers, school administrators and health care policy experts to implement strategies to improve the short-term and long-term mental and physical health of adolescent athletes in the months and years to come as we transition from the CoVID-19 pandemic.

The purpose of this study was to measure changes in the mental and physical health of Wisconsin adolescent athletes manifested during the CoVID-19 related school closures and sport cancellations. To measure these changes, we compared athlete self-report data on depression, physical activity and HRQoL that we collected during-CoVID-19 (DuringCoVID-19) with Wisconsin data from a series of previous investigations carried out by the study team during normal high school sport operations in the years 2015-2018 (PreCoVID-19).³⁰⁻³⁴ We hypothesize that adolescent student athletes will report significantly worse mental health, as

well as lower quality of life and physical activity scores during the CoVID-19 pandemic shutdown, than has previously been reported prior to CoVID-19.

METHODS

This study was approved by the University of Wisconsin-Madison Health Sciences Institutional Review Board in April of 2020. Wisconsin adolescent athletes (male and female, grade: 9–12, age:13-19) were recruited via social media (Facebook, Twitter) to participate in the study by completing an anonymous online survey in May 2020. To ensure widespread distribution and recruitment for the study, the study team provided the social media links to sports medicine provider colleagues across Wisconsin, the Wisconsin Interscholastic Athletic Association (WIAA) and the Wisconsin Athletic trainers' Association (WATA), which provided the links to high school athletes throughout Wisconsin.

The survey was composed of 69 items and included a section to describe the demographics for the participant, which was followed by 3 segments that included validated instruments used to measure depression symptoms, physical activity and HRQoL in adolescents. Demographic information included items for the participants' age, grade, school size, school funding (private or public) and zip code as well as a list of all the high school and club (not affiliated with their school) sports in which they competed during the previous 12 months. The remainder of the survey consisted of an assessment of mental health, physical activity level and HRQoL.

Mental health

The Patient Health Questionnaire-9 Item (PHQ-9) survey was used to evaluate depression symptoms. This 9-item questionnaire asks participants to rate the frequency of depression symptoms experienced in the past two weeks.³⁵ The PHQ-9 scores range from 0-27 with a higher score indicating a greater level of depression symptoms. The PHQ-9 has demonstrated high sensitivity and specificity for depression screening in adolescent patients aged 13 to 17 years. In addition to the total score, PHQ-9 categorical

scores of 0–4, 5–9, 10–14, 15–19 and ≥ 20 correspond to minimal or none, mild, moderate, moderately severe and severe depression symptoms, respectively.^{36,37}

Physical Activity

Physical activity level was assessed with the Hospital for Special Surgery Pediatric Functional Activity Brief Scale (PFABS).^{38,39} This validated 8-item instrument was designed to measure the physical activity of children between 10 and 18 years old for the past month. Scores range from 0 to 30 with a higher score indicating greater physical activity.^{38,39}

Health Related Quality of Life

HRQoL was measured with Pediatric Quality of Life Inventory 4.0 (PedsQL). The 23-item PedsQL questionnaire assesses HRQoL for the previous 7 days. The PedsQL has been validated for use in children ages 2 to 18.^{40,41} Physical and psychosocial subscale summaries, as well as the total scores, range from 0 to 100 with a higher score indicating greater HRQoL. The PedsQL survey has been used to measure HRQoL in samples of healthy and injured adolescent athletes.⁴²

Statistical analyses

Statistical analyses were performed for participants who provided a valid, complete survey. Participants were excluded if they did not complete the entire survey, were not in grades 9 -12, or indicated they did not participate in club or interscholastic sports within the previous 12 months.

Demographic variables and individual sport participation were summarized (mean (SD) or N (%)) overall and by study group (PreCoVID-19 athlete cohort and During-CoVID-19 athlete cohort) stratified by sex. Univariable comparisons of these variables between groups were made via t-tests or chi-square tests. Means and 95% confidence intervals (CI) for each group were estimated by survey weighted ANOVA models separately for depression, physical activity and quality of life measures. A weighted

ordinal logistic regression (OLR) model was used to estimate the percentages of level of depression (PHQ-9) for each group. A group by sex interaction was modeled for all sex-specific estimates.

Sample survey weights were derived based on sport and sex using the 2018-2019 Wisconsin high school sport participation statistics compiled annually by the National Federation of High Schools (NFHS).¹ Survey weights for multiple sport athletes were based on the most common sport played. Inverse propensity score weights (IPSW) were also calculated to account for design-based differences in the two groups.⁴³ Propensity score weights were assigned using logistic regression which was weighted based on the sample survey weights. Group assignment was the dependent variable, and age, sex, and high school individual sport status were predictor variables.

The PHQ9, PFABS and PedsQL scores were compared with historical data of Wisconsin adolescent athletes collected during normal school operations in the years 2015–2018.²⁵⁻²⁹ Final models comparing the During-CoVID-19 group to the PreCoVID-19 group included weights as the multiplication of the sampling weights and IPSWs.⁴⁴

RESULTS

A total of 3,243 Wisconsin adolescent athletes (age = 16.2±1.2 yrs., female = 58%) completed the survey in May 2020 (During-CoVID-19) with responses obtained from participants residing in 71 (99%) of the 72 counties in Wisconsin. Due to the convenience sampling design, information regarding the response rate was unavailable. A total of 2,730 (84%) participants indicated they attended a publicly funded school with the median [IQR] school enrollment = 802 [401, 1248]. Nine hundred seventy (29.9%) competed in a single sport while the remaining 2,273 (70.1%) competed in two or more sports for their school in the prior 12 months. During-CoVID-19 participants were most likely to compete in track (32.8%), basketball (29.1%) and volleyball (21.2%). In addition to competing for their high school teams, 1,962 (60.5%) competed as an individual or as part of a club team outside their school setting.

The PreCoVID-19 cohort consisted of 5,231 athletes (age = 15.7 ± 1.2 , female = 65%) that were enrolled in cohort and longitudinal studies in 2015-2018.³⁰⁻³⁴ A summary of the number of athletes by sex, age and grade for both the During-CoVID-19 and PreCoVID-19 cohorts are found in Table 1. The PreCoVID-19 participants resided in 56 (78%) of Wisconsin counties. A total of 4,399 (84%) of these participants attended a publicly funded school with the median [IQR] school enrollment = 654 [299, 1017]. The PreCoVID-19 participants were most likely to compete in volleyball (55.7%), basketball (43.2%) and football (23.3%). A summary of the number of athletes and percentages of the sports played for both the During-CoVID-19 and PreCoVID-19 cohorts are found in Supplementary Table 1.

Mental health

The number of females that reported a mild level of depression in the During-CoVID-19 athletes was 43% higher compared to the PreCoVID-19 females (During-CoVID-19 = 35%, PreCoVID-19 = 24%, $p < 0.001$). Females in the During-CoVID-19 group reported a three-fold increase in moderate, moderately severe or severe levels of depression compared to the PreCoVID-19 athletes (During-CoVID-19 = 37%, PreCoVID-19 = 11%, $p < 0.001$). The prevalence of depression levels for females in both the PreCoVID-19 and During-CoVID-19 groups are found in Figure 1. The number of males in the During-CoVID-19 group that reported a mild level of depression was 130% higher than compared to PreCoVID-19 males ($p < 0.001$). The proportion of males in the During-CoVID-19 group that reported moderate, moderately severe or severe levels of depression was 4.5 times higher than males in the PreCoVID-19 group (During-CoVID-19 = 27%, PreCoVID-19 = 6%, $p < 0.001$). The prevalence of depression levels for males in the During-CoVID-19 and PreCoVID-19 group is found in Figure 2.

Overall, the PHQ-9 scores for athletes During-CoVID-19 were 2.5 times higher (worse) than the athletes in the PreCoVID-19 group. The During-CoVID-19 females reported a (mean [95%CI]) PHQ-9 score that was over twice the PHQ-9 score for females in the PreCoVID-19

group (8.2 [7.9, 8.5] vs. 3.6 [3.4, 3.8]). Similarly, the During-CoVID-19 males had PHQ-9 scores that were three-times higher than the males in the PreCoVID-19 group (7.8 [7.5, 8.1] vs. 2.6 [2.2, 2.9]. The PHQ-9 scores for both groups are found in Table 1.

Physical Activity

The total PFABS scores (mean [95%CI]) were lower (less physical activity) for athletes in the During-CoVID-19 group compared to the PreCoVID-19 group (12.2 [11.9, 12.5] vs. 27.7 [24.5, 24.9], $p < 0.001$). The During-CoVID-19 females reported PFABS scores that were more than 50% lower (worse) than the PreCoVID-19 females (11.4 [11.1, 11.8] vs. 24.5 [24.2, 24.7], $p < 0.001$). Similarly, males in the During-CoVID-19 group reported PFABS scores approximately 50% lower compared to athletes in the PreCoVID-19 group (13.9 [13.4, 14.4] vs. 25.7 [25.3, 26.1], $p < 0.001$). The PFABS scores for both groups are found in Table 1.

Health Related Quality of Life

Higher (better) PedsQL physical summary, psychosocial summary, and PedsQL total scores were reported for athletes in the PreCoVID-19 group compared to the athletes in the During-CoVID-19 group ($p < 0.001$). The PreCoVID-19 physical health summary scores (mean [95%CI]) for females were 11% higher than those reported in the During-CoVID-19 group (89.6 [89.1, 90.0] vs. 80.9 [80.5, 81.4], $p < 0.001$). Similarly, the PreCoVID-19 physical health summary scores for females were 20% higher than the scores reported in the During-CoVID-19 group (90.1 [89.6, 90.6] vs. 75.0 [74.5, 75.5], $p < 0.001$), while the PreCoVID-19 total PedsQL scores for females were 17% higher than the scores reported in the During-CoVID-19 group (90.0 [89.5, 90.4] vs. 77.1 [76.6, 77.6], $p < 0.001$).

The PreCoVID-19 physical health summary scores (mean [95%CI]) for males were 12% higher than the scores reported by athletes in the During-CoVID-19 group (96.6 [95.9, 97.3] vs. 86.1 [85.4, 86.8], $p < 0.001$). Similarly, the PreCoVID-19 physical health summary scores for males were 14% higher than the scores reported by males in the During-CoVID-19 group (90.9 [90.2, 91.7] vs. 78.6 [77.9, 79.3], $p < 0.001$). The PedsQL total scores reported by the males in

the PreCoVID-19 group were also 14% higher than scores for males in the During-CoVID-19 group (92.9 [92.2, 93.6] vs. 81.2.1 [80.6, 81.9], $p < 0.001$). The PedsQL scores for both groups are found in Table 1.

DISCUSSION

The primary findings of this study are that the mental health, physical activity and HRQoL scores for adolescent athletes were significantly worse during the CoVID-19 pandemic than scores reported by adolescent athletes in prior years. To our knowledge there are no existing data that demonstrates mental health, physical activity and HRQoL *changes* for adolescent athletes immediately following the cancellation of school and athletic seasons during the CoVID-19 pandemic. It has been suggested that the negative psychosocial effects of CoVID-19 related mitigation measures may be manifested in additional health care utilization and spending in the coming months and years.^{25,26} The results of this study can be used to inform stakeholders of the results of quarantine restrictions and how they negatively impact the health of adolescent athletes.

Mental Health

The levels of depression symptoms and total symptom scores reported (as measured by the PHQ-9) in the During-CoVID-19 cohort were significantly higher than our PreCoVID-19 athletes. Additionally, our During-CoVID-19 cohort had higher depression symptoms than previous studies of adolescents that observed severe depression symptoms in approximately 2% of adolescents.⁴⁵ While the prevalence of moderate to severe depressive symptoms was greater among females than males in the During-CoVID-19, both males and females demonstrated dramatically higher rates compared to data collected before the CoVID-19 pandemic.⁴⁵ The depression data reported for participants in the During-CoVID-19 group demonstrate a significant mental health burden among this population of young athletes.

The decrease in mental health is multifaceted and complex and may be related to the decreased socialization, increased family strain, and reduced access to support services.⁴⁶ In Wisconsin, many families self-isolated or severely limited social contact for several weeks to several months in the Spring of 2020. Previous studies have demonstrated that quarantines can negatively impact mental health.^{46,47} Schools play an important role in providing access to mental health services for disadvantaged populations, and health care providers, parents, and policy-makers should be mindful of the mental health strain the current pandemic is placing on adolescents.⁴⁸

Physical Activity

Our study is the first in the United States to demonstrate a significant decrease in physical activity levels in adolescents during the CoVID-19 pandemic. While male athletes reported slightly higher levels of physical activity than females, the decrease in both males and females during the CoVID-19 restrictions was similar, with current levels of physical activity approximately 50% lower than levels reported in a similar population prior to CoVID-19 reported by Fabricant.³⁸ Physical activity is known to have a beneficial effect on a wide range of health outcomes in adolescents, including sleep, academic success, well-being, and mental health.⁴⁹⁻⁵¹ Therefore, it is possible that the identified decrease in mental health in our participants may be at least partly due to the removal of the positive effects that physical activity has for adolescents.

Childhood obesity was a health care crisis prior to the COVID-19 pandemic that is increasing in the US and is projected to become worse due to the pandemic.^{28,29,52} A decrease in physical activity and increase in stress may increase the risk of severe COVID-19 cases in obese children.²⁹ Decreased physical activity in adolescents may also have longer-term negative effects and implications in terms of increased risk for obesity and cardiometabolic disease if these levels remain low for prolonged periods.⁵³ Chronically low levels of physical activity may also compound the mental health consequences of the current crisis.^{49,50} Further,

there is evidence that sport participation during high school is related to improved health and well-being throughout adulthood.⁷⁻¹¹ As time spent being physical active during the school day has decreased, organized sports have been proposed as a means to stabilize long-term physical activity in adolescents and improve adult physical activity.⁴⁹ Though returning adolescents to organized sports is a complex issue and requires careful consideration, stakeholders should consider the promotion of physical activity a top priority during the CoVID-19 pandemic.

Health Related Quality of Life

Several studies have shown that individuals with increased physical activity and or interscholastic sport participation report higher HRQoL scores compared to inactive adolescents and high school non-athletes.^{42,54} Therefore, it is not surprising that HRQoL scores decreased in the During-CoVID-19 cohort compared to the PreCoVID-19 athletes. Interestingly, the overall PedsQL scores observed for the During-CoVID-19 athletes are similar to scores recorded in the general adolescent population (during the pandemic?) but higher than scores reported in adolescents with chronic disease conditions (during the pandemic?).^{40,55}

In our study, females reported higher levels of depression symptoms as well as lower physical activity and HRQoL scores than males. This is consistent with previous research on depression, physical activity and HRQoL in adolescents.^{36-40,50,52,56} Therefore it is unlikely that the sex differences in depression symptoms and HRQoL scores we noted are attributable to the CoVID-19 pandemic directly. Nonetheless, we found dramatic reductions in HRQoL for both males and females during the COVID-19 shutdown, and researchers should continue to monitor HRQoL in adolescents as separation from peers and sports continue during restrictions initiated to limit the spread of CoVID-19.

Limitations

This study has several limitations. First, the data provided were self-reported from online surveys and not the result of a clinical examination conducted by a health care provider.

Nonetheless, our findings include a large sample of athletes and align with reports from experts who have stated that CoVID-19 will impact the health of youth populations.^{13,16,17,25,26} Second, we acknowledge that there may be a response bias for participants. We cannot know for certain if the sample is representative of all Wisconsin adolescent athletes or biased towards athletes who were more likely to respond if they experienced the most profound impacts on their health. Third, due to the survey delivery method, our sample may be biased toward athletes from higher socioeconomic families with easy access to internet services and social media platforms. We could not eliminate this bias, but we did collect results from 99% of Wisconsin counties, which span a wide range of socioeconomic levels. Further, we utilized medical colleagues as well as the Wisconsin Interscholastic Athletic Association and the Wisconsin Athletic Trainers' Association to publicize the study and urge all adolescent athletes to participate. In addition, we used recent high school sport participation statistics from Wisconsin schools to calculate sample weights to mitigate the potential effects of selection bias. This increased the representativeness of our results to the entire adolescent athlete population in Wisconsin. Additionally, it is possible that other variables, such as participation in other co-curricular activities and missing school related events, were not accounted for in our study and could confound the results. Finally, our study did not include a control group consisting of non-athletes. Our existing data sets only include information on athletes, therefore we could only address potential changes over time (Pre-COVID to During-COVID) in athletic populations. We acknowledge it is very likely that the impacts observed in this sample of athletes would be observed across a wide spectrum of youth due the cancellations of most extracurricular activities.

CONCLUSION

The physical activity and mental health of adolescent athletes worsened significantly during the CoVID-19 pandemic compared to PreCoVID-19 values. Our findings suggest that adolescent

athletes have experienced dramatic increases in symptoms of depression along with subsequent decreases in physical activity and HRQoL during this time. Sports medicine and health policy experts should consider the potential negative mental and physical health effects that have become prevalent among adolescent athletes during the CoVID-19 pandemic, and stakeholders should consider systematic steps that can be implemented to reduce these negative changes in the months and years to come.

Word Count = 3433.

List of Tables and Figures

Table 1. Comparison of Depression, Physical Activity and Quality of Life Scores for Wisconsin Adolescent Athlete Historical Data with Data Collected in During-CoVID-19.

Figure 1. Prevalence of Depression Symptoms for Female Adolescent Athletes Prior to the COVID-19 Pandemic (PreCoVID-19) and in During the CoVID-19 Pandemic (DuringCoVID-19).

Figure 2. Prevalence of Depression Symptoms for Female Adolescent Athletes Prior to the COVID-19 Pandemic (PreCoVID-19) and in During the CoVID-19 Pandemic (DuringCoVID-19).

Supplementary Table 1. Distribution of Adolescent Sport Participants for the Historical and During-CoVID-19 Samples

REFERENCES

1. Participation in school athletics. <https://www.childtrends.org/indicators/participation-in-school-athletics>. Accessed July 12,2020.
2. Bailey R. Physical education and sport in schools: a review of benefits and outcomes. *Journal of School Health*. 2006;76(8):397-401. doi:10.1111/j.1746-1561.
3. Eime RM, Young JA, Harvey JT, Charity MJ, Payne WR. A systematic review of the psychological and social benefits of participation in sport for children and adolescents: informing development of a conceptual model of health through sport. *Int J Behav Nutr and Phys Acti*, 2013;10(1):98-104.
4. Gould D, Flett R, Lauer L. The relationship between psychosocial developmental and the sports climate experienced by underserved youth. *Psychology of Sport and Exercise*. 2012;13(1):80.
5. Harrison PA, Narayan G. Differences in behavior, psychological factors, and environmental factors associated with participation in school sports and other activities in adolescence. *Journal of School Health* 2003;73(3):113-20.
6. Marques A, Ekelund U, Sardinha LB. Associations between organized sports participation and objectively measured physical activity, sedentary time and weight status in youth. *J Sci Med in Sport*. 2016;19(2):154-7.
7. Ashdown-Franks G, Sabiston CM, Solomon-Krakus S, O'Loughlin JL. Sport participation in high school and anxiety symptoms in young adulthood. *Mental Health and Phys Activity*. 2017;12:19-24.
8. Dohle S, Wansink B. Fit in 50 years: participation in high school sports best predicts one's physical activity after Age 70. *BMC Public Health* 2013. <https://doi.org/10.1186/1471-2458-13-110>.

9. Easterlin MC, et. al. Association of team sports participation with long-term mental health outcomes among individuals exposed to adverse childhood experiences. *JAMA Pediatr.* 2019; 173 (7) 673- 681.
10. Khan KM, Thompson AM, Blair SN, Sallis JF, Powell KE, Bull FC, Bauman AE. Sport and exercise as contributors to the health of nations. *The Lancet.* 2012;380:59-64.
11. Kniffin KM, Wansink B, Shimizu M. Sports at work: Anticipated and persistent correlates of participation in high school athletics. *Journal of Leadership & Organizational Studies.* 2015;22(2):217-30.
12. Troutman KP, Dufur MJ. From High School Jocks to College Grads: Assessing the Long-Term Effects of High School Sport Participation on Females' Educational Attainment. *Youth & Society.* 2007;38(4):443-462
13. Goldstein E, Wen H, Miller BF. Coronavirus Disease 2019 (COVID-19) and Mental Health for Children and Adolescents *JAMAPeds* Published Online: April 14, 2020.doi:10.1001/jamapediatrics.2020.1456
14. Christakis. School Reopening—The Pandemic Issue That Is Not Getting Its Due Published. *JAMA Pediatr.* Published online May 13, 2020. doi:10.1001/ 2020.2068.
15. Lee. Mental health effects of school closures during COVID-19 . *Lancet Child Adolesc Health* 2020 Published Online April 14, 2020 [https://doi.org/10.1016/S2352-4642\(20\)30109-7](https://doi.org/10.1016/S2352-4642(20)30109-7).
16. Czeisler ME, Lane R, Petrosky E. et al. Mental Health, Substance Use, and Suicidal Ideation During the COVID-19 Pandemic — United States, June 24–30, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:1049-1057.
17. Gassman-Pines A, Ananat EO, Fitz-Henley J. COVID-19 and Parent-Child Psychological Well-being. *Pediatrics.* 2020;146(4):e2020007294.
18. McGuine TA, Biese KM, Petrovska L, Hetzel SJ, Reardon C, Kliethermes S, Bell DR, Brooks A, Watson AM. The health of US adolescent athletes during CoVID-19 related school

- closures and sport cancellations. *Journal of Athletic Training*. 2020 Nov 5;
<https://doi.org/10.4085/1062-6050-0478.20>.
19. Wisconsin Interscholastic Athletic Association. <https://www.wiaawi.org/>. Accessed May 12, 2020.
20. McGuine TA, Post EG, Hetzel S, Pfaffar A, Brooks MA, Bell D. A Prospective Study on the Impact of Sport Specialization on Lower Extremity Injury Rates in High School Athletes. *Am J Sports Med*. 2017;45(12):2706-12.
21. McGuine TA, Pfaffar AY, Kliethermes SA, Schwarz A, Hetzel S, Hammer E & Broglio SP. The impact of sport related concussion injuries on concussion symptoms and health related quality of life in male and female adolescent athletes: A prospective study. *Am J Sports Med*. 2019;47(14):3514–3520.
22. Bell DR, Post EG, Trigsted SM, Schaefer DA, McGuine TA, Watson AM, Brooks MA. Sport specialization characteristics between rural and suburban high school athletes. *Ortho J Sports Med*. 2018;6(1):2325967117751386.
23. Post EG, Bell DR, Trigsted SM, Brooks AM, & McGuine TA. The Association of Volume, Club Sports, and Specialization with Injury History in Youth Athletes. *Sports Health*. 2017;9(6): 518-523 doi: 10.1177/1941738117714160. PMID: 28628419
24. School Closures in Wisconsin https://evers.wi.gov/Documents/COVID-19/English_K-12FAQ.pdf. Accessed May 12 2020.
25. Rajapakse N, Dixit D. Human and novel coronavirus infections in children: a review. *Paediatrics and international child health*. 2020 Jun 26:1-20.
26. Fegert JM, Vitiello B, Plener PL, Clemens V. Challenges and burden of the Coronavirus 2019 (COVID-19) pandemic for child and adolescent mental health: a narrative review to highlight clinical and research needs in the acute phase and the long return to normality. *Child and Adolescent Psychiatry and Mental Health*. 2020 Dec;14:1-1.

27. World Health Organization. Considerations for school-related public health measures in the context of COVID-19: annex to considerations in adjusting public health and social measures in the context of COVID-19, 14 September 2020. *World Health Organization*; 2020.
28. An R. Projecting the impact of COVID-19 pandemic on childhood obesity in the US: A microsimulation model. *Journal of Sport and Health Science*. 2020 May 2.
29. Browne NT, Snethen JA, Greenberg CS, Frenn M, Kilanowski JF, Gance-Cleveland B, Burke PJ, Lewandowski L. When pandemics collide: The impact of COVID-19 on childhood obesity. *Journal of Pediatric Nursing*. 2020 Nov 11.
30. McGuine TA, Pfaller A, Kliethermes S, Schwarz A, Hetzel S, Hammer E, Broglio S. The effect of sport-related concussion injuries on concussion symptoms and health-related quality of life in male and female adolescent athletes: a prospective study. *Am J Sports Med*. 2019 Dec;47(14):3514-20.
31. Donovan L, Hetzel S, Laufenberg CR & McGuine TA. Prevalence and Impact of Chronic Ankle Instability in Adolescent Athletes. *Orthop J Sports Med*. Feb. 2020. 17;8(2):2325967119900962.
32. McGuine TA, Pfaller A, Hetzel S, Broglio SP, Hammer E. A prospective study of concussions and health outcomes in high school football players. *Journal of Athletic Training*. 2020 Oct;55(10):1013-9.
33. Hammer E, Hetzel S, Pfaller A, McGuine T. Longitudinal assessment of depressive symptoms after sport-related concussion in a cohort of high school athletes. *Sports Health*. 2020 Aug 28:1941738120938010.
34. McGuine TA, Biese KM, Lang P, Kliethermes S, Bell DR, Brooks A, Watson AM The Incidence and Risk Factors for Injuries in Girls Volleyball: A Prospective Study of 2072 Players. *Journal of Athletic Training* (in press).

35. Andrews JH, Cho E, Tugendrajch SK, et. al., Evidence-Based Assessment Tools for Common Mental Health Problems: A Practical Guide for School Settings. *Children & Schools*. 2020;42(1):41–52.
36. Kroenke K, Wu J, Yu Z, Bair MJ, Kean J, Stump T, Monahan PO. Patient Health Questionnaire Anxiety and Depression Scale: Initial Validation in Three Clinical Trials. *Psychosom Med*. 2016;78(6):716-27.
37. Richardson LP, McCauley E, Grossman DC, et al. Evaluation of the Patient Health Questionnaire-9 Item for Detecting Major Depression Among Adolescents. *Pediatrics*. 2010;126:1117–1123.
38. Fabricant PD, Robles A, Downey-Zayas T, et al. Development and validation of a pediatric sports activity rating scale: the Hospital for Special Surgery Pediatric Functional Activity Brief Scale (HSS Pedi-FABS). *Am J Sports Med*. 2013;41(10):2421-2429.
39. Fabricant PD, Robles A, McLaren SH, Marx RG, Widmann RF, Green DW. Hospital for Special Surgery Pediatric Functional Activity Brief Scale predicts physical fitness testing performance. *Clin Orthop Relat Res*. 2014;472(5):1610-1616.
40. Varni JW, Seid M, Kurtin PS. PedsQL™ 4.0: Reliability and validity of the Pediatric Quality of Life Inventory™ Version 4.0 Generic Core Scales in healthy and patient populations. *Med Care*. 2001;39(8):800-812.
41. Varni JW, Burwinkle TM, Seid M, Skarr D. The PedsQL™* 4.0 as a pediatric population health measure: feasibility, reliability, and validity. *Ambulatory Pediatrics*. 2003;3(6):329-341.
42. Lam KC, Valier AR, Bay RC, McLeod TC. A unique patient population? Health-related quality of life in adolescent athletes versus general, healthy adolescent individuals. *J Ath Train*. 2013;48(2):233-41.

43. Tan WY. Sampling distributions and robustness of t, F and variance-ratio in two samples and ANOVA models with respect to departure from normality. *Communications in Statistics. Theory and Methods*. 1982; 11:2485-2511
44. Ridgeway G, Kovalchik SA, Griffin BA, & Kabeto MU. Propensity score analysis with survey weighted data. *Journal of Causal Inference*. 2015; 3(2): 237-249.
45. Cyranowski JM, Frank E, Young E, Shear MK. Adolescent Onset of the Gender Difference in Lifetime Rates of Major Depression: A Theoretical Model. *Arch Gen Psychiatry*. 2000;57(1):21–27. doi:10.1001/archpsyc.57.1.2
46. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, Rubin GJ. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *The Lancet*. 2020 Feb 26.
47. Sprang G, Silman M. Posttraumatic stress disorder in parents and youth after health-related disasters. *Disaster Medicine and Public Health Preparedness*. 2013 Feb;7(1):105-10.
48. Ali M. Utilization of Mental Health Services in Educational Setting by Adolescents in the United States *Journal of School Health* March 2019. <https://doi.org/10.1111/josh.12753>.
49. Valkenborghs SR, Noetel M, Hillman CH, Nilsson M, Smith JJ, Ortega FB, Lubans DR. The impact of physical activity on brain structure and function in youth: A systematic review. *Pediatrics*. 2019 Oct 1;144(4):e20184032.
50. Biddle SJ, Ciacconi S, Thomas G, Vergeer I. Physical activity and mental health in children and adolescents: An updated review of reviews and an analysis of causality. *Psychology of Sport and Exercise*. 2019 May 1;42:146-55.
51. Vella SA, Cliff DP, Magee CA, Okely AD. Associations between sports participation and psychological difficulties during childhood: a two-year follow up. *J Sci Med Sport*. 2015; 1;18(3):304-9.
52. Skinner AC, Ravanbakht SN, Skelton JA, Perrin EM, Armstrong SC. Prevalence of obesity and severe obesity in US children, 1999–2016. *Pediatrics*. 2018 Mar 1;141(3).

53. Telama R, Yang X, Viikari J, Välimäki I, Wanne O, Raitakari O. Physical activity from childhood to adulthood: a 21-year tracking study. *Am J Prev Med.* 2005;28(3):267-273.
54. Houston MN, Hoch MC, Hoch JM. Health-related quality of life in athletes: a systematic review with meta-analysis. *J Ath Train.* 2016;51(6):442-53.
55. Varni J., Limbers CA. & Burwinkle TM. Impaired health-related quality of life in children and adolescents with chronic conditions: a comparative analysis of 10 disease clusters and 33 disease categories/severities utilizing the PedsQL™ 4.0 Generic Core Scales. *Health Qual Life Outcomes* **5**, 43 (2007). <https://doi.org/10.1186/1477-7525-5-43>.
56. Dahab K, Potter MN, Provance A, Albright J, Howell DR. Sport Specialization, Club Sport Participation, Quality of Life, and Injury History Among High School Athletes. *J Ath Training.* 2019;54(10):1061-6.
57. Åkesdotter C, Kenttä G, Eloranta S, Franck J. The prevalence of mental health problems in elite athletes. *J Sci Med in Sport.* 2020 Apr 1;23(4):329-35.

List of Tables and Figures

Table 1. Comparison of Depression, Physical Activity and Quality of Life Scores for Wisconsin Adolescent Athlete Historical Data with Data Collected in During-CoVID-19.

Figure 1. Prevalence of Depression Symptoms for Female Adolescent Athletes Prior to the COVID-19 Pandemic (PreCoVID-19) and in During the CoVID-19 Pandemic (DuringCoVID-19).

Figure 2. Prevalence of Depression Symptoms for Female Adolescent Athletes Prior to the COVID-19 Pandemic (PreCoVID-19) and in During the CoVID-19 Pandemic (DuringCoVID-19).

Supplementary Table 1. Distribution of Adolescent Sport Participants for the Historical and During-CoVID-19 Samples

Table 1. Comparison of Depression, Physical Activity and HRQoL Scores for Adolescent Athletes Prior to the CoVID-19 Pandemic (PreCoVID-19) and During the CoVID-19 Pandemic (During-CoVID-19).

Variable	PreCoVID-19 All Participants (n = 5231)	During-CoVID-19 All Participants N = 3243	P value	PreCoVID-19 Females (n = 3402)	During- CoVID-19 Females (n = 1877)	P value	PreCoVID-19 Males (n = 1829)	During-CoVID-19 Males (n = 1366)	P value
Sex – Male [^]	1829 (35.0%)	1366 (42.1%)	< 0.001	--	--		--	--	
Age [^]	15.7 (1.1)	16.2 (1.2)	< 0.001	15.6 (1.1)	16.1 (1.2)	< 0.001	15.8 (1.2)	16.2 (1.2)	< 0.001
Grade [^]			< 0.001			< 0.001			0.003
9	1763 (33.7%)	862 (26.6%)		1177 (34.6%)	499 (26.6%)		586 (32.0%)	363 (26.6%)	
10	1388 (26.5%)	845 (26.1%)		937 (27.5%)	509 (27.1%)		451 (24.7%)	336 (24.6%)	
11	1153 (22.0%)	848 (26.1%)		720 (21.2%)	470 (25.0%)		433 (23.7%)	378 (27.7%)	
12	927 (17.7%)	688 (21.2%)		568 (16.7%)	399 (21.3%)		359 (19.6%)	289 (21.2%)	
PHQ9 Total Score **	3.3 (3.1, 3.5)	8.0 (7.8, 8.2)	< 0.001	3.6 (3.4, 3.8)	8.2 (7.9, 8.5)	< 0.001	2.6 (2.2, 2.9)	7.8 (7.5, 8.1)	< 0.001
PFABS Total Score *	24.7 (24.5, 24.9)	12.2 (11.9, 12.5)	< 0.001	24.5 (24.2, 24.7)	11.4 (11.1, 11.8)	< 0.001	25.7 (25.3, 26.1)	13.9 (13.4, 14.4)	< 0.001
PedsQL ***									
Physical Summary	91.7 (91.3, 92.1)	82.6 (82.2, 83.0)	< 0.001	89.6 (89.1, 90.0)	80.9 (80.5, 81.4)	< 0.001	96.6 (95.9, 97.3)	86.1 (85.4, 86.8)	< 0.001
Psychosocial Summary	90.4 (89.9, 90.8)	76.2 (75.8, 76.6)	< 0.001	90.1 (89.6, 90.6)	75.0 (74.5, 75.5)	< 0.001	90.9 (90.2, 91.7)	78.6 (77.9, 79.3)	< 0.001
PedsQL Total	90.9 (90.5, 91.3)	78.4 (78.0, 78.8)	< 0.001	90.1 (89.6, 90.6)	75.0 (74.5, 75.5)	< 0.001	92.9 (92.2, 93.6)	81.2 (80.6, 81.9)	< 0.001

* Historical group N=1002; ** Historical group N = 2156; *** Historical group N = 5231

[^] Reported as n (%) for Sex and Grade, mean (SD) for Age,

Total scores for the PFABS, PHQ-9 and PedsQL reported as the sample weight * inverse propensity score weighted mean (95% CI)

Figure 1. Prevalence of Depression Symptoms for Female Adolescent Athletes Prior to the COVID-19 Pandemic (PreCoVID-19) and in During the CoVID-19 Pandemic (DuringCoVID-19).

PHQ9 categories are sample weight * inverse propensity score weighted % (95% CI)

Figure 2. Prevalence of Depression Symptoms for Male Adolescent Athletes Prior to the COVID-19 Pandemic (PreCoVID-19) and in During the CoVID-19 Pandemic (DuringCoVID-19).

PHQ9 categories are sample weight * inverse propensity score weighted % (95% CI)

581
582

Supplementary Table 1. Distribution of Adolescent Sport Participants for the PreCoVID-19 and During-CoVID-19 Groups

Sport	PreCoVID-19 All participants = 5,231 n (%)	PreCoVID-19 Females = 3,402 n (%)	During-CoVID-19 Females = 1,877 n (%)	PreCoVID-19 Males = 1,829 n (%)	During-CoVID-19 Males = 1,366 n (%)
Baseball	537 (10.3%)	5 (0.1%)	1 (0.1%)	532 (29.1%)	417 (30.5%)
Basketball	2259 (43.2%)	1590 (46.7%)	502 (26.7%)	669 (36.6%)	437 (32.0%)
Cheer	117 (2.2%)	115 (3.4%)	127 (6.8%)	2 (0.1%)	4 (0.3%)
Cross Country	122 (2.3%)	93 (2.7%)	272 (14.5%)	29 (1.6%)	159 (11.6%)
Football	1218 (23.3%)	6 (0.2%)	14 (0.7%)	1212 (66.3%)	621 (45.5%)
Golf	30 (0.6%)	24 (0.7%)	73 (3.9%)	6 (0.3%)	90 (6.6%)
Gymnastics	48 (0.9%)	48 (1.4%)	31 (1.7%)	0 (0.0%)	0 (0.0%)
Ice Hockey	111 (2.1%)	26 (0.8%)	38 (2.0%)	85 (4.6%)	44 (3.2%)
Lacrosse	91 (1.7%)	16 (0.5%)	19 (1.0%)	75 (4.1%)	15 (1.1%)
Soccer	749 (14.3%)	540 (15.9%)	391 (20.8%)	209 (11.4%)	189 (13.8%)
Softball	848 (16.2%)	846 (24.9%)	504 (26.9%)	2 (0.1%)	3 (0.2%)
Swimming	28 (0.5%)	22 (0.6%)	86 (4.6%)	6 (0.3%)	40 (2.9%)
Tennis	75 (1.4%)	70 (2.1%)	114 (6.1%)	5 (0.3%)	52 (3.8%)
Track	1117 (21.4%)	907 (26.7%)	588 (31.3%)	210 (11.5%)	468 (34.3%)
Volleyball	2916 (55.7%)	2907 (85.4%)	651 (34.7%)	9 (0.5%)	28 (2.0%)
Wrestling	123 (2.4%)	3 (0.1%)	30 (1.6%)	120 (6.6%)	174 (12.7%)
Other ^{\$}	62 (1.2%)	55 (1.7%)	94 (5.0%)	7 (0.4%)	51 (3.7%)
Total Sports Played [^]	2.0 (0.8)	2.1 (0.8)	1.9 (0.8)	1.7 (0.7)	2.0 (0.8)

583 [^]: reported as Mean (SD)

584 ^{\$} Other sports: Archery, Bowling, Field Hockey, Rugby, Snow Skiing (Alpine or Nordic), Power lifting, Trap Shooting, Ultimate
585 Frisbee

586