Title: Correlates of and changes in aerobic physical activity and strength training before and after the onset of COVID-19 pandemic in the UK – findings from the HEBECO study.

Authors: Aleksandra Herbec^{1,2*}, Verena Schneider¹, Abi Fisher¹, Dimitra Kale¹, Lion Shahab¹, Phillippa Lally¹.

1 Department of Behavioural Science and Health, University College London, UK

2 Clinical, Educational and Health Psychology, University College London, UK

*contact: Aleksandra Herbec a.herbec@ucl.ac.uk

Pre-registration: <https://osf.io/spdtb>

Tables are at the end of the manuscript

Abstract

Objectives: Understanding changes in moderate to vigorous aerobic physical activity (MVPA) and strength training (MSA) from before to after (pre-/post-) the onset of the Covid-19 pandemic in the UK (first lockdown) and their correlates can inform interventions.

Methods: Cross-sectional analysis of retrospective and concurrent data on MVPA/MSA pre- and post-Covid-19 (until 14th June 2020) among 2,657 UK adults. The associations between socio-demographic and health characteristics, MVPA/MSA pre-Covid-19, living and exercise conditions and meeting WHO recommended levels for MVPA/MSA/both (vs meeting neither), and changes in MVPA/MSA from pre- to post-Covid-19 following stratification for pre-Covid-19 MVPA/MSA levels were evaluated.

Results: A third of adults maintained (30.4%), decreased (36.2%) or increased (33.4%) their MVPA levels post-Covid-19. For MSA, the percentages were 61.6%, 18.2%, and 20.2%, respectively. MVPA increased or decreased by an average of 150min/week, and MSA by 2 days/week. Meeting both MSA+MVPA recommendations during lockdown (vs. meeting neither) was positively associated with meeting MVPA+MSA pre-lockdown (aOR=16.11,95%CI=11.24-23.07), and post-16-years of age education (aOR=1.57,1.14-2.17), and negatively associated with being obese (aOR=0.49,0.33-0.73), older age (65+ vs ≤34; aOR=.53,.32-.87), and annual household income <50.000GBP (vs ≥50.000GBP; aOR=0.65,0.46-0.91). The odds for decreasing MVPA were significantly lower for white ethnicity, post-16-years of age education, access to garden/balcony, and higher for those who were in total isolation. The odds for decreasing MSA were significantly higher for those who were overweight or obese.

Conclusion: Aerobic and strength training were differently impacted during the first UK lockdown, with poorer outcomes associated with older age, lower education, and higher body mass index.

Keywords: aerobic activity, strength training, covid-19, UK, cross-sectional, changes, MSA, MVPA

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

What are the new findings:

- At the start of the Covid-19 pandemic, a third of UK adults decreased and a third increased moderate-to-vigorous physical activity; the corresponding proportions for muscle strength training were 18% and 20%.
- Older adults were more likely to maintain inactivity from before to start of the pandemic.
- Adults with higher body mass index were at risk of not meeting WHO recommendations for weekly levels of strength and aerobic training, and at decreasing muscle strength training in this period.
- Those with lower incomes, no post-16-years of age education, ethnic minority background and living in total isolation had poorer physical activity outcomes.

Introduction

Physical activity has been negatively affected by the Covid-19 pandemic, which could have major implications for general health and Covid-19 outcomes^{1,2,3}. As the Covid-19 pandemic continues and new lockdowns are introduced it is paramount to understand how aerobic and muscle-strength training were affected during the pandemic so far, and which group in the society may require most support.

Adults who engage in less than 30min of moderate physical activity per week are considered to be inactive^{4,5}. For best health outcomes adults are recommend to engage in moderate-to-vigorous intensity aerobic physical activity (MVPA; i.e. activities that increase heart rate and make one feel warmer) for at least 150 minutes/week, as well as in muscle-strengthening activity (MSA; e.g., strength/resistance training) for at least two days/week⁶. MVPA and MSA both independently and combined lower morbidity and mortality^{7-11,12}.

The Covid-19 pandemic and the social distancing measures introduced in the UK on 23rd March 2020 have affected many opportunities to engage in physical activity¹³. Many sports centres (e.g. gyms) were temporarily closed and team sports banned, while access to outdoors spaces (e.g. remote national parks) was greatly reduced. On the other hand, some lockdown measures could facilitate exercising, including the introduction of remote working. Furthermore, exercising outdoors was listed as one of the few activities that were still permitted even during the strictest lockdown in the UK, alongside shopping for essential items or medicine and going to work.

Physical activity during the Covid-19 pandemic declined¹⁴, but a small minority of adults increased their activity¹⁵. Female gender, lower income, older age, health conditions, perceived risks and poorer mental health were associated with lower physical activity levels¹⁵⁻¹⁸. However, research to date has focused primarily on MVPA or failed to distinguish between MVPA or MSA^{15,19,20} and few studies accounted for confounders that could be relevant during the pandemic, such as having access to indoor and outdoor exercising space or the household makeup¹⁵.

Aims

We characterised MVPA and MSA levels during the periods *before* and *after* (pre- /post-) the Covid-19 pandemic had started in the UK, with the latter covering the period of the first strictest UK lockdown. The research questions (RQs) were:

- RQ1: What were the levels and changes in MVPA and MSA among UK adults from pre- to post- Covid-19?
- What socio-demographic, environmental and health factors were associated with:
 - RQ2: meeting WHO recommendations for MVPA and MSA post-Covid-19?
 - RQ3: (i) decrease in MVPA among those who were active, and (ii) increase in MVPA among those who were inactive from pre- to post-Covid-19?
 - RQ4: (i) decrease in MSA among those who were active, and (ii) increase in MSA among those who were inactive, from pre- to post-Covid-19?

Methods

Study Design

The study involved cross-sectional data analysis from the baseline survey of the HEalth BEhaviours during the COvid-19 pandemic (HEBEco) study (<https://osf.io/sbgru/>). Data were collected using REDCap at UCL (Harris et al, 2019). The study protocol was preregistered (<https://osf.io/eighs/>). Departures from the protocol are explained in Supplementary Materials 1 (<https://osf.io/w7vbf/>). The results from all the pre-registered analyses not reported here are available on Open Science Framework: <https://osf.io/2ujxq/>. The reporting follows the STROBE guidelines.²¹

Participants

Recruitment involved a UK-wide campaign including paid and unpaid posts on social media and information shared through the networks of Cancer Research UK, Public Health England, other charities, sports clubs, universities, and local authorities. Participants with complete data who enrolled into the baseline between 5th May 2020 (start of the data collection for all study variables) until 14th June 2020 (prior to easing of restrictions of the first UK lockdown) were included.

Measures

For the wording of all measures see <https://osf.io/bja7g/> and Supplementary Materials 2. At baseline participants indicated the time when their lives started to be affected by the Covid-19 pandemic in any way, which was used as an anchor to assess before/since (pre-/post-Covid-19) MVPA and MSA. Almost half (44.0%, weighted) of adults selected the second half of March, followed by 26.9% who selected the first half of March.

Meeting WHO recommendations for MVPA and MSA

The pre-/post-Covid-19 MVPA and MSA were measured using the same set of questions^{12,22}. For MVPA, participants were first asked “In the month Before/Since Covid-19, on average, HOW MANY TIMES PER WEEK did you do at least 15 MINUTES or more of moderate or vigorous AEROBIC PHYSICAL ACTIVITY? Examples: brisk walk, jogging, dancing, cycling for recreation or commute, swimming, team or racket sports.” The answers were capped at 14+, equivalent to at least twice per day. The 15 minute period was selected as it corresponds to the minimum amount of PA needed for mortality reduction and extending lifespan²³. Those who engaged in at least one such session were asked “In the month BEFORE/Since COVID-19, HOW LONG (in MINUTES) was your average session of moderate or vigorous AEROBIC PHYSICAL ACTIVITY? Do NOT include strength training.” Total weekly MVPA was computed by multiplying the session number by the minutes.

MSA activity was assessed by a single question: “Before/Since Covid-19, on average, on HOW MANY DAYS PER WEEK did you do STRENGTH TRAINING? Examples: Pilates, push-ups, squats, yoga, and exercises involving free weights, weight machines or elastic band.” The answers were capped at 4+ given that the recommendations were for 2 days of MSA per week.

Data on MVPA and MSA were used to categorise participants, separately for pre-/post-Covid-19, into four groups based on whether they met the WHO recommendations for MSA (at least 2 days/week) and MVPA (≥ 150 minutes/week)⁶: meeting neither, meeting MSA only, meeting MVPA only, or meeting recommendations for both MSA and MVPA.

Change in MVPA and MSA

Changes in MVPA (RQ 3) and MSA (RQ 4) were calculated by subtracting the pre-Covid-19 levels of MSA or MVPA from the post-Covid-19 levels. Participants were categorised into: maintenance (change in MVPA < 20 min; change in MSA = 0), decrease (MVPA by ≥ 20 min; MSA by ≥ 1 day) or increase in activity (MVPA by ≥ 20 min; MSA by ≥ 1 day). Dichotomous variables were then created: decrease (vs maintain/increase) and increase (vs maintain/decrease) in MVPA and MSA.

To minimise the bias due to ceiling and floor effects when assessing changes to MVPA and MSA, the analyses for RQs 3 and 4 were conducted after stratifying for pre-Covid-19 MVPA and MSA. Participants were categorised into: inactive MVPA (< 30 min of MVPA/week^{4,5}) or MSA (0 days of MSA/week) and active (≥ 30 min/week of MVPA; and ≥ 1 day/week of MSA) pre-Covid-19.

Participants who were active post-Covid-19 were also asked “Do you do the same FORM of exercise as you did before the COVID-19? (even if it is in a different location)?” The answer options were: None of the same/some of the same/about half the same/mostly the same/exactly the same.

Explanatory factors and correlates

Socio-demographic characteristics and living conditions assessed were: gender (female/other), ethnicity (white/non-white); education (post-16-years of age/other); employed (yes/no); furloughed/laid off (yes/no); income (low-middle < 50 000 GBP/high ≥ 50 000 GBP/prefer-not-to-say);

age (in decades was entered as a continuous variable where it met assumptions, or as a 3-level categorical variable: ≤34, 35-64, 65+).

We assessed health behaviours, health and living conditions that could impact on MVPA or MSA levels: self-reports of any condition that limited physical activity (yes/no); body mass index (BMI²²: normal and underweight ≤24.99/overweight 25-29.99/ obese ≥30); smoking status (current smoker vs not)²⁴; weekly frequency of alcohol drinking²⁵; deterioration in psychological wellbeing from pre- to post-Covid-19 (yes/no); isolation status (total isolation/other); participants' perceived risk of Covid-19 to their health ('no or minor risk'/other); access to a garden or balcony big enough to exercise comfortably (yes/no); access to a public park/green space that is within a walking distance and open during Covid-19 (yes/no); living with children aged ≤15 (yes/no); living with vulnerable persons (persons over the age of 70, in poor health, or who may be vulnerable to Covid-19; yes/not).

Model included two time co-variables to account for weather changes: enrolment time (up until 15th May, the second half of May, the first half June), and time when Covid-19 started to affect individuals (before mid-March/after).

Analysis

Analyses were conducted in SPSS 26 with the data weighted using the 2018 Census and APS mid-year estimates for age, gender, ethnicity, country of living and household income. The analysis used weights trimmed to top 98th percentile to minimise the impact of extremely high weights²⁶. Differences between the included and excluded participants were assessed with chi-squared for categorical and t-tests for continuous data.

For RQ1, descriptive statistics were computed to characterise the levels of MVPA, MSA, and inactivity pre-/post-Covid-19 and changes in exercise form. For RQ2 univariate and fully adjusted multinomial regression models were computed. For RQ3 and RQ4 the sample was stratified using pre-Covid-19 MVPA or MSA levels to assess outcomes of interest using univariate and adjusted logistic regression models. Sensitivity analyses involved replicating the analyses using unweighted data to check for the robustness of the results, and for RQ3 using different cut-off values of 15mins and 30mins.

Family-wise error was corrected for by using the Benjamini-Hochberg procedure separately for each research question (Benjamini and Hochberg, 1995).

Results

The final analytic sample included 2,657 adults with complete study data (weighted N=2,442). Table 1 present comparisons of the excluded and included sample. The included weighted sample comprised 52% females, 90.5% of white ethnicity, and 67.0% with high school education or higher.

Below are results for the weighted fully adjusted analyses. Results from univariable analyses are reported in Supplementary Materials 3, and from sensitivity analyses in Supplementary Materials 4 and 5, and have not changed the conclusions.

Research Question 1: Changes in MVPA and MSA pre- to post-Covid-19.

Table 2 presents data on physical activity levels before and since Covid-19. Pre-Covid-19 17.6% of adults engaged in no MVPA and MSA, 55.3% engaged in no MSA, and 19.4% engaged in no MVPA. Just under 15% of adults met the recommended levels of both MVPA and MSA pre- and post- Covid-19. The proportion of those who had no MVPA or MSA activity increased minimally (17.4% to 22.1%) which was primarily driven by declines in MVPA. Among those who were active post-Covid-19, 41.5% continued with the mostly or exactly activity form as pre-Covid-19.

From before to since Covid-19 similar proportions of adults maintained (30.4%), decreased (36.2%) or increased (33.5%) their weekly MVPA. Among those who were active pre-Covid-19, 46.7% decreased MVPA. The decrease was of on average 219min/week. Among those who were inactive pre-Covid-19, 29.7% increased MVPA, with an average increase of 144min/week.

Maintenance of MSA was relatively more common (61.6%), with 18.2% decreasing and 20.2% increasing the number of days they engaged in MSA. Among those who were active pre-Covid-19 (40.2%), 45.1% decreased MSA, with an average decrease of 1.9 days/week. Among those who engaged in no MSA pre-Covid-19 (59.8%), 16.4% increased, with an average increase of 2.5 days/week.

Research Question 2: Predictors of meeting WHO recommendations

Table 3 presents results from fully adjusted models. Being aged 65+, having a lower pre-Covid-19 household income, having a condition limiting physical activity, being obese, living in total isolation and deterioration in psychological wellbeing were associated with lower odds, and having at least high school education or meeting the WHO recommended levels of MVPA, MSA or both pre-Covid-19 was associated with greater odds of meeting both MVPA and MSA WHO guidelines since Covid-19 compared with not meeting either.

Having a condition that limited physical activity, being in total isolation and deterioration in psychological wellbeing was associated with lower odds, and meeting the WHO recommended levels of MVPA or both MVPA and MSA was associated with greater odds of meeting MVPA only post-Covid-19. Being aged 35-64, and having obesity (BMI $\geq$ 30) was associated with lower odds, while higher education, and meeting both MVPA and MSA or only MSA recommendations pre-Covid-19 was associated with meeting the recommendations for MSA only post-Covid-19.

RQ 3: Associations with changes in MVPA

Older, white participants with at least high school education and access to a garden/balcony to exercise comfortably in were less likely to decrease MVPA activity (Table 4). Those with conditions limiting PA, living in total isolation and experiencing a deterioration in psychological wellbeing during lockdown were more likely to decrease MVPA activity from pre-Covid levels.

White ethnicity, being employed during Covid-19, and living in total isolation were associated with lower odds of increasing MVPA among this group. At least high school education and living with children was associated with higher odds of increasing MVPA activity.

RQ 4: Associations with changes in MSA

Being employed during lockdown and being overweight (BMI 25-29.99) or obese (BMI ≥ 30) was associated with higher odds of decreasing MSA activity. Older age (35-64, and 65+) and deterioration in psychological wellbeing were associated with lower odds of increasing MSA. See Table 5 for details.

Discussion

This study shows that the first Covid-19 lockdown in the UK affected differently the MVPA and MSA levels, with different factors associated with engagement in and changes in these two activity types.

Meeting WHO recommendations for MVPA and MSA post-Covid-19

Adherence to WHO recommendations for both MVPA and MSA pre- and post-Covid-19 was low – at about 15%. Older adults (aged 65+), adults who were obese, those with pre-Covid-19 household income below 50,000 GBP and who had conditions limiting physical activity were at risk of not meeting the WHO recommendations for both MVPA and MSA. These groups should be target by future interventions.

Moreover, meeting the recommendations for MVPA and MSA before Covid-19 was strongly predictive of meeting them post-Covid-19. Additionally, adults who remained active tended to maintain at least similar levels and forms of the exercises as pre-pandemic. Thus, as UK adults remain relatively consistent in their behaviour even during the lockdown, this study highlights the need to especially support those with low activity levels to develop healthy exercising routines.

Changes to MVPA and MSA levels from pre- to the post-Covid-19 pandemic start

Although the group level data suggest little change in physical activity pre- to post-Covid-19, especially MSA, about a third of adults decreased and another third increased their MVPA from pre- to post-Covid-19. Those who changed MVPA decreased or increased their MVPA levels by over three and two hours per week, respectively. Those who changed MSA either gained or lost two days of MSA per week.

Decreases in MVPA was found among other adults samples in the UK¹⁵, Italy¹⁹ and Spain²⁰. However, the findings that over 70% of UK adults maintained or increased their MVPA could be at least partially attributed to the UK government's consistent lockdown policies that allowed leaving the house for exercising, which placed UK in contrast to other countries^{20,27}.

The maintenance of MSA levels could be at least partially explained by the low pre-Covid-19 MSA levels. Additionally, many MSA exercises can be performed with no equipment and at one's home (e.g. Pilates, push-ups) and thus may be less affected by social distancing measures. This is further supported by the present findings that total isolation (i.e. not leaving the house for any reason) was associated with MVPA but not MSA levels.

Factors associated with changes and maintenance of MVPA and MSA post-Covid-19

Older adults were more likely to maintain their pre-Covid-19 MVPA and MSA levels, including inactivity. Due to more established routines and lower baseline physical activity this group might have been less affected by the lockdown restrictions, such as gym closures or team sports bans²⁸.

MVPA inactivity was more commonly maintained by those who were employed or living in total isolation. Such adults likely had fewer opportunities or flexibility to exercise. Additionally, those who had a health condition limiting physical activity and those living in total isolation tended to decrease MVPA. Factors that were associated with maintaining MVPA activity post-Covid-19 were white ethnicity, higher education, and having a garden or balcony large enough to exercise.

Increasing MVPA levels among those who were previously inactive was most common among those with higher education, which is in line with prior studies²⁹, and those who lived with children. Caring for children might have promoted some forms of MVPA, including outdoors, particularly as parents became responsible for children's physical education during school lockdown closures.

In terms of changes in MSA levels, older adults were at higher risk of maintaining MSA inactivity, while adults with higher BMI levels were at greater risk of decreasing MSA from pre- to post-Covid-19. Therefore, both older adults and those with higher BMI should be among priority groups targeted with MSA interventions, particularly as MSA can bring important clinical benefits to these two groups³⁰⁻³³. Other priority groups for MSA training are those with lower household income and lower educational attainment.

Finally, as found previously¹⁸, psychological wellbeing was associated with physical activity levels. In the present study deterioration in psychological wellbeing was associated with not meeting guidelines for MVPA or decreasing MVPA levels, but not with MSA. Due to the cross-sectional design causality cannot be assumed, as poorer psychological wellbeing (e.g. lower mood, anxiety, stress), can be both a predictor and a consequence of low exercise levels^{34,35}. However, interventions aimed at improving physical activity are likely to improve mental health as well³⁴⁻³⁶.

Strength and limitation

The study is among a few that assessed the levels and changes to both MVPA and MSA during the first UK lockdown. The study drew on previously used measures of MVPA and MSA³⁷, which were supplemented by images to clarify exercise types. This study also benefits from a large list of correlates and covariates being measured to control for confounding. A number of sensitivity analyses were conducted to test the robustness of findings. The key limitations are that this was a cross-sectional study among self-selected sample that relied on self-report and recall that are prone to bias. Finally, the unfolding of the Covid-19 pandemic has coincided with season change from winter to spring. Without a true baseline from the same period in 2019 it is not possible to tease apart the effect of weather change from that of the pandemic.

Conclusions

The findings point to social inequalities in how the first lockdown in the UK has affected physical activity levels, with differential impact on aerobic and strength training. Dedicated interventions are needed to support MVPA and MSA especially among those who have low activity levels, those who are older, have lower income, and have higher body mass index.

Footnotes

- **Contributors** AH conceived the idea in consultation with PL, AB and LS. AH and VS analysed the data. All authors interpreted the data. AH drafted the first version of the manuscript. All authors critically revised the manuscript and approved the final draft before submission.
- **Funding** This project is partially funded by an ongoing Cancer Research UK Programme Grant to UCL Tobacco and Alcohol Research Group (C1417/A22962) and by SPECTRUM, a UK Prevention Research Partnership Consortium (MR/S037519/1).
- **Competing interests** None declared.
- **Patient and public involvement** Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.
- **Patient consent for publication** Obtained.
- **Ethics approval** The study was approved by the by UCL Research Ethics Committee at the UCL Division of Psychology and Language Sciences (PaLS) (CEHP/2020/579; as part of the larger programme of work: The optimisation and implementation of interventions to change behaviours related to health and the environment).
- **Acknowledgements:** We are grateful to all participants who have been supporting our research. We would like to thank Public Health England, and particularly members of the Behavioural Insights at Public Health England for providing feedback on the survey wording. We would like to thank Public Health England, Cancer Research UK, local authorities, Mayors' offices, as well as charities and other organisations in the UK, including the Asthma UK and British Lung Foundation Partnership, for supporting our recruitment campaign.

References

1. Warburton DER, Bredin SSD. Health benefits of physical activity: a systematic review of current systematic reviews. *Curr. Opin. Cardiol.* 2017;32(5):541-556.
2. Woods JA, Hutchinson NT, Powers SK, et al. The COVID-19 pandemic and physical activity. *Sports Medicine and Health Science.* 2020;2(2):55-64.
3. Shur NF, Johns D, Kluzek S, Peirce N. Physical inactivity and health inequality during coronavirus: a novel opportunity or total lockdown? *BMJ Open Sport & Exercise Medicine.* 2020;6(1):e000903.
4. Public Health England. Physical inactivity levels in adults aged 40 to 60 in England. 2017; Accessed on 20th September 2020 from: <https://www.gov.uk/government/publications/physical-inactivity-levels-in-adults-aged-40-to-60-in-england/>.
5. Health Survey for England. Health Survey for England 2016 - Physical activity in adults. 2017; Accessed on 20th Sept 2020 from: <http://healthsurvey.hscic.gov.uk/media/63730/HSE16-Adult-phy-act.pdf>
6. WHO. Physical activity. . 2018; Accessed on 20th November 2020 from <https://www.who.int/news-room/fact-sheets/detail/physical-activity>.
7. Bennie JA, Shakespear-Druery J, De Cocker K. Muscle-strengthening Exercise Epidemiology: a New Frontier in Chronic Disease Prevention. *Sports medicine - open.* 2020;6(1):40.
8. Kamada M, Shiroma EJ, Buring JE, Miyachi M, Lee IM. Strength Training and All-Cause, Cardiovascular Disease, and Cancer Mortality in Older Women: A Cohort Study. *Journal of the American Heart Association.* 2017;6(11).
9. Stamatakis E, Lee IM, Bennie J, et al. Does Strength-Promoting Exercise Confer Unique Health Benefits? A Pooled Analysis of Data on 11 Population Cohorts With All-Cause, Cancer, and Cardiovascular Mortality Endpoints. *American journal of epidemiology.* 2018;187(5):1102-1112.
10. Zhao G, Li C, Ford ES, et al. Leisure-time aerobic physical activity, muscle-strengthening activity and mortality risks among US adults: the NHANES linked mortality study. *Br. J. Sports Med.* 2014;48(3):244-249.
11. Zhao M, Veeranki SP, Magnussen CG, Xi B. Recommended physical activity and all cause and cause specific mortality in US adults: prospective cohort study. *BMJ (Clinical research ed).* 2020;370:m2031.
12. Bennie JA, De Cocker K, Teychenne MJ, Brown WJ, Biddle SJH. The epidemiology of aerobic physical activity and muscle-strengthening activity guideline adherence among 383,928 U.S. adults. *The international journal of behavioral nutrition and physical activity.* 2019;16(1):34.
13. Ricci F, Izzicupo P, Moscucci F, et al. Recommendations for Physical Inactivity and Sedentary Behavior During the Coronavirus Disease (COVID-19) Pandemic. *Frontiers in public health.* 2020;8:199.
14. Ammar A, Brach M, Trabelsi K, et al. Effects of COVID-19 Home Confinement on Eating Behaviour and Physical Activity: Results of the ECLB-COVID19 International Online Survey. *Nutrients.* 2020;12(6).
15. Rogers NT, Waterlow N, Brindle HE, et al. Behavioural change towards reduced intensity physical activity is disproportionately prevalent among adults with serious health issues or self-perception of high risk during the UK COVID-19 lockdown. . *MedRxiv* 2020.
16. Meyer J, McDowell C, Lansing J, et al. Changes in physical activity and sedentary behaviour due to the COVID-19 outbreak and associations with mental health in 3,052 US adults. *Cambridge Open Engage.* 2020.
17. Smith L, Jacob L, Butler L, et al. Prevalence and correlates of physical activity in a sample of UK adults observing social distancing during the COVID-19 pandemic. *BMJ Open Sport & Exercise Medicine.* 2020;6(1):e000850.

18. Jacob L, Tully MA, Barnett Y, et al. The relationship between physical activity and mental health in a sample of the UK public: A cross-sectional study during the implementation of COVID-19 social distancing measures. *Mental health and physical activity*. 2020;19:100345.
19. Maugeri G, Castrogiovanni P, Battaglia G, et al. The impact of physical activity on psychological health during Covid-19 pandemic in Italy. *Heliyon*. 2020;6(6):e04315.
20. López-Bueno R, Calatayud J, Andersen LL, et al. Immediate Impact of the COVID-19 Confinement on Physical Activity Levels in Spanish Adults. *Sustainability*. 2020;12(14):5708.
21. von Elm E, Altman DG, Egger M, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ (Clinical research ed)*. 2007;335(7624):806-808.
22. Churilla JR, Johnson TM, Richardson MR, Williams BD, Rariden BS, Boltz AJ. Mode of physical activity participation by body mass index: 2015 behavioural risk factor surveillance system. *Res. Sports Med*. 2018;26(2):147-157.
23. Wen CP, Wai JPM, Tsai MK, et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *The Lancet*. 2011;378(9798):1244-1253.
24. Kaczynski AT, Manske SR, Mannell RC, Grewal K. Smoking and Physical Activity: A Systematic Review. *American Journal of Health Behavior*. 2008;32(1).
25. Leasure JL, Neighbors C, Henderson CE, Young CM. Exercise and Alcohol Consumption: What We Know, What We Need to Know, and Why it is Important. *Frontiers in psychiatry*. 2015;6:156.
26. Potter FA. Study of Procedures to Identify and Trim Extreme Sample Weights. . *Proceedings of the Survey Research Methods Section, American Statistical Association*. 1990:225-230.
27. DW. Coronavirus: What are the lockdown measures across Europe? 2020; Accessed 20th Sept 2020 from: <https://www.dw.com/en/coronavirus-what-are-the-lockdown-measures-across-europe/a-52905137>.
28. Sun F, Norman IJ, While AE. Physical activity in older people: a systematic review. *BMC public health*. 2013;13:449.
29. Shaw BA, Spokane LS. Examining the association between education level and physical activity changes during early old age. *J. Aging Health*. 2008;20(7):767-787.
30. Kraschnewski JL, Sciamanna CN, Poger JM, et al. Is strength training associated with mortality benefits? A 15year cohort study of US older adults. *Prev Med*. 2016;87:121-127.
31. Guizelini PC, de Aguiar RA, Denadai BS, Caputo F, Greco CC. Effect of resistance training on muscle strength and rate of force development in healthy older adults: A systematic review and meta-analysis. *Exp. Gerontol*. 2018;102:51-58.
32. Said MA, Abdelmoneem M, Almaqhawi A, Hamid Kotob AA, Alibrahim MC, Bougmiza I. Multidisciplinary approach to obesity: Aerobic or resistance physical exercise? *Journal of exercise science and fitness*. 2018;16(3):118-123.
33. Willis LH, Slentz CA, Bateman LA, et al. Effects of aerobic and/or resistance training on body mass and fat mass in overweight or obese adults. *J Appl Physiol (1985)*. 2012;113(12):1831-1837.
34. Sharma A, Madaan V, Petty FD. Exercise for mental health. *Primary care companion to the Journal of clinical psychiatry*. 2006;8(2):106.
35. Mikkelsen K, Stojanovska L, Polenakovic M, Bosevski M, Apostolopoulos V. Exercise and mental health. *Maturitas*. 2017;106:48-56.
36. Callow DD, Arnold-Nedimala NA, Jordan LS, et al. The Mental Health Benefits of Physical Activity in Older Adults Survive the COVID-19 Pandemic. *Am. J. Geriatr. Psychiatry*. 2020;28(10):1046-1057.
37. Bennie JA, Pedisic Z, van Uffelen JG, et al. The descriptive epidemiology of total physical activity, muscle-strengthening exercises and sedentary behaviour among Australian adults--results from the National Nutrition and Physical Activity Survey. *BMC public health*. 2016;16:73.

Table 1. Sample characteristics (weighted).

	Included sample N=2442	Excluded sample N=350	<i>p</i>
Female	52.6%	48.7%	0.179
Age; M (SD)	50.0 (16.0)	42.7 (18.9)	<0.001
White ethnicity	90.5%	82.4%	<0.001
Household income ≥50,000	18.1%	15.5%	0.095
Income: <50,000 GBP	73.5%	73.1%	
Income: Prefer not to say	8.3%	11.5%	
Education: Highschool or higher	67.0%	69.1%	0.432
Employed	48.3%	40.7%	0.008
Laid-off/Furloughed	12.7%	15.2%	0.197
Condition limiting PA	16.3%	21.2%	0.035
BMI ≤24	43.2%	27.3%	<0.001
BMI: 25-29.99	35.1%	16.8%	
BMI: Obese: 30+	21.7%	8.4%	
BMI don't know/prefer not to say ^a	0%	47.6%	
Total isolation	7.8%	8.8%	0.541
Minor/no Covid-19 risk percept	33.3%	38.6%	0.069
Living with children	17.0%	15.7%	0.549
Living with vulnerable	15.1%	21.8%	0.001
Access garden/balcony	72.2%	62.6%	<0.001
Access green space	59.5%	52.4%	0.012
Time life affected by Covid-19 mid- March or sooner ^b	44.8%	53.3%	0.003
Enrolled from 1 st June (ref)	4.4%	5.4%	0.404
Enrolled up until 15 th May	50.1%	46.7%	
Enrolled second half of May	45.4%	47.9%	
Smoker	24.1%	38.9%	<0.001
Weekly alcohol drinking, none (ref)	21.8%	21.8%	0.920
Weekly or less	27.9%	26.9%	
More than weekly	50.3%	51.3%	
Deteriorated psychological wellbeing	54.1%	52.8%	0.708
Meeting WHO PA rec (meeting both=ref)	15.6%	17.3%	0.218
Meeting only MVPA before	23.0%	22.0%	
Meeting only MSA before	13.0%	7.7%	
Meeting neither before	48.4%	53.0%	

^aAs per protocol, the participants who did not provide BMI data (e.g. they selected 'prefer not to say' or 'don't know' on the question on height or weight) but who provided other study data were to be included in the analyses, but were ultimately excluded due to having missing data on other variables. ^bTime that participant's life started to be affected by Covid-19 in any way

Table 2. Aerobic physical activity (MVPA) and strength training (MSA) before and since COVID-19.

	Before Covid-19 ¹	Since Covid-19 ¹
No activity (0 sessions MVPA and 0 days MSA) % (N)	17.6 (430)	22.1 (540)
MVPA		
MVPA 0 sessions % (N)	19.4 (473)	24.8 (604)
MVPA min/week among active; Median (IQR)	125 (210.0)	150 (220.0)
Mean (SD)	238.1 (424.0)	230.9 (398.8)
MVPA min/week among entire sample; Median (IQR)	90 (180.0)	90 (224.5)
Mean (SD)	196.2 (395.4)	179.8 (364.7)
MSA		
MSA % (N)		
0 das/week	55.3 (1557)	56.2 (1584)
1 day/week	11.1 (314)	9.6 (271)
2 days/week	10.4 (293)	7.9 (223)
3 days/week	9.0 (253)	7.6 (213)
4+ days/week	7.0 (197)	11.3 (320)
MSA Median (IQR)	0 (2)	0 (2)
Meet guidelines % (N)		
Neither MVPA nor MSA	45.1 (1271)	44.6 (1256)
Meet MVPA (≥150min/week) only	21.3 (599)	21.2 (598)
Meet MSA (≥2 days/week) only	11.7 (330)	11.9 (336)
Meets both MVPA and MSA	14.5 (410)	14.8 (416)
Change in form of exercise among those who exercise % (N)		
None of the same	-	17.1 (331)
Some of the same	-	32.7 (632)
About half the same	-	8.7 (168)
Mostly the same	-	26.0 (503)
Exactly the same	-	15.5 (299)

HEBECO Aerobic Strength Covid-19 Pandemic

Table 3. Predictors of meeting both, MVPA only and MSA only WHO guidelines since Covid-19 in comparison to not meeting WHO guidelines for MVPA and MSA (reference, n=1178). Results from fully adjusted models on weighted data with BH FDR correction of p-values (significant in bold).

	Both MVPA and MSA (n=388)				MVPA only (n=570)				MSA only (n=306)			
	aOR	95% CI		p	aOR	95% CI		p	aOR	95% CI		p
Female	1.08	0.82	1.41	0.588	0.79	0.62	1.00	0.049	1.02	0.77	1.35	0.906
Age ≤34												
Age: 35-64	0.70	0.49	0.99	0.047	1.11	0.78	1.56	0.566	0.56	0.38	0.82	0.003
Age: 65+	0.53	0.32	0.87	0.012	1.10	0.71	1.70	0.676	0.99	0.60	1.61	0.961
White ethnicity	1.61	1.03	2.52	0.036	1.57	0.99	2.49	0.054	0.72	0.47	1.10	0.130
Household income ≥50.000												
Income: <50 000 GBP	0.65	0.46	0.91	0.011	0.71	0.52	0.96	0.028	1.13	0.76	1.67	0.556
Income: Prefer not to say	0.44	0.25	0.77	0.004	0.63	0.38	1.03	0.065	0.88	0.49	1.59	0.666
Education: Highschool or higher	1.57	1.14	2.17	0.006	1.04	0.81	1.35	0.745	1.74	1.24	2.45	0.002
Employed	0.76	0.55	1.05	0.094	1.09	0.82	1.43	0.562	1.06	0.75	1.49	0.734
Laid-off/Furloughed	0.79	0.51	1.23	0.295	1.05	0.71	1.54	0.813	1.14	0.71	1.83	0.584
Condition limiting physical activity	0.44	0.28	0.71	0.001	0.43	0.29	0.63	<0.001	0.97	0.65	1.45	0.881
BMI ≤24												
BMI: 25-29.99	0.81	0.60	1.10	0.176	1.01	0.78	1.32	0.926	0.72	0.53	0.99	0.043
BMI: Obese: 30+	0.49	0.33	0.73	<0.001	0.87	0.64	1.19	0.394	0.37	0.24	0.56	<0.001
Total isolation	0.44	0.23	0.83	0.011	0.26	0.14	0.50	<0.001	0.83	0.49	1.40	0.478
Minor/no Covid-19 risk percept	1.00	0.74	1.34	0.977	1.20	0.93	1.55	0.167	0.94	0.68	1.30	0.718
Living with children	0.81	0.57	1.16	0.254	0.75	0.55	1.03	0.079	0.66	0.44	0.99	0.045
Living with vulnerable	1.17	0.82	1.68	0.395	0.86	0.61	1.20	0.366	0.87	0.58	1.30	0.499
Access garden/balcony to exercise comfortably in	1.10	0.81	1.51	0.533	1.02	0.79	1.33	0.861	0.80	0.58	1.09	0.159
Access green space within walking distance	1.27	0.96	1.69	0.096	1.31	1.03	1.66	0.028	0.91	0.68	1.23	0.548
Smoker	0.72	0.51	1.02	0.062	0.93	0.70	1.22	0.593	0.80	0.57	1.14	0.218
Weekly alcohol drinking, none (ref)												
Weekly or less	1.00	0.68	1.49	0.988	1.43	1.02	2.00	0.038	1.06	0.71	1.59	0.782
More than weekly	1.04	0.73	1.50	0.811	1.28	0.94	1.74	0.121	0.89	0.61	1.29	0.528
Deteriorated psychological wellbeing	0.56	0.43	0.73	<0.001	0.71	0.56	0.89	0.003	0.72	0.55	0.96	0.025
Meeting WHO PA ^a rec pre-Covid (meeting neither=ref)												
Meeting both before	16.11	11.24	23.07	<0.001	2.31	1.58	3.39	<0.001	3.72	2.45	5.62	<0.001
Meeting only MVPA before	3.88	2.64	5.70	<0.001	7.57	5.82	9.84	<0.001	1.47	0.94	2.30	0.092
Meeting only MSA before	6.38	4.26	9.54	<0.001	0.74	0.44	1.25	0.264	9.61	6.70	13.77	<0.001

a=meeting World Health Physical activity recommendations before the first Covid-19 lockdown in the UK. Notes: All models were also adjusted for the time of enrolment and time when Covid-19 started to affect individuals in any way. Following the BH correction the highest p-value that met the threshold for significance was p=0.019

HEBECA PA analysis baseline

Table 4. Independent associations of change in MVPA. Segmented analyses of active sample (pre-Covid-19 MVPA activity ≥ 30 min; predicting decrease by ≥ 20 min) and inactive sample (pre-Covid-19 MVPA activity < 30 min; predicting increase by ≥ 20 min).

	Sample active (≥ 30 min/week) pre-Covid-19 (weighted n=1857)				Sample inactive (< 30 min/week) pre-Covid-19 (weighted n=585)			
	Decrease (n=868) vs not				Increase (n=174) vs not			
	aOR	95% CI	p		aOR	95% CI	p	
Female	1.22	1.00	1.49	0.047	1.56	1.02	2.38	0.040
Age (in decades)	0.91	0.85	0.98	0.015	0.72	0.61	0.84	<0.001
White ethnicity	0.62	0.44	0.86	0.005	1.01	0.44	2.33	0.982
Household income $\geq 50,000$								
Income: $< 50,000$ GBP	1.30	1.00	1.67	0.048	0.49	0.26	0.93	0.030
Income: Prefer not to say	1.22	0.81	1.85	0.335	0.36	0.14	0.93	0.034
Education: Highschool or higher	0.73	0.58	0.91	0.006	2.26	1.40	3.63	0.001
Employed	0.86	0.69	1.07	0.180	0.51	0.31	0.85	0.009
Laid-off/Furloughed	1.05	0.76	1.47	0.750	1.30	0.72	2.37	0.385
Condition limiting PA	1.74	1.27	2.39	0.001	0.49	0.27	0.89	0.019
BMI ≤ 24								
BMI: 25-29.99	1.03	0.82	1.28	0.816	1.54	0.94	2.53	0.088
BMI: Obese: 30+	0.87	0.66	1.14	0.317	0.74	0.42	1.31	0.307
Total isolation	3.81	2.33	6.23	<0.001	0.29	0.12	0.73	0.008
Minor/no Covid-19 risk percept	0.99	0.80	1.23	0.926	1.20	0.76	1.92	0.436
Living with children	0.84	0.65	1.09	0.187	2.32	1.27	4.24	0.006
Living with vulnerable	0.78	0.59	1.02	0.074	0.90	0.51	1.59	0.716
Access garden/balcony to exercise comfortably in	0.75	0.60	0.94	0.011	0.72	0.46	1.13	0.157
Access green space within walking distance	0.82	0.67	1.01	0.059	1.10	0.72	1.67	0.665
Smoker	1.09	0.85	1.40	0.504	1.15	0.73	1.80	0.551
Weekly alcohol drinking, none (ref)								
Weekly or less	0.78	0.59	1.04	0.087	0.77	0.43	1.38	0.378
More than weekly	0.78	0.60	1.02	0.068	0.88	0.52	1.50	0.644
Deteriorated psychological wellbeing	1.40	1.15	1.71	0.001	1.09	0.72	1.66	0.682

Notes: All models were also adjusted for the time of enrolment and time when Covid-19 started to affect individuals in any way. Following the BH correction the highest p-value that met the threshold for significance was $p=0.019$.

HEBECO PA analysis baseline

Table 5. Research question 4: Independent associations of change in MSA. Segmented analyses of active sample (pre-Covid-19 MSA activity ≥ 1 day/week; predicting decrease by ≥ 1 day/week) and less inactive sample (pre-Covid-19 MSA activity 0 days/week; predicting increase by ≥ 1 day/week). Findings from fully adjusted logistic regression models on weighted data and using BH FDR adjustment (significant in bold).

	Sample active (≥ 1 day/week) pre-Covid-19 (weighted n=985)				Sample inactive (0 days/week) pre-Covid-19 (weighted n=1456)			
	Decrease (n=444) vs not				Increase (n=239) vs not			
	aOR	95% CI		p	aOR	95% CI		p
Female	0.94	0.71	1.23	0.634	0.96	0.70	1.31	0.792
Age (in decades)	1.03	0.94	1.14	0.516	-	-	-	-
Age ≤ 34								
Age: 35-64	-	-	-	-	0.22	0.15	0.33	<0.001
Age: 65+	-	-	-	-	0.34	0.20	0.58	<0.001
White ethnicity	0.75	0.51	1.11	0.154	0.71	0.41	1.23	0.216
Household income $\geq 50,000$								
Income: <50,000 GBP	1.11	0.80	1.55	0.535	0.67	0.45	0.99	0.047
Income: Prefer not to say	0.94	0.54	1.62	0.818	0.56	0.29	1.08	0.084
Education: Highschool or higher	0.82	0.59	1.15	0.260	1.20	0.85	1.70	0.310
Employed	1.81	1.34	2.46	<0.001	1.08	0.75	1.55	0.671
Laid-off/Furloughed	1.34	0.85	2.12	0.205	1.30	0.80	2.10	0.289
Condition limiting PA	1.11	0.72	1.72	0.632	0.64	0.38	1.05	0.078
BMI ≤ 24.99								
BMI: 25-29.99	1.88	1.39	2.55	<0.001	0.89	0.63	1.26	0.512
BMI: Obese: 30+	3.38	2.23	5.14	<0.001	0.64	0.42	0.97	0.037
Total isolation	1.87	1.02	3.42	0.042	0.80	0.42	1.52	0.499
Minor/no Covid-19 risk percept	1.43	1.06	1.92	0.019	0.93	0.65	1.31	0.669
Living with children	1.01	0.71	1.44	0.964	1.48	0.94	2.31	0.088
Living with vulnerable	1.17	0.79	1.74	0.437	0.95	0.63	1.44	0.809
Access garden/balcony to exercise comfortably in	1.02	0.74	1.41	0.891	0.84	0.60	1.17	0.301
Access green space within walking distance	1.07	0.79	1.43	0.667	0.87	0.64	1.18	0.370
Smoker	0.63	0.43	0.91	0.013	1.04	0.74	1.46	0.828
Weekly alcohol drinking, none (ref)								
Weekly or less	1.11	0.74	1.65	0.620	1.13	0.73	1.76	0.580
More than weekly	1.26	0.86	1.85	0.237	1.06	0.71	1.59	0.786
Deteriorated psychological wellbeing	1.27	0.97	1.66	0.082	0.62	0.46	0.83	0.002

Notes: All models were also adjusted for the time of enrolment and time when Covid-19 started to affect individuals in any way. Following the BH correction the highest p-value that met the threshold for significance for was $p=0.019$.