

# **Appendix**

## *Appendix 1 – Search Strategy for MEDLINE and EMBASE*

1. exp Malnutrition/
2. exp Protein-Energy Malnutrition/
3. exp Child Nutrition Disorders/
4. exp Wasting Syndrome/
5. "moderate acute malnutrition".mp.
6. (moderate\* adj2 malnutrition).mp.
7. (moderate\* adj2 malnourish\*).mp.
8. (moderate\* adj2 undernutrition).mp.
9. (moderate\* adj2 wast\*).mp.
10. "mid\* upper arm circumference".mp.
11. (mid\* upper arm circumference adj2 "125").mp.
12. "MUAC".mp.
13. 1 OR 2 OR 3 OR 4 OR 5 OR 6 OR 7 OR 8 OR 9 OR 10 OR 11 OR 12
14. exp Infant/
15. exp Child/
16. exp Child, Preschool/
17. 14 OR 15 OR 16
18. exp Dietary Supplements/
19. exp Food, Fortified/
20. exp Food, Formulated/
21. ((fortif\* or enrich\* or formulat\* or supplement\* or lipid\* or blend\*) adj2 (food\* or spread\* or diet\* or feed\* or product\*)).mp.
22. "ready to use food\*".mp.
23. "ready to use supplementary food".mp.
24. "RUSF".mp.
25. "lipid-based nutrient supplement".mp.
26. (lipid-based adj2 supplement\*).mp.
27. "LNS".mp.
28. Nutributter.mp.
29. "ready-to-use therapeutic food".mp.

30. "RUTF".mp.
31. (therapeutic adj2 food).mp.
32. plumpy\*.mp.
33. "FBF".mp.
34. "fortified blended flour".mp.
35. super-cereal\*.mp.
36. "corn soy\* blend\*".mp.
37. "wheat soy\* blend\*".mp.
38. "rice milk blend\*".mp.
39. "milk rice blend\*".mp.
40. "pea wheat blend\*".mp.
41. "cereal pulse blend\*".mp.
42. "CSB".mp.
43. "CSB+".mp.
44. 18 OR 19 OR 20 OR 21 OR 22 OR 23 OR 24 OR 25 OR 26 OR 27 OR 28 OR 29 OR 30 OR 31 OR 32 OR 33  
OR 34 OR 35 OR 36 OR 37 OR 38 OR 39 OR 40 OR 41 OR 42 OR 43
45. exp Education/
46. exp Health Education/
47. exp Counseling/
48. "standard care".mp.
49. "basic care".mp.
50. "control".mp.
51. (education adj2 nutrition).mp.
52. (counseling adj2 nutrition).mp.
53. ((local or home) adj2 (food\* or feed\* or diet\*)).mp.
54. 45 OR 46 OR 47 OR 48 OR 49 OR 50 OR 51 OR 52 OR 53
55. 13 AND 17 AND 44 AND 54
56. Limit 55 to (English language and humans and yr="2006-Current")

*Appendix 2 – Reasons for exclusion following screening of full-text*

<b>Study</b>	<b>Reason for Exclusion</b>
<b>Aboud 2008</b>	Enrolled children did not have MAM
<b>Aburto 2010</b>	Investigated effect of micronutrient supplementation on physical activity
<b>Ackatia-Armah 2012</b>	Full text not available
<b>Addo 2020</b>	Investigated improvements in gross motor and communication scores
<b>Ahmed 2019</b>	Study protocol for ongoing trial
<b>Akhter 2011</b>	Preventative management of undernutrition
<b>Amagloh 2012</b>	Investigated composition of sweet potato based complementary food
<b>Annan 2014</b>	Technical brief
<b>Arimond 2015</b>	Preventative management of undernutrition
<b>Ashton 2011</b>	Preventative supplementation
<b>Ashworth 2009</b>	Investigated the likelihood of effect of nutrition education programmes
<b>Batra 2016</b>	Enrolled children did not have MAM
<b>Bazzano 2017</b>	Literature review
<b>Borg 2017</b>	Enrolled children did not have MAM
<b>Borg 2019</b>	Lessons learned when testing a novel supplementary food
<b>Borg 2020</b>	Preventative supplementation
<b>Briend 2011</b>	Population based mathematical model
<b>Brown 2009</b>	Literature review
<b>Chanani 2019</b>	No comparator treatment group.
<b>Chang 2013</b>	Follow-up study post treatment of MAM
<b>Choudhury 2016</b>	Defined undernutrition using weight-for-age
<b>Cohuet 2012</b>	Investigated acceptability and intra-household use of supplements
<b>Daniel 2019</b>	Investigated dietician impact in hospital setting
<b>Das 2019</b>	Literature review
<b>Das 2020</b>	Systematic review

<b>Delchevalerie 2015</b>	Defined MAM using Department for Health and Human Statistics
<b>Desai 2014</b>	Retrospective cohort study
<b>Dido 2011</b>	Full text not available
<b>Egata 2014</b>	Investigated predictors of acute undernutrition
<b>Fatima 2018</b>	Enrolled mildly malnourished children (WHZ>-2 to <-1)
<b>Fernald 2016</b>	Investigated impact of supplementation on stunting
<b>Fitriyanto 2019</b>	No comparator treatment group.
<b>Flax 2013</b>	Investigated feeding behaviours with supplementation
<b>Flax 2015</b>	Investigated dietary intake during supplementation
<b>Friis 2015</b>	Literature review
<b>Gaffey 2013</b>	Investigated management of moderate diarrhoea
<b>Gigante 2007</b>	Enrolled school-aged children
<b>Hossain 2012</b>	Conference proceeding - full text unavailable
<b>Huybregts 2012</b>	Preventative management of undernutrition. Undernutrition defined using NCHS references.
<b>Huybregts 2017</b>	Preventative management of undernutrition
<b>Iannotti 2015</b>	Enrolled children aged 3-13 years.
<b>Ickes 2015</b>	Diet adequacy assessment
<b>Ikawati 2019</b>	Defined undernutrition using weight-for-age
<b>Inayati 2012</b>	Enrolled children did not have MAM - had mild wasting (defined as a WHZ>-1.5 to <-1)
<b>Isanaka 2010</b>	Preventative supplementation
<b>luel-Brockdorf 2017</b>	Investigated home behaviours during supplementation
<b>James 2016</b>	Prospective cohort study
<b>Juliana 2017</b>	Investigated zinc deficiency
<b>Kajjura 2019</b>	Investigated hygiene practice and knowledge of mothers following nutrition education
<b>Karakochuk 2015</b>	Defined MAM using NCHS references
<b>Kekalih 2019</b>	Study protocol for ongoing trial
<b>Khan 2013</b>	No comparator treatment group. Enrolled children aged 6 months to 8 years
<b>Khatib 2010</b>	Descriptive study of a vulnerable group
<b>Kimani-Murage 2013</b>	Full text not available - enrolment of well nourished mother-child pairs

<b>Kulwa 2014</b>	Study protocol for ongoing trial
<b>Langendorg 2014</b>	Preventative supplementation
<b>Lanou 2019</b>	Enrolled children did not have MAM
<b>Lazzerini 2013</b>	Systematic review
<b>Lelijveld 2020</b>	Follow-up study 4 months post treatment of MAM
<b>Linneman 2007</b>	No comparator treatment group.
<b>Magnin 2017</b>	No comparator treatment group.
<b>Maheswari 2012</b>	Used weight to define undernutrition
<b>Manary 2020</b>	Literature review
<b>Mangani 2014</b>	Investigated morbidity in well nourished children
<b>Marron 2015</b>	Full text not available
<b>Matondo 2016</b>	Defined MAM using NCHS references
<b>Maust 2015</b>	Combined protocol for MAM and SAM treatment. Results published do not differentiate between MAM and SAM
<b>Moramarco 2016</b>	Retrospective observational study
<b>Moramarco 2018</b>	Retrospective observational study
<b>Mostafa 2020</b>	Proof-of-concept study - full text not available - study not complete
<b>Nackers 2010</b>	Defined MAM using NCHS references
<b>Nane 2019</b>	Study protocol for ongoing trial
<b>Olney 2017</b>	Preventative supplementation - primary outcome assessed anaemia
<b>Pereira 2020</b>	Investigated glucose levels and Hb with supplementation
<b>Perez-Exposito 2009</b>	Literature review
<b>Phuka 2009</b>	Measured mean anthropometric changes. Primary outcome was reduction in stunting.
<b>Phuka 2011</b>	Investigated acceptability of LNS in Malawi - amount consumed, time to eat and maternal rating
<b>Prado 2016</b>	Enrolled children did not have MAM
<b>Puett 2013</b>	Cost-effectiveness analysis
<b>Pulakka 2017</b>	Enrolled children did not have MAM - measured impact of supplementation on physical activity
<b>Ragini 2014</b>	Primary outcome focused on mothers' continuation of nutrition education practices
<b>Rogers 2017</b>	Behaviour change and cost-effectiveness analysis

<b>Ruel 2008</b>	Compared preventative supplementation with recuperative
<b>Sachdeva 2014</b>	Severe acute malnutrition
<b>Saleem 2014</b>	Enrolled children did not have MAM
<b>Sayyad-Neerkorn 2015</b>	Preventative supplementation
<b>Scherbaum 2015</b>	Recovery defined as WHZ>-1.5
<b>Schlossman 2015</b>	Defined undernutrition using weight-for-age and height-for-age
<b>Schlossman 2017</b>	Defined undernutrition using weight-for-age and height-for-age
<b>Schlossman 2018</b>	Enrolled school-aged children
<b>Segre 2017</b>	Investigated economic impacts of local vs offshore therapeutic supplements
<b>Sguassero 2012</b>	Preventative supplementation
<b>Shen 2017</b>	Cost-effectiveness analysis
<b>Siega-Riz 2014</b>	Supplementation effect on micronutrient status
<b>Somasse 2013</b>	Follow-up study post treatment of MAM
<b>Somasse 2016</b>	Management of severe acute malnutrition
<b>Steenkamp 2015</b>	No comparator treatment group.
<b>Stobaugh 2017</b>	Investigated children post recovery from MAM
<b>Stobaugh 2017</b>	Investigating sustaining recovery post treatment of MAM
<b>Stobaugh 2018</b>	Investigated factors associated with relapse post MAM treatment
<b>Tandon 2018</b>	Investigated effect of supplementation on neutropenia in cancer patients with acute malnutrition
<b>Taye 2016</b>	Descriptive study
<b>Thakwalakwa 2015</b>	Defined MAM using NCHS references
<b>Trehan 2015</b>	Investigated an extended course of treatment for MAM post recovery
<b>Van der Kam 2016</b>	Preventative supplementation
<b>Van Waardenburg 2009</b>	Investigated critically ill infants with bronchiolitis
<b>Verna 2012</b>	Conference proceeding - full text unavailable. Did not use WHO growth standards to define MAM
<b>Vray 2018</b>	Study protocol for ongoing trial
<b>Wang 2013</b>	Investigated acceptability and feeding practices in MAM treatment

<b>Wilner 2017</b>	Lessons learned / communication whilst implementing a supplementary feeding programme
<b>Zhang 2019</b>	Systematic review

*Appendix 3 – Nutrient compositions of interventions by study and including WHO Recommendations for children aged 6-59 months*

Study	Intervention	Composition				
		Dry mass of food (g)	Energy (kcal)	Energy Density (kcal/g)	Protein (g)	Fat (g)
<b>WHO Recommendations</b>	per 1000kcal of Supplement (70% of daily energy intake)	-	1000	0.8	20-43	25-65
<b>Matilsky 2009</b>	CSB	200	749	3.7	34	-
	Milk/peanut FS & Soy/peanut FS	136	749	5.5	19	-
<b>LaGrone 2012</b>	CSB+	143	563	3.9	21	13
	Soy RUSF	104	563	5.4	17	40
	Soy/whey RUSF	103	563	5.5	15	38
<b>Nikiema 2014</b>	CSB++	65	273	4.2	10	6
	RUSF	50	258	5.2	9	17
<b>Medoua 2015</b>	CSB+, oil	72	320	5	10	11
	RUSF	61	320	5.2	10	21
<b>Amegovu 2015</b>	CSB+	269	1,200	4	46	30
	SPB	269	1,228	5	40	48
<b>Ackatia-Armah 2015</b>	RUSF	92	500	5.4	13	33
	CSB++	127	501	3.9	18.4	12
	Misola	125	500	4	18.5	14
	LMF + MNP	129	500	3.9	15.5	14
<b>Stobaugh 2016</b>	Soy RUSF	105	560	5.3	17	37
	Whey RUSF	105	516	4.9	11	36
<b>Kajjura 2019</b>	MSBP	150	675	4.5	26	6.6
	CSB+	158	600	3.8	25	9.5
<b>Kohlmann 2019</b>	A-RUTF	100	560	5.6	14.5	29.2
	S-RUTF	100	559	5.6	15.8	33



<b><i>Azimi 2020</i></b>	RUSF	95	475	5	15	23
<b><i>Roediger 2020</i></b>	HiPro-RUSF	100	530	5.3	13.5	32.8
	C-RUSF	100	537	5.4	13.4	32.9
<b><i>Bailey 2020</i></b>	RUSF	-	500	-	-	-
	RUTF	-	500	-	-	-