

Supplemental Tables

Table e-1

Cognitive domain	Assessments
Memory	Three sub-scores derived from a 12-word five trial list-learning task: list learning total score, delayed recall score, and delayed recognition score
Language	Picture naming (modified Boston Naming) test, Category fluency (Animal Naming) test, Phonemic fluency (C, F, L in English speakers and P, S and V in Spanish speakers)
Processing Speed	Color Trails test Form 1 [¥] , Grooved Pegboard task (non-dominant hand) [¥]
Executive Function	Difference in time to complete the Color Trails test Form 1 [¥] and Form 2 [¥] , Sum of the Odd-Man-Out subtests 2 and 4

[¥] For the participants who did not complete the tasks within the maximum time allowed, “imputed” time was calculated by adding penalty time to maximum time allowed. The penalty time was obtained by multiplying the number of incomplete items by “expected time per correct items completed” estimated using Kaplan-Meier.

Table e-2: immune markers selected by Least Absolute Shrinkage and Selection Operator procedure as linearly associated with global cognition score and each domain score

Global Cognitive Score	Memory	Language	Processing Speed	Executive Function
CCL11	CCL11	CCL11	CCL11	CCL11
CXCL9	CXCL9	CXCL9	CXCL9	CXCL9
HGF	HGF	HGF	HGF	HGF
Serpin_E1	Serpin_E1	Serpin_E1	Serpin_E1	Serpin_E1
sFasl	sFasl	sFasl	sFasl	Leptin
SCF	CSF2	Leptin	CSF2	CCL3
Leptin	SCF	CCL3	SCF	CCL2
CCL3	CXCL12a	IL12p40	CCL2	PDGFBB
	IL21		IL1a	
	CCL4		IFNb	
	TGFa			

CCL11 = C-C Motif Chemokine Ligand 11; CXCL9 = C-X-C Motif Chemokine Ligand 9; HGF = hepatocyte growth factor; sFasL = soluble Fas ligand; SCF = stem cell factor; CSF2 = Colony stimulating factor 2; CCL2 = C-C Motif Chemokine Ligand 2; CCL3 = C-C Motif Chemokine Ligand 3; CCL4 = C-C Motif Chemokine 4; IL12p40 = interleukin-12 p40; IL21 = interleukin 21; IL1a = interleukin 1a; CCL4 = C-C Motif Chemokine Ligand 4; IFNb = interferon beta; PDGFBB = Platelet derived growth factor-BB; TGFa = transforming growth factor α ; TRAIL = Tumor necrosis factor-related apoptosis-inducing ligand

Table e-3. Association of immune markers and interactions with memory domain*

Immune marker [†]	Estimate	Standard Error	Pr > ChiSq
CXCL9	-0.047	0.03	0.115
HGF	-0.082	0.029	0.005
CCL11	0.005	0.025	0.844
SCF	-0.050	0.027	0.061
CSF2	0.057	0.029	0.051
Serpin_E1	0.057	0.025	0.021
IL21	-0.011	0.074	0.880
sFasL	0.029	0.026	0.262
CCL4	0.126	0.066	0.055
TGFa	0.019	0.028	0.504
TGFa*CXCL10	-0.011	0.045	0.814
CCL11*CCL5	0.031	0.023	0.179
CCL2*IL12p40	0.007	0.018	0.693
CXCL10*TNFa	0.015	0.025	0.547
sFasL*PDGFBB	-0.042	0.021	0.043
TRAIL*VCAM1	-0.062	0.026	0.015
CXCL9*CXCL10	-0.026	0.018	0.144
HGF*Leptin	-0.035	0.024	0.145
CXCL12a	-0.041	0.070	0.556
CXCL10*IFNb	0.033	0.044	0.456
IL12p40*TRAIL	0.039	0.022	0.071

*Adjusted for age, sex, race-ethnicity, education, medical insurance type, diabetes, smoking status, systolic blood pressure, diastolic blood pressure, high-density lipoprotein, low-density lipoprotein, estimated glomerular filtration rate

[†] Immune markers were log 2 transformed and standardized

CCL11 = C-C Motif Chemokine Ligand 11; CXCL9 = C-X-C Motif Chemokine Ligand 9; HGF = hepatocyte growth factor; sFasL = soluble Fas ligand; SCF = stem cell factor; CSF2 = Colony stimulating factor 2; CCL2 = C-C Motif Chemokine Ligand 2; CCL3 = C-C Motif Chemokine Ligand 3; CCL4 = C-C Motif Chemokine 4; IL12p40 = interleukin-12 p40; IL21 = interleukin 21; IL1a = interleukin 1a; CCL4 = C-C Motif Chemokine Ligand 4; IFNb = interferon beta; PDGFBB = Platelet derived growth factor-BB; TGFa = transforming growth factor α ; TRAIL = Tumor necrosis factor-related apoptosis-inducing ligand

Supplement Table e-4. Association of immune markers and interactions with language domain*

Immune marker [†]	Estimate	Standard Error	p
IL12p40	-0.066	0.029	0.024
HGF	-0.045	0.029	0.126
CCL11	-0.038	0.021	0.073
Serpin_E1	0.012	0.026	0.641
sFasL	0.055	0.022	0.012
VCAM1	0.044	0.027	0.100
Leptin	0.046	0.024	0.058
CXCL9	-0.015	0.024	0.546
CCL3	0.020	0.023	0.398
IL7*PDGFBB	-0.016	0.024	0.490
CXCL9*CXCL10	-0.009	0.016	0.566
PDGFBB*IL13	-0.013	0.025	0.598
Serpin_E1*CCL5	-0.022	0.018	0.233
IL12p40*TRAIL	0.027	0.017	0.114
CCL3*TNFb	-0.003	0.012	0.834
CXCL10*CCL5	0.033	0.020	0.102

*Adjusted for age, sex, race-ethnicity, education, medical insurance type, diabetes, smoking status, systolic blood pressure, diastolic blood pressure, high-density lipoprotein, low-density lipoprotein, estimated glomerular filtration rate

[†] Immune markers were log 2 transformed and standardized

CCL11 = C-C Motif Chemokine Ligand 11; CXCL9 = C-X-C Motif Chemokine Ligand 9; HGF = hepatocyte growth factor; sFasL = soluble Fas ligand; SCF = stem cell factor; CSF2 = Colony stimulating factor 2; CCL2 = C-C Motif Chemokine Ligand 2; CCL3 = C-C Motif Chemokine Ligand 3; CCL4 = C-C Motif Chemokine 4; IL12p40 = interleukin-12 p40; IL21 = interleukin 21; IL1a = interleukin 1a; CCL4 = C-C Motif Chemokine Ligand 4; IFN β = interferon beta; PDGFBB = Platelet derived growth factor-BB; TGF α = transforming growth factor α ; TRAIL = Tumor necrosis factor-related apoptosis-inducing ligand

Supplement Table e-5. Association of immune markers and interactions with processing speed*

Immune marker[†]	Estimate	Standard Error	p
CXCL9	-0.034	0.032	0.297
CCL11	-0.045	0.026	0.086
HGF	-0.071	0.031	0.023
CSF2	0.010	0.031	0.739
CCL2	0.000	0.026	0.999
SCF	-0.007	0.029	0.804
IL1a	0.011	0.033	0.740
IFN β	0.046	0.029	0.119
sFasL	0.019	0.028	0.492
Serpin_E1	0.033	0.024	0.181
HGF*Leptin	-0.060	0.027	0.028
CCL11*IL1RA	-0.049	0.017	0.005
Serpin_E1*CCL5	-0.022	0.022	0.313
Leptin*VEGFD	-0.047	0.026	0.066
Leptin*CCL4	-0.032	0.031	0.289
CXCL9*IL10	-0.030	0.023	0.190
CSF2*IL17F	0.092	0.063	0.142
VEGFD*CXCL10	0.056	0.024	0.021
CCL11*sICAM1	0.039	0.028	0.157
Leptin*CXCL10	0.053	0.050	0.294
HGF*CXCL12a	0.028	0.020	0.159
SCF*BDNF	-0.002	0.026	0.931
CCL2*BDNF	0.022	0.023	0.352
IFN β *PDGFBB	0.017	0.025	0.500
IL10*BDNF	0.004	0.036	0.908
Serpin_E1*IL31	0.054	0.025	0.028
IL10*PDGFBB	0.026	0.031	0.404
IL1RA*BDNF	0.012	0.029	0.669
CCL11*CCL5	0.023	0.026	0.372
CCL3*TNF β	-0.012	0.016	0.458
IFN β *CXCL10	-0.011	0.028	0.685
CSF2*CXCL10	-0.027	0.054	0.620
CSF2*IFN β	-0.045	0.058	0.438

*Adjusted for age, sex, race-ethnicity, education, medical insurance type, diabetes, smoking status, systolic blood pressure, diastolic blood pressure, high-density lipoprotein, low-density lipoprotein, estimated glomerular filtration rate[†] Immune markers were log 2 transformed and standardized

CCL11 = C-C Motif Chemokine Ligand 11; CXCL9 = C-X-C Motif Chemokine Ligand 9;
HGF = hepatocyte growth factor; sFasL = soluble Fas ligand; SCF = stem cell factor;
CSF2 = Colony stimulating factor 2; CCL2 = C-C Motif Chemokine Ligand 2;
CCL3 = C-C Motif Chemokine Ligand 3; CCL4 = C-C Motif Chemokine 4;

IL12p40 = interleukin-12 p40; IL21 = interleukin 21; IL1a = interleukin 1a; CCL4 = C-C Motif Chemokine Ligand 4; IFN β = interferon beta; PDGFBB = Platelet derived growth factor-BB; TGF α = transforming growth factor α ; TRAIL = Tumor necrosis factor-related apoptosis-inducing ligand

Supplement Table e-6. Association of immune markers with executive function domain*

Immune marker [†]	Estimate	Standard Error	p
CCL11	-0.056	0.023	0.014
Leptin	-0.005	0.026	0.859
HGF	-0.075	0.026	0.004
CXCL9	0.026	0.026	0.319
CCL2	-0.011	0.022	0.619
CCL3	0.050	0.025	0.047
Serpine1	0.030	0.022	0.172
PDGFBB	0.058	0.021	0.006
CCL11*SCF	-0.039	0.021	0.057
Serpine1*CCL5	-0.038	0.018	0.037
CCL4*VCAM1	-0.050	0.030	0.099
CCL5*CCL4	0.012	0.028	0.656
Leptin*CXCL12a	-0.016	0.028	0.562
Leptin*IFN α 2	-0.027	0.023	0.248
sFasL*IL21	0.001	0.015	0.924
IL21*IL7	0.017	0.019	0.376
CCL4*VEGFA	-0.016	0.010	0.098
VEGFA*IL13	-0.040	0.017	0.023
CCL11*IL1RA	-0.013	0.016	0.425
PDGFBB*BDNF	-0.023	0.020	0.250
IL5*CXCL10	-0.065	0.087	0.458
PDGFBB*VEGFD	0.043	0.022	0.048
CCL3*TNF β	-0.001	0.018	0.950
CXCL9*PDGFBB	0.040	0.029	0.055
CCL3*sICAM1	0.052	0.025	0.039
Serpine1*IL17F	0.025	0.020	0.215
sICAM1*TNF α	0.008	0.024	0.730
CXCL10*IL22	-0.001	0.044	0.974
CXCL10*IL2	0.078	0.081	0.336
CCL3*IL5	0.042	0.022	0.056

*Adjusted for age, sex, race-ethnicity, education, medical insurance type, diabetes, smoking status, systolic blood pressure, diastolic blood pressure, high-density lipoprotein, low-density lipoprotein, estimated glomerular filtration rate

[†] Immune markers were log 2 transformed and standardized

CCL11 = C-C Motif Chemokine Ligand 11; CXCL9 = C-X-C Motif Chemokine Ligand 9; HGF = hepatocyte growth factor; sFasL = soluble Fas ligand; SCF = stem cell factor; CSF2 = Colony stimulating factor 2; CCL2 = C-C Motif Chemokine Ligand 2; CCL3 = C-C Motif Chemokine Ligand 3; CCL4 = C-C Motif Chemokine 4; IL12p40 = interleukin-12 p40; IL21 = interleukin 21; IL1a = interleukin 1a; CCL4 = C-C Motif Chemokine Ligand 4; IFN β = interferon beta; PDGFBB = Platelet derived growth factor-BB;

TGF α = transforming growth factor α ; TRAIL = Tumor necrosis factor-related apoptosis-inducing ligand