Original Article

Changes in healthcare workers' knowledge, attitudes, practices, and stress during the COVID-19 pandemic

Running title: Changing KAP among HCWs during COVID-19

Mohamad-Hani Temsah, MD, FRCPCH^{1,2,3*}, Abdullah Al Huzaimi, MD^{1,4*}, Abdulkarim Alrabiaah, MD^{1,2}, Nurah Alamro, MD, DrPH^{1,5,6}, Fahad Al-Sohime, MD^{1,2}, Ayman Al-Eyadhy, MD^{1,2}, Khalid Alhasan, MD^{1,2}, Jameela A Kari, MD, FRCPCH⁷, Ali Alhaboob, MD^{1,2}, Amro Al Salmi, MD², Wejdan AlMuhanna, MD², Ibrahim Almaghlouth, MD, MSc, FRCPCH^{1,8}, Fadi Aljamaan, MD^{1,9}, Rabih Halwani, PhD¹⁰, Mazin Barry, MD, FRCPCH, FACP, DTM&H^{1,11}, Fahad Al-Zamil, MD^{1,2}, Ahmad Al-Hadi, MD^{1, 12, 13}, Sarah Al-Subaie, MD^{1,2}, Amr Jamal, MD^{1,5,14#}, Ali Mohammed Somily, MD, FRCPCH^{1,15}

¹*College of Medicine, King Saud University, Riyadh, Saudi Arabia*

²*Department of Pediatrics, King Saud University Medical City, Riyadh, Saudi Arabia*

³*Prince Abdullah Ben Khaled Coeliac Disease Chair, Faculty of Medicine, King Saud University, Riyadh, Saudi Arabia*

⁴*Cardiac Science Department, King Saud University Medical City, Riyadh, Saudi Arabia*

⁵*Department of Family and Community Medicine, King Saud University Medical City, Riyadh, Saudi Arabia*

⁶*Prince Sattam bin Abdulaziz Research Chair for Epidemiology and Public Health, King Saud University, Riyadh, Saudi Arabia*

⁷*Paediatric Nephrology Centre of excellence, Department of Paediatrics, King Abdulaziz University, Jeddah, Saudi Arabia*

⁸*College of Medicine Research Center, King Saud University, Riyadh, Saudi Arabia*

⁹*Adult Critical Care Department, King Saud University, King Saud University Medical City/King Khalid University Hospital, Riyadh, Saudi Arabia*

¹⁰*Sharjah Institute of Medical Research, College of Medicine, University of Sharjah, Sharjah, United Arab Emirates*

¹¹*Infectious Disease Unit, Department of Internal Medicine, King Saud University, Riyadh, Saudi Arabia*

¹²*Department of Psychiatry, College of Medicine, King Saud University and King Saud University Medical City, Riyadh, Saudi Arabia*

¹³*SABIC Psychological Health Research and Applications Chair, College of Medicine, King Saud University, Riyadh, Saudi Arabia*

¹⁴*Evidence-Based Health Care & Knowledge Translation Research Chair, King Saud University, Riyadh, Saudi Arabia*

¹⁵*Department of Pathology and Laboratory Medicine, College of Medicine, King Saud University and King Saud University Medical City, Riyadh, Saudi Arabia*

*These authors contributed equally to this work.

#Corresponding author:

Amr Jamal

College of Medicine, King Saud University, King Saud University Medical City

PO Box 2925, Riyadh 11461, Saudi Arabia

Email: amrjamal@ksu.edu.sa

Tel: +96611-467-0836, fax: +96611-467-1967

Abstract

Introduction: Coronavirus disease 2019 (COVID-19) has caused an unprecedented health crisis around the world, not least because of its heterogeneous clinical presentation and course. The new information on the pandemic emerging daily has made it challenging for healthcare workers (HCWs) to stay current with the latest knowledge, which could influence their attitudes and practices during patient care.

Methods: This study is a follow-up evaluation of changes in HCWs' knowledge, attitudes, and practices as well as anxiety levels regarding COVID-19 since the beginning of the pandemic. Data were collected through an anonymous, predesigned, self-administered questionnaire that was sent online to HCWs in Saudi Arabia.

Results: The questionnaire was sent to 1500 HCWs, with a 63.8% response rate (N=957). The majority of respondents were female (83%), and the most common age group was 31–40 years (52.2%). Nurses constituted 86.3% of the respondents. HCWs reported higher anxiety during the COVID-19 pandemic which increased from 4.91 ± 2.84 to 8.6 ± 2.27 on an 11-point Likert scale compared to other viral outbreaks. HCWs believed that their own preparedness as well as that of their hospital's intensive care unit (ICU) or emergency room (ER) was higher during the COVID-19 pandemic than during the Middle East respiratory syndrome coronavirus pandemic (2012–2015). About 58% of HCWs attended one or more simulations concerning the management of COVID-19 patients in their ICU/ER, and nearly all had undergone N95 mask fit testing. The mean score of HCWs' knowledge of COVID-19 was 9.89/12. For most respondents (94.6%), the perception of being at increased risk of infection was the main cause of anxiety related to COVID-19; the mean score of anxiety over COVID-19 increased from 4.91 ± 2.84 before to 8.6 ± 2.27 during the pandemic in Saudi Arabia.

Conclusions: HCWs' anxiety levels regarding COVID-19 have increased since a pandemic was declared. It is vital that healthcare facilities provide more emotional and psychological support for all HCWs.

Keywords: COVID-19, changing KAP, healthcare worker.

Introduction

Coronavirus disease 2019 (COVID-19) has caused an unprecedented health crisis around the world and has surprised healthcare workers (HCWs) because of its wide dynamic and heterogeneous clinical presentation, from asymptomatic to critical illness leading to hospitalization and death.^[1] The rapid evolution of available literature and new information on the pandemic emerging daily has made it challenging for HCWs to stay current with the latest knowledge, which could influence their attitudes and practices during patient care. Additionally, basic principles of infection prevention and control (IPC) are more widely recognized and implemented by HCWs in Saudi Arabia, including hand hygiene, personal protective equipment (PPE) compliance, and N95 fit testing.^[2] The previous experience of Middle East respiratory syndrome coronavirus (MERS-CoV) outbreaks at several hospitals^[3-9] has enhanced IPC knowledge and practices among HCW, and commissioning bodies such as the Saudi Central Board for Accreditation of Healthcare Institutions have updated their definitions and standards for coronaviruses and acute respiratory syndromes to include COVID-19.^[10] As such, hospitals in Saudi Arabia were quick to adapt to and prepare for the pandemic before the first cases appeared in the country.^[11]

In the present study, we assessed the anxiety levels and knowledge base of HCWs in a tertiary care hospital in Saudi Arabia during the pandemic as compared to its very beginning in order to determine the degree of preparedness of the healthcare system to not only to manage patients but also to meet the needs of HCWs, and identify potential deficiencies that can be improved by focused education and training.

Methods

Data collection

This was the second study for serial cross-sectional surveys among HCWs in Saudi Arabia during the COVID-19 pandemic. The survey was a pilot-validated, self-administered questionnaire sent online to HCWs that was adapted from our previously published study.^[12] We used multiple professional social media groups and email lists from our previous surveys during the MERS-CoV outbreak in 2014 and the early phase of the COVID-19 pandemic in early 2020.

Data were collected over 2 weeks (April 13–27, 2020). The first part of the survey addressed respondents' characteristics (sex, age, etc.) and sources of information used during the pandemic.

We assessed the following domains for each participant: (1) knowledge concerning COVID-19; (2) attitudes toward IPC measures; and (3) changes in hygiene practices. Knowledge on COVID-19 was assessed using 6 questions consisting of true or false answers, with a score of 0–12 points assigned (Supplementary Table S1). The degree of change in HCWs' attitudes was measured using a series of questions on a Likert-like scale from 1–5 (Supplementary Table S2). Hygiene practices were similarly measured to assess the degree of behavioral change in HCWs on scale of 1–4 (Supplementary Table S3).

HCW anxiety and preparedness with respect to COVID-19 were evaluated based on self-reported perceived anxiety over COVID-19 vs MERS-CoV vs seasonal influenza (11-point Likert scale), along with the level of worry over COVID-19 infection in the previous 2 weeks. HCWs' perceptions of their own preparedness as well as that of their hospital with respect to the COVID-19 pandemic vs the previous MERS-CoV outbreak were compared using an 11-point Likert scale. Additionally, the availability of a psychological support system, HCWs' adherence to seasonal influenza vaccine, and participation in COVID-19 simulation training, N95 mask fit testing and sources of anxiety for each HCWs were evaluated. Data collected from HCWs through the online survey were entered into a secure spreadsheet in SPSS (SPSS Inc, Chicago, IL, USA). All data was treated with strict confidentiality, with the anonymity of respondents maintained throughout the study.

Statistical analysis

Descriptive statistics approaches with mean, median, and standard deviation were applied to continuous variables, while percentages were used for dichotomous variables. The 2-sample t test was used to evaluate continuous scores and the Z-test was used to compare proportions. The multiple response dichotomy analysis was applied to describe the healthcare workers sources of covid19 Information's.

A multivariate logistic regression model was used to explore the associations between the outcome variable the level of anxiety from COVID-19 and demographic characteristics of HCWs, attitude and hygiene practice scores, and sources of anxiety, the associations between predictors and the outcome was expressed as Odds Ratio and 95% Confidence Interval. The SPSS IBM Version21 was used for the data analysis, the excel program was used for creating figures and depictions, the p-value statistical significance was considered at 0.050 level. The Institutional Review Board of King Saud University Medical City approved the study. HCWs gave informed consent prior to participation.

Results

Of the 1500 HCWs who were contacted, 1453 agreed to participate in the study. Participants with information missing on more than 50% of the questionnaire were excluded.

Demographic and sources of information data

Data from the questionnaire were completed by 957 (63.8%) HCWs and were included in the final analysis. Most respondents were employed at public governmental tertiary hospitals in Riyadh, Saudi Arabia; the majority were female (83%) and between 31–40 years of age (52.2%). Nurses constituted 86.3% of the study population. Most respondents worked in general hospital wards (26.5%), followed by outpatient clinics (18.8%) and adult emergency rooms (ERs) (14.7%) (Table 1). HCWs obtained information on COVID-19 from multiple sources (Figure 1).

HCW anxiety and preparedness with respect to COVID-19 during the pandemic

HCWs' self-reported anxiety levels regarding COVID-19, MERS-CoV, and seasonal influenza were compared (Figure 2). The highest levels of anxiety were experienced during the COVID-19 pandemic compared to other viral outbreaks. HCWs believed that their own preparedness and those of their hospital's ER or intensive care unit (ICU) was higher during the COVID-19 pandemic than during the 2012–2015 MERS-CoV outbreaks (Table 2); however, 88% felt more worried about the former.

About 97% of respondents had undergone N95 mask fit testing in 2020 or in the previous year, but only 58% had attended at least one simulation session related to managing COVID-19 patients in their hospital's ICU/ER. About one-quarter of HCWs worked at hospitals where no psychological support system was available during the COVID-19 pandemic.

Knowledge, attitudes, and practices (KAP) of HCWs with respect to COVID-19

The mean score for HCWs' knowledge pertaining to COVID-19 was 9.89 out of 12. HCWs' attitude toward infection control measures were measured on a 5-point Likert-like scale, with a mean score of 4.42 ± 1.1 (Table 3), which indicated that most participants agreed or strongly agreed with level of importance of implementing their hospital's IPC measures (Supplementary Table S2).

The mean hygiene practice score was 3.8 ± 0.32 , representing a moderate-to-large change in self-rated compliance with hygiene practices and behaviors (e.g., hand hygiene) (Supplementary Table S3).

Sources of anxiety among HCWs

Sources of perceived anxiety related to COVID-19 in HCWs are shown in Figure 3. Most HCWs (94.6%) were concerned about the risk of acquiring COVID-19 during the pandemic, while more than half of the group was worried about the depletion of PPE at their hospital in general or in specifically high-risk departments such as the ICU or ER. On the other hand, the HCWs described several coping interventions to handle the stress during the pandemic, as demonstrated in the "word cloud" representation in Figure 4.

Factors associated with HCWs' perceived anxiety with respect to COVID-19

Factors contributing to HCWs' perceived anxiety over COVID-19 during the pandemic were analyzed using a multivariate logistic regression model (Table 4). The model was significant overall ($f[17,939]=5.98$, $P<0.001$).

Neither sex nor age was correlated with perceived anxiety over COVID-19. However, the clinical role and work location of the HCW showed positive associations

with perceived anxiety level during the pandemic: resident physicians and nurses had lower mean perceived anxiety than HCWs in other roles (e.g. consultant or assistant consultant) (both $P < 0.010$). Additionally, HCWs at general hospitals (including in COVID-19 wards) and pediatric ER had higher mean perceived anxiety than their colleagues in other departments ($P = 0.018$ and 0.016 , respectively).

Hospital setting and sector were not correlated with mean perceived anxiety of HCWs during the COVID-19 pandemic; however, HCWs at hospitals equipped with a psychological support system dedicated to staff during the pandemic reported a lower mean perceived anxiety than those working in hospitals without such a system ($P = 0.001$).

HCWs who believed that the medical staff were at higher risk of contracting COVID-19 and for whom the possibility of shortage of medical supplies and disposable materials (e.g., ventilator circuits) were a source of anxiety had a significantly higher mean perceived anxiety level than other HCWs ($P = 0.015$ and 0.010 , respectively), although potential shortages in PPEs or equipment such as ventilators were not a significant source of anxiety. In addition, the fear of staff shortage during the pandemic did not significantly affect HCWs' mean perceived anxiety.

HCWs' improvement in hygiene practice score was positively and significantly correlated with mean perceived anxiety over COVID-19 ($P < 0.001$); additionally, their attitude toward hygiene practices was associated with a significantly higher mean anxiety ($P = 0.037$).

Additional Bivariate analysis were done and revealed a significantly higher mean anxiety level over COVID-19 in HCWs at hospitals without support systems than in those at hospitals where such systems were available (9.02 ± 1.88 vs 8.46 ± 2.37 , $P < 0.001$) consistent with multivariate linear regression analysis. We also found that HCWs who used social networks as a source of information had a higher mean anxiety over COVID-19 than those who did not use social media (8.72 ± 2.23 vs 8.39 ± 2.33 , $P = 0.032$) (Table S4).

HCW anxiety and KAP before vs during the COVID-19 pandemic

We compared the results of the present study (Phase II), which was carried out during the active COVID-19 pandemic in Saudi Arabia (April 13–27, 2020), with those of our previous study (Phase I; February 5–16, 2020), which was conducted by the same research team immediately before the first case of COVID-19 was reported in Saudi Arabia.

HCWs' mean scores of perceived anxiety over COVID-19 increased from 4.91 ± 2.84 to 8.6 ± 2.27 on an 11-point Likert scale (Table 5). The degree of anxiety over other viral outbreaks, including MERS-CoV and seasonal influenza, similarly increased, as did concern over contracting COVID-19.

The Knowledge of COVID-19 in the second survey exceeded that measured in February 2020 ($t=33.32$, $df=1537$, $p<0.0001$ (2-sample t test), which is expected based on the elapsed time and exposure of the respondents to COVID-19 within that time frame. Likewise, scores for HCWs' hygiene practices and attitudes toward hygiene increased significantly in the current survey compared to those measured in February ($P<0.001$).

The proportion of HCWs who received the annual influenza vaccine this year increased significantly during the pandemic as compared to before ($Z=-6.4183$, $P<0.0001$). The proportion of HCWs who intended to reschedule their annual leave also increased from 15.3% in February to 45.7% in April 2020 ($Z=-12.1816$, $P<0.001$).

Discussion

The participants were predominantly female (83%) and most were nurses (86.3%), which is comparable to the demographic profile of the previous studies conducted in early 2020 prior to the COVID-19 outbreak in Saudi Arabia.^[13–16] Most respondents were between the ages of 31 and 40 years; some studies have shown that medical staff aged 31–40 years were more worried about becoming infected with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2, the causative agent of COVID-19) and infecting their families compared to other age groups.^[13, 15] A large proportion of the study participants were from acute care areas (42.42%), which is similar to other published work.

The current COVID-19 pandemic has been accompanied by an overabundance of information that has made it difficult to obtain accurate facts as most individuals rely on social media for this purpose;^[6] this contributes to higher levels of anxiety.^[17] Classic forms of media (i.e., television and newspapers) as well as social media affect the attitudes of both the public and health professionals, while consulting reliable sources of information is a basic condition for IPC. Compared to before the COVID 19 pandemic, hospital announcements were the top source of information used by HCWs during the pandemic (77.8% vs 86.7%), as was the case for the MERS-CoV outbreak.^[12] This finding highlights the importance of having a dedicated team to provide accurate information from trusted sources. News on social networks was a source of information for 61.7% of HCWs before and 66.3% during the pandemic, which may have contributed to greater anxiety in our study participants as bivariate analysis showed a higher mean anxiety over COVID-19 found that HCWs who used social networks as a source of information, although the use of unofficial and nonscientific sources of data was discouraged in awareness campaigns conducted by hospitals. Use of the Ministry of Health and World Health Organization websites as a source of information by HCWs increased from 42.3% to 71.6% and from 50% to 70.8%, respectively, between the 2 surveys.

Healthcare simulation tests have been useful for managing the COVID-19 pandemic. However, there are also practical constraints on running in situ simulations during a pandemic, such as the need for physical distancing, rigorous sanitization measures for mannequins and training equipment, and heightened anxiety among simulation participants; indeed, the fear of acquiring the infection could explain why 41.7% did not attend the simulation tests.^[18] In contrast, 97.65% of HCWs underwent N95 mask fit testing; this high percentage compared to the simulation test may be explained by the fact that the former was performed individually and therefore elicited less anxiety in HCWs in terms of the risk of acquiring COVID-19, and that such testing has been mandated for all HCW.

Surveys are commonly used to identify knowledge gaps and behavioral patterns in order to implement effective measures for improvement of processes and practices. We observed that knowledge scores were higher among participants in Phase II (during

the COVID-19 pandemic) than in Phase I (pre-pandemic); the trend is similar to that observed during the previous MERS-CoV outbreak.^[12, 19] Most study participants had a high level of knowledge concerning SARS-CoV-2 infection and modes of transmission; this is expected given their profession as healthcare workers. However, previously published data from a KAP survey in the general population in China during the COVID-19 pandemic revealed a high rate of correct responses in the knowledge questionnaire, which the authors attributed to the high educational level of the participants and the intensified public health education programs.^[20] Other studies have reported variable findings regarding KAP among HCWs during Ebola and Zika virus outbreaks. For example, satisfactory knowledge of Ebola virus disease without corresponding good practices was reported among Nigerian HCWs.^[21] This increase in HCWs knowledge could improve perceptions and positive attitudes that can translate into good practices, aiding in the prevention and management of infectious diseases. In our previous phase 1 study, the knowledge score was significantly associated with positive attitude and practice scores: that is, HCWs with a high level of knowledge had more positive attitudes and perceptions toward preventive measures and engaged to a greater extent in IPC practices. Others have reported similar associations in KAP surveys of COVID-19 and other infectious diseases.^[22-24]

The HCWs in our study reported a high level of anxiety with respect to COVID-19, with 88% having more anxiety as compared to the previous MERS-CoV outbreak. Similarly, both local and international studies have found that anxiety levels were high among HCWs during both pandemics.^[25-27] A recent systematic review of 115 publications concluded that all coronavirus outbreaks (SARS, MERS-CoV, and COVID-19) had a substantial impact on the physical and mental health of HCWs.^[27]

The availability of hospital-based psychological support was associated with decreased levels of anxiety among HCWs in our survey. Psychological support is critical for the well-being of frontline HCWs.^[28] However, establishing psychological support services in Hospitals might not be enough to guarantee their efficacy, for instance in the case where there are obstacles to their access by HCWs who need them.^[29] One study proposed specific measures that healthcare managers should implement to protect the mental health of healthcare staff: managers must be honest about current and future

situations and have regular meetings to discuss protocols and HCWs' wellbeing; and after the crisis has passed, they must actively monitor, support, and—where necessary—provide treatment to HCWs.^[30] In accordance with this issue, the United Nations issued a policy report on the importance of HCWs' mental health, and encouraged all parties to facilitate HCWs' access to mental health services.^[31]

Risk of being infected with SARS-CoV-2 was the highest source of anxiety among HCWs in our study, followed by depletion of PPE at their hospital in general or more specifically in the ICU/ER departments. These findings are similar to those of a study of 69 HCWs who participated in multiple listening sessions. There were 7 sources of anxiety, ranging from fear of not having enough equipment and of contracting COVID-19 or transmitting it to their loved ones, to fear of not performing well when they were needed in areas beyond their expertise.^[32] Such anxiety could affect the confidence of HCWs in themselves as well as general trust in the healthcare system. In China, HCWs who were interviewed reported that they did not require psychological help but needed enough protective supplies and more uninterrupted rest time.^[29] Another study examining factors related to HCWs' psychological difficulties found that infection of colleagues and family members, protective measures, and medical violence were among the main concerns of HCWs in areas affected by COVID-19.^[33-34]

This study had some limitations. Recall bias may have affected some participants' responses, and most participants were nurses and females. Another issue is that many HCWs in our cohort had experienced the previous MERS-CoV epidemic; therefore, additional research in other settings are needed in order to determine whether our findings are generalizable.

Conclusion

HCWs' anxiety levels over COVID-19 increased after the pandemic was declared. Healthcare facilities need to provide more emotional and psychological support for all HCWs, including psychological first aid delivered through webinars to each unit on topics such as dealing with anxiety and insomnia, peer support, practicing self-care, and support for moral distress; as well as providing individual support sessions.

Ethics approval and consent to participate: Institutional Review Board, King Saud University, IRB # 20/0064/IRB

Consent for publication: Granted upon manuscript acceptance

Competing interests: The authors of this work have nothing to disclose.

Funding: The authors are grateful to the Deanship of Scientific Research, King Saud University for funding through Vice Deanship of Scientific Research Chairs.

References

- [1] Eckerle I, Meyer B. SARS-CoV-2 seroprevalence in COVID-19 hotspots. *Lancet*. 2020;396:10250:514–5.
- [2] Ministry of Health (MOH). Middle East Respiratory Syndrome Coronavirus; Guidelines for Healthcare Professionals - April 2018 - v 5.1. <https://www.moh.gov.sa/CCC/healthp/regulations/Documents/MERS-CoV%20Guidelines%20for%20Healthcare%20Professionals%20-%20May%202018%20-%20v5.1%20%281%29.pdf>. Accessed 31 Aug 2020.
- [3] Barry M, Phan MV, Akkielah L, et al. Nosocomial outbreak of the Middle East Respiratory Syndrome coronavirus: A phylogenetic, epidemiological, clinical and infection control analysis. *Travel Med Infect Dis*. 2020;26:101807.
- [4] Drosten C, Muth D, Corman VM, et al. An observational, laboratory-based study of outbreaks of Middle East respiratory syndrome coronavirus in Jeddah and Riyadh, kingdom of Saudi Arabia, 2014. *Clin Infect Dis*. 2015;60:369–77.
- [5] Assiri A, McGeer A, Perl TM, et al. Hospital outbreak of Middle East respiratory syndrome coronavirus. *N Engl J Med*. 2013;369:407–16.
- [6] Al-Omari A, Rabaan AA, Salih S, Al-Tawfiq JA, Memish ZA. MERS coronavirus outbreak: Implications for emerging viral infections. *Diagn Microbiol Infect Dis*. 2019;93:265–85.
- [7] Fagbo SF, Skakni L, Chu DK, et al. Molecular epidemiology of hospital outbreak of Middle East respiratory syndrome, Riyadh, Saudi Arabia, 2014. *Emerg Infect Dis*. 2015;21 Suppl 11:1981–8.
- [8] Oboho IK, Tomczyk SM, Al-Asmari AM, et al. 2014 MERS-CoV outbreak in Jeddah – a link to health care facilities. *N Engl J Med*. 2015;372:846–54.
- [9] Balkhy HH, Alenazi TH, Alshamrani MM, et al. Description of a hospital outbreak of Middle East respiratory syndrome in a large tertiary care hospital in Saudi Arabia. *Infect Control Hosp Epidemiol*. 2016;37:1147–55.
- [10] Saudi Central Board for Accreditation of Healthcare Institutions. National Standards for Hospitals. <https://portal.cbahi.gov.sa/english/cbahi-standards/NHS-3rd-Edition-2016>. Accessed 31 Aug 2020.

- [11] Barry M, Ghonem L, Alsharidi A, et al. Coronavirus disease-2019 pandemic in the Kingdom of Saudi Arabia: Mitigation measures and hospital preparedness. *JNSM*. 2020;3:155–8.
- [12] Alsubaie S, Hani Temsah M, Al-Eyadhy AA, et al. Middle East Respiratory Syndrome Coronavirus epidemic impact on healthcare workers' risk perceptions, work and personal lives. *J Infect Dev Ctries*. 2019;13:920–6.
- [13] Cai H, Tu B, Ma J, Chen L, Fu L, Jiang Y, et al. Psychological impact and coping strategies of frontline medical staff in Hunan between January and March 2020 during the outbreak of coronavirus disease 2019 (COVID-19) in Hubei, China. *Med Sci Monit*. 2020;26:e924171.
- [14] Kang L, Li Y, Hu S, et al. The mental health of medical workers in Wuhan, China dealing with the 2019 novel coronavirus. *Lancet Psychiatry*. 2020;7:e14.
- [15] Lai J, Ma S, Wang Y, et al. Factors associated with mental health outcomes among health care workers exposed to coronavirus disease 2019. *JAMA Netw Open*. 2020;3:e203976.
- [16] Mohindra R, Ravaki R, Suri V, Bhalla A, Singh SM. Issues relevant to mental health promotion in frontline health care providers managing quarantined/isolated COVID19 patients. *Asian J Psychiatr*. 2020;51:102084.
- [17] Barry M, Al Amri M, Memish ZA. COVID-19 in the shadows of MERS-CoV in the Kingdom of Saudi Arabia. *J Epidemiol Glob Health*. 2020;10:1–3.
- [18] Brazil V, Lowe B, Ryan L, et al. Translational simulation for rapid transformation of health services, using the example of the COVID-19 pandemic preparation. *Adv Simul (Lond)*. 2020;5:9.
- [19] Bukhari EE, Temsah MH, Aleyadhy AA, et al. Middle East respiratory syndrome coronavirus (MERS-CoV) outbreak perceptions of risk and stress evaluation in nurses. *J Infect Dev Ctries*. 2016;10:845–50.
- [20] Zhong BL, Luo W, Li HM, et al. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: A quick online cross-sectional survey. *Int J Biol Sci*. 2020;16:1745–52.

- [21] Oladimeji AM, Gidado S, Nguku P, et al. Ebola virus disease - gaps in knowledge and practice among healthcare workers in Lagos, August 2014. *Trop Med Int Health*. 2015;20:1162–70.
- [22] ul Haq N, Hassali MA, Shafie AA, Saleem F, Farooqui M, Aljadhey H. A cross sectional assessment of knowledge, attitude and practice towards Hepatitis B among healthy population of Quetta, Pakistan. *BMC Public Health*. 2012;12:692.
- [23] Huynh G, Nguyen T, Tran V, Vo K, Vo V, Pham L. Knowledge and attitude toward COVID-19 among healthcare workers at District 2 Hospital, Ho Chi Minh City. *Asian Pac J Trop Dis*. 2020;13:260–5.
- [24] Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK. Novel coronavirus (COVID-19) knowledge and perceptions: A survey of healthcare workers. *medRxiv*. 2020. doi: <https://doi.org/10.1101/2020.03.09.20033381>.
- [25] Alkhamees AA, Alrashed SA, Alzunaydi AA, Almohimeed AS, Aljohani MS. The psychological impact of COVID-19 pandemic on the general population of Saudi Arabia. *Compr Psychiatry*. 2020;102:152192.
- [26] Temsah MH, Al-Sohime F, Alamro N, et al. The psychological impact of COVID-19 pandemic on health care workers in a MERS-CoV endemic country. *J Infect Public Health*. 2020;13:877–82.
- [27] Salazar de Pablo G, Vaquerizo-Serrano J, Catalan A, et al. Impact of coronavirus syndromes on physical and mental health of health care workers: Systematic review and meta-analysis. *J Affect Disord*. 2020;275:48–57.
- [28] Liu YE, Zhai ZC, Han YH, Liu YL, Liu FP, Hu DY. Experiences of front-line nurses combating coronavirus disease-2019 in China: A qualitative analysis. *Public Health Nurs*. 2020. <https://doi.org/10.1111/phn.12768>.
- [29] Chen Q, Liang M, Li Y, et al. Mental health care for medical staff in China during the COVID-19 outbreak. *Lancet Psychiatry*. 2020;7:e15–6. [https://doi.org/10.1016/s2215-0366\(20\)30078-x](https://doi.org/10.1016/s2215-0366(20)30078-x).
- [30] Greenberg N, Docherty M, Gnanapragasam S, Wessely S. Managing mental health challenges faced by healthcare workers during covid-19 pandemic. *BMJ*. 2020;368:m1211.

- [31] Guterres A. Launch of the policy brief on COVID-19 and mental health 2020.
Available at: <https://www.un.org/en/coronavirus/mental-health-services-are-essential-part-all-government-responses-covid-19>. 31 August 2020.
- [32] Shanafelt T, Ripp J, Trockel M. Understanding and addressing sources of anxiety among health care professionals during the COVID-19 pandemic. JAMA. 2020 Apr 7. doi: 10.1001/jama.2020.5893.
- [33] Liu Q, Luo D, Haase JE, et al. The experiences of health-care providers during the COVID-19 crisis in China: A qualitative study. Lancet Glob Health. 2020;8:e790–8.
- [34] Dai Y, Hu G, Xiong H, Qiu H, Yuan X. Psychological impact of the coronavirus disease 2019 (COVID-19) outbreak on healthcare workers in China. medRxiv. 2020. doi: <https://doi.org/10.1101/2020.03.03.20030874>.

Table 1. Sociodemographic and professional characteristics of healthcare workers
(N=957)

	Frequency	Percentage
Sex		
Male	163	17
Female	794	83
Age (years), mean (SD)		36.69 (7.79)
21–30 years	192	20.1
31–40 years	500	52.2
41–50 years	206	21.5
≥51 years	59	6.2
Clinical role		
Consultant	69	7.2
Assistant consultant	22	2.3
Registrar	13	1.4
Resident	27	2.8
Nurse	826	86.3
Department		
Adult ICU	139	14.5
Pediatric ICU	75	7.8
Adult ER	141	14.7
Pediatric ER	51	5.3
Isolation ward	34	3.6
General ward	254	26.5
OPD	180	18.8
Operation room	40	4.2
Other units (dialysis, radiology, other services)	43	4.5
Hospital sector		
Private	32	3.3
Public/governmental	925	96.7
Hospital setting		

Tertiary	850	88.8
Non-tertiary (secondary etc)	107	11.2
Hospital location		
Riyadh	907	94.8
Other city	50	5.2

ER, emergency room; ICU, intensive care unit; OPD, outpatient department; SD, standard deviation.

Table 2. Descriptive analysis of healthcare workers' perceived stress, anxiety, and preparedness practices during the COVID-19 pandemic

	Frequency (%) / Scale mean (SD)
Anxiety	
A) Rate the anxiety you have experienced during the following epidemic/pandemics: (11-point Likert scale)	
Current COVID-19 outbreak/pandemic	8.56 (2.27)
MERS-CoV	6.40 (2.35)
Seasonal flu	4.94 (2.65)
B) How worried were you in the past week about contracting COVID-19? (5-point Likert scale)	3.44 (1.10)
C) Compared to the MERS-CoV epidemic in 2012–2015, how do you feel about the recent COVID-19 (2019-NCOV) pandemic?	
Same level of stress during both pandemics	73 (7.6%)
I have felt more stressed during the COVID-19 (2019-NCOV) pandemic	843(88.1%)
I felt more stressed during the previous MERS-CoV epidemic	41(4.3%)
Preparedness	
Your preparedness for the COVID-19 pandemic (11-point Likert scale)	8.84 (2.03)
Your preparedness for the previous MERS-CoV outbreak (11-point Likert scale)	8.53 (2.35)
Your ICU/ER department's preparedness during the COVID-19 pandemic (11-point Likert scale)	9.10 (2.02)
Your ICU/ER department's preparedness during the MERS-CoV epidemic (11-point Likert scale)	8.83 (2.19)
Have you received the influenza vaccine (flu shot) this season?	
Yes, I receive it every year	0757 (79.1%)
Yes, only this year	50 (5.2%)

No	150(15.7%)
Did you undergo an N95 mask fit test?	
Yes, in 2020	471 (49.2%)
Yes, before this year	463 (48.4%)
Never	23 (2.4%)
Did you have a simulation about possible COVID-19 patients in your hospital's ICU/ER?	
Yes, once	252 (26.3%)
Yes, several simulation scenarios	306 (32%)
No	399 (41.7%)
Does your hospital have psychological support services to support HCWs during COVID-19?	
Yes	709 (74.1%)
No	248 (25.9%)

2019-NCoV, 2019 novel coronavirus; COVID-19, coronavirus disease 2019; ER, emergency room; HCW, healthcare worker; ICU, intensive care unit; MERS-CoV, Middle East respiratory syndrome coronavirus; SD, standard deviation.

Table 3. Descriptive statistics of HCWs' perceptions of COVID-19

	Mean (SD)
Knowledge of COVID-19 (0 to 12 points)	9.89 (1.98)
Attitudes toward infection control measures (level of agreement, 5-point Likert scale)	4.42 (1.1)
Hygiene practice improvement score (level of behavior change, 4-point Likert scale)	3.82 (0.32)

COVID-19, coronavirus disease 2019; SD, standard deviation.

Table 4. Multivariate linear regression analysis of healthcare workers' perceived anxiety caused by COVID-19 (N=956)

	Beta coefficient	95% CI beta coefficient		P value
		Lower Bound	Upper Bound	
(Constant)	4.144	1.640	6.648	0.001
Sex: female	-0.027	-0.502	0.447	0.910
Age (years)	-0.001	-0.020	0.018	0.899
Clinical role				
Resident physician	-1.382	-2.356	-0.407	0.006
Registered nurse	-1.031	-1.643	-0.420	0.001
Work location				
Pediatric emergency room worker	0.773	0.145	1.401	0.016
General wards worker	0.388	0.068	0.709	0.018
Hospital setting				
Public hospital	-0.018	-0.811	0.775	0.965
Tertiary hospital	0.287	-0.165	0.738	0.213
Psychological support services available at hospital: Yes	-0.568	-0.894	0-.242	0.001
Plans to reschedule annual leave from work: Yes	0.506	0.226	0.787	<0.001
Belief causing anxiety				
Increased risk of transmission among healthcare workers	0.734	0.143	1.325	0.015
Risk of depletion of hospital's personal protective equipment	0.216	-0.127	0.559	0.216
Lack of some types of equipment during pandemic (eg, ventilator shortages)	-0.061	-0.450	0.328	0.759

Depletion of disposable medical materials (eg, ventilator circuits, endotracheal tubes)	0.526	0.125	0.927	0.010
Shortage of staff and increased demand on all staff	-0.914	-2.001	0.174	0.099
Improvement in hygiene practices	0.895	0.445	1.346	<0.001
Attitudes toward infection control measures	0.136	0.008	0.264	0.037

SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

Table 5. Bivariate comparisons with 2-sample mean and proportion differences in healthcare workers' perceptions and behaviors before and during the COVID-19 pandemic

	Phase I* N=582	Phase II N=957	Test statistic	P value**
	Mean (SD)	Mean (SD)		
Anxiety over various viral outbreaks				
MERS-CoV (11-point Likert scale)	4.44 (2.48)	6.40 (2.35)	t=15.54, df=1537	<0.0001
Current COVID-19 pandemic (11-point Likert scale)	4.91 (2.84)	8.6 (2.27)	t=28.070, df=1537	<0.0001
Seasonal flu (11-point Likert scale)	3.22 (2.57)	4.94 (2.65)	t=12.50, df=1537	<0.0001
Worry about contracting COVID-19 (5-point Likert scale)	2.57 (1.071)	3.44(1.1)	t=15.20, df=1537	<0.0001
Knowledge of COVID-19 score	79.7 (13.7)	82.50 (16.54)	t=33.32, df=1537	<0.0001
HCW hygiene practice improvement score (4-point Likert scale)	2.82 (0.88)	3.82 (0.32)	t=31.90, df=1537	<0.0001
HCW attitudes toward importance of hygiene (5-point Likert scale)	4.10 (1.02)	4.42 (1.10)	t=5.69, df=1537	<0.0001
Received annual flu immunization/vaccination: Yes, n (%)	411 (70.6%)	807 (84.3%)	Z=-6.4183, 437	<0.0001
Leave rescheduling: Yes, n (%)	89 (15.3%)	437 (45.7%)	Z=-12.181, 6	<0.0001

*First survey was conducted before any cases were reported in Saudi Arabia **Two tailed.

COVID-19, coronavirus disease 2019; HCW, healthcare worker; MERS-CoV, Middle East respiratory syndrome coronavirus; SD, standard deviation.

Table S1. Healthcare workers' knowledge on COVID-19

	Incorrectly answered, n (%)	Correct answered, n (%)
COVID-19 can be transmitted by contacting patients' stools and then touching the nose or eyes (T)	46 (4.8)	911 (95.2)
COVID-19 can be transmitted through contact with packages shipped from China (F)	425 (44.4)	532 (55.6)
COVID-19 causes severe respiratory symptoms (T)	17 (1.8)	940 (98.2)
COVID-19 causes severe neurologic symptoms (F)	419 (43.8)	538 (56.2)
There is no vaccine against SARS-CoV-2 virus infection (T)	41 (4.3)	916 (95.7)
There is no specific treatment for COVID-19 (T)	57 (6)	900 (94)
COVID-19, coronavirus disease 2019; F, false; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; T, true		

Table S2. Healthcare workers' attitudes toward SARS-CoV-2 infection control measures

Attitude toward domestic hygienic practices	Level of importance*, mean (SD)
It is important to wear face masks at work to prevent SARS-CoV-2 infection	4.38 (1.22)
It is important to maintain good indoor ventilation to the spread of COVID-19	4.22 (1.23)
It is important to avoid going to crowded places during flu season	4.52 (1.17)
Hand hygiene is very important to protect against SARS-CoV-2 infection	4.57 (1.18)

*Likert scale score of 1 to 5 (1, strongly disagree; 2, disagree; 3, neutral; 4, agree; 5, strongly agree).

COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; SD, standard deviation

Table S3. Changes in healthcare workers' hygiene practices

Hygiene practice	Change in behavior*, mean (SD)
Compliance with hand hygiene at the hospital	3.82 (0.46)
Compliance with universal precautions (eg, masks and gloves)	3.81 (0.48)
Avoidance of contact with people showing flu-like symptoms	3.78 (0.51)
Avoidance of social visits (eg, visiting friends)	3.87 (0.41)
Avoidance of handshaking	3.88 (0.43)
Compliance with the N95 fit test at the hospital	3.75 (0.57)

*Scored on a scale of 1 to 4 (1, no change; 2, little change; 3, moderate change; 4, large change).

SD, standard deviation.

Table S4: Bivariate analysis of the HCW's current perceived Anxiety from COVID19 disease across the levels of their demographic & professional characteristics and perceptions.

		Mean (SD)- Worry		
		level from	test statistic	p-value
		COVID19		
Sex				
	Male	8.89 (1.99)	t(260.66)=1.95	0.052
	Female	8.55 (2.32)		
Age (years)-Groups				
	21-30 years	8.42 (2.17)	f(3,953)=0.81	0.486
	31-40 years	8.70 (2.23)		
	41-50 years	8.53 (2.47)		
	>=51 years	8.63 (2.20)		
Clinical Role				
	Consultant	9.10 (1.73)	f(4,46.12)=4.10	0.007
	Assistant Consultant	9.73 (1.64)		
	Registrar	8.69 (2.53)		
	Resident	8.00 (2.47)		
	Nurse	8.55 (2.30)		
Working department				
	Other (please specify)	8.56 (2.33)	f(8,224.80)=2.50	0.013
	Pediatric ICU	8.89 (2.54)		
	Adult ICU	8.56 (2.24)		
	Pedia ER	9.24 (1.66)		
	Adult ER	8.12 (2.27)		
	Isolation ward	8.68 (2.18)		
	General ward	8.87 (2.10)		
	OPD	8.38 (2.47)		
	Operative Room	8.38 (2.29)		
Working hospital sector				

Private	8.91 (2.10)	t(955)=0.77	0.444
Public/Governmental	8.59 (2.28)		
Hospital settings			
Non-Tertiary (Secondary...)	8.39 (2.47)	t(955)=1.02	0.307
Tertiary	8.63 (2.24)		
Hospital location			
Other cities	9.16 (1.75)	t(58.62)=2.26	0.027
Riyadh city	8.57 (2.29)		
Does your hospital have psychological support services to support your HCWs during COVID?			
Yes	8.46 (2.37)	t(539.50)=3.80	<0.001
No	9.02 (1.88)		
Will you consider rescheduling/changing your work time to avoid contact with COVID19 Patients?			
Yes	8.88 (2.13)	t(949.4)=3.50	0.001
No	8.38 (2.35)		
Have you taken the Influenza vaccine (Flu Shot) this season:			
Yes, I do that every year	8.54 (2.32)	f(2,954)=1.52	0.219
Yes, only this year	8.92 (2.10)		
No	8.83 (2.05)		
Do you think a dedicated psychological support system could help HCWs in the ER/ICUs during the pandemic?			
Strongly Disagree	9.20 (2.01)	f(4,952)=1.93	0.104
Disagree	7.36 (1.69)		
Neither agree nor disagree	8.74 (2.10)		

Agree	8.46 (2.22)		
Strongly agree	8.67 (2.35)		
Did you receive N95 mask fit test?			
Yes, in 2020	8.51 (2.39)	f(2,59.30)=0.90	0.422
Yes, before this year	8.70 (2.12)		
Never	8.43 (2.64)		
Use of various sources of information :			
Hospital announcements			
No	8.91 (2.10)	t(955)=0.30	0.799
Yes	8.59 (2.28)		
Official statements			
No	8.42 (2.28)	t(955)=1.23	0.219
Yes	8.65 (2.27)		
MOH website			
No	8.59 (2.21)	t(955)=0.10	0.922
Yes	8.61 (2.29)		
WHO website.			
No	8.62 (2.26)	t(955)=0.18	0.86
Yes	8.60 (2.27)		
CDC website			
No	8.60 (2.27)	t(955)=0.054	0.957
Yes	8.61 (2.27)		
Other Internet sources			
No	8.52 (2.29)	t(955)=0.20	0.234
Yes	8.69 (2.25)		
YouTube			
No	8.52 (2.24)	t(955)=1.34	0.179
Yes	8.72 (2.31)		
Social Networks			
No	8.39 (2.33)	t(955)=2.14	0.032

Yes

8.72 (2.23)

Figure legends

Figure 1. Sources of information regarding COVID-19 used by HCWs.

Figure 2. HCWs' self-rated mean anxiety related to various viral outbreaks (11-point Likert-like scale).

Figure 3. Sources of concern and anxiety related to COVID-19 among HCWs.

Figure 4. Word cloud representation of what best made healthcare workers cope with the stress during the pandemic (N = 957)

hygiene Pray god safe covid patients people PPEs stay safe protect COVID19
updating knowledge well cases Hand washing eating protocol handwashing knowledge
hygiene social distance will update covid given using ppe avoid wear PPE home staffs
exercise Team work precautions eat healthy talking family
Proper work sleep Family support Faith
information psychological support social distancing rest
Awareness prevent good pandemic updates
Proper information hospital wear mask Pray enough PPE
Prayer preventive measures Stay home MOH
follow stress family family friends hand hygiene help
support infection control proper Relax Reading Self PPE
online covid food always availability ppe Stay eat healthy foods knowing watch
time vigilant proper PPE keep social distance prevention management
proper hand hygiene Compliance providing health practice Talk Education
decrease watching movies compliance infection control take hand washing social keep infected
aware meditation patients virus