Genetic drift versus regional spreading dynamics of COVID-19

R. Di Pietro ¹, M. Basile ¹, L. Antolini ² and S. Alberti ³ *

¹ Department of Medicine and Aging Sciences, Section of Biomorphology, G. d'Annunzio University of Chieti-Pescara, Italy.

² Center for Biostatistics, Department of Clinical Medicine, Prevention and Biotechnology, University of Milano-Bicocca, Monza, Italy.

³ Unit of Medical Genetics, Department of Biomedical Sciences - BIOMORF, University of Messina, via Consolare Valeria, Messina, Italy.

* Address reprint requests to Prof. Alberti at the Unit of Medical Genetics, Department of Biomedical Sciences - BIOMORF, University of Messina, via Consolare Valeria, Messina, Italy, or at salberti@unime.it

Drs. Di Pietro and Basile contributed equally to this article.

Running title: COVID-19 transmission determinants

Word count of the abstract: 234

Word count of the text: 2,742

24 **Abstract**

25 **Background**

26 Current propagation models of COVID-19 are poorly consistent with existing epidemiological data
27 and with evidence that the SARS-CoV-2 genome is mutating, for potential aggressive evolution of
28 the disease.

29 **Methods**

30 We challenged regional versus genetic evolution models of COVID-19 at a whole-population level,
31 over 168,089 laboratory-confirmed SARS-CoV-2 infection cases in Italy, Spain and Scandinavia.
32 Diffusion data in Germany, France and UK provided a validation dataset of 210,239 additional cases.

33 **Results**

34 The mean doubling time of COVID-19 cases was 6.63 days in Northern versus 5.38 days in Southern
35 Italy. Spain extended this trend of faster diffusion in Southern Europe, with a doubling time of 4.2
36 days. Slower doubling times were observed in Sweden (9.4 days), Finland (10.8 days), Norway (12.95
37 days). COVID-19 doubling time in Germany (7.0 days), France (7.5 days) and UK (7.2 days) supported
38 the North/South gradient model. Clusters of SARS-CoV-2 mutations upon sequential diffusion across
39 distinct geographic areas were not found to clearly correlate with regional distribution dynamics.

40 **Conclusions**

41 Acquisition of mutations, upon SARS-CoV-2 spreading across distinct geographic areas, did not
42 distinctly associate to enhanced virus aggressiveness, and failed to explain regional diffusion
43 heterogeneity at early phases of the pandemic. Our findings indicate that COVID-19 transmission
44 rates associate to a sharp North/South climate gradient, with faster spreading in Southern regions.
45 Thus, warmer climate conditions may not limit SARS-CoV-2 infectivity. Very cold regions may be
46 better spared by recurrent courses of SARS-CoV-2 infection.

47
48
49 **Keywords:** COVID-19, pandemic, spreading dynamics, mutation rates.

Introduction

Studies on early dynamics of COVID-19 [1] revealed that the epidemic doubled in size every 6.4 [2] to 7.4 [1] days, with a reproductive number (R_0) of infectious cases from 2.2 [1] to 2.7 [2]. Later investigations followed disease spreading to Singapore [3], Germany [4], France and Finland (www.ecdc.europa.eu/en/covid-19-pandemic) [5-7]. However, major uncertainties remained on SARS-CoV-2 transmission dynamics [7]. Considerable effort across major research institutions was invested into modelling SARS-CoV-2 spreading determinants. Models were generated, which took into account, among others, global traveling, population density, demographic characteristics, age distribution, social dynamics, governmental policies, air pollution, virus infectious capacity, SARS-CoV-2 containment procedures, together with economical and healthcare factors [7-12] (10.21203/rs.3.rs-82122/v1). However, limited, if any, regional heterogeneity in COVID-19 transmission could be identified using such diffusion models. We reasoned that fundamental variables were missing from current analyses, and went on for identifying such missing factor(s).

SARS-CoV-2 was suggested to be sensitive to temperature and humidity, which may affect diffusion across diverse climate areas [13] (papers.ssrn.com/sol3/papers.cfm?abstract_id=3550308; ssrn.com/abstract=3556998; www.medrxiv.org/content/10.1101/2020.02.22.20025791v1). Accordingly, initial climate-dependent propagation models predicted a limited impact of COVID-19 in the Southern hemisphere, during seasons that were infection-prone in the Northern hemisphere (papers.ssrn.com/sol3/papers.cfm?abstract_id=3550308; ssrn.com/abstract=3556998). However, early foci of infection were detected in Australia and New Zealand (Figure 1). Outbreaks were also revealed in South America and extended to Central America and Mexico. Further infection foci were revealed in Saudi Arabia and Africa, and extended to sub-Saharan countries (Tables S1, S2), questioning simple models of climate-dependent SARS-CoV-2 transmission.

Viral evolution processes [14] may mimic regional COVID-19 spreading dynamics [15-18]. SARS-CoV-2 possesses a single-strand RNA genome [19], prone to acquire genomic mutations (nextstrain.org/ncov/; www.gisaid.org/). However, the SARS-CoV-2 RNA polymerase has error-correcting capacity, and shows replication error rates >10-fold lower than other RNA viruses [15]. Correspondingly, SARS-CoV-2 sequence diversity is low [17]. Spike proteins, in particular, show very few mutations overall [17]. Never the less, SARS-CoV-2 bearing distinct sets of mutations were isolated in different regions of the world (nextstrain.org/ncov/global), leaving the question open on whether viral genetic drift may drive evolution toward increased aggressiveness.

We thus went on to challenge regional versus genetic evolution models of COVID-19 at a population-wide level. Best chances for detecting basic transmission determinants of SARS-CoV-2 were expected before any large-scale defensive approach was implemented. Corresponding strategies

were adopted to determine the impact of UV light exposure on SARS-CoV-2 diffusion [20]. Western Europe provided a vast terrain for this approach, because of the large-scale outbreaks of COVID-19 early-on in the course of the pandemic. Further advantage was provided by Europe's high healthcare management and data collection standards (Bloomberg Global Health Index, 2018, www.bloomberg.com/; WHO, www.who.int/whr/en/; worldpopulationreview.com/countries/best-healthcare-in-the-world/; [21]), which supported a robust detection of basic diffusion parameters of COVID-19.

Broadly diverse climate regions around the CET longitude (15°E), were severely exposed to COVID-19. Spain and Italy were the countries with the highest early incidence of COVID-19 in Europe (Figures 1, S1, Table S3). The heaviest initial casualties in Italy were suffered by Lombardy and Veneto, i.e. cold and humid areas during wintertime. Markedly warmer and drier climate conditions prevail in Southern regions of the country. A further shift toward warmer/drier conditions occurs in Spain. Scandinavian countries provided a reference for cold winter temperatures, over a Sweden-Finland-Norway axis (Table S4). This offered unique opportunities to assess a climate-dependent coronavirus infection model. Such analysis was performed at a whole-population level on 86,498 laboratory-confirmed SARS-CoV-2 infection cases in Italy, 64,095 in Spain, and 17,496 cases in Scandinavia (github.com/pcm-dpc/COVID-19) (Supplemental Appendix). Diffusion data in France (Table S5), Germany (Table S6), and UK (Table S7) provided a validation dataset, encompassing 210,239 COVID-19 cases. This model was then merged with that of coronavirus genetic evolution (Figure 2), for detecting signs of positive selection for increased aggressiveness across the analyzed regions.

Methods

Incidence data

Data on laboratory-confirmed SARS-CoV-2 infection cases in Europe were collected at early time-points of the pandemic/peak diffusion rates from the following sources: Italy (github.com/pcm-dpc/COVID-19), France (dashboard.covid19.data.gouv.fr/vue-d-ensemble?location=FRA), UK (www.nhs.uk/), Germany (corona.rki.de), Spain (RTVE - Ministry of Health; www.rtve.es/noticias/20200415/mapa-del-coronavirus-espana/2004681.shtml), Sweden (Public Health Agency of Sweden; www.folkhalsomyndigheten.se/smittskydd-beredskap/utbrott/aktuella-utbrott/covid-19), Finland (National Institute for Health and Welfare THL; thl.fi/en/web/thlfi-en), Norway (Norwegian Institute of Public Health; www.fhi.no/sv/smittsomme-sykdommer/corona/dags--og-ukerapporter/dags--og-ukerapporter-om-koronavirus).

As current transmission models did not lead to predict regional heterogeneity [7-12] (10.21203/rs.3.rs-82122/v1), we set our search to identify additional variables that were not taken into account by pre-existing models.

Key considerations of our strategy were:

- I. An explosive diffusion of SARS-CoV-2 in Western Europe occurred early along the course of the pandemic, providing a vast number of infection cases, over largely overlapping calendar time-frames.
- II. High-quality disease-reporting procedures allowed whole-population-level analyses, with inclusion of 378,328 laboratory-confirmed SARS-CoV-2 infection cases across continental Europe and UK.
- III. The North-South span of the European regions involved in the early phases of COVID-19, provided vast diversity of climatic zones. The null hypothesis we challenged was that COVID-19 transmission velocity would not have been different across climate areas, quantitatively categorized as an independent variable.
- IV. As viral evolution processes in specific geographic areas [14] can effectively mimic climate region-associated spreading [15-18], these processes were analyzed accordingly.
- V. A potent confounding factor in disease transmission analyses is the founder effect, i.e. the date of first moving of infectious cases to a geographic site. To prevent this bias, the doubling number of COVID-19 cases, rather than their absolute number, was considered [20].
- VI. A further confounding factor was that of large-scale virus diffusion containment measures. Hence, we directed our search toward the initial period of explosive diffusion of the virus, and ended our observations at the time of first reduction of infectious case incidence rates, upon implementation of containment measures, in a region-by-region manner [20].

Disease severity was then classified as (a) hospitalized cases, (b) intensive-care unit patients, (c) recovered cases, (d) deaths. These findings were presented as cumulative incidence by region.

The cumulative incidence of COVID-19 cases was then linked to Köppen–Geiger climate classification maps (koeppen-geiger.vu-wien.ac.at/present.htm). These were computed as mean parametrization of data collected between 1980 and 2016 [22]. The tripartite classification by country areas was compounded as an independent variable versus COVID-19 doubling time (Table 1).

SARS-CoV-2 mutation analysis

SARS-CoV-2 genomic RNA sequences and country-correlated data were obtained from nextstrain.org/ncov/global. Each data-point was represented as a bead, whereby each bead

corresponded to a specific set of virus mutations (mutation haplotype) (Figures 2, S2-9). ‘Beads-on-a-string’ plots were then generated, that represented linked series of individual mutation haplotypes, that acquired subsequent mutations over time. Phylogeny trees for such mutation clusters were then obtained, for drawing distinct evolutionary branches of SARS-CoV-2 (nextstrain.org/ncov/europe?branchLabel=aa) (Figures 2, S2-9).

Statistical analysis

The cumulative incidence of COVID-19 cases versus calendar dates was acquired at province/county level [23, 24]. Corresponding plots acted as a smoother, for accurate determination of infection curve parameters. They also served to average urban versus countryside population dynamics/events on a province-by-province basis. Disease cumulative incidence graphs were found to largely follow a peculiar linear growth pattern [25]. This allowed to rigorously apply linear regression methodology for determining case-incidence rates.

At subsequent time points, deviations from linearity, with flattening of disease incidence curves, were recorded, following implementation of country-wide restrictions in traveling and social interactions (www.gazzettaufficiale.it/eli/gu/2020/03/08/59/sg/pdf). These inflection points were taken as landmark dates, and marked the end of the observation period. From each of these dates, the doubling time for cumulative number of diagnoses was calculated backward for each province as follows. Two dates were identified: the maximum date, at which the cumulative number of diagnoses were lower than a half of the cumulative number of diagnoses at the landmark time, and the minimum date, with a cumulative number of diagnoses greater than a half of the cumulative number of diagnoses at the landmark date. The fraction of days from the minimum date to achieve half of the cumulative number of diagnoses at the landmark date were obtained by a linear assumption for the cumulative incidence between the two dates. Correspondingly, distinct calendar dates were applied to data-collection in different provinces, regions and countries, according to the spreading sequence of the pandemic. Of note, each of these estimates corresponded to the fastest spreading velocity of COVID-19 in each region.

Coefficients, standard error, 95% confidence intervals were computed. Percentile distribution boxplots of COVID-19 cases doubling times were drawn. Median, maximum value, minimum value and distribution outliers were estimated. The correlation between COVID-19 spreading rates versus normalized climate-area values was computed by Anova.

Software

184 Stata software version 16 was used for data importing, manipulation and graphics (StataCorp.
185 2019. *Stata Statistical Software: Release 16*. College Station, TX: StataCorp LLC).

187 **Results**

188 **COVID-19 case doubling-time by geographic area**

189 Infection transmission rates were computed for:

190 Italy: on COVID-19 cases from March 3rd to March 27th (n=86,498) ([Supplemental Appendix](#))
191 ([Figures S10-12](#)).

192 Spain: on COVID-19 cases from February 25th to March 27th 2020 (n=64,095) ([Figure S13](#)).

193 Norway: on data (>50 infection case outbreaks) obtained from February 21st to April 14th 2020
194 (n=6,676) ([Figure S14](#)).

195 Finland: on COVID-19 cases from March 1st to April 7th 2020 (n=2,646) ([Figure S15](#)).

196 Sweden: on data (>50 infection case outbreaks) obtained from February 26th to April 9th 2020
197 (n=8,995) ([Figure S16](#)).

198 France: on COVID-19 cases from February 25th to April 4th 2020 ([Figure S17](#)).

199 UK: on COVID-19 cases from February 1st to April 9th 2020 ([Figure S17](#)).

200 Germany: on COVID-19 cases from February 24th to April 2nd 2020 ([Figure S17](#)).

202 **COVID-19 doubling time versus climate region**

203 Quantitative climate assessments are affected by complex, interdependent sets of variables [26]. Up
204 to 89 distinct parameters are required for meteorological classification alone
205 (apps.ecmwf.int/datasets/data/interim-full-moda/levtype=sfc/). Discrete humidity measures,
206 temperature profiles, (papers.ssrn.com/sol3/papers.cfm?abstract_id=3556998) [13, 27] and weather
207 structure, intertwine with lifestyle, social and occupational determinants [27]
208 (www.medrxiv.org/content/10.1101/2020.03.23.20040501v4). Hence, fundamental sources of
209 uncertainty are associated to climate modeling [11]. We thus resorted to utilizing the Köppen–Geiger
210 climate classification (koeppen-geiger.vu-wien.ac.at/present.htm), as drawn over 30+ years of
211 observations, and robustly validated [22, 26, 28, 29]. This was summarized as a tripartite
212 classification of country/region/province, which was compounded as an independent variable versus
213 COVID-19 spreading velocity ([Table 1](#)).

214 Cumulative numbers of COVID-19 cases versus calendar dates were normalized to the highest
215 case incidence in each area ([Figures S10-17](#)). Pandemic doubling times were correspondingly
216 computed ([Table S3](#)) [20] and grouped by geographic region. The average doubling time for Northern
217 Italy was 6.63 (SD=1.94) days; 5.87 (SD=1.08) days in Central regions; 5.38 (SD=2.31) days in

218 Southern areas, for significantly shorter doubling time in Southern regions ($P=0.02$ versus Northern
219 Italy) (Table S3, Figures 3, S10-12). The mean COVID-19 doubling-time for the whole country was
220 6.06 (SD=1.95) days.

221 With a doubling time of 4.2-days, Spain extended such a trend (Figure S13). At the opposite
222 end of the climate spectrum, Scandinavia showed longer COVID-19 doubling times, over a Sweden-
223 Finland-Norway axis, with a doubling time of 9.4 days (SD=1.2) for Sweden ($P<0.0001$ versus
224 Northern Italy), 10.8 days for Finland, 12.95 days (SD=0.52) for Norway ($P<0.0001$ versus Northern
225 Italy) (Figures 3, S14-16). This depicted a distinct North-South gradient of COVID-19 spreading
226 velocity (Anova $P<0.0001$) (Figure 4, Table 1).

227 This climate model was challenged versus a validation data-set of 210,239 laboratory-
228 confirmed COVID-19 cases in Germany, France and UK. Average climate areas for all three
229 countries fell in between classification classes of Northern Italy and Southern Sweden. Pandemic
230 doubling time was computed to be 7.0 days in Germany (Figure S17). In sharp consistency,
231 corresponding doubling times in France and UK were 7.5 and 7.2 days, respectively. This fell in
232 between Northern Italy and Sweden data, as predicted by our model.

233 Disease severity as classified hospitalization, intensive-care unit and fatality rates were
234 compounded as cumulative incidence by region (Figure S12). However, analysis of neither disease
235 onset severity nor outcome provided correlation with parameters of regional diffusion heterogeneity.
236 It should be noted that data on recoveries and deaths are not consistently classified across all the
237 regions under study, and are considered less reliable than those of confirmed COVID-19 cases [20].
238

239 SARS-CoV-2 genetic-drift driven diffusion

240 Sequence mutation analysis revealed different branches of acquired mutations, i.e. distinct groups of
241 viral genome mutations (haplotypes), at sites of major diffusion in Europe
242 (nextstrain.org/ncov/europe) (Figures 2, S2-9). Each of these branches was observed to acquire
243 additional mutations over time, in an uneven manner among different geographic areas. We searched
244 this data-set for potential indicators of positive selection for specific virus mutation(s). One virus
245 mutation, i.e. the spike D614G amino acid change, was associated with increased COVID-19
246 aggressiveness [17]. The D614G mutation appeared very early in Europe (inferred date, Jan. 6, 2020)
247 (nextstrain.org/ncov/; www.gisaid.org/) [17] and spread evenly across European countries.

248 Among descendants of D614G viruses, we looked for evidence of positive selection over
249 additional mutations. It should be noted that most viral mutations may not have phenotypic effects,
250 most of them being probably neutral or near neutral. Further, while some mutations may become
251 dominant over time, the overall diversity of SARS-CoV-2 genomes will continue to increase due to

genetic drift. Never the less, if positive selection for one or more virus mutation had been at work, deviation from even distributions of virus descendants across regions had to be expected. Among flags of such unevenness, we looked for (a) mutation-correlated increase of disease severity over time (b) prevalence of such mutation(s) in hardest-hit countries, (c) progressively broader diffusion of more aggressive virus genotypes along the course of the pandemic.

The highest numbers of accumulated mutations were revealed in SARS-CoV-2 in Wales and Senegal isolates (Figure 2B). Hence, they most likely represented late correlates of viral genetic drift over time. Of interest, the lowest number of accumulated mutations was recorded in Italy, a country with high disease severity in Europe. This appeared poorly consistent with a progressive increase of disease severity upon accumulation of novel mutations, suggesting instead correlation with a short SARS-CoV-2 evolution time. Large mutation loads were observed in Spain (n=14), the second hardest-hit country in Europe. However, a similar mutation load was observed in Sweden (n=13), a country with much more limited COVID-19 transmission and severity, further supporting correlation with genetic drift. Consistent, large mutation loads were observed in late-disease-insurgence countries, such as France and Belgium (n=16), supporting a slow SARS-CoV-2 genomic evolution, along the course of the disease (Figures S3-9).

Discussion

Large efforts have gone into modelling COVID-19 transmission, according to global and local population dynamics, demographics, governmental policies and infectious ability of the virus [7-12] (10.21203/rs.3.rs-82122/v1). Most models, though, showed inadequate capacity for predicting regional diffusion dynamics of the pandemic [11].

We speculated that fundamental variables associated to COVID-19 uneven diffusion remained to be identified, and set a search for discovering such factor(s). We went on to perform a population-wide analysis, on 378,328 laboratory-confirmed SARS-CoV-2 infection cases in continental Europe and UK. A robust determination required collecting epidemiological data [23, 24], before intervention with disease-containment measures. We thus went on to identify landmark dates, as inflection points of disease incidence curves associated to disease-taming procedures. This led us to first identify a quasi-universal pattern of linear growth of COVID-19 cases over time, consistent with a unique diffusion mode of SARS-CoV-2 [25]. Distinct COVID-19 transmission rates were then identified as associated to different geographic regions.

Still, accumulation of mutations of SARS-CoV-2 may have led to distinct selection for disease progression over different regions. An indicator of selective pressure for viral evolution has been that of progressively larger prevalence across different geographic locations [17], as indicated for the

spike D614G mutation. However, D614G was associated with higher upper respiratory tract viral loads [16-18], but not with increased disease severity. This appeared puzzling [15], as higher viral loads have been associated to worse disease course [30] and to increased mortality [31]. The hypothesis of positive selection of spike D614G was further investigated in the United Kingdom using more than 25,000 whole-genome SARS-CoV-2 sequences. This indicated 614G increases in frequency relative to 614D as consistent with selective advantage, but not in all cases [16]. This suggested that “a combination of evolutionary selection for G614 and the founder’s effects of being introduced into highly mobile populations may have together contributed in part to its rise” [17]. These findings were recapitulated “as a slow genetic drift .. of a highly stable [SARS-CoV-2] genome” [15, 32], during the analyzed time-frames.

Never the less, we looked for evidence of selection for viral evolution, utilizing broader indicators of predominance of specific mutation(s) versus disease severity. Four major mutation groups/haplotypes were revealed in all examined European countries. The highest number of accumulated mutations was revealed in Wales and Senegal SARS-CoV-2 isolates, suggesting correlation with genetic drift, at late stages of the disease. The lowest number of accumulated mutations was recorded in Italy, the country that first showed severe disease outbreaks in Europe, and similar mutation loads were observed in Spain, the second hardest-hit country in Europe, and Sweden, a country with much less explosive COVID-19 transmission, suggesting correlation with disease duration, rather than with selection for higher disease severity. The larger mutation loads were revealed in France and Belgium, both late-disease-insurgence countries, further supporting a relationship between mutation acquisition and length of disease course. Taken together, our findings add evidence to a model of SARS-CoV-2 genetic drifting during the early course of the pandemic, rather than to selective pressure for increased COVID-19 aggressiveness.

As COVID-19 spreading models based on population demographics and socio-economic factors could not account for regional diffusion heterogeneity, neither could an evolving infectious capacity of the virus, our findings supported a climate dependency of COVID-19 transmission capacity. This showed a sharp North-South gradient, with the shortest COVID-19 doubling times in Southern Italy and Spain. At the opposite end of the climate spectrum, Scandinavia showed the longest COVID-19 doubling times, over a Sweden-Finland-Norway axis. This climate model was verified in validation datasets of COVID-19 cases in Germany, France and UK, which included 210,239 laboratory-confirmed SARS-CoV-2 infection cases, and showed pandemic doubling times that were intermediate between Northern and Southern regions, in sharp consistency with the climate-area Köppen–Geiger model [22].

Findings of more efficient coronavirus spreading in warmer regions are consistent with resilience of coronaviruses to environmental conditions [33-35]. Of note, the Middle East Respiratory Syndrome (MERS) was first reported in Saudi Arabia (www.cdc.gov/coronavirus/mers). MERS is caused by the MERS-CoV, which is structurally and genetically related to SARS-CoV, indicating that at least some coronavirus strains may better propagate in high-temperature climate conditions (www.cdc.gov/coronavirus/mers/risk.html).

However, climate areas are associated to much more complex sets of variables, among them indoor versus outdoor temperature profiles, specific/relative/absolute humidity [13, 26, 27], UV exposure versus daily time/season/latitude [20, 27], weather structure and ventilation, together with social behavior, inter-individual distancing, indoor-crowding, lifestyle, outside physical activity, (papers.ssrn.com/sol3/papers.cfm?abstract_id=3556998) [27], versus community structure, socio-economical and healthcare factors (www.medrxiv.org/content/10.1101/2020.03.23.20040501v4). Recent studies have begun to dissect such determinants, indicating impact of UV exposure on COVID-19 transmission [20], while suggesting no role of temperature and humidity (SSRN 3567840, 2020 - papers.ssrn.com) [20, 29]. Our own research confirms these findings, and the lack of impact of temperature on COVID-19 progression over the areas analyzed (unpublished observations). Additional work is expected to bring-in further insight over climate area-associated virus diffusion determinants.

Taken together, our findings suggest higher SARS-CoV-2 resilience in warmer regions than previously predicted, and caution that high environmental temperatures may not efficiently tame SARS-CoV-2 infectiousness [12]. Cold regions may be better spared by recurrent courses of COVID-19.

Acknowledgments

We are much indebted to all the information curators we cite, and to the website providers, the article data and graphic primers have been downloaded from.

Role in the article

All authors contributed to literature search, figures, study design, data collection, data analysis, data interpretation. S.A and R.DiP. wrote the manuscript draft. All authors contributed to discussing and writing the final text. R.DiP. and M.B. contributed equally to this work.

351 **Footnote page**

352 Conflict of interest

353 The authors do not have a commercial or other association that might pose a conflict of interest.

354

355 Funding

356 The University of Messina provided funding to this work.

357

358 Meeting presentation

359 These findings have not yet been presented to meetings.

360

361 Corresponding author contact information

362 Prof. Saverio Alberti, Head, Medical Genetics, Department of Biomedical Sciences, University of

363 Messina, via Consolare Valeria, 98100 Messina, Italy, Phone: (+39) 090-221.3375, E-mail:

364 salberti@unime.it

365 **References**

- 366 1. Li, Q. et al. (2020) Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected
367 Pneumonia. *New England Journal of Medicine* 382 (13), 1199-1207.
- 368 2. Wu, J.T. et al. (2020) Nowcasting and forecasting the potential domestic and international spread
369 of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *The Lancet* 395 (10225),
370 689-697.
- 371 3. Pung, R. et al. (2020) Investigation of three clusters of COVID-19 in Singapore: implications for
372 surveillance and response measures. *The Lancet* 395 (10229), 1039-1046.
- 373 4. Hoehl, S. et al. (2020) Evidence of SARS-CoV-2 Infection in Returning Travelers from Wuhan,
374 China. *New England Journal of Medicine* 382 (13), 1278-1280.
- 375 5. Sun, J. et al. (2020) COVID-19: Epidemiology, Evolution, and Cross-Disciplinary Perspectives.
376 *Trends in Molecular Medicine*.
- 377 6. McMichael, T.M. et al. (2020) Epidemiology of Covid-19 in a Long-Term Care Facility in King
378 County, Washington. *New England Journal of Medicine*.
- 379 7. Lipsitch, M. et al. (2020) Defining the Epidemiology of Covid-19 — Studies Needed. *New England*
380 *Journal of Medicine* 382 (13), 1194-1196.
- 381 8. Pearson, C.A. et al. (2020) Projected early spread of COVID-19 in Africa through 1 June 2020.
382 *Euro surveillance : bulletin Europeen sur les maladies transmissibles = European communicable*
383 *disease bulletin* 25 (18), 2000543.
- 384 9. Hernandez-Vargas, E.A. and Velasco-Hernandez, J.X. (2020) In-host Mathematical Modelling of
385 COVID-19 in Humans. *Annual reviews in control*, 10.1016/j.arcontrol.2020.09.006.
- 386 10. Kennedy, D.M. et al. (2020) Modeling the effects of intervention strategies on COVID-19
387 transmission dynamics. *Journal of clinical virology : the official publication of the Pan American*
388 *Society for Clinical Virology* 128, 104440-104440.
- 389 11. Baker, R.E. et al. (2020) Susceptible supply limits the role of climate in the early SARS-CoV-2
390 pandemic. *Science*, eabc2535.
- 391 12. Kissler, S.M. et al. (2020) Projecting the transmission dynamics of SARS-CoV-2 through the
392 postpandemic period. *Science*, eabb5793.
- 393 13. Sundell, N. et al. (2016) A four year seasonal survey of the relationship between outdoor climate
394 and epidemiology of viral respiratory tract infections in a temperate climate. *Journal of Clinical*
395 *Virology* 84, 59-63.
- 396 14. Eletreby, R. et al. (2020) The effects of evolutionary adaptations on spreading processes in
397 complex networks. *Proceedings of the National Academy of Sciences of the United States of America*
398 117 (11), 5664-5670.

399 15. Rausch, J.W. et al. (2020) Low genetic diversity may be an Achilles heel of SARS-CoV-2.
400 Proceedings of the National Academy of Sciences 117 (40), 24614.

401 16. Volz, E. et al. (2021) Evaluating the Effects of SARS-CoV-2 Spike Mutation D614G on
402 Transmissibility and Pathogenicity. Cell 184 (1), 64-75.e11.

403 17. Korber, B. et al. (2020) Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases
404 Infectivity of the COVID-19 Virus. Cell 182 (4), 812-827.e19.

405 18. Li, Q. et al. (2020) The Impact of Mutations in SARS-CoV-2 Spike on Viral Infectivity and
406 Antigenicity. Cell 182 (5), 1284-1294 e9.

407 19. Mousavizadeh, L. and Ghasemi, S. (2020) Genotype and phenotype of COVID-19: Their roles in
408 pathogenesis. Journal of Microbiology, Immunology and Infection.

409 20. Carleton, T. et al. (2021) Global evidence for ultraviolet radiation decreasing COVID-19 growth
410 rates. Proceedings of the National Academy of Sciences 118 (1), e2012370118.

411 21. Lozano, R. et al. (2020) Measuring universal health coverage based on an index of effective
412 coverage of health services in 204 countries and territories, 1990–2019: a systematic analysis
413 for the Global Burden of Disease Study 2019. The Lancet 396 (10258), 1250-1284.

414 22. Beck, H.E. et al. (2018) Present and future Köppen-Geiger climate classification maps at 1-km
415 resolution. Scientific data 5, 180214-180214.

416 23. Ambroggi, F. et al. (2006) Molecular subtyping of breast cancer from traditional tumor marker
417 profiles using parallel clustering methods. Clin Cancer Res 12 (3 Pt 1), 781-90.

418 24. Cimoli, G. et al. (2004) Meta-analysis of the role of p53 status in isogenic systems tested for
419 sensitivity to cytotoxic antineoplastic drugs. Biochim Biophys Acta 1705 (2), 103-20.

420 25. Turner, S. et al. (2020) A network-based explanation of why most COVID-19 infection curves
421 are linear. Proceedings of the National Academy of Sciences 117 (37), 22684-22689.

422 26. Babin, S. (2020) Use of Weather Variables in SARS-CoV-2 Transmission Studies. International
423 journal of infectious diseases : IJID : official publication of the International Society for Infectious
424 Diseases 100, 333-336.

425 27. Sajadi, M.M. et al. (2020) Temperature, Humidity, and Latitude Analysis to Estimate Potential
426 Spread and Seasonality of Coronavirus Disease 2019 (COVID-19). JAMA network open 3 (6),
427 e2011834-e2011834.

428 28. Chakrabarti, S.S. et al. (2020) COVID-19 in India: Are Biological and Environmental Factors
429 Helping to Stem the Incidence and Severity? Aging and disease 11 (3), 480-488.

430 29. Briz-Redón, Á. and Serrano-Aroca, Á. (2020) A spatio-temporal analysis for exploring the effect
431 of temperature on COVID-19 early evolution in Spain. The Science of the total environment 728,
432 138811-138811.

433 30. Yu, F. et al. (2020) Quantitative Detection and Viral Load Analysis of SARS-CoV-2 in Infected
434 Patients. *Clinical Infectious Diseases*.

435 31. Westblade, L.F. et al. (2020) SARS-CoV-2 Viral Load Predicts Mortality in Patients with and
436 Without Cancer Who Are Hospitalized with COVID-19. *Cancer Cell*.

437 32. Dearlove, B. et al. (2020) A SARS-CoV-2 vaccine candidate would likely match all currently
438 circulating variants. *Proceedings of the National Academy of Sciences* 117 (38), 23652.

439 33. Chin, A.W.H. et al. (2020) Stability of SARS-CoV-2 in different environmental conditions. *The*
440 *Lancet Microbe* 1 (1), e10.

441 34. Kampf, G. et al. (2020) Persistence of coronaviruses on inanimate surfaces and their inactivation
442 with biocidal agents. *Journal of Hospital Infection* 104 (3), 246-251.

443 35. van Doremalen, N. et al. (2020) Aerosol and Surface Stability of SARS-CoV-2 as Compared with
444 SARS-CoV-1. *New England Journal of Medicine*.

445

446

447 **Table 1: COVID-19 doubling time versus climate area.**

Country/region	COVID-19 doubling time (days)	Climate area	Lab-confirmed case numbers *
Spain	4.2	Csa/Csb/Bsk	64,095
Southern Italy	5.38	Csa/Csb	5,322
Central Italy	5.87	Csa/Cfa/Cfb	10,842
Northern Italy	6.63	Cfa/Cfb	70,334
Germany	7.0	Cfb	73,522
France	7.5	Cfb	68,665
UK	7.2	Cfb	68,052
Sweden	9.4	Dfc/Cfb	11,321
Finland	10.8	Dfc/Dfb	2,646
Norway	12.95	Dfc/Dfb/ET	5,855

448
449 *: Laboratory-confirmed SARS-CoV-2 infection cases in Europe cases were retrieved by country at peak diffusion rates,
450 before the landmark dates indicated: Italy (github.com/pcm-dpc/COVID-19, March 27th 2020), France
451 (dashboard.covid19.data.gouv.fr/vue-d-ensemble?location=FRA; April 4th 2020), UK (www.nhs.uk/; April 9th
452 2020), Germany (corona.rki.de; April 2nd 2020), Spain (RTVE - Ministry of Health;
453 www.rtve.es/noticias/20200415/mapa-del-coronavirus-espana/2004681.shtml; March 27th 2020), Sweden (Public Health
454 Agency of Sweden; www.folkhalsomyndigheten.se/smittskydd-beredskap/utbrott/aktuella-utbrott/covid-19; April 9th
455 2020), Finland (National Institute for Health and Welfare THL; thl.fi/en/web/thlfi-en; April 7th 2020), Norway
456 (www.fhi.no/sv/smittsomme-sykdommer/corona/dags--og-ukerapporter/dags--og-ukerapporter-om-koronavirus; April
457 14th 2020).

458

Figure 1. Worldwide progression of COVID-19 cases.

(A) COVID-19 cumulative case incidence across the world, as of March 21st, 2020; numbers are color-coded and are proportional to circle diameter (www.healthmap.org/covid-19/).

(B) COVID-19 cumulative case incidence, as in (A), zoomed over Central Europe.

(C) COVID-19 incidence of active cases, as of March 31st, 2020; numbers are proportional to circle diameter (Johns Hopkins University, JHU; coronavirus.jhu.edu/map.html).

(D) Coronavirus spreading around the world as of April 4th. Overall confirmed cases by country since February 24th (JHU, public.flourish.studio/visualisation/1694807/).

Region	Coef.	Std. Err.	t	P>t	[95% Conf. Interval]	
intercept Northern Italy	6,63	0,33			5,97	7,29
Southern vs Northern Italy	-1,25	0,53	-2,35	0,02	-2,31	-0,20
Central vs Northern Italy	-0,76	0,55	-1,39	0,166	-1,84	0,32
Sweden vs Northern Italy	2,77	0,72	3,86	<0.0001	1,35	4,19
Norway vs Northern Italy	6,32	0,77	8,25	<0.0001	4,80	7,83

Figure 3. COVID-19 diffusion across geographic areas.

(top) Distribution boxplots of SARS-CoV-2 infected cases doubling times in the areas/countries analyzed. Upper horizontal line: 75th percentile; lower horizontal line: 25th percentile; horizontal bar within box: median; upper horizontal bar outside box: maximum value; lower horizontal bar outside box: minimum value. Dots: distribution outliers.

(bottom) SARS-CoV-2 infected cases doubling times versus central intercept (benchmark=Northern Italy). Coef.: coefficient; Std. Err.: standard error; 95% confidence intervals are shown. P>t: 0.002 Southern versus Northern Italy; <0.0001 Sweden versus Northern Italy; <0.0001 Norway versus Northern Italy.

Figure 4. The COVID-19 North-South gradient.

The North-South gradient of SARS-CoV-2 infected cases doubling times across countries is depicted. Country values were plotted according to their classification by climate zone (Table 1). The Anova P-value for Cartesian numerals association is shown. Vertical arrows: COVID-19 doubling times in validation datasets (Germany, France, UK).